

Modeling and Simulation — Lesson 2

Feedback in the Nature


Dynamics of Predator-Prey Interactions

Václav B. Petrák

Faculty of Biomedical Engineering
Czech Technical University

March 5, 2026

Predator-Prey



- ▶ The Lotka-Volterra equations, also known as the predator-prey equations, model the dynamics of biological systems in which two species interact, one as a **predator** and the other as **prey**.
- ▶ They were proposed independently by Alfred J. Lotka in 1925 and Vito Volterra in 1926.
- ▶ Model is characterized by **oscillations** in the population size of both predator and prey,
- ▶ Peak of the predator's oscillation lags behind the peak of the prey's oscillation.

Assumptions

The model makes several simplifying assumptions but captures the essential dynamics of predator-prey interactions:

Assumptions

The model makes several simplifying assumptions but captures the essential dynamics of predator-prey interactions:

- ▶ the prey population will grow exponentially when the predator is absent (unrealistic in a long term but may be precise for a long period)

Assumptions

The model makes several simplifying assumptions but captures the essential dynamics of predator-prey interactions:

- ▶ the prey population will grow exponentially when the predator is absent (unrealistic in a long term but may be precise for a long period)
- ▶ the predator population will starve in the absence of the prey population — there is no option of switching to another type of prey (may be realistic for some circumstances)

Assumptions

The model makes several simplifying assumptions but captures the essential dynamics of predator-prey interactions:

- ▶ the prey population will grow exponentially when the predator is absent (unrealistic in a long term but may be precise for a long period)
- ▶ the predator population will starve in the absence of the prey population — there is no option of switching to another type of prey (may be realistic for some circumstances)
- ▶ predators can consume any quantities of prey (unrealistic for large quantities)

Assumptions

The model makes several simplifying assumptions but captures the essential dynamics of predator-prey interactions:

- ▶ the prey population will grow exponentially when the predator is absent (unrealistic in a long term but may be precise for a long period)
- ▶ the predator population will starve in the absence of the prey population — there is no option of switching to another type of prey (may be realistic for some circumstances)
- ▶ predators can consume any quantities of prey (unrealistic for large quantities)
- ▶ both populations are moving randomly through a homogeneous, static, environment (may be approximately true for some environments)

The Mathematical Model

The Lotka-Volterra equations are a pair of first-order, nonlinear, differential equations given by:

$$\frac{dN}{dt} = rN - \beta NP \quad (1)$$

$$\frac{dP}{dt} = m\beta NP - dP \quad (2)$$

The Mathematical Model

The Lotka-Volterra equations are a pair of first-order, nonlinear, differential equations given by:

$$\frac{dN}{dt} = rN - \beta NP \quad (1)$$

$$\frac{dP}{dt} = m\beta NP - dP \quad (2)$$

where:

- ▶ N is population number of prey (e.g., rabbits),

The Mathematical Model

The Lotka-Volterra equations are a pair of first-order, nonlinear, differential equations given by:

$$\frac{dN}{dt} = rN - \beta NP \quad (1)$$

$$\frac{dP}{dt} = m\beta NP - dP \quad (2)$$

where:

- ▶ N is population number of prey (e.g., rabbits),
- ▶ P is the number of predators (e.g., foxes),
- ▶ r is the prey growth rate

The Mathematical Model

The Lotka-Volterra equations are a pair of first-order, nonlinear, differential equations given by:

$$\frac{dN}{dt} = rN - \beta NP \quad (1)$$

$$\frac{dP}{dt} = m\beta NP - dP \quad (2)$$

where:

- ▶ N is population number of prey (e.g., rabbits),
- ▶ P is the number of predators (e.g., foxes),
- ▶ r is the prey growth rate
- ▶ d is the death rate of predators

The Mathematical Model

The Lotka-Volterra equations are a pair of first-order, nonlinear, differential equations given by:

$$\frac{dN}{dt} = rN - \beta NP \quad (1)$$

$$\frac{dP}{dt} = m\beta NP - dP \quad (2)$$

where:

- ▶ N is population number of prey (e.g., rabbits),
- ▶ P is the number of predators (e.g., foxes),
- ▶ r is the prey growth rate
- ▶ d is the death rate of predators
- ▶ m is the proportionality constant

The Mathematical Model

The Lotka-Volterra equations are a pair of first-order, nonlinear, differential equations given by:

$$\frac{dN}{dt} = rN - \beta NP \quad (1)$$

$$\frac{dP}{dt} = m\beta NP - dP \quad (2)$$

where:

- ▶ N is population number of prey (e.g., rabbits),
- ▶ P is the number of predators (e.g., foxes),
- ▶ r is the prey growth rate
- ▶ d is the death rate of predators
- ▶ m is the proportionality constant
- ▶ β attack efficiency of the predator

Prey equations

The prey population is given by:

$$\frac{dN}{dt} = rN - \beta NP \quad (3)$$

Prey equations

The prey population is given by:

$$\frac{dN}{dt} = rN - \beta NP \quad (3)$$

In the absence of predators, the equation becomes:

$$\frac{dN}{dt} = rN \quad (4)$$

which represents the Malthusian model. The prey population would grow exponentially with the growth rate r .

Prey equations

The prey population is given by:

$$\frac{dN}{dt} = rN - \beta NP \quad (3)$$

In the absence of predators, the equation becomes:

$$\frac{dN}{dt} = rN \quad (4)$$

which represents the Malthusian model. The prey population would grow exponentially with the growth rate r .

1em The reduction in prey population due to predators is captured by the expression:

$$-\beta NP \quad (5)$$

Prey equations

The prey population is given by:

$$\frac{dN}{dt} = rN - \beta NP \quad (3)$$

In the absence of predators, the equation becomes:

$$\frac{dN}{dt} = rN \quad (4)$$

which represents the Malthusian model. The prey population would grow exponentially with the growth rate r .


The reduction in prey population due to predators is captured by the expression:

$$-\beta NP \quad (5)$$

Here, NP is proportional to the number of encounters. An increased number of predators, an increased prey population, and greater searching efficiency of the predators would all lead to a higher removal rate of prey. β is attack efficiency of the predator. Higher β means higher probability of prey being killed.


Predator-prey encounters

Picture below illustrates the concept of predators-prey encounters as $N \cdot P$ product.



Predator-prey encounters

Picture below illustrates the concept of predators-prey encounters as $N \cdot P$ product.



The concept would not work for predators hunting in packs or for prey staying in herds.

Predator equations

The predator population is given by:

$$\frac{dP}{dt} = m\beta NP - dP \quad (6)$$

Predator equations

The predator population is given by:

$$\frac{dP}{dt} = m\beta NP - dP \quad (6)$$

If the prey population N is 0, the equation becomes:

$$\frac{dN}{dt} = -dN \quad (7)$$

which represents dying of prey at exponential rate d .

Predator equations

The predator population is given by:

$$\frac{dP}{dt} = m\beta NP - dP \quad (6)$$

If the prey population N is 0, the equation becomes:


$$\frac{dN}{dt} = -dN \quad (7)$$


which represents dying of prey at exponential rate d .

The growth of predator population growth by eating of prey


$$+m\beta NP \quad (8)$$

Here we see again βNP . Proportionality constant m tells how many prey individuals are required for increase of predator population by one.





Concept of feedback



1

What is a Vector Field?

- ▶ In mathematics and physics, a **vector field** assigns a vector to every point in a given space.

What is a Vector Field?

- ▶ In mathematics and physics, a **vector field** assigns a vector to every point in a given space.
- ▶ For a 2D dynamical system, the state is represented by coordinates (x, y) . The differential equations define the rate of change at any point:

$$\vec{v}(x, y) = \left(\frac{dx}{dt}, \frac{dy}{dt} \right) \quad (9)$$

What is a Vector Field?

- ▶ In mathematics and physics, a **vector field** assigns a vector to every point in a given space.
- ▶ For a 2D dynamical system, the state is represented by coordinates (x, y) . The differential equations define the rate of change at any point:

$$\vec{v}(x, y) = \left(\frac{dx}{dt}, \frac{dy}{dt} \right) \quad (9)$$

- ▶ Visually, this is represented as a grid of arrows. The **direction** of the arrow shows where the system is heading, and its **length** (magnitude) shows how fast it is changing.
- ▶ Vector fields allow us to understand the qualitative, long-term behavior of a system without needing to analytically solve the complex differential equations.

Vector Field in Predator-Prey Phase Space

- ▶ In the Lotka-Volterra model, our 2D phase space is defined by the prey population N and the predator population P .

Vector Field in Predator-Prey Phase Space

- ▶ In the Lotka-Volterra model, our 2D phase space is defined by the prey population N and the predator population P .
- ▶ At any given state (N, P) , the instantaneous rate of change is the vector:

$$\vec{v} = (rN - \beta NP, m\beta NP - dP) \quad (10)$$

Vector Field in Predator-Prey Phase Space

- ▶ In the Lotka-Volterra model, our 2D phase space is defined by the prey population N and the predator population P .
- ▶ At any given state (N, P) , the instantaneous rate of change is the vector:

$$\vec{v} = (rN - \beta NP, m\beta NP - dP) \quad (10)$$

- ▶ The arrows in this specific vector field indicate how both populations will shift from that exact moment.
- ▶ By following the continuous flow of these arrows, we can trace the closed, periodic trajectories of the populations over time.

Equilibrium Points in the Vector Field

- **Equilibrium points** (or fixed points) occur where the vector magnitude is exactly zero. At these points, the system is completely at rest ($\frac{dN}{dt} = 0$ and $\frac{dP}{dt} = 0$).

Equilibrium Points in the Vector Field

- ▶ **Equilibrium points** (or fixed points) occur where the vector magnitude is exactly zero. At these points, the system is completely at rest ($\frac{dN}{dt} = 0$ and $\frac{dP}{dt} = 0$).
- ▶ For the Lotka-Volterra equations, the vector field reveals two such points:
 1. **Extinction:** $(0, 0)$ — both populations are zero.

Equilibrium Points in the Vector Field

- ▶ **Equilibrium points** (or fixed points) occur where the vector magnitude is exactly zero. At these points, the system is completely at rest ($\frac{dN}{dt} = 0$ and $\frac{dP}{dt} = 0$).
- ▶ For the Lotka-Volterra equations, the vector field reveals two such points:
 1. **Extinction:** $(0, 0)$ — both populations are zero.
 2. **Coexistence:** $\left(\frac{d}{m\beta}, \frac{r}{\beta}\right)$ — populations perfectly balance each other out and remain static.

Equilibrium Points in the Vector Field

- ▶ **Equilibrium points** (or fixed points) occur where the vector magnitude is exactly zero. At these points, the system is completely at rest ($\frac{dN}{dt} = 0$ and $\frac{dP}{dt} = 0$).
- ▶ For the Lotka-Volterra equations, the vector field reveals two such points:
 1. **Extinction:** $(0, 0)$ — both populations are zero.
 2. **Coexistence:** $\left(\frac{d}{m\beta}, \frac{r}{\beta}\right)$ — populations perfectly balance each other out and remain static.
- ▶ The vector field arrows orbit continuously around the coexistence equilibrium point, mathematically illustrating the perpetual lagging and catching-up dynamics of the two species.

Biological Implications

- ▶ The model predicts oscillations in the populations of predator and prey.
- ▶ The fluctuation of the two species is periodic,
- ▶ The amplitude and period of these oscillations depend on the initial conditions and the parameters of the model.
- ▶ The *average* numbers of the two species are constant
- ▶ If we destroy individuals of both species proportionately to their number, the average number of individuals of the prey grows and the average number of predators diminishes.

Conclusion

- ▶ The Lotka-Volterra predator-prey model provides a framework for understanding the dynamics of biological systems with predator-prey interactions.
- ▶ Despite its simplicity, it offers valuable insights into the oscillatory nature of ecological populations and has motivated further research into more complex and realistic models.

Modeling Approaches

Modeling Approach 1: The Euler Method

- ▶ The Euler method is the simplest numerical procedure for solving ordinary differential equations (ODEs).
- ▶ **How it works:** We break time into discrete, tiny steps (Δt). At each step, we calculate the current rate of change and assume it remains constant for that brief interval.
- ▶ **Discrete Equations:**

$$N_{t+\Delta t} = N_t + \Delta t(rN_t - \beta N_t P_t) \quad (11)$$

$$P_{t+\Delta t} = P_t + \Delta t(m\beta N_t P_t - dP_t) \quad (12)$$

- ▶ **Pros:** Transparent; excellent for learning how continuous equations translate to discrete steps.
- ▶ **Cons:** Prone to accumulating errors. If Δt is too large, the inherent closed loops artificially spiral outwards.

Euler Method in Python

Demonstrating the explicit loop, mapping directly to the discrete equations:

```
1 import numpy as np
2
3 r, beta, m, d = 1.0, 0.1, 0.05, 0.5
4 dt = 0.01 # Small time step is crucial for stability
5 steps = 2000
6
7 N, P = np.zeros(steps), np.zeros(steps)
8 N[0], P[0] = 40, 9 # Initial conditions
9
10 for i in range(1, steps):
11     dN_dt = r * N[i-1] - beta * N[i-1] * P[i-1]
12     dP_dt = m * beta * N[i-1] * P[i-1] - d * P[i-1]
13
14     N[i] = N[i-1] + dN_dt * dt
15     P[i] = P[i-1] + dP_dt * dt
```

Modeling Approach 2: ODE Solvers

- ▶ For practical applications and research, manual loops are often insufficient. We use dedicated mathematical solvers, such as SciPy's `solve_ivp`.
- ▶ **How it works:** These solvers act as a "black box" utilizing advanced algorithms (like Runge-Kutta). You provide the differential equations, time span, and initial conditions.
- ▶ **Adaptive Step Size:** The solver automatically shrinks Δt when populations change rapidly (for accuracy) and expands it when stable (to save computing power).
- ▶ **Pros:** Highly accurate, stable, and correctly handles the periodic nature of predator-prey dynamics without artificial spiraling.
- ▶ **Cons:** Less transparent for beginners; the integration loop is hidden behind the scenes.

SciPy Solver in Python

Define the system as a function, allowing the solver to handle integration:

```
1 from scipy.integrate import solve_ivp
2
3 def lotka_volterra(t, state, r, beta, m, d):
4     N, P = state
5     dN_dt = r * N - beta * N * P
6     dP_dt = m * beta * N * P - d * P
7     return [dN_dt, dP_dt]
8
9 params = (1.0, 0.1, 0.05, 0.5)
10 initial_state = [40, 9]
11 t_span = (0, 20)
12
13 # 'RK45' is the default Runge-Kutta method
14 solution = solve_ivp(fun=lotka_volterra, t_span=t_span,
15                      y0=initial_state, args=params)
16
17 time = solution.t
18 prey, predators = solution.y[0], solution.y[1]
```