

# Modeling and Simulation — Lesson 1

## **The Limits to Growth: Basic Population Models**

Václav B. Petrák

Faculty of Biomedical Engineering  
Czech Technical University in Prague

April 16, 2026

## Introduction

- Course information

- Introduction to Modeling

## Section 1: Fibonacci Sequence

- Background

- Exercise

## Section 2: Malthusian Growth Model

- Background

- Exercises

## Section 3: Logistic Growth Model

- Background

- Exercise

## Section 4: Leslie Matrix Model

- Background

- Population Growth Rate ( $\lambda$ )

- Exercise

# Introduction

**Lecturer:** Václav B. Petrák

**Email:** [vaclav.petrak@fbmi.cvut.cz](mailto:vaclav.petrak@fbmi.cvut.cz)

**Phone/WhatsApp:** +420 725 878 390

## Consultations:

- ▶ In person: By individual request. Please email or text me.
- ▶ In person: Friday, May 2<sup>nd</sup>, 2025 (Poster-related problems).
- ▶ Via email: Anytime.

- ▶ **Advance Notice Required:** All absences must be excused in advance whenever possible.
- ▶ **Communication Procedure:** Contact me as soon as possible by email with the date of absence and the reason.
- ▶ **Responsibility for Missed Work:** You are responsible for making up missed assignments and catching up on course materials.

- ▶ We will cover 5 topics in modeling and simulation, one per session.
- ▶ The last lecture will be focused on student presentations.

There is no exam.

Your grade will be based on your independent work.  
There are three outputs used for grading:

- 1. Journal Club Presentation**
- 2. Capstone Poster Presentation**
- 3. Project Review Discussion**

# Journal Club is a discussion of research papers

A journal club is a group that meets to critically evaluate articles in academic literature. Journal clubs:

- ▶ Encourage critical thinking.
- ▶ Help in understanding the studied concepts and methods.
- ▶ Help keep up with the latest research.
- ▶ Improve communication and presentation skills.
- ▶ Simulate the experience of a scientist.

**The main purpose in this course is to explore how the methods we learned are applied in current research.**



# Your Task for Journal Club

- ▶ You will read a research paper related to the topic of your capstone poster.
- ▶ Your task is to read the paper and prepare a 15-minute presentation, followed by a 5-minute discussion, to explain the paper.
- ▶ The presentation dates will be arranged individually.

# The Journal Club presentation will cover:

- Introduction:** Provide a high-level overview of the background of the research topic and the main objectives of the paper.
- Methods:** Give an overview of the methodology used in the paper.
- Results:** Highlight the key results and conclusions.
- Discussion:** Discuss the practical implications of the research, its applications in real-world situations, and any limitations.
- Conclusions:** Summarize the main contributions and overall findings of **the paper**. Reiterate its key takeaways.
- Q&A:** Answer questions from the audience (lecturer).

# Grading Criteria for Journal Club Presentation

## Presentation Design and Delivery (50%)

- ▶ **Design:** The presentation is well-organized and easy to follow.
- ▶ **Timing:** The presenter adhered to the 15-minute format.
- ▶ **Delivery:** Clear and confident speech with appropriate pacing.
- ▶ **Questions:** Convincing response to questions.


# Grading Criteria for Journal Club Presentation

## Content of the Presentation (50%)

- ▶ **Introduction:** Provides a high-level overview, explaining the motivation and research question.
- ▶ **Methods:** Explains the experimental approach used by the authors. Unfamiliar methods are explained without excessive detail.
- ▶ **Results:** The key results from the paper are explained.
- ▶ **Critical Evaluation:** Critique of methods, ideas for improvement, and significance of the results.

# Capstone poster is your independent project

- ▶ You will work on an independent project that builds on and extends course content.
- ▶ The output of your work will be a poster and the code.
- ▶ Presentation dates will be arranged individually.



A **capstone** (or keystone) is the wedge-shaped stone at the apex of an arch. ***It is the final piece placed during construction and locks all the stones into position, allowing the arch to bear weight.***

# The Capstone Poster presentation will cover:

Walk others through your poster in about 5 minutes, explaining its content and key messages.

**Introduction:** Provide a high-level overview of the background and main objectives of **your poster**.

**Methods:** Give an overview of the methodology used in **your poster**.

**Results:** Highlight the key results and conclusions.

**Discussion:** Discuss the practical implications of **your findings** and their applications. Discuss any limitations.

**Conclusions:** Summarize very briefly the main contributions and overall findings. Reiterate key takeaways.

# Grading Criteria for Capstone Poster

## Poster Design and Delivery (50%)

- ▶ **Design:** The poster is well-organized and easy to follow. Make sure that the poster reads well.
- ▶ **Plots:** Clear plots, of simulation results that address poster objectives and show model behavior.
- ▶ **Delivery:** Clear and confident speech with appropriate pacing.
- ▶ **Questions:** Convincing response to questions.

# Grading Criteria for Capstone Poster

## Model Implementation (50%)

- ▶ **Correctness:** Correct functionality of fundamental models.
- ▶ **Understanding:** Demonstrated understanding of core concepts.
- ▶ **Extension:** Successful and meaningful implementation of additional features (if applicable).
- ▶ **Discussion:** Insightful discussion on how parameters/initial conditions affect outcomes. Identification and explanation of behaviors.



# During Project Review we will discuss your work

Project Review assesses **your understanding** of the Capstone Poster, including the methodology, results, and the quality, design, and documentation of your code.

- ▶ **Primary Focus:** The review covers the entire Capstone Poster, with code as one component.
- ▶ **Interactive Session:** An interactive walk-through where you explain your project's methodology, design choices, and code functionality.
- ▶ **Demonstrating Ownership:** An opportunity to demonstrate your understanding, justify development decisions, and articulate the reasoning behind your approach.
- ▶ **Goal:** Ensure the correctness of your project while demonstrating your grasp of the implemented concepts.

# Grading Criteria for Project Review

## Readability and Maintainability (20%)

- ▶ Clean, well-formatted, and easy-to-follow code.
- ▶ Consistent naming conventions and style.
- ▶ Logical organization: functions that break code into manageable components where necessary.
- ▶ Sufficient and meaningful comments for complex/non-obvious code.

# Grading Criteria for Project Review

## Project Understanding (80%)

- ▶ Ability to explain the modeled system: what it represents, what assumptions were made, and why.
- ▶ Ability to interpret simulation results in terms of real-world behavior of the system.
- ▶ Ability to explain how model parameters and initial conditions influence outcomes.
- ▶ Ability to suggest meaningful extensions, alternative approaches, or improvements to the model.

# Avoid using AI to solve entire exercises

- ▶ Consider a math problem:
  - *Given that two trains are traveling on parallel tracks in opposite directions with speeds of 56 km/h and 52 km/h... after what duration will they cross each other?*
- ▶ The primary goal is **not** to find the answer but to practice your reasoning and calculation skills.
- ▶ Similarly, the aim of modeling and simulation exercises is to learn and practice new skills.
- ▶ Avoid using generative AI like ChatGPT or Gemini to solve the **entire problem** for you.
  - It spoils the learning process for you.
  - It may disrupt the pacing of the lesson.

During lessons, you may use AI to:

- ▶ Ask clarifying questions.
- ▶ Debug your code (after you've made your own attempts).
- ▶ Get ideas for further work.

**Let's discuss what constitutes fair use of generative AI in the capstone poster.**

# A model is a representation of a **system**

- ▶ A model is a representation of a **system** that enables us to investigate its properties and **predict** future outcomes.
  - A **system** is a set of components that interact with each other within a boundary to function as a whole (e.g., Solar system, rabbit and fox populations).
- ▶ In modeling, we aim to identify components, relationships, and behaviors to predict system dynamics.



# A model is a representation of a **system**

- ▶ A model is a representation of a **system** that enables us to investigate its properties and **predict** future outcomes.
  - A **system** is a set of components that interact with each other within a boundary to function as a whole (e.g., Solar system, rabbit and fox populations).
- ▶ In modeling, we aim to identify components, relationships, and behaviors to predict system dynamics.
- ▶ Modeling always requires simplification (abstraction).

Example of a model:  
SIR (susceptible,  
infectious, and recovered)  
model for dynamics of  
infectious diseases



# A model is a representation of a **system**

- ▶ A model is a representation of a **system** that enables us to investigate its properties and **predict** future outcomes.
  - A **system** is a set of components that interact with each other within a boundary to function as a whole (e.g., Solar system, rabbit and fox populations).
- ▶ In modeling, we aim to identify components, relationships, and behaviors to predict system dynamics.
- ▶ Modeling always requires simplification (abstraction).
- ▶ A **mathematical model** uses **mathematical equations to describe a system**.

SIR model equations:

$$\frac{dS}{dt} = -\beta SI$$

$$\frac{dI}{dt} = \beta SI - \gamma I$$

$$\frac{dR}{dt} = \gamma I$$



All models are approximations. Assumptions, whether implied or clearly stated, are never exactly true.

**All models are wrong, but some models are useful.**

So the question you need to ask is not *"Is the model true?"*

It never is.

But *"Is the model good enough for this particular application?"*

— George E. P. Box


## Section 1: Fibonacci Sequence

# A very simple population model

The Italian mathematician Leonardo Fibonacci (1170 - 1250) considered the following conditions when calculating the number of rabbits born over the course of one year:

- ▶ A single newly born pair of rabbits (one male, one female) is put in a field.
- ▶ Rabbits are able to mate at the age of one month, so that at the end of their second month, a female can produce another pair of rabbits.
- ▶ Rabbits never die, and a mating pair always produces one new pair (one male, one female) every month from the second month on.

# Graphical representation of the first 6 months



# Fibonacci sequence

This problem leads to a sequence of numbers called the **Fibonacci sequence**:

1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, ...

The first two numbers are given, and subsequent numbers are found by adding the two preceding numbers. Thus, we have the following recursive definition (a definition that defines elements in a set in terms of other elements in the set) for Fibonacci numbers:

$$F_1 = F_2 = 1$$


$$F_n = F_{n-1} + F_{n-2}$$

The ratio of two consecutive Fibonacci numbers converges to a number known as the **golden ratio**:

$$\psi = \frac{1 + \sqrt{5}}{2} \approx 1.618... \quad (1)$$


# Fibonacci sequence

A spiral created by drawing circular arcs connecting the opposite corners of squares in a tiling where side lengths correspond to successive Fibonacci numbers.




# Fibonacci sequence in nature

- **Spiral Patterns:** The growth chambers of a nautilus shell create a spiral that approximately follows the Fibonacci sequence and the golden ratio.



# Fibonacci sequence in nature

- ▶ **Spiral Patterns:** The growth chambers of a nautilus shell create a spiral that approximately follows the Fibonacci sequence and the golden ratio.
- ▶ **Branches:** Tree branches sometimes follow the Fibonacci sequence. The main branch splits into two, then one of those branches splits while the other stays dormant.






# Fibonacci sequence in nature

- ▶ **Spiral Patterns:** The growth chambers of a nautilus shell create a spiral that approximately follows the Fibonacci sequence and the golden ratio.
- ▶ **Branches:** Tree branches sometimes follow the Fibonacci sequence. The main branch splits into two, then one of those branches splits while the other stays dormant.
- ▶ **Flower Petals:** The number of petals on many flowers corresponds to Fibonacci numbers: lilies (3 petals), buttercups (5 petals), and daisies (often 34 or 55 petals).






# Fibonacci sequence in nature

- ▶ **Spiral Patterns:** The growth chambers of a nautilus shell create a spiral that approximately follows the Fibonacci sequence and the golden ratio.
- ▶ **Branches:** Tree branches sometimes follow the Fibonacci sequence. The main branch splits into two, then one of those branches splits while the other stays dormant.
- ▶ **Flower Petals:** The number of petals on many flowers corresponds to Fibonacci numbers: lilies (3 petals), buttercups (5 petals), and daisies (often 34 or 55 petals).
- ▶ **Why?** Fibonacci patterns in nature are likely related to optimal growth and packing efficiency.
- ▶ **While the Fibonacci sequence appears in nature, it is not a universal rule.**



# The Fibonacci sequence is extensively used in the arts

## Golden ratio in famous logos



# Fibonacci sequence tasks as Python refresher

We will use the Fibonacci sequence to review some basic concepts in Python. Basic experience with programming is expected.

## We will cover:

- ▶ Loop control statements
- ▶ Vector and matrix indexing (using NumPy)
- ▶ Functions
- ▶ Plotting (using Matplotlib)

# Task 1.1: Loop control statements

With loop control statements, you can repeatedly execute a block of code. There are two main types of loops:

1. **for** statements loop a specific number of times (often iterating over a sequence).
2. **while** statements loop as long as a condition remains true.

Your tasks:

1. Generate the first 20 elements of the Fibonacci sequence using a **for** loop.
2. Generate all Fibonacci numbers smaller than 500 using a **while** loop.

*Note: Save generated numbers to a list or array; we will need them for the next task.*

## Solution 1.1.1: First 20 Fibonacci numbers

Generate the first 20 elements of the Fibonacci sequence using a **for** loop.

```
1 import numpy as np
2
3 n = 20
4 fib_seq = [1, 1]
5
6 for i in range(2, n):
7     fib_seq.append(fib_seq[i-1] + fib_seq[i-2])
8
9 print(fib_seq)
```

## Solution 1.1.2: Fibonacci numbers smaller than 500

Generate all Fibonacci numbers smaller than 500 using a **while** loop.

```
1 fib_seq = [1, 1]
2 max_value = 5000
3 i = 1 # Python uses 0-based indexing
4
5 while (fib_seq[i] + fib_seq[i-1]) < max_value:
6     fib_seq.append(fib_seq[i] + fib_seq[i-1])
7     i += 1
8
9 print(fib_seq)
```

## Task 1.2: Vector Indexing

Vector indexing allows you to access and manipulate specific elements within an array. We will use the **NumPy** library for this.

### ► Basic Indexing (0-based)

- Access elements by their position: `x[2]` gets the *third* element of `x`.
- Negative indexing: `x[-1]` gets the last element.
- Slicing: `x[1:5]` gets elements from index 1 up to (but not including) index 5.

### ► Boolean (Logical) Indexing:

- Create a boolean mask to filter elements: `x[x > 10]` extracts elements greater than 10.



## Task 1.2: Vector Indexing

**Your tasks** (Assume `fib_seq` is a NumPy array):

- ▶ Divide the 6<sup>th</sup> and 5<sup>th</sup> element of the array.
- ▶ Divide the last and 2nd to last element of the array.
- ▶ Get all Fibonacci numbers smaller than 50.
- ▶ Get all Fibonacci numbers between 50 and 500.
- ▶ With the help of indexing and the modulus operator, replace all even Fibonacci numbers in the array with 0.

## Task 1.2: Solution

- ▶ Divide 6<sup>th</sup> and 5<sup>th</sup> element of the array.

```
fib_seq[5] / fib_seq[4]
```

- ▶ Divide last and 2nd last element of the array.

```
fib_seq[-1] / fib_seq[-2]
```

- ▶ Get all Fibonacci numbers smaller than 50.

```
fib_seq[fib_seq < 50]
```

- ▶ Get all Fibonacci numbers between 50 and 500.

```
fib_seq[(fib_seq > 50) & (fib_seq < 500)]
```

- ▶ Replace all even Fibonacci numbers in the vector with 0.

```
fib_seq[fib_seq % 2 == 0] = 0
```

# Matrix indexing is similar to vector indexing

- ▶ Define matrix (using NumPy)

```
M = np.array([[1,2,3], [4,5,6], [7,8,9]])
```

- ▶ Extract the element in row 2, column 3

```
M[1, 2]
```

- ▶ One or both of the row and column subscripts can be slices:

```
M[0:3, 1:3]
```

- ▶ A ":" selects the entire dimension

```
M[:, 1:3]
```

- ▶ To treat M as a flat vector (similar to linear indexing), use flatten:

```
M.flatten()[7]
```

- ▶ You can use a boolean array as a mask.

```
N = np.array([[1.3, 2.0], [4.8, 5.0]])
```

```
N[N % 1 == 0]
```

```
N[N > 3]
```

## Task 1.3: Functions background

- ▶ Functions are reusable blocks of code that take inputs, perform calculations, and return outputs. They are essential for code organization and modularity.
- ▶ Functions in Python are defined using the `def` keyword.

```
1  # declare function
2  def circle_area(radius):
3      pi = 3.14159 # approx. value of pi
4      area = pi * radius**2
5      return area
6
7  # call function
8  my_circle = circle_area(5)
```

# Task 1.3: Functions

## Your tasks

1. Write a function to return a list/array with generated  $n$  Fibonacci numbers.
2. Write a function to return a random Fibonacci number.
3. Write a function to check if a number is a Fibonacci number.

*Think about your solution first; after that feel free to search for hints online.*

## Solution 1.3.1: Fibonacci generator

- ▶ We will reuse our previous Task 1.1 code.
- ▶ Handling edge cases ( $n=0$ ,  $n=1$ ) is important.
- ▶ **What are the limitations of this code?**

```
1 def generate_fibonacci(n):  
2     fib_seq = [1, 1]  
3     for i in range(2, n):  
4         fib_seq.append(fib_seq[i-1] + fib_seq[i-2])  
5     return fib_seq[:n]  
6  
7 print(generate_fibonacci(10))
```

## Solution 1.3.2: Random Fibonacci number

```
1 import random
2
3 def random_fibonacci(n):
4     fib_seq = [1, 1]
5     for i in range(2, n):
6         fib_seq.append(fib_seq[i-1] + fib_seq[i-2])
7
8     fib_seq = fib_seq[:n]
9     return random.choice(fib_seq)
10
11 print(random_fibonacci(10))
```

## Solution 1.3.3: Check if number is Fibonacci

One possible approach: Check if the last generated number before overstepping the threshold is the number in question.

```
1 def check_fibonacci(val):
2     a, b = 1, 1
3
4     while b < val:
5         memory = b
6         b = a + b
7         a = memory
8
9     return b == val
10
11 # Example usage:
12 # print(check_fibonacci(8)) # True
```



## Task 4: Plotting

Plotting allows you to represent your data graphically. Python uses the library **Matplotlib** for this.

Your tasks:

1. Plot the first 15 elements of the Fibonacci sequence.
2. Calculate and plot the ratio of  $n[i+1] / n[i]$  for the first 15 elements of the Fibonacci sequence.

# Solutions 1.4

1. Plot the first 15 elements of the Fibonacci sequence:

```
1 plt.plot(fib_seq_15, '-s')
2 plt.show()
3
4 # Calculate ratio ( $F_n / F_{n-1}$ )
5 ratio = fib_seq_15[1:] / fib_seq_15[:-1]
6 print(ratio)
```

2. Calculate and plot the ratio of  $n[i+1] / n[i]$  in the first 15 elements.

```
1 plt.figure()
2 plt.plot(ratio, '-s')
3 plt.show()
```

## Section 2: Malthusian Growth Model

# Malthusian Growth Model

- ▶ The Malthusian growth model is a mathematical model of population growth.
- ▶ It essentially describes exponential population growth. [Image of exponential growth curve]
- ▶ The model is named after Thomas Malthus, who wrote *An Essay on the Principle of Population* (1798), one of the first and most influential books on population dynamics.



Thomas Robert Malthus (1766 - 1834), an English economist and scholar.

# Key assumptions of Thomas Malthus

- ▶ **Population grows exponentially when resources are abundant.**
- ▶ Malthus further assumed that agricultural production increases only linearly (arithmetically). [Image of Malthusian catastrophe graph showing exponential population vs linear resources]
- ▶ Population growth is limited by available resources and can be reduced by famine, disease, or war.

## Malthusian Catastrophe

A situation where the population exceeds the agricultural production of its environment. This results in severe consequences like poverty, famine, disease, and war. Malthus believed a Malthusian catastrophe was inevitable unless population growth was controlled.

# Malthus, Darwin, and Natural Selection

- ▶ Before reading Malthus, Darwin thought that living things reproduced just enough individuals to keep populations stable.
- ▶ Malthus helped him realize that as populations breed beyond their means, advantageous traits become more prevalent due to competition for resources.
- ▶ This led to the development of the theory of natural selection.

*"In October 1838... I happened to read for amusement Malthus on Population, and being well prepared to appreciate the struggle for existence... **it at once struck me that under these circumstances favourable variations would tend to be preserved, and unfavourable ones to be destroyed.** The result of this would be the formation of a new species."*

— Charles Darwin

# Neo-Malthusianism

- ▶ A revival of Malthusian concerns in the 20<sup>th</sup> century, often focused on environmental limits.
- ▶ Neo-Malthusians advocate for population control due to concerns about overpopulation, resource depletion, and environmental degradation.
- ▶ Widely discussed in the 1960s; a famous example is Paul Ehrlich's book *The Population Bomb* (1968).
- ▶ Their influence can be seen in family planning initiatives and advocacy for sustainability.
- ▶ Modern environmentalism acknowledges the impact of population growth on resource use, pollution, and environmental strain.



# Malthusian Growth Model Equation

The change in population  $dP/dt$  is proportional to the current population size  $P$  at time  $t$ . The parameter  $r$  is the intrinsic rate of increase (the difference between birth rate and death rate).

$$\frac{dP}{dt} = rP$$

To solve the differential equation, we separate the variables. We move terms with  $P$  to one side and terms with  $t$  to the other side.

$$\frac{dP}{P} = rdt$$



# Malthusian Growth Model Equation

We integrate both sides of the equation. Integration introduces a constant of integration,  $C$ .

$$\int \frac{dP}{P} = \int r dt$$
$$\ln |P| = rt + C$$

We exponentiate both sides. We remove the absolute value as the population size is always positive.

$$e^{\ln P} = e^{rt+C}$$
$$P = e^C e^{rt}$$

The integration constant  $e^C$  represents the initial population  $P_0$ . Substitution leads to the final explicit form:

$$P(t) = P_0 e^{rt}$$

# Numerical vs Explicit Solution

- ▶ Numerical and explicit solutions are two methods used to solve mathematical problems in modeling and simulation.
- ▶ **Explicit (Analytical) Solution:** A mathematically exact solution expressed as a formula.
  - Provides a precise value for any time  $t$ .
  - Often allows for a deeper understanding of the underlying mathematical relationships.
  - It can be very difficult or impossible to find explicit solutions for complex, non-linear problems.

# Numerical vs Explicit Solution

- ▶ Numerical and explicit solutions are two methods used to solve mathematical problems in modeling and simulation.
- ▶ **Numerical Solution:** Uses **iterative methods to approximate** a solution using computational steps. [Image of Euler method steps visualization]
  - Allows solving a much wider range of complex problems (e.g., where parameters change over time).
  - Solutions are only approximate.
  - Solutions can be affected by cumulative calculation errors (discretization error).
  - Difficult to validate without real-world data or an explicit solution for comparison.

# Numerical vs Explicit Solution for Malthusian Growth

## Explicit Solution

Calculation using the formula:

$$P(t) = P_0 e^{rt}$$

Where:

- ▶  $P_0$  is the initial population at time 0
- ▶  $r$  is the growth rate
- ▶  $t$  is the time

## Numerical Solution (Euler's Method):

$$\frac{dP}{dt} \approx \frac{\Delta P}{\Delta t}$$

Algorithm:

### 1. Discretization:

$$t_{i+1} = t_i + \Delta t$$

### 2. Iteration (Update Rule):

$$P(t_{i+1}) = P(t_i) + rP(t_i)\Delta t$$

Where:

- ▶  $P(t_i)$  is the population at time step  $i$
- ▶  $\Delta t$  is the time step size

## Task 2.1: Population at discrete times

Calculate population at certain times of interest using the explicit formula.

- ▶ Initial population size ( $P_0$ ): 2.5 million
- ▶ Yearly growth rate ( $r$ ): 2.7%
- ▶ Equation:  $P(t) = P_0 e^{rt}$
- ▶ Time in years ( $t$ ):
  - 1
  - 5
  - 7.6
  - 25.78
- ▶ Note: Population size and growth rate correspond to the population of Senegal in 1950.



## Solution 2.1: Population at discrete times

```
1 import numpy as np
2
3 P0 = 2.5    # Initial population
4 r = 0.027   # Growth rate 2.7%
5
6 t_task1 = np.array([1, 5, 7.5, 25.78])
7
8 print("Years of interest:")
9 print(t_task1)
10 print("Calculated population size:")
11 print(P0 * np.exp(r * t_task1))
```

## Solution 2.1: Population at discrete times

- ▶ The results show population size at times of 1, 5, 7.5, and 25.78 years.
- ▶ The explicit solution allows us to easily calculate population predictions at any given specific time without iterating.

```
1 Years of interest:
2   [1, 5, 7.5, 25.78]
3 Calculated population size (Millions):
4   [2.568, 2.861, 3.061, 5.014]
```

## Task 2.2: Continuous modeling with explicit solution

- ▶ The population has the same parameters as in Task 2.1.
- ▶ Calculate the population for the next 70 years.
- ▶ The time step is 1 year.
- ▶ Plot the results.




## Solution 2.2: Continuous modeling with explicit solution

```
1 import numpy as np
2 import matplotlib.pyplot as plt
3
4 P0 = 2.50
5 r = 0.027
6 t_start, t_end, t_diff = 0, 70, 1
7
8 t_expl = np.arange(t_start, t_end + t_diff, t_diff)
9 P_expl = P0 * np.exp(r * t_expl)
10
11 plt.plot(t_expl, P_expl, linewidth=2)
12 plt.xlabel("Time (years)")
13 plt.ylabel("Population (millions)")
14 plt.ylim(0, 20)
15 plt.show()
```

## Solution 2.2: Continuous modeling with explicit solution

Population increases exponentially.




## Task 2.3: Numerical approach

- ▶ Use the numerical approach (Euler's method) to calculate population from year 0 to 70.
- ▶ The time step should be 1 year.
- ▶ Plot the result.

## Solution 2.3: Numerical approach

```
1 import matplotlib.pyplot as plt
2
3 P0, r = 2.50, 0.027
4 t_start, t_end, t_diff = 0, 70, 1
5
6 t_num = [t_start]
7 P_num = [P0]
8
9 # Numerical calculation loop
10 for i in range(int(t_end/t_diff)):
11     P_new = P_num[i] + r * P_num[i] * t_diff
12     t_new = t_num[i] + t_diff
13     P_num.append(P_new)
14     t_num.append(t_new)
15
16 plt.plot(t_num, P_num, color="red")
17 plt.xlabel("Time (years)")
18 plt.ylabel("Population (millions)")
19 plt.ylim(0, 20)
20 plt.show()
```

## Solution 2.3: Numerical approach



## Task 2.4: Difference in solutions

- ▶ Create a plot that shows results from both explicit and numerical approaches and the difference (error).
- ▶ Add a start year of 1950 to the plot to represent the correct timeline.
- ▶ Compare the difference for time steps ( $\Delta t$ ) of 1 year, 1 month, and 1 day.

## Solution 2.4: The difference in solutions

```
1 import numpy as np
2 import matplotlib.pyplot as plt
3
4 # MODELING PARAMETERS
5 P0, r = 2.50, 0.027
6 start_year = 1950
7 t_start, t_end, t_diff = 0, 70, 1
8
9 # EXPLICIT SOLUTION
10 t_expl = np.arange(t_start, t_end + t_diff, t_diff)
11 P_expl = P0 * np.exp(r * t_expl)
12
13 # NUMERICAL SOLUTION
14 t_num = [t_start]
15 P_num = [P0]
16 for i in range(int(t_end/t_diff)):
17     P_num.append(P_num[i] + r * P_num[i] * t_diff)
18     t_num.append(t_num[i] + t_diff)
19 P_num = np.array(P_num)
20 t_num = np.array(t_num)
```



## Solution 2.4: The difference in solutions

```
1 # PLOTTING
2 fig, (ax1, ax2) = plt.subplots(1, 2, figsize=(12, 5))
3
4 # Plot 1: Explicit vs Numerical
5 ax1.plot(t_expl + start_year, P_expl, color="red",
6          label="Explicit")
7 ax1.plot(t_num + start_year, P_num, color="blue",
8          label="Numerical")
9 ax1.set_title("Explicit and numerical solution")
10 ax1.set_ylim(0, 20)
11 ax1.set_xlim(1950, 2020)
12 ax1.legend()
13
14 # Plot 2: Difference
15 ax2.plot(t_expl + start_year, P_expl - P_num, color="black")
16 ax2.set_title("The difference")
17 ax2.set_ylim(0, 0.5)
18 ax2.set_xlim(1950, 2020)
19
20 plt.tight_layout()
21 plt.show()
```





## Solution 2.4: The difference in solutions

Step size of 1 year leads to substantial error.





## Solution 2.4: The difference in solutions

Step size of 1 month reduces the error significantly.



## Solution 2.4: The difference in solutions

Step size of 1 day leads to negligible error.



## Task 2.5: Real world data

Compare the prediction modeled by the explicit formula with real data of the Senegalese population.

```
1 import numpy as np
2 import matplotlib.pyplot as plt
3
4 # Growth rate
5 r = 0.027
6
7 # Start year
8 start_year = 1950
9
10 # Time axis (years)
11 t_sen = np.array([0, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50,
12                  55, 60, 65, 70, 72, 73, 74])
13
14 # Senegalese population (millions)
```

## Solution 2.5: Real world data



```
1 import numpy as np
2 import matplotlib.pyplot as plt
3
4 # Growth rate
5 r = 0.027
6
7 # Start year
8 start_year = 1950
9
10 # Time axis (years)
11 t_sen = np.array([0, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50,
12                  55, 60, 65, 70, 72, 73, 74])
13
14 # Senegalese population (millions)
15 P_sen = np.array([2.5, 2.8, 3.3, 3.8, 4.4, 5.0, 5.7, 6.5,
16                  7.5, 8.6, 9.7, 11.0, 12.5, 14.4, 16.4, 17.3, 17.8,
17                  18.2])
18
19 # EXPLICIT SOLUTION
20 # Initial population is the first value: P_sen[0]
21 P_expl = P_sen[0] * np.exp(r * t_sen)
```

## Solution 2.5: Real world data

```
1
2 # PLOTTING
3 fig, (ax1, ax2) = plt.subplots(1, 2, figsize=(12, 5))
4
5 # Tile 1: Real data and model
6 ax1.plot(t_sen + start_year, P_expl, color="red",
7          label='Model')
8 ax1.plot(t_sen + start_year, P_sen, color="blue",
9          label='Real Data')
10 ax1.set_title("Real data and model")
11 ax1.set_ylim(0, 20)
12 ax1.legend()
13
14 # Tile 2: The difference
15 ax2.plot(t_sen + start_year, P_expl - P_sen, '-o',
16          color="black")
17 ax2.set_title("The difference")
18
19 plt.tight_layout()
20 plt.show()
```

## Solution 2.5: Real world data

Even a very simple model can capture reality very well.



## Task 2.6: Real world data — USA

- ▶ Compare the prediction modeled by the explicit formula for the USA with real data.
- ▶ Adjust the time span, start year, and initial population.
- ▶ Set growth rate to 2.75% (historical average for that specific start period).

```
1  # Parameters
2  r = 0.0275
3  start_year = 1790
4
5  # Time axis: equivalent to 0:10:230
6  t_usa = np.arange(0, 240, 10)
7
8  # US population data (Millions)
9  P_usa = np.array([
10     3.9, 5.3, 7.2, 9.6, 12.9, 17.1, 23.2, 31.4,
11     38.6, 50.2, 63.0, 76.2, 92.2, 106.0, 123.2, 132.2,
12     151.3, 179.3, 203.3, 226.5, 248.7, 281.4, 308.7, 331.4
13 ])
```



## Solution 2.6: Real world data — USA

```
1 import numpy as np
2 import matplotlib.pyplot as plt
3
4 # Parameters
5 r = 0.0275
6 start_year = 1790
7
8 # Time axis: equivalent to 0:10:230
9 t_usa = np.arange(0, 240, 10)
10
11 # US population data (Millions)
12 P_usa = np.array([
13     3.9, 5.3, 7.2, 9.6, 12.9, 17.1, 23.2, 31.4,
14     38.6, 50.2, 63.0, 76.2, 92.2, 106.0, 123.2, 132.2,
15     151.3, 179.3, 203.3, 226.5, 248.7, 281.4, 308.7, 331.4
16 ])
17
18 # EXPLICIT SOLUTION
19 #  $P(t) = P_0 * e^{(rt)}$ 
20 P_expl = P_usa[0] * np.exp(r * t_usa)
```



## Solution 2.6: Real world data — USA

```
1  # PLOTTING
2  years = t_usa + start_year
3  fig, (ax1, ax2) = plt.subplots(1, 2, figsize=(12, 5))
4
5  # Tile 1: Real data and model
6  ax1.plot(years, P_expl, color="red", label="Model")
7  ax1.plot(years, P_usa, color="black", label="Real Data")
8  ax1.set_title("Real data and model")
9  ax1.set_ylim(0, 350)
10 ax1.legend()
11
12 # Tile 2: The difference
13 ax2.plot(years, P_expl - P_usa, '-o', color="black")
14 ax2.set_title("The difference")
15
16 plt.tight_layout()
17 plt.show()
```

## Solution 2.6: Real world data — USA

The model is reasonably precise for the first one hundred years, then it diverges.

**Why?** (The growth rate slowed down; resources/space became less "unlimited").



## Section 3: Logistic Growth Model

# Logistic growth

The **logistic growth model** represents how populations change **over time when resources are limited**. The logistic model incorporates a carrying capacity—the maximum population size a specific environment can sustainably support.

# Logistic growth

The **logistic growth model** represents how populations change over time when resources are limited. The logistic model incorporates a carrying capacity—the maximum population size a specific environment can sustainably support.

## Key Concepts

- ▶ **Carrying Capacity ( $K$ ):** A ceiling that limits population growth due to limited resources (food, space, water, etc.).

# Logistic growth

The **logistic growth model** represents how populations change over time when resources are limited. The logistic model incorporates a carrying capacity—the maximum population size a specific environment can sustainably support.

## Key Concepts

- ▶ **Carrying Capacity ( $K$ ):** A ceiling that limits population growth due to limited resources (food, space, water, etc.).
- ▶ **Intrinsic Growth Rate ( $r$ ):** The rate at which a population would grow if there were unlimited resources.

# Logistic growth

The **logistic growth model** represents how populations change over time when resources are limited. The logistic model incorporates a carrying capacity—the maximum population size a specific environment can sustainably support.

## Key Concepts

- ▶ **Carrying Capacity ( $K$ ):** A ceiling that limits population growth due to limited resources (food, space, water, etc.).
- ▶ **Intrinsic Growth Rate ( $r$ ):** The rate at which a population would grow if there were unlimited resources.
- ▶ **S-Shaped Curve:** The logistic growth model plots as an S-shaped curve. Growth is initially exponential, then slows as it approaches the carrying capacity.



# Equation of logistic growth

Differential equation of logistic growth

$$\frac{dP}{dt} = rP \left( 1 - \frac{P}{K} \right)$$

# Equation of logistic growth

Differential equation of logistic growth

$$\frac{dP}{dt} = rP \left( 1 - \frac{P}{K} \right)$$

Iterative Form:

$$P(t_{i+1}) = P(t_i) + rP(t_i) \left( 1 - \frac{P(t_i)}{K} \right) \Delta t$$

# Equation of logistic growth

Differential equation of logistic growth

$$\frac{dP}{dt} = rP \left( 1 - \frac{P}{K} \right)$$

Iterative Form:

$$P(t_{i+1}) = P(t_i) + rP(t_i) \left( 1 - \frac{P(t_i)}{K} \right) \Delta t$$

Explicit solution

$$P(t) = \frac{K}{1 + \left( \frac{K - P_0}{P_0} \right) e^{-rt}}$$

# Equation of logistic growth

Differential equation of logistic growth

$$\frac{dP}{dt} = rP \left( 1 - \frac{P}{K} \right)$$

Iterative Form:

$$P(t_{i+1}) = P(t_i) + rP(t_i) \left( 1 - \frac{P(t_i)}{K} \right) \Delta t$$

Explicit solution

$$P(t) = \frac{K}{1 + \left( \frac{K - P_0}{P_0} \right) e^{-rt}}$$

where

- $P(t_i)$  is the population at time  $t_i$

# Equation of logistic growth

Differential equation of logistic growth

$$\frac{dP}{dt} = rP \left( 1 - \frac{P}{K} \right)$$

Iterative Form:

$$P(t_{i+1}) = P(t_i) + rP(t_i) \left( 1 - \frac{P(t_i)}{K} \right) \Delta t$$

Explicit solution

$$P(t) = \frac{K}{1 + \left( \frac{K - P_0}{P_0} \right) e^{-rt}}$$

where

- ▶  $P(t_i)$  is the population at time  $t_i$
- ▶  $P_0$  is the initial population

# Equation of logistic growth

Differential equation of logistic growth

$$\frac{dP}{dt} = rP \left( 1 - \frac{P}{K} \right)$$

Iterative Form:

$$P(t_{i+1}) = P(t_i) + rP(t_i) \left( 1 - \frac{P(t_i)}{K} \right) \Delta t$$

Explicit solution

$$P(t) = \frac{K}{1 + \left( \frac{K - P_0}{P_0} \right) e^{-rt}}$$

where

- ▶  $P(t_i)$  is the population at time  $t_i$
- ▶  $P_0$  is the initial population
- ▶  $r$  is the growth rate

# Equation of logistic growth

Differential equation of logistic growth

$$\frac{dP}{dt} = rP \left( 1 - \frac{P}{K} \right)$$

Iterative Form:

$$P(t_{i+1}) = P(t_i) + rP(t_i) \left( 1 - \frac{P(t_i)}{K} \right) \Delta t$$

Explicit solution

$$P(t) = \frac{K}{1 + \left( \frac{K - P_0}{P_0} \right) e^{-rt}}$$

where

- ▶  $P(t_i)$  is the population at time  $t_i$
- ▶  $P_0$  is the initial population
- ▶  $r$  is the growth rate
- ▶  $K$  is the carrying capacity

# Equation of logistic growth

Differential equation of logistic growth

$$\frac{dP}{dt} = rP \left( 1 - \frac{P}{K} \right)$$

Iterative Form:

$$P(t_{i+1}) = P(t_i) + rP(t_i) \left( 1 - \frac{P(t_i)}{K} \right) \Delta t$$

Explicit solution


$$P(t) = \frac{K}{1 + \left( \frac{K - P_0}{P_0} \right) e^{-rt}}$$

where



- ▶  $P(t_i)$  is the population at time  $t_i$
- ▶  $P_0$  is the initial population
- ▶  $r$  is the growth rate
- ▶  $K$  is the carrying capacity
- ▶  $t$  is time,  $\Delta t$  is the time step





# Logistic growth of population over time



# Example — French and English railway construction





# Example — French and English railway construction



## Topics for discussion

- Why do you think that the French railway construction follows logistic growth but the English does not?



# Example — French and English railway construction



## Topics for discussion

- ▶ Why do you think that the French railway construction follows logistic growth but the English does not?
- ▶ How would you explain **carrying capacity** in the context of railroad construction?

# Example — French and English railway construction



## Topics for discussion

- ▶ Why do you think that the French railway construction follows logistic growth but the English does not?
- ▶ How would you explain **carrying capacity** in the context of railroad construction?
- ▶ What may affect the carrying capacity for railways? Is the capacity constant or dynamic?

# Closer look at logistic growth equation

Differential equation of logistic growth

$$\frac{dP}{dt} = rP \left( 1 - \frac{P}{K} \right)$$

**Let's explore the equation**

- ▶ Try to figure out how the carrying capacity affects population growth. Try different combinations of P and K.

# Closer look at logistic growth equation

Differential equation of logistic growth

$$\frac{dP}{dt} = rP \left( 1 - \frac{P}{K} \right)$$

**Let's explore the equation**

- ▶ Try to figure out how the carrying capacity affects population growth. Try different combinations of  $P$  and  $K$ .
- ▶ What are the equilibrium solutions where there is no population growth?

# Derivation of Equilibrium Solutions

To find equilibrium solutions, we set  $\frac{dP}{dt} = 0$ :

$$0 = rP \left( 1 - \frac{P}{K} \right)$$



# Derivation of Equilibrium Solutions

To find equilibrium solutions, we set  $\frac{dP}{dt} = 0$ :

$$0 = rP \left( 1 - \frac{P}{K} \right)$$

This equation is satisfied when either:

- The population is zero

$$P = 0$$

# Derivation of Equilibrium Solutions

To find equilibrium solutions, we set  $\frac{dP}{dt} = 0$ :

$$0 = rP \left( 1 - \frac{P}{K} \right)$$

This equation is satisfied when either:

- The population is zero

$$P = 0$$

- The growth rate is zero

$$r = 0$$

# Derivation of Equilibrium Solutions

To find equilibrium solutions, we set  $\frac{dP}{dt} = 0$ :

$$0 = rP \left( 1 - \frac{P}{K} \right)$$

This equation is satisfied when either:

- The population is zero

$$P = 0$$

- The growth rate is zero

$$r = 0$$

- The population size equals the carrying capacity

$$1 - \frac{P}{K} = 0$$

$$1 = \frac{P}{K}$$

$$P = K$$


- ▶ **Pierre Verhulst:** A Belgian mathematician who first proposed the logistic model in the 1830s and 1840s. His model sought to refine the exponential population growth model of Malthus.

# History

- ▶ **Pierre Verhulst:** A Belgian mathematician who first proposed the logistic model in the 1830s and 1840s. His model sought to refine the exponential population growth model of Malthus.
- ▶ Verhulst got inspiration from the concept of intraspecific competition—the idea that as populations get larger, individuals compete more intensely for limited resources.

# History

- ▶ **Pierre Verhulst:** A Belgian mathematician who first proposed the logistic model in the 1830s and 1840s. His model sought to refine the exponential population growth model of Malthus.
- ▶ Verhulst got inspiration from the concept of intraspecific competition—the idea that as populations get larger, individuals compete more intensely for limited resources.
- ▶ This equation was published after Verhulst had read Thomas Malthus' *An Essay on the Principle of Population*.



Pierre

*Verhulst and the first plot of the logistic growth curve*

# The logistic growth model has extensive applications

- ▶ **Biology:** Used extensively in modeling animal, plant, and bacterial population growth.

# The logistic growth model has extensive applications

- ▶ **Biology:** Used extensively in modeling animal, plant, and bacterial population growth.
- ▶ **Epidemiology and medicine:** Helps analyze how diseases spread within a population or tumor growth.



# The logistic growth model has extensive applications

- ▶ **Biology:** Used extensively in modeling animal, plant, and bacterial population growth.
- ▶ **Epidemiology and medicine:** Helps analyze how diseases spread within a population or tumor growth.
- ▶ **Economics:** Modeling market adoption of new products or technologies—a diffusion of innovation.

# The logistic growth model has extensive applications

- ▶ **Biology:** Used extensively in modeling animal, plant, and bacterial population growth.
- ▶ **Epidemiology and medicine:** Helps analyze how diseases spread within a population or tumor growth.
- ▶ **Economics:** Modeling market adoption of new products or technologies—a diffusion of innovation.
- ▶ **Linguistics:** An innovation that is at first marginal begins to spread more quickly with time, and then more slowly as it becomes more universally adopted.

# The logistic growth model has extensive applications

- ▶ **Biology:** Used extensively in modeling animal, plant, and bacterial population growth.
- ▶ **Epidemiology and medicine:** Helps analyze how diseases spread within a population or tumor growth.
- ▶ **Economics:** Modeling market adoption of new products or technologies—a diffusion of innovation.
- ▶ **Linguistics:** An innovation that is at first marginal begins to spread more quickly with time, and then more slowly as it becomes more universally adopted.
- ▶ **Statistics and machine learning:** Logistic functions are often used in artificial neural networks to introduce nonlinearity in the model—but it is mostly for the shape.

# Tasks overview

## 1. Numerically solve the logistic growth equation and plot:

- Population size vs. time
- Population increase vs. time
- Per capita growth rate vs. population size
- Population increase vs. population size

# Tasks overview

## 1. Numerically solve the logistic growth equation and plot:

- Population size vs. time
- Population increase vs. time
- Per capita growth rate vs. population size
- Population increase vs. population size

## 2. Adjust the appearance of the plot:

- Add a plot title and axis labels
- Change line thickness and appearance
- Experiment with other changes

# Tasks overview





1. Numerically solve the logistic growth equation and plot:
  - Population size vs. time
  - Population increase vs. time
  - Per capita growth rate vs. population size
  - Population increase vs. population size
2. Adjust the appearance of the plot:
  - Add a plot title and axis labels
  - Change line thickness and appearance
  - Experiment with other changes
3. Experiment with model parameters:
  - Try to increase the initial population over the capacity of the environment
4. Model the population\* of the USA and compare it with real data.

\* For the population model of the USA, set the step size to 1 month, the initial population size to 5.2 million in 1790, the growth rate to 0.0237, and the capacity of the environment to 460 million.

## Solution 3.1: Logistic growth model

```
1 import numpy as np
2 import matplotlib.pyplot as plt
3
4 # Parameters and Time
5 P0, r, K = 3.9, 0.0255, 430
6 t = np.arange(0, 400, 1/12)
7
8 # Vectorized Logistic Growth (Analytical Solution)
9 #  $P(t) = K / (1 + ((K - P0) / P0) * \exp(-rt))$ 
10 P = K / (1 + ((K - P0) / P0) * np.exp(-r * t))
11 P_diff = r * P * (1 - P / K) * (1/12)
12
13 # Plotting
14 fig, axs = plt.subplots(2, 2, figsize=(10, 8))
15 plots = [
16     (t, P, "Population vs Time"),
17     (t, P_diff, "Growth Amount vs Time"),
18     (P, P_diff / P, "Per Capita Growth vs Pop"),
19     (P, P_diff, "Growth Amount vs Pop")
20 ]
21
22 for ax, (x, y, title) in zip(axs.flat, plots):
23     ax.plot(x, y)
24     ax.set_title(title)
```

## Solution 3.1: Logistic growth model









## Solution 3.2: Adjusted plot

```
1 import numpy as np
2 import matplotlib.pyplot as plt
3
4 # Params & Time
5 P0, r, K, dt = 3.9, 0.0255, 430, 1/12
6 t = np.arange(0, 400 + dt, dt)
7 P = np.zeros(len(t))
8 P[0] = P0
9
10 for i in range(len(t) - 1):
11     P[i+1] = P[i] + r * P[i] * (1 - P[i] / K) * dt
12
13 # Derived Data
14 P_diff = np.concatenate([[np.nan], np.diff(P)])
15 plots = [(t, P, "Pop vs Time"), (t, P_diff, "Growth vs
16         Time"), (P, P_diff/P, "Per Capita vs Pop"), (P,
17         P_diff, "Growth vs Pop")]
18
19 fig, axs = plt.subplots(2, 2, figsize=(10, 7))
20 for ax, (x, y, title) in zip(axs.flat, plots):
21     ax.plot(x, y)
22     ax.set_title(title)
23
24 plt.tight_layout()
25 plt.show()
```


# Solution 3.2: Adjusted plot



## Solution 3.3: US population

```
1 import numpy as np
2 import matplotlib.pyplot as plt
3 # Parameters
4 r, P0, K, start_year = 0.0237, 5.2, 460, 1790
5 t_usa = np.arange(0, 240, 10)
6 P_usa = np.array([3.9, 5.3, 7.2, 9.6, 12.9, 17.1, 23.2,
7                   31.4, 38.6, 50.2, 63.0, 76.2, 92.2, 106.0, 123.2,
8                   132.2, 151.3, 179.3, 203.3, 226.5, 248.7, 281.4, 308.7,
9                   331.4])
10
11 dt = 1/12
12 t_num = np.arange(0, 230 + dt, dt)
13 P_num = np.zeros(len(t_num))
14 P_num[0] = P0
15
16 for i in range(len(t_num) - 1):
17     P_num[i+1] = P_num[i] + r * P_num[i] * (1 - P_num[i] /
18         K) * dt
19
20 # Plotting
21 plt.plot(t_num + start_year, P_num, 'r-', label='Logistic
22     Model')
23 plt.plot(t_usa + start_year, P_usa, 'ko', label='Real Data')
24 plt.axis([1790, 2020, 0, 350])
25 plt.legend(): plt.title("USA Population: Logistic Growth
```

## Solution 3.3: US population



## Section 4: Leslie Matrix Model

# Limitations of previous models

- ▶ So far, our models (Malthusian, Logistic) assumed all individuals in a population are identical.
- ▶ **In reality, age matters:**
  - **Reproduction:** Young individuals (larvae, babies) cannot reproduce yet.
  - **Survival:** Survival rates often depend on age (e.g., higher mortality in very young or very old individuals).
- ▶ To capture this, we need an **age-structured model**.
- ▶ The most common discrete-time model for this is the **Leslie Matrix**.

# The Leslie Matrix Model

Developed by Patrick H. Leslie in 1945.

- ▶ It tracks the number of females in different age classes at discrete time steps.
- ▶ **Variables:**
  - $n_i(t)$ : Number of individuals in age class  $i$  at time  $t$ .
  - $F_i$ : **Fecundity** (fertility) rate. Average number of female offspring produced by a female in age class  $i$ .
  - $S_i$ : **Survival** rate. Probability that a female in age class  $i$  survives to age class  $i + 1$ .

# Mathematical Formulation

If we have 3 age classes, the population vector at time  $t + 1$  is calculated as:

$$n_1(t + 1) = F_1 n_1(t) + F_2 n_2(t) + F_3 n_3(t) \quad \text{New offspring (born from all classes)}$$

$$n_2(t + 1) = S_1 n_1(t) \quad \text{Survivors from class 1 moving to 2}$$

$$n_3(t + 1) = S_2 n_2(t) \quad \text{Survivors from class 2 moving to 3}$$

This can be written in matrix form:


$$\begin{bmatrix} n_1 \\ n_2 \\ n_3 \end{bmatrix}_{t+1} = \begin{bmatrix} F_1 & F_2 & F_3 \\ S_1 & 0 & 0 \\ 0 & S_2 & 0 \end{bmatrix} \begin{bmatrix} n_1 \\ n_2 \\ n_3 \end{bmatrix}_t$$

$$\mathbf{n}_{t+1} = \mathbf{L} \cdot \mathbf{n}_t$$



# Life Cycle with Age-Class Structure

Individuals are grouped into discrete age classes of equal duration.



- **Circles:** Represent specific age classes or ages (1 to 4).
- **Horizontal Arrows ( $S_x$ ):** Represent **survival probabilities**. Probability that an individual of age  $x$  survives to age  $x + 1$ . There is no survival beyond group 4.
- **Curved Arrows ( $F_x$ ):** Represent **births (fecundity)**. Arrows lead back to Age 1 because newborns enter the first class. Arrows emerging from ages 2, 3, and 4 indicate these ages are reproductive.

# Matrix Representation of the Life Cycle

The life cycle diagram translates directly into a **projection matrix** (Leslie Matrix).

**The Equation:**

$$n_{t+1} = L \cdot n_t$$

**The Matrix Expansion:**

$$\begin{bmatrix} n_1 \\ n_2 \\ n_3 \\ n_4 \end{bmatrix}_{t+1} = \begin{bmatrix} 0 & F_2 & F_3 & F_4 \\ S_1 & 0 & 0 & 0 \\ 0 & S_2 & 0 & 0 \\ 0 & 0 & S_3 & 0 \end{bmatrix} \begin{bmatrix} n_1 \\ n_2 \\ n_3 \\ n_4 \end{bmatrix}_t$$

# Structure of the Leslie Matrix

$$L = \begin{bmatrix} F_1 & F_2 & F_3 & \dots & F_k \\ S_1 & 0 & 0 & \dots & 0 \\ 0 & S_2 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \dots & \vdots \\ 0 & 0 & \dots & S_{k-1} & 0 \end{bmatrix}$$

- ▶ **Top Row:** Fertility rates (production of new offspring).
- ▶ **Sub-diagonal:** Survival rates (transition to next age group).
- ▶ **All other entries:** Zero (you cannot skip an age or get younger).

What happens if we simulate this for many time steps?

- ▶ **Population Growth:** The total population eventually grows (or declines) at a constant rate  $\lambda$ .
- ▶ **Stable Age Distribution:** The *proportion* of individuals in each age class becomes constant.
  - Example: 60% juveniles, 30% adults, 10% seniors.
  - This happens regardless of the initial population state!

*Note for linear algebra enthusiasts:  $\lambda$  is the dominant eigenvalue of  $L$ , and the stable age distribution is the corresponding eigenvector.*

# The Finite Rate of Increase ( $\lambda$ )

As the population approaches the Stable Age Distribution, it begins to grow (or decline) at a constant rate every time step.

We call this rate  $\lambda$  (lambda).

$$N_{total}(t + 1) = \lambda \cdot N_{total}(t)$$

## Biological Interpretation:

- ▶  $\lambda > 1$ : The population is **growing** exponentially.
- ▶  $\lambda = 1$ : The population is **stable** (stationary).
- ▶  $\lambda < 1$ : The population is **declining** to extinction.

# Importance of $\lambda$

Why is calculating  $\lambda$  so critical in conservation and ecology?

- ▶ **Conservation Status:** Determines if an endangered species is recovering ( $\lambda > 1$ ) or needs intervention ( $\lambda < 1$ ).
- ▶ **Pest Control:** To eradicate a pest, we must lower survival/fertility until  $\lambda < 1$ .
- ▶ **Harvesting:** In fisheries or forestry, we want to harvest the "excess" without driving  $\lambda$  below 1.
- ▶ **Comparative Analysis:** Allows us to compare population health across different habitats regardless of current population size.

# How to Calculate $\lambda$

There are two main ways to find  $\lambda$ :

## 1. The Analytical Method (Eigenvalues)

- ▶  $\lambda$  is the **dominant eigenvalue** (largest absolute value) of the Leslie Matrix  $L$ .
- ▶ Requires linear algebra:  $\det(L - \lambda I) = 0$ .

## 2. The Numerical Method (Simulation)

- ▶ If we simulate the population long enough, the ratio between consecutive years converges to  $\lambda$ .
- ▶  $\lambda \approx \frac{N(t+1)}{N(t)}$  (for large  $t$ ).

## Task 4.1: Defining the Model

### Scenario: A population of hypothetical birds

- ▶ **Class 1 (Chicks, 0-1 yr):** Do not reproduce ( $F_1 = 0$ ). 60% survive to next year ( $S_1 = 0.6$ ).
- ▶ **Class 2 (Young Adults, 1-2 yr):** Moderate reproduction ( $F_2 = 1.5$ ). 50% survive ( $S_2 = 0.5$ ).
- ▶ **Class 3 (Older Adults, 2-3 yr):** High reproduction ( $F_3 = 2.0$ ). They die after this year ( $S_3 = 0$ ).

### Your Task:

1. Construct the Leslie Matrix  $L$  in Python (using NumPy).
2. Define an initial population vector  $n_0$ : 100 chicks, 0 adults.



## Solution 4.1: Defining the Model

```
1 import numpy as np
2
3 # Define parameters
4 F1 = 0;   F2 = 1.5; F3 = 2.0
5 S1 = 0.6; S2 = 0.5
6
7 # Construct Leslie Matrix
8 L = np.array([[F1, F2, F3],
9               [S1, 0,  0],
10              [0,  S2, 0]])
11
12 # Initial population (100 chicks, 0 young adults, 0 old
13   adults)
14 n0 = np.array([100, 0, 0])
```

## Task 4.2: Simulation Loop

### Your Task:

- ▶ Simulate the population for 20 years.
- ▶ Store the population vector for each year.
- ▶ *Hint: You will need a matrix to store results, where columns represent time steps.*

## Solution 4.2: Simulation Loop

```
1 years = 20
2 # Pre-allocate matrix (3 rows, years+1 columns)
3 pop_history = np.zeros((3, years + 1))
4
5 # Set initial year (Python is 0-indexed)
6 pop_history[:, 0] = n0
7
8 # Loop through time
9 for t in range(years):
10     #  $n(t+1) = L @ n(t)$ 
11     # The @ operator performs matrix multiplication
12     pop_history[:, t+1] = L @ pop_history[:, t]
```

## Task 4.3: Analyzing Results

### Your Task:

1. Calculate the **total population** for each year (sum of all age classes).
2. Plot the Total Population over time.
3. Plot the specific age classes over time.

## Solution 4.3: Plotting

```
1 import matplotlib.pyplot as plt
2
3 # Calculate total population (sum each column, axis=0)
4 total_pop = np.sum(pop_history, axis=0)
5 t = np.arange(years + 1)
6
7 # Plot
8 plt.figure(figsize=(10, 6))
9
10 plt.subplot(2,1,1)
11 plt.plot(t, total_pop, 'k-', linewidth=2)
12 plt.title('Total Population Growth'); plt.grid(True)
13
14 plt.subplot(2,1,2)
15 plt.plot(t, pop_history[0, :], 'b-o', label='Chicks')
16 plt.plot(t, pop_history[1, :], 'r-o', label='Young')
17 plt.plot(t, pop_history[2, :], 'g-o', label='Old')
18 plt.legend(); plt.grid(True)
19 plt.show()
```

## Task 4.4: Stable Age Distribution (Advanced)

We claimed earlier that the *proportion* of age groups becomes constant. **Your Task:**

- ▶ Normalize the population data at each step (divide each column by the total population of that column).
- ▶ Inspect the last few columns of this normalized matrix.
- ▶ Are the proportions changing?

## Solution 4.4: Stable Age Distribution

```
1 # Normalize to find proportions
2 # axis=0 means we sum down the columns
3 proportions = pop_history / np.sum(pop_history, axis=0)
4
5 # Display the proportions for the last year (-1 index)
6 print("Final Age Distribution:")
7 print(proportions[:, -1])
8
9 # Check the previous year to see if it stabilized
10 print("Previous Year Distribution:")
11 print(proportions[:, -2])
```

*You should see that the vector values stabilize, confirming the Stable Age Distribution theorem.*

## Task 4.5: Calculating $\lambda$ in Python

### Your Task:

1. **Method A (Simulation):** Calculate the growth rate using the total population from the *last two years* of your simulation.

$$\lambda_{sim} = \frac{\text{Total Pop (Year 20)}}{\text{Total Pop (Year 19)}}$$

2. **Method B (Linear Algebra):** Use NumPy's linear algebra library to calculate the eigenvalues of matrix  $L$  directly.
3. Compare the two values.



## Solution 4.5: Calculating $\lambda$

```
1 # --- Method A: From Simulation ---
2 # Total population at last step / Total population at 2nd
  to last step
3 lambda_sim = total_pop[-1] / total_pop[-2]
4 print(f"Simulated Lambda: {lambda_sim:.4f}")
5
6 # --- Method B: Eigenanalysis ---
7 # Calculate eigenvalues
8 eigenvalues = np.linalg.eigvals(L)
9
10 # The dominant eigenvalue is the maximum real value
11 lambda_eig = np.max(np.real(eigenvalues))
12 print(f"Theoretical Lambda: {lambda_eig:.4f}")
13
14 # They should be nearly identical!
```

Note: `'np.linalg.eigvals'` returns complex numbers. We use `'np.real'` and `'np.max'` to find the dominant growth rate.