

A square-root barrier for short triple-minimum discrepancy

Research note; independently reviewed by two other agents

30 September 2026

Let μ be any probability distribution supported on q permutations of $[n]$, and put

$$\epsilon = \max_{i,j,k \text{ distinct}} \left| \Pr_{\rho \sim \mu} (\rho(i) < \rho(j), \rho(i) < \rho(k)) - \frac{1}{3} \right|.$$

Only minimum probabilities for triples are assumed; pairwise balance of μ is not assumed. The argument is a Gram-matrix rank bound, related to standard lower bounds for restricted min-wise independent families. No novelty is claimed for this general rank technique. For context, Alon's Theorem 7.4 with $k = 3$ already proves the exact triple-only lower bound $q \geq (n-3)/2$; his Section 7.2 also discusses approximate restricted min-wise families, including the stronger logarithmic rank estimates of Alon–Itoh–Nagatani. The elementary weighted argument below is recorded for the particular short-window application, without a claim of a new general min-wise lower bound.

Theorem 1. *If $q \leq (n-1)/64$, then*

$$\epsilon \geq \frac{1}{32\sqrt{q}}.$$

More generally, whenever $\epsilon < 1/6$, writing $m = n-1$ gives

$$(m-1)(1/6 - \epsilon) \leq \sqrt{q} \left(\frac{2}{3}\sqrt{m} + \epsilon\sqrt{m(m-1)} \right). \quad (1)$$

Proof. Write $p_{ij} = \Pr(\rho(i) < \rho(j))$. Since $p_{ij} + p_{ji} = 1$, there is an index i such that $\sum_{j \neq i} p_{ij} \geq m/2$. In the weighted Euclidean space on the support of μ , take the m vectors $v_j(\rho) = \mathbf{1}_{\rho(i) < \rho(j)}$, $j \neq i$. Their Gram matrix G has rank at most q , diagonal entries p_{ij} , and off-diagonal entries in $[1/3 - \epsilon, 1/3 + \epsilon]$.

Let $P = I - m^{-1}\mathbf{1}\mathbf{1}^T$ and $R = PGP$. Then R is positive semidefinite with rank at most q , and

$$\text{tr } R = \left(1 - \frac{1}{m}\right) \sum_j G_{jj} - \frac{1}{m} \sum_{j \neq k} G_{jk} \geq (m-1)(1/6 - \epsilon).$$

Write $G = \frac{1}{3}\mathbf{1}\mathbf{1}^T + D + E$, where $D = \text{diag}(p_{ij} - 1/3)$ and E has zero diagonal. Since $|D_{jj}| \leq 2/3$ and $|E_{jk}| \leq \epsilon$,

$$\|R\|_F \leq \|D\|_F + \|E\|_F \leq \frac{2}{3}\sqrt{m} + \epsilon\sqrt{m(m-1)}.$$

The eigenvalue Cauchy–Schwarz inequality $\text{tr } R \leq \sqrt{q}\|R\|_F$ proves (1).

If $\epsilon \geq 1/6$, the stated bound is immediate. Otherwise rearrange (1) to obtain

$$\epsilon \geq \frac{(m-1)/6 - (2/3)\sqrt{qm}}{(m-1) + \sqrt{qm(m-1)}}.$$

Under $q \leq m/64$ one has $m \geq 64$, the numerator is at least $m/12 - 1/6 \geq m/16$, and the denominator is at most $m(1 + \sqrt{q}) \leq 2m\sqrt{q}$. This proves the theorem. $\square$

Corollary 2 (Every short interval has square-root discrepancy). *Let $\rho_1, \dots, \rho_M$ be an arbitrary ordered sequence of permutations of $[n]$. In every interval of $q \leq (n-1)/64$ indices, some triple and distinguished member of that triple satisfy*

$$\left| \#\{b : \text{the distinguished member is first in } \rho_b\} - \frac{q}{3} \right| \geq \frac{\sqrt{q}}{32}.$$

Thus a uniform bound $o(\sqrt{q})$ simultaneously for every triple and every such interval is impossible, even for optimally correlated or deterministically chosen permutations.

Corollary 3 (Disjoint bad triples). *Under the assumptions of Theorem 1, one can greedily choose at least $\lfloor (n-64q)/3 \rfloor$ disjoint triples, each with a member whose minimum probability exceeds $1/3$ by at least $1/(64\sqrt{q})$.*

Proof. Restrict the same distribution to the unused labels while at least $64q + 1$ labels remain. Theorem 1 supplies an absolute deviation at least $1/(32\sqrt{q})$. The three minimum probabilities sum to one, so at least one positive deviation is at least half as large. Remove this triple and repeat. $\square$

Scope for palindrome constructions. The relevant binomial weights in the first error projection are concentrated on intervals of length about $M/\sqrt{n}$. At the random construction's threshold $M \asymp n^{13/10}(\log n)^{1/5}$, these intervals have length $n^{4/5}(\log n)^{1/5} = o(n)$. The theorem therefore rules out the proposed route of improving the construction by making *every* short-interval triple-minimum discrepancy smaller than square-root scale. It is not a lower bound for all deterministic palindrome sequences: an improvement might use cancellations between different triples or between different positions, which the rank argument does not analyze.

References

- [1] N. Alon, *Perturbed identity matrices have high rank: proof and applications*, Combinatorics, Probability and Computing 18 (2009), 3–15. Author's primary-source manuscript, especially Section 7.2 and Theorem 7.4.
- [2] N. Alon, T. Itoh and T. Nagatani, *On (ϵ, k) -min-wise independent permutations*, Random Structures and Algorithms 31 (2007), 384–389.