

Scheduling with Uniform Tests: Information and Optimization

Václav Rozhoň

August 2026

Abstract

We study single-machine scheduling with unit-time tests and total completion time. A test reveals a job's unknown processing time; without a test, a job may either be unavailable, run for a common upper bound u , or run blindly for its actual duration, depending on the model.

For deterministic algorithms we determine the exact size-asymptotic ratio for every common upper bound $u \in (0, \infty]$. The answer is a continuous six-piece curve whose cap-free obligatory endpoint is $1.5705740966\dots$ and whose global maximum is $1.8667603992\dots$. Against an oblivious adversary, we prove the exact randomized finite-cap curve: it has four pieces, global maximum $1.6257523846\dots$, and equals $4/3$ for every $u \geq 25/4$. The curve is the worst-case envelope of a stronger instance-optimal value for every processing-time multiset. A separate uniform argument proves the same sharp $4/3$ value for obligatory testing with unbounded, unannounced processing times.

For blind execution we prove a stronger instance-optimal statement at the quadratic scale. A universal policy learns, from a sublinear pilot, the same explicit fluid value attainable with advance knowledge of the multiset; a matching oblivious lower bound holds even for announced adaptive policies. The proofs combine pairwise delay accounting, completion-density thresholds, fractional-knapsack envelopes, random relabelling, and sampling without replacement.

Contents

1	Introduction and proof intuition	3
1.1	The common core: obligatory testing	3
1.2	Beyond a worst-case ratio: instance optimality	4
1.3	The information view: blind execution	5
1.4	The optimization view: a common upper bound	5
1.5	Techniques	6
1.6	Related work	7
2	OLD intro	8
2.1	The exact answers	8
2.2	Why total completion time leads to pair accounting	10
2.3	Lower-bound intuition: hide the short jobs until the end	11
2.4	Upper-bound intuition: three policies and one common bank	12
2.5	Randomized intuition: completion density and sampling	13
2.6	Blind execution: from tests to a fractional knapsack	14
2.7	Why the infinite endpoint needs a uniform proof	14
2.8	How to read the proofs	14

3	Models, benchmarks, and common preliminaries	15
3.1	Information and adversaries	16
3.2	Offline optima and pair accounting	16
3.3	Competitive-ratio conventions	17
3.4	Test-order adversaries	17
4	The cap-free endpoint $u = \infty$	18
4.1	Harmonic core and matching lower bound	18
4.2	Upper bound: a self-contained proof	23
5	Finite caps: the exact curve	31
5.1	The binary stopping game and its lower envelope	33
5.2	Matching upper bounds through the binary regime	35
5.3	The unrestricted mixed lower bound	39
5.4	The large-upper-limit lower bound by cap-free transfer	41
5.5	Completion of the curve and consistency at the joins	42
5.6	Matching upper bound for the common-upper model	42
6	Randomized common upper bounds: the exact four-piece curve	49
6.1	Offline and stationary fluid coefficients	49
6.2	Flattening the survival function	50
6.3	The two scalar maximizations	52
6.4	Learning the announced policy	53
6.5	Binary oblivious lower bounds	53
7	Randomized obligatory testing: the sharp ratio is $4/3$	56
7.1	The normalized offline objective	56
7.2	Stationary prefixes	56
7.3	Learning the prefix from a sublinear sample	58
7.4	Good samples, fallback, and bad-event excess	59
7.5	The balanced binary lower bound	60
8	Blind execution and instance-optimal two-phase policies	61
8.1	The distributional benchmark	61
8.2	The fixed-fraction completion envelope	63
8.3	Adaptive first touches are a uniform urn	64
8.4	The announced-multiset lower bound	65
8.5	A growing grid and arbitrary bounded multisets	66
8.6	Removing the announcement	66
8.7	Why the medium block is necessary	67
9	Machine-checked verification	68
9.1	Deterministic exact ratios	68
9.2	Randomized obligatory testing	68
9.3	Blind-execution instance optimality	68
10	Discussion and open questions	69
11	Conclusion	70
	Acknowledgements	70

A	Finite rational checkpoint for the cap-free lower bound	71
B	Independent certificate by bubbling caps to a prefix	72
C	Machine-checked deterministic theorem and its correspondence	74
C.1	Objects on the two sides	74
C.2	Expansion of the lower and upper fields	75
C.3	Correspondence with the two deterministic theorems	76
C.4	Proof boundary	76

1 Introduction and proof intuition

Suppose that several jobs are waiting for one machine. A job can be *tested* for one unit of time, after which its processing time becomes known. Testing sounds helpful, but its cost is paid at the worst possible place: while the machine tests one job, every unfinished job waits. Our question is how much information or speed-up is worth buying before the machine should simply start completing jobs.

There are two natural interpretations. In the *information* view, an untested job may be run blindly for its true duration; testing only tells us that duration before we commit to the job. In the *optimization* view, an unoptimized job takes a common time u , whereas a test unlocks an optimized duration $p_i \leq u$. Requiring every job to be tested is the clean common core: it isolates the conflict between learning the instance and finishing it. Throughout the paper there is one nonpreemptive machine, all jobs are available at time zero, and the objective is total completion time.

The three models differ only in what can be done before a test:

model	untested action	role of a test
obligatory	none	make the job available
blind execution	run for the true p_i	learn p_i before running
common upper u	run for time u	unlock processing at the revealed p_i

Our results give exact worst-case ratios in the obligatory and common-upper models. They also give a stronger, instance-dependent answer for randomized algorithms: after a sublinear pilot sample, advance knowledge of the entire multiset is worth only $o(n^2)$.

1.1 The common core: obligatory testing

We begin with the most constrained model. Every job must first undergo its unit test. A positive outcome p_i then requires another operation of length p_i ; a zero job completes with its test. After seeing an outcome, the scheduler must decide whether to process it now or let it wait while more jobs are tested.

Competitive analysis compares the online cost with the clairvoyant optimum, which knows all processing times in advance. We study the leading quadratic coefficient: an additive $o(n^2)$ loss is ignored. For deterministic algorithms this is the usual adversarial game. For randomized algorithms the input is fixed before the private random bits are drawn—an *oblivious* adversary.

Theorem 1.1 (Obligatory testing, informal). *The exact size-asymptotic deterministic competitive ratio is $1.5705740966\dots$. Against an oblivious adversary, the exact randomized ratio is $4/3$, even when processing times are unbounded and their multiset is not announced.*

The two optimal algorithms look rather different. The deterministic algorithm tests jobs in a fixed order and uses a threshold that changes with the observed history. Short outcomes are completed immediately; long ones are deferred and sorted in a final SPT tail. The threshold remembers how many positive jobs have appeared and how much work has already been promised to the tail. Its analysis uses a bank account that pays for every pair of jobs delayed in the wrong order.

The randomized algorithm first samples a sublinear number of jobs. From the sample it learns the processing-time prefix with the best *completion density*: completed mass divided by testing and immediate processing work. It then tests the remaining jobs in random order, completes outcomes below the learned threshold immediately, and defers the rest. A fixed cutoff prevents a missed enormous job from contaminating the discovery phase; if the learned density is too small, the algorithm safely falls back to testing everything and sorting afterwards.

The lower bounds explain why these choices are necessary. Against a deterministic algorithm, decreasing positive levels are revealed from long to short and a large block of zeros is hidden until the end. Harmonic gaps balance the cost of processing a positive job too early against the cost of deferring it. Against a randomized algorithm, the hard multiset is much simpler: half the jobs have length zero and half have length two, with a uniformly random labelling. Until many labels have been inspected, every unit of work completes only about half a job; integrating this completion envelope gives the factor $4/3$. The precise statements and proofs appear in Theorems 2.1 and 2.4 and Sections 4 and 7.

1.2 Beyond a worst-case ratio: instance optimality

A single competitive ratio deliberately forgets the instance. This can be misleading here. On some multisets testing everything and sorting is nearly optimal; on others testing is almost useless. An algorithm certified only as 1.5706- or $4/3$ -competitive does not say that it automatically adapts to these easy cases.

We therefore add a second comparison. For a multiset M , imagine the best randomized online policy that is told M but not its hidden labelling. Its leading cost is an explicit fluid value $n^2\Phi(D_M)$, where D_M is the empirical distribution. We ask whether one universal policy, not told M , can attain the same value. This is an “outer” version of online analysis: on each instance we compare not only with the offline optimum, but also with every other online policy, including policies given more information.

Theorem 1.2 (Randomized instance optimality, informal). *For every fixed common upper bound u , there is an explicit benchmark $\Phi_u(D)$ and one randomized policy, not told the processing-time multiset, whose cost on every multiset $M \subseteq [0, u]$ is*

$$n^2\Phi_u(D_M) + o_u(n^2).$$

No randomized policy—even one told M —can improve the leading term under a uniformly hidden labelling; consequently some fixed oblivious labelling witnesses the same lower bound.

The analogous statement holds in the blind-execution model for every fixed bound $p_i \leq L$, with a different explicit benchmark $\Phi_{\text{blind}}(D_M)$ and error $o_L(n^2)$.

Randomization is essential to the universal part of this statement. A deterministic policy has a fixed pattern of labels from which it tries to learn, and an oblivious labelling can make that pattern unrepresentative. A random pilot instead estimates the histogram of any fixed multiset. Once the histogram is known approximately, the policy solves a one-dimensional quadratic optimization and executes the resulting threshold schedule.

The boundedness qualifications in the two parts should not be confused with the obligatory theorem above. The blind theorem assumes $p_i \leq L$, and the common-upper theorem already has $p_i \leq u$. They use these bounds to make the pilot and grid errors uniformly $o(n^2)$. Randomized obligatory testing, by contrast, allows arbitrary processing times and uses its cutoff/fallback argument instead.

1.3 The information view: blind execution

We now drop the obligation to test. An untouched job may be run blindly for its actual duration p_i ; only after it finishes is that duration known. Thus a test does not make the job faster. It buys the option to postpone a long outcome and to move a short outcome forward.

Here a worst-case ratio to the clairvoyant optimum can be infinite. A zero-heavy multiset illustrates the obstruction. The optimum puts all zero jobs first at no processing cost, whereas an online policy cannot identify those jobs without either testing many labels or occasionally running a positive job before many hidden zeros. As the positive fraction tends to zero, the online leading cost can exceed the clairvoyant leading cost by an arbitrarily large factor. Instance optimality remains meaningful: it asks for the best cost possible under the same hidden-labelling uncertainty, rather than dividing by a benchmark that may be subquadratic.

For a distribution D , write $\mu = \mathbb{E}P$ and choose the testing threshold τ from

$$\mathbb{E}(\tau - P)^+ = 1.$$

Testing one unit of job mass and immediately completing outcomes below τ behaves like one divisible completion module. For any fixed tested fraction q , all remaining work is optimally arranged by increasing cost per completion. The resulting canonical policy is

$$\begin{aligned} \text{threshold tests} &\longrightarrow \text{known values below } \mu \longrightarrow \text{blind jobs} \\ &\longrightarrow \text{known values above } \mu. \end{aligned}$$

Its normalized cost $F_D(q)$ is quadratic in q , and $\Phi_{\text{blind}}(D) = \min_{0 \leq q \leq 1} F_D(q)$. If the test module is no denser than blind execution, the minimizer is $q = 0$ and the policy simply runs every job blindly. Otherwise a genuine interior stopping fraction can be optimal. A three-point example in Section 8 shows that the known middle block cannot in general be omitted.

The universal policy learns this distributional schedule from a sublinear blind pilot. Conversely, a completion-envelope argument rules out any better leading cost, even for an adaptive policy that knows the multiset. This is the blind part of the preceding informal theorem; its precise statement is Theorem 2.5.

1.4 The optimization view: a common upper bound

In the second interpretation, raw execution of every untouched job takes a known common time u . A unit test reveals an optimized time $p_i \leq u$ and allows the scheduler to use it. Small u makes optimization unattractive; large u makes raw execution expensive. Letting u grow recovers obligatory testing, but the finite values are not a monotone interpolation: at intermediate u , the offline optimum can exploit raw execution in ways that are particularly difficult to imitate online.

Theorem 1.3 (Common-upper curves, informal). *For every fixed $u > 0$, the exact size-asymptotic deterministic competitive ratio is a continuous six-piece function $V(u)$. Its*

global maximum is 1.8667603992..., and for every u beyond 4.5052... it is already equal to the obligatory value 1.5705740966....

Against an oblivious adversary, the exact randomized ratio is a continuous four-piece function $R_{\text{rand}}(u)$. Its global maximum is 1.6257523846..., and it is exactly $4/3$ for every $u \geq 25/4$.

The deterministic curve is supported by three algorithmic shapes. For small u , run every job raw. For intermediate u , test jobs and combine a forced immediate prefix with a fixed short/long threshold. For large u , use the history-dependent obligatory threshold, with a temporary reserve that pays for capped outcomes. The matching adversaries move through binary zero/cap instances, capped versions of the harmonic construction, and finally the cap-free harmonic instance. These changes of optimal policy and hard instance create the six branches.

The randomized result is best understood through the stronger benchmark Φ_u . For a fixed empirical distribution, the optimal leading policy tests a fraction q , immediately completes outcomes below its density threshold, processes the tested residual stock in SPT order, and finally runs untouched jobs raw. Dividing its value by the clairvoyant coefficient $\Omega_u(D)$ gives the instance-specific ratio

$$\alpha_u(D) = \frac{\Phi_u(D)}{\Omega_u(D)}.$$

The exact oblivious curve is the outer maximization

$$R_{\text{rand}}(u) = \sup_D \alpha_u(D).$$

This infinite-dimensional optimization collapses to binary distributions. Different zero/cap mixtures give the two middle branches, and the balanced $\{0, 2\}$ distribution gives the final $4/3$ plateau. Thus the curve is not merely a worst-case calculation: it is the envelope of a per-multiset instance-optimal theorem. Exact formulas are given in Theorems 2.2 and 2.3.

1.5 Techniques

Four ideas recur across the models.

Pairwise delay accounting. An operation contributes to the completion time of its own job and once more for every other job still unfinished. The objective is therefore a sum of diagonal terms and one delay term per pair of jobs. The diagonal is only $O(n)$ after normalization, while the $\Theta(n^2)$ pair terms determine the exact asymptotic constants. Deterministic lower bounds make many pairs wait in the wrong order; deterministic upper bounds charge those errors to an offline pair through a bank potential.

Completion density and envelopes. A test together with the outcomes completed immediately can be viewed as a module: it spends some work and completes some job mass. Maximizing mass per unit work yields a threshold. Once these modules, known processing classes, and raw or blind jobs are divisible, their best order is fractional knapsack. The area below the unfinished-mass curve gives both the explicit quadratics Φ_u and Φ_{blind} and a lower bound against arbitrary adaptive stopping rules.

Random relabelling and learning. The algorithm privately permutes labels and uses an $o(n)$ pilot to estimate the empirical distribution. For lower bounds, first-touch causality gives a complementary fact: the adaptive order in which a deterministic policy first touches uniformly labelled occurrences is itself a uniform permutation. The test/raw decision is predictable from the earlier revealed values. A single without-replacement concentration event can therefore control every prefix, after which the realized stopping fraction is inserted pathwise into the fluid envelope.

From instance values to exact curves. For randomized common-upper scheduling we first solve every empirical distribution, obtaining $\Phi_u(D)$. Only then do we maximize $\Phi_u(D)/\Omega_u(D)$. Survival-function flattening reduces the outer optimization to scalar binary families. This separation between the inner algorithmic problem and the outer worst-case distribution is what makes the four-piece curve transparent.

1.6 Related work

Scheduling with tests is part of optimization with explorable uncertainty, where uncertain data can be queried at a cost; the query viewpoint goes back at least to Kahan [Kah91]. Levi, Magnanti, and Shaposhnik study a stochastic scheduling model in which tests diagnose random job attributes and derive prefix-testing policies followed by expected-Smith-ratio order [LMS19, LMS24]. Our four-block blind policy is a unit-weight fluid analogue, but our multiset is adversarial, need not converge, is not announced to the universal policy, and is hidden by an oblivious labelling.

The adversarial common-upper model was introduced by Dürr et al. [DEMM20]. For one machine and total completion time they obtained deterministic bounds, including the $1.8668\dots$ UTE guarantee for binary values under a common upper bound. Our deterministic theorem closes the common-upper gap for arbitrary $p_i \in [0, u]$ and every fixed u . Our randomized theorem gives a separate exact four-piece curve against an oblivious adversary. Related variants allow nonuniform test times [AE21, LLWZ23, KLL⁺26]. Other work considers testing budgets [DKL⁺23], multiple machines [GCH24], or weights [BS26].

Dogeas, Erlebach, and Liang introduced obligatory-test scheduling for total completion time [DEL24]. For uniform tests they gave the SIDLE upper bound $1.58451\dots$ and the deterministic lower bound $\sqrt{2}$. Liang and Liang subsequently raised the single-machine lower bound to $3/2$ through a completion-threshold framework [LL26]. Our harmonic construction and adaptive threshold meet at $1.5705740966\dots$, while our randomized result resolves the oblivious single-machine value at $4/3$.

The processing-time-oracle model of Dufossé et al. [DDN⁺22] is close to blind execution: an untested job takes its true duration, but only two possible durations are allowed. We permit an arbitrary bounded multiset and obtain a distribution-sensitive benchmark and four-block policy. If no tests are available and processing times are revealed only at completion, one obtains the classical nonclairvoyant model of Motwani, Phillips, and Torng [MPT94]. These semantics differ from common-upper raw execution, which always lasts the declared upper bound, regardless of the hidden optimized value.

The rest of the paper first fixes the models and offline benchmarks in Section 3. It then proves the deterministic obligatory endpoint and finite common-upper curve, the randomized common-upper and obligatory results, and the blind instance-optimal theorem. The machine-checked scope is summarized in Section 9.

2 OLD intro

2.1 The exact answers

The obligatory endpoint had a gap between the best known deterministic lower and upper bounds [DEL24, LL26]. The finite common-upper problem likewise had no exact answer for arbitrary revealed values [DEMM20]. We close both gaps in the natural size-asymptotic convention.

Let $z_\star > 1$ be the unique solution of

$$z_\star - 3 = \log z_\star, \quad (1)$$

where all logarithms are natural, and put

$$R_\star = 1 + \frac{2}{z_\star - 1} = 1.570574096649395\dots, \quad z_\star = 4.505241495792883\dots \quad (2)$$

Theorem 2.1 (Obligatory endpoint). *The following matching statements hold for $u = \infty$.*

- (i) *For every deterministic online algorithm $\mathcal{B}$ and every $\varepsilon > 0$, there are arbitrarily large instances I for which*

$$\text{ALG}_{\mathcal{B}}(I) \geq (R_\star - \varepsilon) \text{OPT}(I).$$

- (ii) *There is one deterministic algorithm $\mathcal{A}$ and one absolute constant C such that every n -job instance satisfies*

$$\text{ALG}_{\mathcal{A}}(I) \leq R_\star \text{OPT}(I) + Cn.$$

Consequently the optimal deterministic size-asymptotic value at $u = \infty$ is exactly $R_\star$.

For the finite curve, let $\varphi = (1 + \sqrt{5})/2$ and define

$$u_\diamond = \frac{1 + \sqrt{3 + 2\sqrt{5}}}{2}, \quad u_0 = 1 + s_0, \quad s_0^3 = (s_0 + 1)^2,$$

where $s_0 > 1$. For $s = u - 1$, let $\rho_I(u)$ be the positive root of

$$s\rho^2 + (s^3 + s^2 + 1)\rho - (s^2 + s + 1)(s + 2) = 0.$$

On $\varphi + 2 \leq u \leq z_\star$, define $C_{\text{mix}}(u) = 1 + c$ by the unique pair (c, m) satisfying

$$\text{artanh } m - m = 1 - \frac{1}{c} + \frac{1}{2} \log \left(1 + \frac{2}{c} \right), \quad u = 1 + \frac{2}{c} - m, \quad (3)$$

with $2/(z_\star - 1) \leq c \leq 1/\varphi$ and $0 \leq m \leq 1/\varphi$.

Theorem 2.2 (Finite common-upper curve). *For every fixed $u > 0$, the optimal deterministic size-asymptotic competitive ratio is*

$$V(u) = \begin{cases} 1, & 0 < u \leq 1, \\ u, & 1 \leq u \leq u_\diamond, \\ \rho_I(u), & u_\diamond \leq u \leq u_0, \\ 1 + 1/\sqrt{u-1}, & u_0 \leq u \leq \varphi + 2, \\ C_{\text{mix}}(u), & \varphi + 2 \leq u \leq z_\star, \\ R_\star, & u \geq z_\star. \end{cases} \quad (4)$$

Every lower branch holds against arbitrary deterministic algorithms on arbitrarily large fixed instances. Every upper branch is attained up to $O_u(n)$. On the finite part of the last branch, one algorithm and one absolute $O(n)$ constant work simultaneously for all $u \geq z_*$. Together with Theorem 2.1, the same witness covers the endpoint $u = \infty$.

The deterministic theorem is only the first part of the landscape. Against an oblivious adversary, private randomization gives the following exact four-piece curve. Let $u_r > 1$ be the unique root of

$$u_r^3 - 6u_r^2 + 5u_r - 1 = 0, \quad u_r = 5.048917339522306 \dots \quad (5)$$

Theorem 2.3 (Randomized finite common-upper curve). *For every fixed $u > 0$, the optimal randomized size-asymptotic competitive ratio in the common-upper model against an oblivious adversary is*

$$R_{\text{rand}}(u) = \begin{cases} 1, & 0 < u \leq 1, \\ \frac{u^3}{u^2 + (u-1)^3}, & 1 \leq u \leq u_r, \\ 1 + \frac{1}{2(\sqrt{u}-1)}, & u_r \leq u \leq 25/4, \\ \frac{4}{3}, & u \geq 25/4. \end{cases} \quad (6)$$

More precisely, there are randomized nonanticipating algorithms $\mathcal{A}_{n,u}$ and numbers $\varepsilon_u(n) \rightarrow 0$ such that every fixed labelled vector $p \in [0, u]^n$ satisfies

$$\mathbb{E} \text{ALG}_{\mathcal{A}_{n,u}}(p) \leq R_{\text{rand}}(u) \text{OPT}_u(p) + \varepsilon_u(n)n^2. \quad (7)$$

Conversely, for every randomized online algorithm and every $\varepsilon > 0$, there are arbitrarily large n and a fixed vector $p \in [0, u]^n$ for which

$$\mathbb{E} \text{ALG}(p) \geq (R_{\text{rand}}(u) - \varepsilon) \text{OPT}_u(p). \quad (8)$$

The curve is continuous at all three junctions. Its global maximum is attained at $u = (3 + \sqrt{3})/2$ and equals

$$\max_{u>0} R_{\text{rand}}(u) = \frac{27 + 6\sqrt{3}}{23} = 1.625752384583185 \dots \quad (9)$$

Thus randomization improves every nontrivial part of the deterministic finite-cap curve. The last branch already has value $4/3$, but passing to $u = \infty$ still requires a proof uniform over unbounded processing times. That endpoint remains sharp even when the processing-time multiset is not announced.

Theorem 2.4 (Randomized obligatory testing). *There is a universal randomized nonanticipating algorithm, depending only on n , such that every fixed vector $p \in [0, \infty)^n$ satisfies*

$$\mathbb{E} \text{ALG}(p) \leq \frac{4}{3} \text{OPT}(p) + 20378n^{7/4} \leq \left(\frac{4}{3} + 40756n^{-1/4} \right) \text{OPT}(p). \quad (10)$$

Conversely, for every randomized algorithm and every sufficiently large even n , some fixed labelling of $n/2$ zero jobs and $n/2$ jobs of processing time two satisfies

$$\mathbb{E} \text{ALG}(p) \geq n^2 - O(n^{3/2}), \quad \text{OPT}(p) = \frac{3}{4}n^2 + n. \quad (11)$$

Hence the exact randomized size-asymptotic competitive ratio for obligatory testing against an oblivious adversary is $4/3$.

Our third result uses a stronger, instance-dependent comparison. In the blind-execution model, sparse zero-heavy instances may have clairvoyant cost $o(n^2)$, so an additive $o(n^2)$ theorem is not a conventional competitive ratio. It instead says that the algorithm learns, at negligible leading cost, everything that advance knowledge of the multiset could contribute to the oblivious minimax value.

Theorem 2.5 (Blind-execution instance optimality). *Fix $L < \infty$. There are a randomized nonanticipating policy $\mathcal{A}^*$, which knows n, L but not the processing-time multiset, and a deterministic sequence $\varepsilon_L(n) \rightarrow 0$ with the following properties. For every multiset $M \subseteq [0, L]$ of size n and every fixed labelling σ ,*

$$\mathbb{E}C_{\mathcal{A}^*}(\sigma(M)) \leq n^2\Phi(D_M) + \varepsilon_L(n)n^2. \quad (12)$$

For every randomized policy $\mathcal{A}$, even one told M , some fixed oblivious labelling σ satisfies

$$\mathbb{E}C_{\mathcal{A}}(\sigma(M)) \geq n^2\Phi(D_M) - \varepsilon_L(n)n^2. \quad (13)$$

The benchmark Φ , defined in Section 8.1, is the minimum of an explicit one-variable quadratic. One may take $\varepsilon_L(n) = O_L(n^{-1/6}\sqrt{\log(n+2)})$ analytically; the formal proof also provides a slower explicit vanishing rate using only finite second-moment estimates.

model	adversary / knowledge	benchmark	result
common upper u	deterministic	clairvoyant OPT	exact six-piece curve $V(u)$
common upper u	randomized oblivious; unknown M	clairvoyant OPT	exact four-piece curve $R_{\text{rand}}(u)$
obligatory	deterministic	clairvoyant OPT	exact R_*
obligatory	randomized oblivious; unknown M	clairvoyant OPT	exact asymptotic ratio $4/3$
blind, $p \leq L$	randomized oblivious; unknown M	announced minimax value	$\Phi(D_M) \pm o_L(1)$ after normalization by n^2

Table 1: The three models and their principal guarantees. The last row is instance optimality at the quadratic scale, not a competitive-ratio claim on instances with subquadratic optimum.

The finite curve is continuous, rises from one to its global maximum

$$V(u_\diamond) = u_\diamond = 1.866760399173862\dots,$$

and then decreases to the obligatory value. In fact it does more than converge:

$$V(u) = V(\infty) = R_* \quad \text{for every } u \geq z_*. \quad (14)$$

This is equality of size-asymptotic game values, not equality of the finite problems. At any finite u , the optimum may still save a test on a job with $p \geq u - 1$.

2.2 Why total completion time leads to pair accounting

The most useful way to think about total completion time is pairwise. Every operation of job i contributes once to C_i , and it contributes once more for each other job that remains unfinished while that operation is executed. Equivalently, the objective consists of diagonal job terms and one delay term for every unordered pair of jobs. The offline pair term is




Figure 1: The exact deterministic size-asymptotic ratio for finite common upper limit u . The curve stabilizes at the obligatory endpoint value; dotted lines mark its five regime changes.

simply the shorter effective length. The online pair term records whether an earlier revealed job was processed immediately or left waiting.

This representation explains both sides of our proof. A lower-bound adversary tries to make many pairs wait in the wrong order. An upper-bound algorithm tries to ensure that every expensive local decision can be charged to a corresponding offline pair. Since there are $\Theta(n^2)$ pairs but only n diagonal terms, controlling the quadratic pair coefficient gives the exact size-asymptotic ratio; bounded rounding errors of $O(1)$ per job become only $O(n)$.

2.3 Lower-bound intuition: hide the short jobs until the end

At the obligatory endpoint, the adversary reveals equal blocks of positive jobs from longer to shorter values and reveals a large zero block last. The algorithm is forced to choose between two risks. Processing a revealed job early may delay many still-hidden zeros. Deferring it may leave a large tail that delays everything completed later. The adaptive revelation is frozen after the interaction, so a standard replay argument produces one fixed instance for the chosen deterministic algorithm.

The processing times are separated by harmonic gaps. This choice is not a guess about the final constant. The zero mass and the number of shorter positive blocks give two multiplicities; their sum exactly cancels the denominator of each harmonic gap. After this cancellation, every possible postponement made by the algorithm has a nonnegative coefficient and every own-phase choice contributes the same completed square. All algorithmic decisions disappear from the leading term. What remains is a one-variable ratio describing the relative positive and zero masses, and its stationary equation is (1).

For finite u , two additional adversarial pictures appear. When u is small or moderate, binary values $p \in \{0, u\}$ already suffice: the adversary watches how much work the algorithm completes before revealing whether the next jobs are zeros or caps. For the transition to the obligatory value, the hard instance is a capped prefix followed by a scaled

copy of the harmonic core. The prefix exploits the optimum’s ability to skip tests; the core exploits uncertainty in the test order. As u grows, the optimal cap mass decreases continuously to zero, leaving exactly the obligatory construction.

2.4 Upper-bound intuition: three policies and one common bank

The exact upper curve uses three simple algorithmic shapes. Their formal pseudocode appears later; here we describe why they are natural.

approximate range	upper-bound policy at a glance
$u \lesssim 1.87$	Do not test: execute every job raw.
$1.87 \lesssim u \lesssim 3.15$	Test all jobs; always finish an initial fraction, then use a short/long threshold. The initial fraction shrinks as u grows.
$3.15 \lesssim u \lesssim 3.62$	Test all jobs; finish values at most one immediately and sort the rest in a final tail.
$3.62 \lesssim u \lesssim 4.51$	Use the history-dependent threshold; the proof adds a reserve for capped outcomes.
$u \gtrsim 4.51$, including ∞	Use the same adaptive threshold, now with no cap reserve.

Raw policy. When u is very small, information costs at least as much as it can save. The algorithm therefore runs every job raw. Slightly beyond that trivial range, raw execution need no longer be clairvoyantly optimal for each job, but comparing its length u with the optimum’s effective lengths already gives the first nontrivial minimax branch.

Forced-prefix threshold policy. In the next regimes the algorithm tests every job. During an initial fraction of the test order it processes every revealed value immediately. After that prefix it processes only sufficiently short outcomes and sends the rest to a shortest-processing-time tail. The forced prefix guarantees that some jobs finish before too many tests accumulate; the suffix threshold protects the schedule from long revealed jobs. As u increases, the useful prefix shrinks continuously to zero. The algebraic branch is exactly the point where the worst binary mixture is tangent to the target ratio; after the prefix vanishes, balancing a capped tail against a threshold-one tail gives the square-root branch.

Adaptive-threshold policy. For large u , a fixed prefix is too rigid. The algorithm again tests jobs in a fixed order, but its immediate-processing threshold reacts to the observed history. It remembers only the number N of positive outcomes and the number D of those outcomes that were deferred. A normalized balance of the form

$$\frac{D - (1 + c)N}{\text{number of untested jobs}}$$

measures how much future delay has already been promised to the tail. Near a neutral balance the algorithm can afford to process moderately long jobs immediately. Every positive outcome drives the balance downward, while a deferral refunds one unit because future tests will also delay that tail job. When the normalized balance becomes sufficiently negative, the threshold falls toward one. All deferred jobs are sorted only after the tests finish.

The analysis first fixes the symbolic immediate/deferred path. Convexity then reduces every ordinary revealed value to one of four endpoints: a true zero, a positive value tending

to zero, the immediate side of the live threshold, or its deferred side. A nonnegative bank potential pays the excess pair cost at each endpoint. The logarithmic threshold is forced by making the immediate and deferred boundary equations balance simultaneously; the obligatory constant is forced by calibrating the bank's initial value.

Finite u creates exactly one new endpoint: a capped value $p = u$, for which the optimum can use raw execution and avoid the test. On the middle bridge this endpoint can temporarily overdraw the ordinary bank, so we add a reserve whose balance depends only on the observed cap fraction. At the end of the bridge the reserve becomes zero. With $r = R_\star - 1$, the worst residual charge of the cap endpoint is at most $2 - r(u - 1)$. The right-hand side is nonpositive precisely when $u \geq 1 + 2/r = z_\star$. Thus beyond $z_\star$ the obligatory four-endpoint bank already pays for the fifth endpoint. The plateau in Equation (14) is a proof-level fact, not merely a numerical coincidence.

2.5 Randomized intuition: completion density and sampling

The randomized common-upper and obligatory results have a different core. Suppose for a moment that the multiset is announced. Choose a processing-time prefix E , test jobs in a uniformly random order, process a revealed job immediately exactly when it lies in E , and sort the remaining tail after the last test. The expected cost of this stationary policy has a closed pairwise formula. The right prefix maximizes

$$\frac{\text{mass completed during discovery}}{\text{test work plus immediate processing work}}.$$

Writing the reciprocal of this density as θ , the optimizer is a threshold: early values are at most θ and late values are at least θ .

For a finite cap, the algorithm compares this stationary policy with running every job raw. Writing $H(t) = \Pr(P > t)$ and $s = u - 1$, the offline quadratic coefficient is

$$\frac{1}{2} \left(1 + \int_0^s H(t)^2 dt \right).$$

The density threshold satisfies $\int_0^\theta (1 - H(t)) dt = 1$. Splitting according to whether $\theta \geq s$ or $\theta \leq s$ reduces the worst distribution to one of two binary endpoints. The distributions on $\{0, u\}$ give the two middle branches of (6); the balanced $\{0, 2\}$ distribution gives the 4/3 plateau. The first transition occurs when the stationary critical point meets the raw/stationary crossing, which is exactly the cubic in (5); the second occurs at $u = 25/4$.

At the obligatory endpoint, a short algebraic identity writes $4 \text{OPT} - 3\mathbb{E} \text{ALG}$ as a sum of three nonnegative terms. Equality again identifies the hard distribution: half of the mass at zero and half at two.

For a finite cap, bounded support makes the announcement easy to remove: a pilot of size $n^{2/3}$ selects between raw execution and the stationary prefix on a grid of $n^{1/3}$ cells, losing only $O_u(n^{11/6})$. At the unbounded obligatory endpoint, the algorithm instead tests $n^{3/4}$ uniformly random jobs and learns the threshold on a grid of $n^{1/4}$ cells. Closing the chosen sample prefix under its numerical threshold is essential: otherwise a cell missed by the sample could violate the population separation. Values above a fixed cutoff are always deferred. If the learned completion density is too small, the algorithm falls back to test-all-then-SPT; precisely in that case the offline objective is already large enough to pay for the fallback. This gives the upper bound in Equation (10). For the lower bound, a uniformly labelled half-zero, half-two multiset is a balanced random bridge. At every time, the number of completed jobs is at most one half of the work spent, up to the bridge discrepancy. Integrating this envelope gives Equation (11).

2.6 Blind execution: from tests to a fractional knapsack

For a processing-time distribution D , let $\mu = \mathbb{E}P$ and define τ by

$$\mathbb{E}(\tau - P)^+ = 1.$$

A unit mass of tests followed by immediate processing of outcomes below τ completes mass $a = \Pr(P < \tau)$ using work $1 + \mathbb{E}[P \mathbf{1}_{P < \tau}] = a\tau$. It is therefore a divisible completion module of work τ per completed job. Untested blind jobs form a second module of cost μ per completion; tested residual values form ordinary known SPT classes.

For any fixed final tested fraction q , a completion-envelope relaxation turns these objects into a fractional knapsack. Its greedy order is

$$\begin{aligned} \text{test module} &\longrightarrow \text{known residual values below } \mu \longrightarrow \text{blind block} \\ &\longrightarrow \text{known residual values above } \mu. \end{aligned}$$

The area below the corresponding unfinished-mass curve is an explicit quadratic $F_D(q)$; minimizing it defines $\Phi(D)$. This pointwise envelope, rather than a local exchange in an adaptive decision tree, is what makes the lower bound robust.

The remaining issue is to compare a completely adaptive finite policy with that deterministic fluid picture. Distinguish equal-valued occurrences and map each hidden labelling to the order in which the policy first touches them. Prefix causality makes this reveal map a bijection of the finite permutation space. Hence a uniformly labelled input becomes an ordinary without-replacement urn and the test/blind decision at the next touch is predictable from the preceding revealed values. One simultaneous urn event then controls every prefix; only after fixing that event do we substitute the policy's realized stopping fraction q . Rounding to a growing grid and a sublinear blind pilot complete the transfer to an arbitrary bounded, unannounced multiset.

2.7 Why the infinite endpoint needs a uniform proof

For each fixed processing-time vector, the finite- u offline lengths $q_u(p)$ eventually equal $1 + p$. Nevertheless, a competitive-ratio proof cannot blindly send u to infinity: the worst instance may choose $p \approx u$, the algorithm may depend on u , and the hidden constant in an $O_u(n)$ estimate may diverge. We avoid every such interchange. The lower bound uses one bounded harmonic core, which embeds in all sufficiently large finite caps. The high- u upper bound uses one u -independent adaptive algorithm and one absolute linear remainder. Consequently the endpoint $u = \infty$ and the finite plateau are certified by the same witnesses.

The linear remainder is lower order because $\text{OPT} \geq n(n+1)/2$ at $u = \infty$ and, for fixed u , the analogous finite bound is also quadratic. It implies an endpoint coefficient $R_\star + \varepsilon$ with a fixed additive constant for every $\varepsilon > 0$. We do *not* prove the stronger statement $\text{ALG} \leq R_\star \text{OPT} + B$ with one fixed B , nor the unrelaxed inequality $\text{ALG} \leq R_\star \text{OPT}$ on every finite instance.

2.8 How to read the proofs

After the literature review and common framework, Section 4 begins the deterministic part with the cap-free endpoint. Its lower subsection gives the harmonic adversary and a reusable scaled-core lemma. Its upper subsection formally states the adaptive algorithm and proves the ordinary four-endpoint bank. These are matching and may still be read independently.

Section 5 then treats finite caps. The binary stopping game and the two simple policies prove the first four branches. The capped-prefix lower construction reuses the harmonic core, while the middle/high upper bound reuses the ordinary bank and analyzes only the new cap endpoint and its reserve. At $u = z_*$ both cap corrections vanish. Appendices A and B contain an exact rational checkpoint for the cap-free lower construction and an independent cap-bubbling certificate.

Section 6 proves the exact randomized finite-cap curve. It first identifies the announced stationary benchmark, reduces the distributional maximization to a scalar binary game, then removes announcement by a sublinear random pilot and proves the matching oblivious Yao bounds. Section 7 proves the sharp randomized $4/3$ endpoint theorem uniformly for unbounded processing times. It first solves the announced stationary problem, then learns the threshold from a sample, and finally gives the balanced binary Yao lower bound. Section 8 turns to blind execution. It defines the fluid benchmark, proves the fixed- q envelope theorem, transfers the envelope to arbitrary adaptive announced policies, and removes announcement by a blind pilot. Section 9 records the machine-checked scope; detailed correspondence for the deterministic theorem remains in Appendix C.

3 Models, benchmarks, and common preliminaries

There are n jobs, all available at time zero, and one machine. Operations are nonpreemptive, and at most one operation can run at a time. A unit test of job j reveals its processing time $p_j \geq 0$.

We use the following three operation models throughout.

Common-upper execution CU(u).

For finite u , all jobs share the known upper limit $p_j \in [0, u]$. A job is completed either by raw execution for exactly u time, or by a unit test and subsequent processing for p_j . Raw execution reveals no hidden value.

Obligatory testing OT.

Processing times are arbitrary finite nonnegative numbers. Every job must be tested for one unit and subsequently processed for p_j ; the processing may be delayed after the test. This is also the formal endpoint $\text{CU}(\infty)$.

Blind execution BE(L).

Here $p_j \in [0, L]$. A job may be run blindly for its actual duration p_j , at whose completion its duration becomes known, or tested for one unit and subsequently processed for p_j . Thus blind execution never pays an upper bound in place of the actual processing time.

The completion time C_j is the completion time of the processing operation; in tested mode, when $p_j = 0$, this is the instant immediately after the test. In the blind model, a blind zero job completes immediately when selected. The objective is $\sum_j C_j$. The number n and the parameter u are known to a common-upper algorithm; a blind-execution algorithm knows n, L .

We write $\text{ALG}_{\mathcal{A}}(I)$ for the objective value of an online algorithm $\mathcal{A}$ on an instance I , and $\text{OPT}(I)$ for the value of an offline optimum that knows all processing times. We omit subscripts and arguments when they are clear.

3.1 Information and adversaries

The deterministic common-upper results allow the usual adversarial input and use adaptive revelation only as a lower-bound device that is frozen by Lemma 3.3. The randomized results use an *oblivious* adversary: the labelled processing-time vector is fixed before the algorithm draws its private random seed. Unless explicitly stated otherwise, the algorithm is not told the multiset of values. An *announced* policy is additionally given that multiset but not its assignment to labels. Announcing the multiset can only help and is therefore a valid strengthening in lower bounds.

All randomized expectations are over private algorithmic randomness. A uniform random relabelling is private randomness, not a distributional assumption on the adversarial multiset. For lower bounds we use a uniform distribution over fixed labellings as a finite Yao distribution and then select one labelling whose cost is at least the average.

3.2 Offline optima and pair accounting

Define the clairvoyant effective length by

$$q_u(p) := \begin{cases} \min\{u, 1 + p\}, & u < \infty, \\ 1 + p, & u = \infty. \end{cases} \quad (15)$$

Lemma 3.1 (Unified offline order). *Let $q_i = q_u(p_i)$ and let $q_{(1)} \leq \dots \leq q_{(n)}$. For every $u \in (0, \infty]$, an offline optimum completes the jobs in nondecreasing order of q_i and has value*

$$\text{OPT}_u = \sum_{k=1}^n \sum_{i=1}^k q_{(i)}, \quad (16)$$

$$= \sum_i q_i + \sum_{i < j} \min\{q_i, q_j\}. \quad (17)$$

For finite u , it uses raw mode when $u \leq 1 + p_j$ and tested mode otherwise. At $u = \infty$, it tests every job and processes the jobs in nondecreasing p_j order.

Proof. Let $C_{[1]} \leq \dots \leq C_{[n]}$ be the completion times of any feasible schedule. Completing job j consumes at least q_j units: for finite u this is the shorter of its two modes, and for $u = \infty$ it is its compulsory test-process length. By time $C_{[k]}$ the machine has completed some k jobs, so

$$C_{[k]} \geq \sum_{i=1}^k q_{(i)}.$$

The schedule that takes the shorter available mode for every job and executes the resulting blocks in nondecreasing q_i order attains every inequality. Summing and expanding the prefix sums gives both formulas. $\square$

In the blind-execution model, a clairvoyant optimum never tests: it knows all values and can execute each job blindly for p_j . The same shortest-first argument gives

$$\text{OPT}_{\text{BE}}(p) = \sum_i p_i + \sum_{i < j} \min\{p_i, p_j\}. \quad (18)$$

We will not use this quantity as the principal benchmark of Theorem 2.5: it may be $o(n^2)$, while that theorem compares policies at the n^2 scale to the stronger announced-multiset online benchmark $\Phi(D_M)$.

At the cap-free endpoint, the pair formula specializes to

$$\text{OPT}_\infty = \sum_{k=1}^n \left(k + \sum_{i=1}^k p_{(i)} \right), \quad (19)$$

$$= \sum_i (1 + p_i) + \sum_{i < j} (1 + \min\{p_i, p_j\}). \quad (20)$$

In particular,

$$\text{OPT}_\infty \geq \frac{n(n+1)}{2}. \quad (21)$$

For finite u , $q_j \geq \min\{u, 1\}$ gives

$$\text{OPT}_u \geq \min\{u, 1\} \frac{n(n+1)}{2}. \quad (22)$$

3.3 Competitive-ratio conventions

The ordinary competitive ratio of $\mathcal{A}$ is

$$\text{CR}(\mathcal{A}) = \sup_I \frac{\text{ALG}_{\mathcal{A}}(I)}{\text{OPT}(I)}.$$

For the result of this paper, the relevant size-asymptotic ratio is

$$\text{CR}_\infty(\mathcal{A}) = \limsup_{n \rightarrow \infty} \sup_{|I|=n} \frac{\text{ALG}_{\mathcal{A}}(I)}{\text{OPT}(I)}. \quad (23)$$

The additive-constant convention calls a coefficient c admissible for $\mathcal{A}$ if there exists $B < \infty$ such that

$$\text{ALG}_{\mathcal{A}}(I) \leq c \text{OPT}(I) + B \quad \text{for every } I.$$

We write

$$\text{CR}_{\text{add}}(\mathcal{A}) = \inf\{c : c \text{ is admissible for } \mathcal{A}\}. \quad (24)$$

The optimal deterministic values in the two conventions are obtained by taking the infimum of $\text{CR}_\infty(\mathcal{A})$ and $\text{CR}_{\text{add}}(\mathcal{A})$, respectively, over deterministic algorithms. For any fixed u , an estimate $\text{ALG} \leq c \text{OPT} + O_u(n)$ fixes Equation (23) at most c and, by Equations (21) and (22), makes every coefficient $c + \varepsilon$ admissible. It need not make the endpoint coefficient c admissible.

Remark 3.2 (No hidden interchange of limits). The notation $u = \infty$ identifies one model exactly. We never infer an endpoint upper bound merely by taking $u \rightarrow \infty$ in an estimate whose algorithm or additive term may depend on u . The endpoint estimate is proved directly in Section 4.2; Section 5.6 then verifies that the same algorithm and certificate work uniformly for every finite $u \geq z_\star$.

3.4 Test-order adversaries

The lower bound will assign processing times according to the order in which tests finish. This is only a proof device.

Lemma 3.3 (Replay). *Fix a deterministic online algorithm. Suppose all untested jobs have the same known parameters, and an adversary assigns processing times deterministically from the transcript and test-completion order. The resulting interaction is reproduced by a fixed input instance.*

Proof. Run the interaction once and record the value revealed for every tested job label. Assign those values to the corresponding labels in advance. An induction over operation completions shows that the deterministic algorithm chooses the same tests, processing operations, and idle intervals on the fixed instance. If the algorithm fails to test or complete some job forever, its competitive ratio is already infinite; otherwise every relevant label is recorded. $\square$

4 The cap-free endpoint $u = \infty$

This section contains the two estimates needed for the cap-free endpoint theorem and isolates the harmonic and four-endpoint cores reused later. The first subsection applies to an arbitrary deterministic online algorithm; the second analyzes one explicit algorithm. Each restates the local notation it uses.

4.1 Harmonic core and matching lower bound

This subsection proves the lower bound in one self-contained calculation. Its notation is local to this subsection. The construction has many equally large groups of positive jobs, revealed from longest to shortest, followed by one large group of zero jobs. The only design choice is the gap between two consecutive positive processing times. Choosing these gaps harmonically will make all decisions of the online algorithm disappear from the leading term.

Theorem 4.1 (Lower bound). *For every deterministic online algorithm $\mathcal{B}$ and every $\varepsilon > 0$, there are arbitrarily large fixed instances I with unit tests and rational processing times such that*

$$\text{ALG}_{\mathcal{B}}(I) \geq (R_{\star} - \varepsilon) \text{OPT}(I).$$

The revelation experiment. Fix an integer K , a rational number $\xi > 0$, and a rational slack $\gamma \in (0, 1)$; one may keep $\gamma = 1/2$ throughout. For a large scaling integer H , the instance will contain H jobs of each of K positive types $0, 1, \dots, K-1$ and ξH zero jobs. We choose H to clear all denominators.

The adversary assigns processing times according to the order in which tests are started. The first H tested jobs receive processing time p_0 , the next H receive p_1 , and so on; after the K positive blocks, every remaining job receives processing time zero. Thus

$$p_0 > p_1 > \dots > p_{K-1} > 1,$$

so the algorithm sees the positive types from longest to shortest and learns about the zero jobs only at the end.

This adaptive description is only a convenient way to construct the input. After running the interaction against a fixed deterministic algorithm, assign to every job label the value revealed when that label was tested. Replaying the algorithm on this fixed assignment produces exactly the same transcript. If the algorithm never tests or completes some job, its cost is infinite and the lower bound is immediate; otherwise every job label is recorded. Hence it is enough to lower-bound the cost in the revelation experiment.

Why each phase can be relaxed to shortest processing time. Let T_{ℓ} be the start of the first test in positive block ℓ , and let T_K be the start of the first test in the zero block. Phase ℓ contains the completions in $(T_{\ell}, T_{\ell+1}]$; a completion exactly at T_{ℓ} belongs

to the preceding phase. Each T_ℓ is the boundary between two nonpreemptive operations. Consequently, a previously tested job that is unfinished at T_ℓ has not started processing and still needs its entire processing time.

We repeatedly use the following elementary observation. Suppose jobs with remaining work $v_1, \dots, v_m$ complete after some time T . If their completion order is $q_1, \dots, q_m$, then the r th relative completion time is at least $\sum_{h \leq r} v_{q_h}$. Summing these prefix inequalities shows that their total relative completion time is minimized by arranging the v_i in nondecreasing order. This is merely a lower bound; the online algorithm need not actually use that order.

For normalized group masses $q_1, \dots, q_m$ and remaining lengths $v_1 \leq \dots \leq v_m$, this shortest-first bound has leading coefficient

$$\mathcal{S}((q_g, v_g)_{g=1}^m) = \sum_{g=1}^m \left(\frac{v_g q_g^2}{2} + q_g \sum_{h < g} v_h q_h \right). \quad (25)$$

More precisely, the corresponding contribution for groups of size $q_g H$ is $H^2 \mathcal{S} + O(H)$. The quadratic term is simply the familiar sum $v_g(1 + \dots + q_g H)$ inside one group; the cross term is the waiting caused by all shorter groups.

We shall ensure that

$$p_0 - p_{K-1} < 1. \quad (26)$$

In phase ℓ , an unfinished job of an older type $j < \ell$ therefore has remaining length p_j , whereas a type- ℓ job completed in its own phase needs both its test and processing, of total length $1 + p_\ell$. Their order in the shortest-first relaxation is

$$p_{\ell-1} < p_{\ell-2} < \dots < p_0 < 1 + p_\ell.$$

At T_K , an untested zero job has remaining length 1, so the relaxed tail starts with all zero jobs and then uses the positive types from shortest to longest. These are the only two places where Equation (26) is used.

The finite accounting identity. We now record the online algorithm's possible behavior. Normalize all job counts by H . Let

- $s_j \in [0, 1]$ be the mass of type- j jobs completed in their own phase;
- $d_{j,\ell} \geq 0$ be the mass of type- j jobs completed in the later phase $\ell > j$; and
-

$$X_{j,\ell} = s_j + \sum_{r=j+1}^{\ell-1} d_{j,r} \leq 1 \quad (\ell > j)$$

be the mass of type j completed before phase ℓ . Thus $X_{j,\ell+1}$ is the mass completed by the end of phase ℓ .

At time T_ℓ , the machine has already performed the tests of the ℓH jobs in earlier positive blocks and has processed every earlier job that has completed. Thus

$$\frac{T_\ell}{H} \geq \ell + \sum_{j < \ell} p_j X_{j,\ell}. \quad (27)$$

For each phase, charge its completions their common start time from Equation (27), and charge their relative completion times using Equation (25). Do the same at T_K for the

remaining positive jobs and the ξH zero jobs. Expanding and collecting equal variables gives

$$\begin{aligned} \frac{\text{ALG}}{H^2} &\geq C_K + \sum_{j=0}^{K-1} \left(\frac{s_j^2}{2} - \theta_j s_j \right) \\ &\quad + \sum_{j < \ell} d_{j,\ell} \left[\ell - j - \theta_j - \sum_{j < k < \ell} (p_j - p_k) X_{k,\ell+1} \right] - O(1/H), \end{aligned} \quad (28)$$

where K, ξ, γ are fixed while $H \rightarrow \infty$, and

$$\theta_j = 1 - \xi(p_j - 1) - \sum_{k > j} (p_j - p_k - 1), \quad (29)$$

$$C_K = \frac{\xi^2}{2} + 2\xi K + K^2 + P_K, \quad P_K = \sum_{j=0}^{K-1} \left(j + \frac{1}{2} \right) p_j. \quad (30)$$

For completeness, here is how to audit Equation (28) without guessing its origin. In phase ℓ , the masses in the shortest-first relaxation are

$$d_{\ell-1,\ell}, d_{\ell-2,\ell}, \dots, d_{0,\ell}, s_\ell$$

with respective lengths

$$p_{\ell-1}, p_{\ell-2}, \dots, p_0, 1 + p_\ell.$$

Multiply their total mass by the right-hand side of Equation (27) and add the functional Equation (25). In the final phase use masses

$$\xi, 1 - X_{K-1,K}, \dots, 1 - X_{0,K}$$

and lengths $1, p_{K-1}, \dots, p_0$. The constant terms sum to C_K . The terms containing only s_j are $s_j^2/2 - \theta_j s_j$; the coefficient of each $d_{j,\ell}$ is precisely the bracket in Equation (28). Thus the displayed identity is just the phasewise shortest-first bound with the phase start times retained. No optimality assumption about the online schedule is hidden in it.

The meaning of the expression is useful. Completing an s_j fraction of a type in its own phase creates a quadratic completion cost but saves later waiting cost, producing $s_j^2/2 - \theta_j s_j$. A completion postponed to a later positive phase is represented by $d_{j,\ell}$ and its bracket. We next choose the processing times so that every such bracket is nonnegative and all θ_j are equal.

The harmonic choice and its telescoping cancellation. Set

$$p_{K-1} = 1 + \frac{\gamma}{\xi}, \quad p_i = p_{i+1} + \frac{1}{\xi + K - 1 - i} \quad (0 \leq i < K - 1). \quad (31)$$

In other words, measured upward from the shortest type, the consecutive gaps are

$$\frac{1}{\xi + 1}, \frac{1}{\xi + 2}, \dots, \frac{1}{\xi + K - 1}.$$

This is the key design choice. If $m = K - 1 - j$, then

$$p_j - 1 = \frac{\gamma}{\xi} + \sum_{r=1}^m \frac{1}{\xi + r},$$

$$\sum_{k>j} (p_j - p_k) = \sum_{r=1}^m \frac{r}{\xi + r}.$$

Adding the first line multiplied by ξ to the second line makes each denominator cancel:

$$\xi(p_j - 1) + \sum_{k>j} (p_j - p_k) = \gamma + \sum_{r=1}^m \frac{\xi + r}{\xi + r} = \gamma + m. \quad (32)$$

Substitution in Equation (29) gives the same value for every type,

$$\theta_j = 1 - \gamma. \quad (33)$$

This is why the increments in Equation (31) are harmonic: the factor ξ coming from the zero block and the multiplicity r coming from the r shorter positive types add up to the denominator $\xi + r$.

The delayed-completion terms are now harmless as well. Since $X_{k,\ell+1} \leq 1$, their brackets are at least

$$\begin{aligned} \ell - j - (1 - \gamma) - \sum_{j < k < \ell} (p_j - p_k) \\ = \gamma + \sum_{j < k < \ell} (1 - (p_j - p_k)) > 0, \end{aligned}$$

where the final inequality follows from Equation (26). We may therefore drop every term containing $d_{j,\ell}$. The remaining minimization is not a multivariable optimization problem: complete one square for each j ,

$$\frac{s_j^2}{2} - (1 - \gamma)s_j = \frac{1}{2}(s_j - (1 - \gamma))^2 - \frac{1}{2}(1 - \gamma)^2.$$

Because $1 - \gamma \in (0, 1)$, this minimum is feasible for $s_j \in [0, 1]$. Hence every online algorithm obeys

$$\liminf_{H \rightarrow \infty} \frac{\text{ALG}}{H^2} \geq Q_K := C_K - \frac{K}{2}(1 - \gamma)^2. \quad (34)$$

It remains to compute the offline benchmark. Knowing all processing times, the optimum runs the zero jobs first and the positive types from shortest to longest. Equivalently, the pairwise contribution of two jobs is $1 + \min\{p_i, p_j\}$. Its leading coefficient is

$$D_K = \frac{\xi^2}{2} + \xi K + \frac{K^2}{2} + P_K, \quad \lim_{H \rightarrow \infty} \frac{\text{OPT}}{H^2} = D_K. \quad (35)$$

The four terms respectively account for zero-zero pairs, zero-positive pairs, the unit-test part of positive-positive pairs, and the processing part of positive-positive pairs. Thus the finite- K lower-bound certificate is simply Q_K/D_K .

Passing to a smooth curve. Write

$$\alpha = \frac{K}{\xi}$$

for the ratio of positive mass to zero mass, and keep α fixed while $K \rightarrow \infty$. We restrict to $0 < \alpha < e - 1$. Then

$$p_0 - p_{K-1} = \sum_{r=1}^{K-1} \frac{1}{\xi + r} < \log\left(1 + \frac{K}{\xi}\right) = \log(1 + \alpha) < 1,$$

which verifies Equation (26).

The processing contribution P_K has the exact form

$$P_K = \frac{K^2}{2} \left(1 + \frac{\gamma}{\xi}\right) + \frac{1}{2} \sum_{h=1}^{K-1} \frac{(K-h)^2}{\xi+h}. \quad (36)$$

Indeed, the common baseline $1 + \gamma/\xi$ contributes the first term, and swapping the order of summation shows that the increment $1/(\xi+h)$ receives total weight $(K-h)^2/2$. After division by ξ^2 , the sum becomes a Riemann integral:

$$\begin{aligned} \frac{P_K}{\xi^2} &\longrightarrow I(\alpha) = \frac{\alpha^2}{2} + \frac{1}{2} \int_0^\alpha \frac{(\alpha-t)^2}{1+t} dt \\ &= \int_0^\alpha (\alpha-t)(1+\log(1+t)) dt \\ &= \frac{1}{2}(1+\alpha)^2 \log(1+\alpha) - \frac{\alpha}{2} - \frac{\alpha^2}{4}. \end{aligned} \quad (37)$$

The second integral form follows either by integration by parts or by summing the limiting processing-time curve $1+\log(1+t)$ against the remaining positive mass $\alpha-t$. The subtraction in Equation (34) is only $O(K)$, whereas $C_K, D_K = \Theta(K^2)$. Since $C_K - D_K = \xi K + K^2/2$, we obtain

$$R(\alpha) := \lim_{K \rightarrow \infty} \frac{Q_K}{D_K} = 1 + \frac{\alpha + \alpha^2/2}{\frac{1}{2} + \frac{\alpha}{2} + \frac{\alpha^2}{4} + \frac{1}{2}(1+\alpha)^2 \log(1+\alpha)}. \quad (38)$$

This last one-variable maximization is where the constant in the theorem comes from. Put $z = (1+\alpha)^2$. Then

$$R(z) = 1 + \frac{2(z-1)}{1+z+z \log z}. \quad (39)$$

The derivative of the fraction has the sign of

$$3 - z + \log z.$$

For $z > 1$ this expression is strictly decreasing, is positive at $z = 1$, and tends to $-\infty$. Hence it has one zero $z_\star$, characterized by

$$z_\star - 3 = \log z_\star.$$

The derivative is positive before $z_\star$ and negative afterward, so this is the global maximum. At the root,

$$1 + z_\star + z_\star \log z_\star = (z_\star - 1)^2,$$

and therefore

$$R(z_\star) = 1 + \frac{2}{z_\star - 1} = R_\star.$$

Moreover, $\sqrt{z_\star} - 1 = 1.12255\dots < e - 1$, so the maximizer lies in the range where Equation (26) holds.

Proof of Theorem 4.1. Choose a rational α sufficiently close to $\alpha_\star = \sqrt{z_\star} - 1$, and choose any rational $\gamma \in (0, 1)$. For a sufficiently large K , set $\xi = K/\alpha$ and use Equation (31); then $Q_K/D_K > R_\star - \varepsilon/2$. All parameters are rational. Let H tend to infinity through multiples of their common denominators. Equations (34) and (35) make the actual ratio exceed $R_\star - \varepsilon$ for all sufficiently large H . Finally, the replay argument at the beginning of the proof converts the revelation experiment into a fixed instance for the given deterministic algorithm. $\square$

Lemma 4.2 (Scaled harmonic core). *Fix a total normalized mass $s > 0$ and a parameter $t \in [1, e)$. Against every deterministic algorithm for the cap-free endpoint, there are scales $M_\nu \rightarrow \infty$ and finite rational multilevel instances with $sM_\nu + o(M_\nu)$ jobs such that*

$$\frac{\text{OPT}}{M_\nu^2} \rightarrow O_0 = \frac{s^2}{4}(1 + t^{-2} + 2 \log t), \quad \liminf_{\nu \rightarrow \infty} \frac{\text{ALG}}{M_\nu^2} \geq A_0^{\text{lb}} := O_0 + \frac{s^2}{2}(1 - t^{-2}). \quad (40)$$

Their normalized total positive processing mass converges to

$$\frac{1}{M_\nu} \sum_{j: p_j > 0} p_j \rightarrow L = s \log t, \quad (41)$$

and their largest processing time converges to $1 + \log t < 2$. The adaptive revelation is converted by replay into a fixed instance.

Proof. The case $t = 1$ is obtained by continuity from instances with vanishing positive mass, so suppose $1 < t < e$. In the construction above set $\alpha = t - 1 = K/\xi$. Before rescaling, the total normalized mass is $K + \xi = \xi t$. Give every unit of normalized mass the common factor $s/(\xi t)$.

The phase accounting, harmonic cancellation, and completion of squares in Equations (28), (32) and (34) are homogeneous of degree two in the job masses. The Riemann-sum limit Equation (37) substituted into Equation (35) therefore gives

$$O_0 = \frac{s^2}{4}(1 + t^{-2} + 2 \log t).$$

Likewise $C_K - D_K = \xi K + K^2/2$, while the $O(K)$ square-completion loss in Equation (34) disappears after division by the quadratic scale. After the same rescaling this gives the guaranteed lower benchmark

$$A_0^{\text{lb}} - O_0 = \frac{s^2}{2}(1 - t^{-2}).$$

Finally, summing the limiting processing-time curve $1 + \log(1 + x)$ over the positive mass gives $L = s \log t$, and Equation (31) gives $p_0 \rightarrow 1 + \log t$. Rational approximation, clearing denominators, and the replay argument used in the theorem produce a diagonal sequence of arbitrarily large fixed rational instances with the stated limit and liminf bounds. $\square$

4.2 Upper bound: a self-contained proof

We prove the following statement from scratch in this subsection.

Theorem 4.3 (Upper bound). *Let $z > 1$ be the solution of $z - 3 = \log z$, and set*

$$R = 1 + \rho, \quad \rho = \frac{2}{z - 1} = 0.5705740966 \dots$$

There is a deterministic online algorithm and an absolute constant C such that every instance with n jobs satisfies

$$\text{ALG} \leq R \text{OPT} + Cn.$$

The proof has three ingredients. First, pairwise accounting gives an exact formula for the cost of a schedule. Second, a simple convexity argument says that it is enough to consider four boundary outcomes for every test. Third, we give a nonnegative *bank balance* W : whenever a boundary outcome creates positive excess cost, the balance drops by at least that amount, up to a constant rounding loss. Thus the proof below uses derivatives only as shorthand for the first-order change of this bank balance; no optimization background is needed.

The algorithm. For a parameter $c > 0$, the formal procedure $\text{ADAPTIVETHRESHOLD}(c)$ is given in algorithm 1. Immediately before a test, let

- x be the number of jobs not yet tested, including the job about to be tested;
- N be the number of earlier tests that revealed a strictly positive processing time; and
- D be the number of these N jobs that the algorithm deferred.

Define the normalized state

$$y = \frac{D - RN}{x}. \quad (42)$$

Because $D \leq N$ and $R > 1$, we always have $y \leq 0$. For later reuse define the parameterized threshold

$$a_c(y) = \begin{cases} 1, & y \leq -1, \\ 1 + \frac{1}{2} \log\left(\frac{c+2}{c-2y}\right), & -1 \leq y \leq 0. \end{cases} \quad (43)$$

The theorem uses $c = \rho$; throughout this subsection we abbreviate $A(y) = a_\rho(y)$ and $R = 1 + \rho$. The defining equation for $z = (\rho + 2)/\rho$ says

$$\log \frac{\rho + 2}{\rho} = \frac{2}{\rho} - 2. \quad (44)$$

Consequently, A is continuous at -1 , increases from $A(-1) = 1$ to $A(0) = 1/\rho = 1.7526207478\dots$, and

$$A'(y) = \frac{1}{\rho - 2y} \quad (-1 < y < 0). \quad (45)$$

Algorithm 1 $\text{ADAPTIVETHRESHOLD}(J, c)$

Input: jobs J and parameter $c > 0$.

- 1: Fix an arbitrary test order $j_1, \dots, j_n$.
 - 2: Set $N \leftarrow 0$, $D \leftarrow 0$, and $Q \leftarrow \emptyset$.
 - 3: **For** $i = 1, \dots, n$ **do**:
 - 4: Set $x \leftarrow n - i + 1$ and $y \leftarrow (D - (1 + c)N)/x$.
 - 5: Set $a \leftarrow a_c(y)$ according to Equation (43).
 - 6: Test j_i for one unit and observe p_{j_i} .
 - 7: **If** $p_{j_i} \leq a$, process j_i immediately.
 - 8: **Otherwise**, add j_i to Q and set $D \leftarrow D + 1$.
 - 9: **If** $p_{j_i} > 0$, set $N \leftarrow N + 1$.
 - 10: Process the jobs of Q in nondecreasing order of their revealed processing times.
-

For $c = \rho$, the state in the pseudocode is exactly (42). A genuine zero increments neither counter and completes at the end of its test. The procedure never uses raw execution; the same pseudocode will therefore apply to the large finite- u regimes.

The numerator $D - RN$ in (42) is best viewed as a balance, not as an estimate of a processing time. Every revealed positive job reduces the balance by R ; deferring it returns one unit, because every later unit test will also delay that deferred job. Dividing by x spreads this balance over the tests that remain. A state close to 0 can afford a threshold larger than one. Once the normalized balance falls below -1 , the proof can only support the conservative threshold one, so A becomes flat.

We will later derive (43), rather than asking the reader to guess why this particular logarithm appears.

Pairwise accounting. For two jobs $i < j$ in test order, charge to their unordered pair the duration of operations of either job performed while the other job is unfinished. If job i is processed immediately, this charge in the algorithm is $1 + p_i$. If i is deferred and j is immediate, it is $2 + p_j$. If both are deferred, it is $2 + \min\{p_i, p_j\}$. Hence

$$\text{ALG} = \sum_i (1 + p_i) + \sum_{i < j} g_{ij}, \quad g_{ij} = \begin{cases} 1 + p_i, & i \text{ immediate,} \\ 2 + p_j, & i \text{ deferred, } j \text{ immediate,} \\ 2 + \min\{p_i, p_j\}, & i, j \text{ deferred.} \end{cases} \quad (46)$$

The offline optimum tests and immediately processes jobs in nondecreasing order of p_i , and therefore

$$\text{OPT} = \sum_i (1 + p_i) + \sum_{i < j} (1 + \min\{p_i, p_j\}). \quad (47)$$

For completeness, the latter claim follows because the k th completed job cannot finish before the machine has performed k tests and the processing of the k shortest jobs. The shortest-first schedule attains all these lower bounds simultaneously.

We shall bound the excess

$$F = \text{ALG} - R \text{OPT}.$$

Why four boundary outcomes suffice. Fix only the symbolic decisions made along a run: for each job, specify whether its value is zero, positive and immediate, or positive and deferred. This fixes all later counters and thresholds. The allowed values of a coordinate are then, respectively,

$$\{0\}, \quad (0, A(y)], \quad (A(y), \infty).$$

If all processing times except p_i are fixed inside such a symbolic run, (46)–(47) express F as an affine function of p_i plus negative multiples of $\min\{p_i, h\}$. Since $p \mapsto \min\{p, h\}$ is concave, F is convex in p_i . A convex function on $(0, A(y)]$ has its supremum at one of the two ends. On $(A(y), \infty)$, the eventual slope is $1 - R = -\rho < 0$; convexity then implies that every earlier slope is also negative, so the supremum is at the lower end.

It is therefore enough to consider the following four formal outcomes.

type	limiting value	algorithmic decision	counter update
Z	0	immediate	none
E	0^+	immediate	$N \leftarrow N + 1$
I	$A(y)$	immediate	$N \leftarrow N + 1$
Q	$A(y)^+$	deferred	$N \leftarrow N + 1, D \leftarrow D + 1$

Here 0^+ is not a new processing time. It means that a positive immediate value tends to zero *after its symbolic decision has been fixed*. Although its processing contribution vanishes, it still increments N and therefore changes all future thresholds. This is why 0 and 0^+ cannot be identified. Similarly, $A(y)^+$ approaches the threshold from the deferred side. Applying the one-coordinate argument successively shows that a uniform bound on all formal four-type words implies the same bound on all genuine instances. Because F is a finite sum of affine and minimum terms on a fixed symbolic path, the simultaneous one-sided limit at every formal vertex exists and is independent of the rates at which its open coordinates approach their endpoints.

One more elementary property will simplify every minimum in the pairwise formulas. Put $u = D - RN \leq 0$. After an outcome of type Z, E or I, or Q, respectively, the pair

(u, x) changes as follows:

$$(u, x - 1), \quad (u - R, x - 1), \quad (u - R, x - 1), \quad (u - \rho, x - 1).$$

Whenever $x \geq 2$, the new quotient is at most u/x . Thus the states, and hence the thresholds, seen by successive tests never increase. When $x = 1$ there is no next test and therefore no next threshold to compare. The nonzero boundary values $A(y)$ consequently arrive in nonincreasing order. In particular, the minimum of the current nonzero boundary value and any earlier one is always the current value.

The proof state and the local excess. The algorithm needs only (x, N, D) . The analysis splits N into two kinds. Among earlier formal outcomes, let

- S be the number of *substantive* positive outcomes I, Q;
- e be the number of infinitesimal outcomes E; and
- d be the number of deferred outcomes Q.

Thus $N = S + e$ and $D = d$. Define

$$\eta = \frac{d - RS}{x}, \quad b = \frac{e}{x}, \quad y = \eta - Rb. \quad (48)$$

We have $d \leq S$, hence $\eta \leq 0$. The variable b records exactly the part of the algorithmic state that is invisible to limiting processing-cost minima: an infinitesimal value affects N , but its minimum with every other processing time tends to zero.

Consider the current boundary outcome at a pre-state (x, S, e, d) , and write $a = A(y)$. Pairwise accounting gives

$$F = -\rho \frac{n(n+1)}{2} + \sum_i \tilde{\ell}_i, \quad (49)$$

where the exact contribution of the current outcome is

$$\tilde{\ell}_Z = \tilde{\ell}_E = d, \quad (50)$$

$$\tilde{\ell}_I = d + a[x + d - RS - R], \quad (51)$$

$$\tilde{\ell}_Q = d + a[d - RS - \rho]. \quad (52)$$

Here is a quick derivation. The current unit test gives one extra unit of algorithmic pair cost for each earlier deferred job, producing d . Its own test and its pairs with all earlier jobs give a universal excess $-\rho(i+1)$ at step i ; summing this term produces the first term in (49). For an I outcome, the current processing operation contributes $a(x+d)$ to the algorithm and $Ra(S+1)$ to the scaled optimum. For a Q outcome, the corresponding quantities are $a(d+1)$ and $Ra(S+1)$. Monotonicity of the thresholds justifies using the current a in every earlier processing-time minimum.

For the potential calculation it is convenient to discard the two favorable diagonal terms $-Ra$ and $-\rho a$. Define the relaxed local rewards

$$\ell_Z = \ell_E = d, \quad (53)$$

$$\ell_I = d + a(x + d - RS) = d + ax(1 + \eta), \quad (54)$$

$$\ell_Q = d + a(d - RS) = d + ax\eta. \quad (55)$$

Then $\tilde{\ell}_c \leq \ell_c$ for all four types, and

$$F \leq -\rho \frac{n(n+1)}{2} + \sum_i \ell_i. \quad (56)$$

How the logarithmic threshold is found. Before giving the full bank balance, it is useful to see where the formula comes from. Ignore 0^+ outcomes for a moment, so $b = 0$, and look in the range $-1 \leq y \leq 0$. Suppose the nonlinear part of the balance has the form $x^2 f(y)$. Asking the immediate boundary I and the deferred boundary Q to be exactly paid by the first-order decrease of this balance gives

$$-2f + (y - R)f' + A(1 + y) = 0, \quad (57)$$

$$1 - 2f + (y - \rho)f' + Ay = 0. \quad (58)$$

Subtracting the equations gives $f' = A - 1$. Differentiating either equation then gives

$$1 + (2y - \rho)A'(y) = 0.$$

With the natural joining condition $A(-1) = 1$, this is exactly (43)–(45).

The universal negative term in (56) has leading part $-\rho n^2/2$. The initial balance will be $n^2 f(0)$, so exact cancellation requires $f(0) = \rho/2$. Since

$$f(y) = \int_{-1}^y (A(t) - 1) dt, \quad (59)$$

direct integration gives

$$f(0) = \frac{1}{2} - \frac{\rho}{4} \log\left(\frac{\rho + 2}{\rho}\right).$$

Thus $f(0) = \rho/2$ is equivalent to (44), or, after setting $z = (\rho + 2)/\rho$, to $z - 3 = \log z$. Thus both the threshold and the constant R are forced by two local balance equations and the initial accounting; they are not the output of an unexplained numerical optimization.

The bank balance. For $-1 \leq y \leq 0$, put

$$f(y) = \int_{-1}^y (A(t) - 1) dt, \quad h(y) = A(y)(1 + y). \quad (60)$$

The term xd in the balance below pays for the fact that each of the x remaining tests will delay each of the d already deferred jobs. The $x^2 f(y)$ term handles substantive outcomes. Finally, the terms involving b repair the discontinuity between a genuine zero and 0^+ .

Define

$$W(x, S, e, d) = xd + x^2 G(y, b), \quad (61)$$

using two formulas for G :

$$G(y, b) = \begin{cases} f(y) + bh(y) + \frac{R}{2}b^2, & y \geq -1, \\ g(\eta), & g(\eta) = \frac{(1 + \eta)_+^2}{2R}, \quad y \leq -1, \end{cases} \quad \eta = y + Rb. \quad (62)$$

The first formula is used while the threshold is changing. The second is used once the threshold has saturated at one. In that flat region the limiting processing-cost terms depend on the substantive balance η , not on the number of 0^+ outcomes separately, which explains why g contains no b .

This piecewise definition is smooth enough for a first-order argument. At $y = -1$, feasibility gives $0 \leq b \leq 1/R$ and $\eta = -1 + Rb$. Since

$$f(-1) = h(-1) = 0, \quad h'(-1) = 1,$$

the first formula in (62) has value $Rb^2/2$ and derivatives $G_y = b$, $G_b = Rb$. The second formula has exactly the same value and derivatives. This also checks the derivative in the raw x coordinate: at fixed (S, e, d) ,

$$\partial_x [x^2 G(y, b)] = x(2G - yG_y - bG_b),$$

which agrees on the two sides because G , G_y , and G_b agree. The remaining raw derivatives are linear combinations of G_y and G_b . The internal join of $(1 + \eta)_+^2$ at $\eta = -1$ also has a continuous first derivative. Hence W is continuously differentiable. Moreover, $W \geq 0$: in the first region $f \geq 0$, $h \geq 0$, and $b, d \geq 0$; in the second this is immediate from (62).

The choice $h = A(1 + y)$ also has a direct interpretation. At $b = 0$, it makes the first infinitesimal outcome 0^+ exactly neutral in the local balance. The quadratic correction $Rb^2/2$ then maintains the balance for all b and is precisely what makes the two regions meet smoothly.

Four local balance checks. We now verify that every formal outcome can be paid from the bank. In raw coordinate order (x, S, e, d) , the four unit-step directions are

$$v_Z = (-1, 0, 0, 0), \quad v_E = (-1, 0, 1, 0), \quad v_I = (-1, 1, 0, 0), \quad v_Q = (-1, 1, 0, 1). \quad (63)$$

For an endpoint type $q \in \{Z, E, I, Q\}$, define its first-order balance

$$J_q = \ell_q + \nabla W \cdot v_q. \quad (64)$$

The inequality $J_q \leq 0$ simply says: local reward plus first-order change in the bank balance is nonpositive.

We need only a few one-variable identities. On $-1 \leq y \leq 0$,

$$f' = A - 1, \quad f(-1) = 0, \quad f(0) = \frac{\rho}{2}, \quad (65)$$

$$-2f + (y - R)f' + A(1 + y) = 0, \quad (66)$$

$$1 - 2f + (y - \rho)f' + Ay = 0, \quad (67)$$

$$-2f + yf' = A(\rho - y) - R \leq 0, \quad (68)$$

$$h + (\rho - y)h' = AR + (\rho - y)(1 + y)A' \geq AR. \quad (69)$$

The first line follows from the definition and (44). For (66)–(67), differentiate the left-hand sides and use $A' = 1/(\rho - 2y)$; both derivatives vanish, and both expressions equal zero at $y = -1$. Equation (68) follows from (66). Its right-hand side is zero at -1 and decreases afterward, because

$$\frac{d}{dy} [A(y)(\rho - y)] = \frac{\rho - y}{\rho - 2y} - A(y) \leq 0.$$

Indeed, $y \leq 0$ gives $(\rho - y)/(\rho - 2y) \leq 1$, while $A(y) \geq 1$. Finally, (69) is obtained by differentiating $h = A(1 + y)$. Notice also that $h, h' \geq 0$.

In the changing-threshold region $y \geq -1$, direct differentiation of (61) first gives, for an arbitrary $G(y, b)$,

$$J_Z/x = -2G + yG_y + bG_b, \quad (70)$$

$$J_E/x = -2G + (y - R)G_y + (b + 1)G_b, \quad (71)$$

$$J_I/x = -2G + (y - R)G_y + bG_b + A(1 + y + Rb), \quad (72)$$

$$J_Q/x = 1 - 2G + (y - \rho)G_y + bG_b + A(y + Rb). \quad (73)$$

Substituting $G = f + bh + Rb^2/2$ and using (66)–(67) reduces the four expressions to

$$J_Z/x = (-2f + yf') + b(-h + yh'), \quad (74)$$

$$J_E/x = b[-h + (y - R)h' + R], \quad (75)$$

$$J_I/x = b[-h + (y - R)h' + AR], \quad (76)$$

$$J_Q/x = b[-h + (y - \rho)h' + AR]. \quad (77)$$

Now the signs are transparent. The Z line is nonpositive by (68), $h, h' \geq 0$, and $y \leq 0$. The Q line is nonpositive by (69). The I line is no larger because $y - R < y - \rho$ and $h' \geq 0$; the E line is no larger again because $A \geq 1$. Thus all four checks hold in the changing-threshold region.

In the flat region $y \leq -1$, we have $A = 1$ and $G = g(\eta)$. The four checks become

$$J_Z/x = J_E/x = -2g + \eta g', \quad (78)$$

$$J_I/x = -2g + (\eta - R)g' + 1 + \eta, \quad (79)$$

$$J_Q/x = 1 - 2g + (\eta - \rho)g' + \eta. \quad (80)$$

If $\eta \leq -1$, then $g = g' = 0$, and these expressions are 0, $1 + \eta$, and $1 + \eta$. If $-1 \leq \eta \leq 0$, then $g' = (1 + \eta)/R$, and they are, respectively,

$$-\frac{1 + \eta}{R}, \quad -\frac{1 + \eta}{R}, \quad 0.$$

They are therefore nonpositive in both cases. We have proved

$$J_Z, J_E, J_I, J_Q \leq 0 \quad (81)$$

at every feasible state.

From infinitesimal checks to unit steps. Equation (81) compares a local reward with the *derivative* of W , whereas one job changes the integer counters by one. This loses only a constant per job. More precisely, there is an absolute constant C_0 such that every admissible transition satisfies

$$W(s + v_q) - W(s) \leq \nabla W(s) \cdot v_q + C_0. \quad (82)$$

We include the bounded-curvature argument because it is what makes the $O(n)$ term uniform over all instances.

Suppose first that $x \geq 2$ and linearly interpolate the unit transition. The numerators of both y and η have the form $u - \kappa t$, where $u \leq 0$ and $\kappa \geq 0$, while the denominator is $x - t$. Their derivatives are therefore nonpositive, so the segment crosses each joining surface at most once. In the changing-threshold region, $y \geq -1$ implies

$$R(S + e) - d = \rho S + (S - d) + Re \leq x.$$

Consequently,

$$0 \leq \frac{S}{x} \leq \frac{1}{\rho}, \quad 0 \leq \frac{e}{x} \leq \frac{1}{R}, \quad 0 \leq \frac{d}{x} \leq \frac{1}{\rho}.$$

After normalizing by x , all reachable states in this region thus lie in a fixed compact set. The balance W is homogeneous of degree two, so its second derivatives in the raw

coordinates are homogeneous of degree zero and are uniformly bounded on this normalized set.

In the flat region the formula is even simpler:

$$W = xd + \frac{(x + d - RS)_+^2}{2R}. \quad (83)$$

Its gradient is globally Lipschitz. Because the two regional formulas have the same first derivative at their interface, an interpolated unit step has a uniformly bounded second derivative except at finitely many harmless joining points. More explicitly, let $\phi(t) = W(s + tv_q)$. The preceding bounds and the C^1 gluing make ϕ' Lipschitz with one uniform constant L along the whole segment, so

$$|\phi(1) - \phi(0) - \phi'(0)| \leq \int_0^1 |\phi'(t) - \phi'(0)| dt \leq \int_0^1 Lt dt = \frac{L}{2}.$$

This proves (82) for $x \geq 2$. When $x = 1$, the changing-threshold pre-states lie in the same compact normalized set, while (83) applies globally in the flat region. At $x = 0$ we define $W = 0$; this agrees with (83), since terminal feasibility gives $d \leq S$ and hence $d - RS \leq 0$.

Combining (64)–(82) gives the actual one-step bank inequality

$$\ell_i + W(s_{i+1}) - W(s_i) \leq C_0. \quad (84)$$

Telescoping the account. Initially $S = e = d = 0$, $x = n$, and $y = 0$. Therefore

$$W(s_0) = n^2 f(0) = \frac{\rho}{2} n^2. \quad (85)$$

The terminal balance is zero. Summing (84) over all n jobs yields

$$\sum_i \ell_i \leq \frac{\rho}{2} n^2 + C_0 n.$$

Substitution into (56) gives

$$\text{ALG} - R \text{OPT} = F \leq -\rho \frac{n(n+1)}{2} + \frac{\rho}{2} n^2 + C_0 n = \left(C_0 - \frac{\rho}{2}\right) n.$$

The constant is independent of the four-type word. Taking $C = \max\{0, C_0 - \rho/2\}$, the boundary reduction therefore extends the same estimate to every vector of nonnegative processing times, proving Theorem 4.3.

Finally, $\text{OPT} \geq n(n+1)/2$, so the additive $O(n)$ term is lower order and

$$\frac{\text{ALG}}{\text{OPT}} \leq R + O(1/n).$$

This proves the desired size-asymptotic upper bound. Notice exactly where the remaining finite-instance loss enters: it is only the uniform Taylor remainder in (82) (together with the deliberately discarded favorable diagonal terms), not any unproved limiting step in the endpoint reduction. In particular, this argument proves a linear additive term; it does not prove one fixed additive constant at the endpoint R .

Corollary 4.4 (Exact asymptotic values). *The optimal deterministic size-asymptotic value and the optimal deterministic additive-constant infimum both equal $R_\star$. Moreover, for every $\varepsilon > 0$, the Adaptive-Threshold algorithm satisfies*

$$\text{ALG}(I) \leq (R_\star + \varepsilon) \text{OPT}(I) + B_\varepsilon \quad (86)$$

for all instances, with B_ε independent of I .

Proof. The upper bound and Equation (21) imply

$$\frac{\text{ALG}}{\text{OPT}} \leq R_\star + \frac{2C}{n+1},$$

so the size-asymptotic ratio is at most $R_\star$. The lower bound is Theorem 4.1. For the additive statement, Young's inequality and Equation (21) give

$$Cn \leq \frac{\varepsilon}{2}n^2 + \frac{C^2}{2\varepsilon} \leq \varepsilon \text{OPT} + \frac{C^2}{2\varepsilon}.$$

Thus one can take $B_\varepsilon = C^2/(2\varepsilon)$, proving that every coefficient above $R_\star$ is admissible for this algorithm. Conversely, fix $c < R_\star$. By Theorem 4.1, every deterministic algorithm has a sequence of instances with $\text{OPT} \rightarrow \infty$ and $\text{ALG} / \text{OPT} > c + \delta$ for some $\delta > 0$. The inequality $\text{ALG} \leq c \text{OPT} + B$ would then force $\delta \leq B / \text{OPT} \rightarrow 0$, a contradiction. This proves the additive-constant claim. $\square$

5 Finite caps: the exact curve

We now move from the cap-free endpoint to finite u . Raw execution becomes available, all jobs have the same known upper limit, and a test reveals $p_j \in [0, u]$. Thus the offline effective length of job j is

$$q_j = \min\{u, 1 + p_j\}, \quad (87)$$

and the offline optimum schedules the q_j in nondecreasing order. We write $V(u)$ for the optimal deterministic size-asymptotic ratio at this fixed common upper limit.

The answer has six branches. We first define their transition points. Put

$$\varphi = \frac{1 + \sqrt{5}}{2}, \quad u_\diamond = \frac{1 + \sqrt{3 + 2\sqrt{5}}}{2} = 1.866760399173862\dots \quad (88)$$

Let $s_0 > 1$ be the root of

$$s_0^3 = (s_0 + 1)^2, \quad u_0 = 1 + s_0 = 3.147899035704787\dots \quad (89)$$

For $s = u - 1$, put

$$T_s(z) = sz^2 + (s^3 + s^2 + 1)z - (s^2 + s + 1)(s + 2), \quad (90)$$

and let $\rho_I(u)$ be the positive root of

$$s\rho^2 + (s^3 + s^2 + 1)\rho - (s^2 + s + 1)(s + 2) = 0. \quad (91)$$

Denote the left-hand side by $T_s(\rho)$. Equivalently,

$$\rho_I(u) = \frac{-1 - u + 2u^2 - u^3 + \sqrt{u^6 - 4u^5 + 10u^4 - 6u^3 - 3u^2 + 6u - 3}}{2(u - 1)}. \quad (92)$$

At the other end, let $z_\star > 1$ solve $z_\star - 3 = \log z_\star$ and define

$$r = \frac{2}{z_\star - 1}, \quad R_{\text{obl}} = 1 + r = 1.570574096649395\dots, \quad z_\star = 4.505241495792883\dots \quad (93)$$

For $\varphi + 2 \leq u \leq z_\star$, define $C_{\text{mix}}(u) = 1 + c$ by the unique solution (c, m) of

$$\text{artanh } m - m = 1 - \frac{1}{c} + \frac{1}{2} \log\left(1 + \frac{2}{c}\right), \quad u = 1 + \frac{2}{c} - m, \quad (94)$$

with $r \leq c \leq 1/\varphi$ and $0 \leq m \leq 1/\varphi$.

Theorem 5.1 (Exact common-upper curve). *For every fixed $u > 0$, the optimal deterministic size-asymptotic ratio in the optional-testing model is*

$$V(u) = \begin{cases} 1, & 0 < u \leq 1, \\ u, & 1 \leq u \leq u_\diamond, \\ \rho_I(u), & u_\diamond \leq u \leq u_0, \\ 1 + \frac{1}{\sqrt{u-1}}, & u_0 \leq u \leq \varphi + 2, \\ C_{\text{mix}}(u), & \varphi + 2 \leq u \leq z_*, \\ R_{\text{obl}}, & u \geq z_*. \end{cases} \quad (95)$$

For every branch there is an explicit deterministic algorithm satisfying

$$\text{ALG} \leq V(u) \text{OPT} + O_u(n).$$

On finite $u \geq z_*$ the remainder is one absolute $O(n)$ term, and Theorem 2.1 supplies the same algorithm at $u = \infty$. Conversely, for every deterministic algorithm $\mathcal{A}$ and every $\epsilon > 0$, there are arbitrarily large fixed instances I such that

$$\text{ALG}_{\mathcal{A}}(I) \geq (V(u) - \epsilon) \text{OPT}(I).$$

The upper-bound witness can be read from the following policy map. The algorithms themselves are specified formally in algorithms 1 to 3.

range	policy	parameter choice
$0 < u \leq u_\diamond$	RAW	none
$u_\diamond < u \leq u_0$	FORCEDPREFIXUTE	$b = b(u)$ from (112)
$u_0 < u \leq \varphi + 2$	FORCEDPREFIXUTE	$b = 0$
$\varphi + 2 < u < z_*$	ADAPTIVETHRESHOLD	$c = c(u)$ from (94)
$z_* \leq u \leq \infty$	ADAPTIVETHRESHOLD	$c = r$

The global-maximum assertion follows directly from the displayed formulas. Only the algebraic branch needs a check. Implicit differentiation of $T_s(\rho_I(u)) = 0$, with $s = u - 1$, gives

$$\rho'_I(u) = -\frac{\rho^2 + (3s^2 + 2s)\rho - 3(s+1)^2}{2s\rho + s^3 + s^2 + 1}. \quad (96)$$

Using $T_s(\rho) = 0$, its numerator factors as

$$\frac{2s^3 + s^2 - 1}{s} \left(\rho - \frac{2s^3 + 3s^2 - 2}{2s^3 + s^2 - 1} \right).$$

On the relevant interval $s \geq u_\diamond - 1 > 2/3$, the prefactor is positive, while the fraction in parentheses is at most $5/3$ because $4s^3 - 4s^2 + 1 > 0$. The sign certificate (116) gives $\rho > 5/3$. Hence $\rho'_I(u) < 0$. The reciprocal branch and the mixed parametrization are also decreasing, so V attains its global maximum only at $u = u_\diamond$.

The lower bounds, including the fact that the middle construction is safe against raw execution, are proved next. We then give self-contained upper bounds through $u = \varphi + 2$. The parameterized upper bound for the last two branches is proved separately in Section 5.6.

5.1 The binary stopping game and its lower envelope

The first four branches already occur on binary instances $p_j \in \{0, u\}$. The stopping point must be hidden from the algorithm; announcing it in advance gives a strictly weaker lower bound.

Lemma 5.2 (Raw-safe hidden stopping). *Fix $u > 1$, a target $\rho \leq u$, and a stopping fraction $\alpha \in (0, 1)$. Against a deterministic algorithm, there is a legal binary adversary whose leading excess at its first crossing is*

$$(u - \rho)(1 - \sigma^2) + \sigma^2 f_{u,\rho}(y, b), \quad (97)$$

where $0 \leq b \leq y \leq 1$ and

$$0 \leq y - \alpha < \frac{1}{n - v}. \quad (98)$$

Moreover,

$$f_{u,\rho}(y, b) = 1 - \rho + 2y + (u - 1)(1 - \rho)y^2 + b^2 + 2b(u - 1 - uy). \quad (99)$$

Here $\sigma \in [0, 1]$ is the fraction of jobs not already executed raw. Consequently, if

$$\min_{0 \leq b \leq \alpha} f_{u,\rho}(\alpha, b) \geq 0,$$

then the adversary gives ratio at least $\rho - O_u(1/n)$.

Proof. Start in a long phase. A fresh job that is tested receives $p = u$, while a fresh job run raw is assigned the hidden value zero. If v jobs have been run raw and L tested-long jobs have been seen, stop at the first touch for which

$$L \geq \alpha(n - v). \quad (100)$$

All untouched jobs are then assigned value zero. Only for the purpose of lower-bounding the remaining cost, we grant the algorithm this future information and allow an optimal continuation; in the actual interaction, values are still revealed only by tests. If the line is never crossed, every job was run raw: if even one job had been tested, then after the last first touch the tested fraction would be one. On the all-zero hidden instance the ratio is $u \geq \rho$.

At a crossing, let e of the L long jobs already be complete, let $d = L - e$ be pending, and let $x = n - v - e - d$ jobs remain untouched. The following favorable canonical schedule lower-bounds the algorithm: run the v raw jobs; test and immediately process the e completed long jobs; perform the d noncompleting long tests; test the x zero jobs; and finally process the d pending long jobs. Indeed, if a noncompleting unit test is immediately before an available length- u completion while h jobs remain, the two orders contribute $h + hu$ and $hu + (h - 1)$, respectively; moving the completion left saves one. Raw completions are available from time zero, and identical long jobs may be relabelled so that the e completed ones are tested first. After the switch, a zero test-completion has length one and therefore precedes a pending operation of length u . These exchanges respect precedence.

The exact costs of this relaxed schedule and of OPT are

$$A_{\text{ex}} = u \left(vn - \frac{v(v-1)}{2} \right) + (1+u) \left(e(n-v) - \frac{e(e-1)}{2} \right)$$

$$+ d(n - v - e) + x(d + x) - \frac{x(x - 1)}{2} + u \frac{d(d + 1)}{2}, \quad (101)$$

$$O_{\text{ex}} = \frac{n(n + 1)}{2} + (u - 1) \frac{(e + d)(e + d + 1)}{2}. \quad (102)$$

Put $\nu = v/n$, $\delta = (v + e + d)/n$, and $\lambda = e/n$. Twice the quadratic coefficients in Equations (101) and (102) are

$$\begin{aligned} \mathcal{A} &= 1 + 2\delta(1 - u\nu) + (u - 1)\delta^2 + 2\nu(\nu + u - 2) \\ &\quad + \lambda^2 + 2\lambda(\nu + u - 1 - u\delta), \\ \mathcal{O} &= 1 + (u - 1)(\delta - \nu)^2. \end{aligned}$$

Now put $\sigma = 1 - \nu$, $y = (e + d)/(n - v)$, and $b = e/(n - v)$. Direct expansion gives exactly Equation (97) after normalizing by $n^2/2$. The first-crossing overshoot in Equation (100) is less than one: a test increases $e + d - \alpha(n - v)$ by one and a raw first touch increases it by α . This proves Equation (98), and in particular $\sigma(y - \alpha) < 1/n$. There is one small domain issue: the actual $b \leq y$ can exceed α . Put $\bar{b} = \min\{b, \alpha\}$. Then $0 \leq b - \bar{b} \leq y - \alpha$. Since $f_{u,\rho}$ is Lipschitz on the unit square,

$$\sigma^2 f_{u,\rho}(y, b) \geq \sigma^2 f_{u,\rho}(\alpha, \bar{b}) - O_u(1/n). \quad (103)$$

Thus the stated minimum over $0 \leq \bar{b} \leq \alpha$ suffices uniformly even when $n - v$ is small. The discarded diagonal terms are $O_u(n)$ in the unnormalized costs. Finally, the adaptive transcript defines a fixed instance by replay: every tested answer was fixed when revealed, and every raw job's zero value stayed hidden. This proves the lemma. $\square$

Minimizing Equation (99) over the algorithm's completion choice gives

$$b_*(y) = \max\{0, uy - (u - 1)\}. \quad (104)$$

The upper constraint $b \leq y$ never binds, because $uy - (u - 1) \leq y$ for $y \leq 1$. There are therefore two elementary quadratic branches. On the positive branch, put

$$A_u = u^2 - u + 1, \quad D_{u,\rho} = A_u + (u - 1)\rho. \quad (105)$$

Then

$$\min_b f_{u,\rho}(y, b) = 1 - \rho - (u - 1)^2 + 2A_u y - D_{u,\rho} y^2. \quad (106)$$

Its maximizer is $\alpha = A_u/D_{u,\rho}$, and tangency to zero is

$$(1 - \rho - (u - 1)^2)D_{u,\rho} + A_u^2 = 0. \quad (107)$$

Expanding Equation (107) gives Equation (91). On the zero- b branch,

$$\min_b f_{u,\rho}(y, b) = 1 - \rho + 2y + (u - 1)(1 - \rho)y^2, \quad (108)$$

and tangency gives

$$\rho = 1 + \frac{1}{\sqrt{u - 1}}, \quad \alpha = \frac{1}{\sqrt{u - 1}}. \quad (109)$$

The latter point belongs to the zero- b branch iff $(u - 1)^3 \geq u^2$, with equality at $u = u_0$.

For $\rho = u$, the maximum in Equation (106) is nonnegative exactly while

$$[u(u - 1)]^2 - u(u - 1) - 1 \leq 0, \quad (110)$$

whose positive equality is $u = u_\diamond$. We have proved the following binary lower envelope.

Proposition 5.3 (Binary lower curve). *Every deterministic algorithm has size-asymptotic ratio at least*

$$B(u) = \begin{cases} 1, & 0 < u \leq 1, \\ u, & 1 \leq u \leq u_\diamond, \\ \rho_I(u), & u_\diamond \leq u \leq u_0, \\ 1 + \frac{1}{\sqrt{u-1}}, & u \geq u_0. \end{cases} \quad (111)$$

The lower bound uses only $p_j \in \{0, u\}$.

Proof. For $1 < u \leq u_\diamond$, use $\rho = u$ and the maximizing point of Equation (106). For $u_\diamond \leq u \leq u_0$, use the positive root $\rho_I(u)$ and $\alpha = A_u/D_{u, \rho_I(u)}$; the tangency identity makes the maximum zero. The minimizing completion mass is positive in the interior and becomes zero at $u = u_0$. For $u \geq u_0$, use the zero- b tangency point from Equation (109). The branch conditions are precisely the two inequalities checked above. Finally, $V(u) \geq 1$ is trivial for $u \leq 1$. In each nontrivial case Lemma 5.2 supplies fixed binary instances by replay. $\square$

5.2 Matching upper bounds through the binary regime

The first policy is formalized in algorithm 2.

Algorithm 2 RAW(J, u)

Input: jobs J and a finite common upper limit u .

- 1: Fix an arbitrary order $j_1, \dots, j_n$.
 - 2: **For** $i = 1, \dots, n$, execute j_i untested for u time.
-

For $u \leq u_\diamond$, use RAW. It is optimal when $u \leq 1$; when $u > 1$, every effective length in Equation (87) is at least one, so the ratio is at most u .

For the next interval, let $\rho = \rho_I(u)$, $s = u - 1$, and define

$$K = s^2 + s + 1, \quad D = K + s\rho, \quad b = \frac{K - s^2\rho}{D}, \quad \chi = \frac{1 - b}{\rho}, \quad \alpha = \frac{K}{D}. \quad (112)$$

The defining equation for ρ gives

$$0 \leq b < \chi < \alpha < 1, \quad \chi = \frac{s(s+1)}{D}, \quad b = (s+1)\alpha - s; \quad (113)$$

For completeness, these inequalities follow without numerical root isolation. Since T_s is increasing on positive arguments,

$$T_s\left(\frac{K}{s^2}\right) = \frac{K}{s^3}(-s^3 + s^2 + 2s + 1) \geq 0 \quad (s \leq s_0), \quad (114)$$

and hence $b = (K - s^2\rho)/D \geq 0$, with equality only at $s = s_0$. Moreover,

$$\alpha - \chi = \frac{1}{D}, \quad \chi - b = \frac{s^2\rho - 1}{D} > 0. \quad (115)$$

For the last sign,

$$9T_s(5/3) = f(s) := 6s^3 - 12s^2 - 2s - 3 < 0. \quad (116)$$

throughout the present interval. Indeed, on $[4/5, 1]$ the polynomial is decreasing and $f(4/5) = -1151/125$; on $[1, s_0]$ it has one interior minimum, while $f(1) = -11$ and $f(s_0) < f(13/6) = -95/36$. Here $s_0 < 13/6$ follows by substituting $13/6$ in $s^3 - (s+1)^2$. Thus $\rho > 5/3$, while $s > 4/5$, and $s^2\rho > 16/15$. Finally $K < D$ gives $\alpha < 1$. In particular, $b = 0$ occurs only at $u = u_0$. The remaining two elementary branches use the single procedure in algorithm 3.

Algorithm 3 FORCEDPREFIXUTE(J, u, b)

- Input:** jobs J , finite $u > 1$, and $b \in [0, 1]$.
- 1: Fix a test order $j_1, \dots, j_n$; set $k \leftarrow \lfloor bn \rfloor$, $\tau \leftarrow \min\{1, u-1\}$, and $Q \leftarrow \emptyset$.
 - 2: **For** $i = 1, \dots, n$ **do**:
 - 3: Test j_i for one unit and observe p_{j_i} .
 - 4: **If** $i \leq k$ or $p_{j_i} \leq \tau$, process j_i immediately.
 - 5: **Otherwise**, add j_i to Q .
 - 6: Process the jobs of Q in nondecreasing revealed processing time.
-

On $u_\diamond \leq u \leq u_0$, use FORCEDPREFIXUTE with b from Equation (112). Rounding bn changes the cost by $O_u(n)$.

Here is the complete binary calculation. Adjacent exchanges place as many long jobs as possible in the forced prefix and put all suffix-long jobs before suffix zeros. If x is the total long mass, then the prefix-long mass is $a = \min\{b, x\}$ and the deferred-long mass is $d = (x - b)_+$. After division by n^2 , the leading excess $G(x) = A - \rho O$ is

$$G(x) = \begin{cases} \frac{1-\rho}{2} + u \left(x - \frac{x^2}{2} \right) - \frac{\rho s}{2} x^2, & 0 \leq x \leq b, \\ \frac{1-\rho}{2} + u \left(b - \frac{b^2}{2} \right) + (1-b)(x-b) + \frac{s}{2}(x-b)^2 - \frac{\rho s}{2} x^2, & b \leq x \leq 1. \end{cases} \quad (117)$$

The second quadratic is strictly concave, has its stationary point at $x = \alpha$, and satisfies $G(\alpha) = 0$; after substituting Equation (112), the latter identity is exactly Equation (91). Since $b < \alpha$, its right derivative at b is positive. The left derivative satisfies $G'_-(b) = G'_+(b) + s(1-b) > 0$, and the first quadratic is concave, so it is increasing on $[0, b]$. Thus $G(x) \leq 0$ for every $x \in [0, 1]$, including the limiting case $b = 0$. This proves the binary upper bound without a hidden local-versus-global step.

For $u_\diamond \leq u \leq 2$, arbitrary values reduce to binary values by independent Bernoulli extremalization, in the following order. First raise *every* suffix-deferred value to u . This cannot lower ALG, while its offline effective length was already u . Only after this deterministic step, replace a prefix value p by u with probability p/u and by zero otherwise, and replace a suffix-immediate value $p \leq u-1$ by u with probability p and by zero otherwise. Notice that the u realization of a suffix-immediate job becomes deferred, so status is not preserved. Nevertheless the online check is pairwise. For jobs i before j , the leading pair contribution is

$$g_{ij} = \begin{cases} 1 + p_i, & i \text{ is immediate,} \\ 2 + p_j, & i \text{ is deferred and } j \text{ is immediate,} \\ 2 + \min\{p_i, p_j\}, & i, j \text{ are deferred.} \end{cases} \quad (118)$$

A rounded prefix job stays immediate and preserves its expected value. If i is suffix-immediate, its rounded contribution is one when it becomes zero and at least two when it

becomes u , hence its expectation is at least $1 + p_i$. If i is deferred and j is suffix-immediate, then i has been raised to u and the expected pair contribution is $2 + up_j \geq 2 + p_j$; a deferred-deferred pair can only increase. Thus expected ALG does not decrease, up to its $O_u(n)$ diagonal terms.

The expected effective lengths are, respectively,

$$1 + \frac{u-1}{u}p \leq q(p), \quad 1 + (u-1)p \leq 1 + p, \quad (119)$$

and u for a deferred value. Since the minimum is jointly concave, $\mathbb{E} \min(q'_i, q'_j) \leq \min(q_i, q_j)$. Thus some binary realization has competitive excess at least that of the original instance.

It remains to justify arbitrary values for $2 \leq u \leq u_0$. The following quadratic lemma records the entire endpoint calculation.

Lemma 5.4 (UTE endpoint game). *For $2 \leq u \leq u_0$, UTE with the parameters Equation (112) satisfies*

$$\text{ALG} \leq \rho_I(u) \text{OPT} + O_u(n). \quad (120)$$

Proof. Fix the symbolic prefix/immediate/deferred decisions. Convexity in one coordinate reduces a prefix job to $p \in \{0, u\}$, a suffix-immediate job to $p \in \{0, 1\}$, and a suffix-deferred job to $p \in \{1^+, u\}$. For the last claim, on $1 < p < u - 1$ the deferred-deferred term has coefficient $1 - \rho < 0$ multiplying a concave minimum; on $[u - 1, u]$, OPT is flat and ALG is nondecreasing. Adjacent exchanges give the canonical endpoint order.

Normalize counts by n . Let a be prefix- u mass, d suffix- u mass, t suffix- 1^+ mass, and m suffix-immediate-one mass. Their constraints are

$$0 \leq a \leq b, \quad d, t, m \geq 0, \quad d + t + m \leq 1 - b. \quad (121)$$

Pairwise accounting using Equation (118) gives

$$A = \frac{1}{2} + (s+1) \left(a - \frac{a^2}{2} \right) + (1-b)(d+t+m) - \frac{m^2}{2} + \frac{s}{2}d^2, \quad (122)$$

$$O = \frac{1}{2} + \frac{(a+d+t+m)^2}{2} + \frac{s-1}{2}(a+d)^2. \quad (123)$$

Moving m to t does not change O and increases A , so $m = 0$. Put $x = a + d$ and write $F(x) = A - \rho O$ after optimizing the other variables. The derivative of the gap in t is $1 - b - \rho(x + t)$, so its maximizing value is $(\chi - x)_+$. At fixed x, t , moving mass from d to a leaves O unchanged and changes A at rate $s(1 - x) + b - a \geq 0$. Hence

$$t = (\chi - x)_+, \quad a = \min\{b, x\}, \quad d = (x - b)_+. \quad (124)$$

For $x \geq \chi$, the gap $A - \rho O$ is the binary concave quadratic already discussed; it is tangent to zero at $x = \alpha$. For $x < \chi$, put

$$C_0 = -\frac{\rho-1}{2} + \frac{(1-b)^2}{2\rho}, \quad \gamma = s - \rho(s-1). \quad (125)$$

Substitution gives

$$A - \rho O = C_0 + a(s + b - sx) - \frac{a^2}{2} + \frac{\gamma x^2}{2}. \quad (126)$$

One has $\gamma > 0$ on $1 \leq s \leq s_0$. Indeed, $T_s(2) = s(s^2 - s + 1) > 0$, so for $s \leq 2$ the equation $T_s(\rho) = 0$ gives $\rho < 2$, and hence $\gamma > 2 - s \geq 0$. For $2 \leq s \leq s_0 < 13/6$,

$$16T_s(7/4) = 12s^3 - 20s^2 + s - 4 > 0; \quad (127)$$

the polynomial is 14 at $s = 2$ and has positive derivative thereafter. Thus $\rho < 7/4$ and $\gamma > 7/4 - 3s/4 > 1/8$.

On $0 \leq x \leq b$, one has $a = x$, and

$$A - \rho O = C_0 + (s + b)x - \frac{s + 1 + \rho(s - 1)}{2}x^2. \quad (128)$$

Its derivative is decreasing, and the derivative at b multiplied by D is

$$\rho(1 + s^2 + \rho s^2(s - 1)) > 0, \quad (129)$$

so the maximum is at b . On $b \leq x \leq \chi$, Equation (126) is convex and its maximum is at b or χ . The endpoint χ lies on the binary branch. Finally,

$$F(b) - F(\chi) = \frac{\chi - b}{2} [2sb - \gamma(b + \chi)], \quad (130)$$

and direct use of Equation (112) gives

$$D[\gamma(b + \chi) - 2sb] = -s + (1 + s - s^3)\rho + s^2(s - 1)\rho^2 > 0. \quad (131)$$

For a rational sign certificate, T_s is increasing on positive arguments and

$$9T_s(5/3) = 6s^3 - 12s^2 - 2s - 3 < 0, \quad (132)$$

so $\rho > 5/3$. If the right-hand side of Equation (131) is denoted by $q_s(\rho)$, then

$$3 \partial_\rho q_s(5/3) = 7s^3 - 10s^2 + 3s + 3 > 0. \quad (133)$$

The last polynomial is increasing for $s \geq 1$, so q_s is increasing for $\rho \geq 5/3$. Finally,

$$9q_s(5/3) = 10s^3 - 25s^2 + 6s + 15 > 0. \quad (134)$$

For the last sign, write $s = 3/2 + z$; the expression becomes $3/2 - 3z/2 + 20z^2 + 10z^3$. It is at least $3/2$ for $-1/2 \leq z \leq 0$, and for $z \geq 0$ it is larger than $3/2 - 9/320$. These inequalities prove that every endpoint gap is nonpositive. Diagonals and prefix rounding contribute $O_u(n)$. $\square$

At $u = u_0$, $b = 0$. The same zero-prefix algorithm remains exact through $u = \varphi + 2$.

Lemma 5.5 (Zero-prefix extension). *For $u_0 \leq u \leq \varphi + 2$, FORCEDPREFIXUTE($J, u, 0$) from algorithm 3 is $(1 + 1/\sqrt{u - 1})$ -competitive up to $O_u(n)$.*

Proof. The same coordinatewise reduction leaves capped-deferred mass d , boundary-deferred mass t , immediate-one mass m , and zeros. With $s = u - 1$, these masses satisfy $d, t, m \geq 0$ and $d + t + m \leq 1$, and the leading costs are

$$A = \frac{1}{2} + d + t + m - \frac{m^2}{2} + \frac{sd^2}{2}, \quad O = \frac{1}{2} + \frac{(d + t + m)^2}{2} + \frac{(s - 1)d^2}{2}. \quad (135)$$

Move m to t and put $y = d + t$. For fixed y , the ratio is fractional-linear in d^2 , so its maximum is on one of two faces:

$$\frac{1 + 2y}{1 + y^2} \leq \varphi \quad (d = 0), \quad 1 + \frac{2y}{1 + sy^2} \leq 1 + \frac{1}{\sqrt{s}} \quad (d = y). \quad (136)$$

The first one-variable function is maximized at $y = 1/\varphi$ and the second at $y = 1/\sqrt{s}$, which gives the two displayed constants. The second value dominates the first exactly when $u \leq \varphi + 2$. Binary instances attain it. $\square$

5.3 The unrestricted mixed lower bound

For $\varphi + 2 < u < z_*$, neither the binary family nor the pure cap-free family is tight. The correct adversary puts a capped block before a scaled harmonic core. Two lemmas ensure that optional raw execution and early processing of caps cannot escape it.

The cap-free construction needed below is exactly the scaled harmonic core from Lemma 4.2; no second phase-accounting proof is needed here. We now prepend capped jobs to that reusable core.

Lemma 5.6 (Cap deferral). *Prepend capped mass m with value $p = u$ to a later cap-free block of mass $s = 1 - m$ and total processing mass L . Suppose every later value satisfies $1 + p \leq u$ and*

$$s(u - 1) - L - m \geq 0. \quad (137)$$

Then processing any capped job before all later jobs are complete cannot decrease the leading online cost.

Proof. A cap completed after its test block but before a later job of value p has mutual cost $1 + u$; leaving it pending until after that job gives $2 + p$. The exchange saves $u - 1 - p \geq 0$. If cap mass q is instead completed inside the cap-test block, its pairs with all later jobs add $q[s(u - 1) - L]$. Its largest possible saving against not-yet-tested caps is $qm - q^2/2$. Relative to leaving every cap pending, the change is therefore at least

$$q[s(u - 1) - L - m] + \frac{q^2}{2} \geq 0. \quad (138)$$

The finite diagonal discrepancy is lower order. $\square$

After cap deferral, pairwise accounting gives the offline coefficient O and the guaranteed online lower benchmark A^{lb} :

$$A^{\text{lb}} = A_0^{\text{lb}} + m(1 - m) + m(1 + L) + \frac{u}{2}m^2, \quad (139)$$

$$O = O_0 + m(1 - m + L) + \frac{u}{2}m^2. \quad (140)$$

Indeed, a cap-later pair contributes $2 + p$ online and $1 + p$ offline, whereas a cap-cap pair contributes $2 + u$ online and u offline; collecting these pairs yields the two cross terms and the cap squares above. In particular, $A^{\text{lb}} - O = (A_0^{\text{lb}} - O_0) + m$.

Lemma 5.7 (Raw-safe quota and replay). *Fix $m \in (0, 1)$ and a later cap-free block satisfying the hypotheses of Lemma 5.6. Against every deterministic optional algorithm there are fixed instances whose asymptotic excess is at least the cap-free hybrid benchmark $A^{\text{lb}} - CO$ from Equations (139) and (140) for every target $C \leq u$.*

Proof. In the initial phase, a tested fresh job receives $p = u$, while a job run raw is assigned hidden value zero. If v raw jobs and ℓ tested caps have been touched, stop at the first crossing

$$\frac{\ell}{n - v} \geq m. \quad (141)$$

With $S = n - v$, the overshoot is less than one, so $\ell = mS + O(1)$. If no crossing occurs, no test can have occurred: after all first touches, any positive ℓ would make the ratio in Equation (141) equal one. Thus every job was run raw and the all-zero hidden instance has ratio $u \geq C$.

At a crossing, scale the prescribed later instance to the remaining $S - \ell$ first touches, absorbing the one-job cap overshoot and integer rounding into its zero group. A later test reveals the next value in this fixed type stream; a later raw action consumes the next value but keeps it hidden from the optional algorithm.

Move every initial raw-zero completion to the start. It has the same length u as cap processing, and crossing a unit test only lowers completion cost. With

$$Z_v = vn - \frac{v(v-1)}{2}, \quad (142)$$

this prefix contributes uZ_v online and Z_v offline. Delete it. The remaining S labels consist, up to one-job rounding, of capped fraction m and the prescribed later cap-free block.

If the optional algorithm later runs a fresh assigned value p raw, a physical cap-free simulation tests and immediately processes that job in time $1+p \leq u$, suppresses the answer from the virtual optional algorithm, and idles until its virtual u -block ends. Inductively every physical completion is no later than its virtual optional completion. Since the finite-cap and cap-free offline lengths are both $1+p$ on the later block,

$$\text{ALG}_{\text{opt}} - C \text{OPT}_{\text{opt}} \geq (u - C)Z_v + (\text{ALG}_{\text{hyb}} - C \text{OPT}_{\text{hyb}}) - O_u(n). \quad (143)$$

The first term is nonnegative. Finally, record every assigned value on its job label. Replaying this fixed vector produces the same transcript because the values learned only by the physical simulation were hidden from the virtual algorithm. This proves legality. $\square$

Apply Lemma 4.2 with $s = 1 - m$. Substitution into Equations (139) and (140) gives the guaranteed two-parameter lower ratio

$$\mathcal{R}_u(m, t) = 1 + \frac{m + \frac{(1-m)^2}{2}(1-t^{-2})}{\frac{(1-m)^2}{4}(1+t^{-2} + 2\log t) + m(1-m)(1+\log t) + \frac{u}{2}m^2}. \quad (144)$$

At an interior stationary point with value $1 + c$, differentiation and the zero-gap equation give

$$\log t = u - 2 - \frac{1}{c}, \quad t^2 = \frac{1-m}{1+m} \left(1 + \frac{2}{c}\right). \quad (145)$$

Eliminating t gives precisely Equation (94). Conversely, those equations make $\mathcal{R}_u(m, t) = 1 + c$.

The branch runs from

$$(u, m, t) = \left(\varphi + 2, \frac{1}{\varphi}, 1\right) \quad \text{to} \quad (u, m, t) = (z_*, 0, \sqrt{z_*}). \quad (146)$$

It satisfies $u \geq \varphi + 2$, $\log t < 1$, and $m \leq 1/\varphi$. Therefore

$$(1-m)(u-1-\log t) > (1-m)\varphi \geq \frac{1}{\varphi} \geq m, \quad (147)$$

so Equation (137) holds; also $1+p < 3 < u$ for every later positive value in a sufficiently fine finite construction. The functions

$$g(c) = 1 - \frac{1}{c} + \frac{1}{2} \log \left(1 + \frac{2}{c}\right), \quad h(m) = \text{artanh } m - m \quad (148)$$

have $g'(c) = 2/[c^2(c+2)] > 0$ and $h'(m) = m^2/(1-m^2) > 0$. Hence $m(c) = h^{-1}(g(c))$ is strictly increasing, while

$$u(c) = 1 + \frac{2}{c} - m(c) \quad (149)$$

is strictly decreasing. The identities $g(r) = h(0) = 0$ and $g(1/\varphi) = h(1/\varphi)$ give the two endpoints in Equation (146); consequently every $u \in [\varphi + 2, z_*]$ has exactly one parameter pair.

To pass to fixed finite instances, first approximate (m, t) rationally while retaining the strict cap inequalities, then choose a sufficiently fine harmonic family, and finally multiply all rational block sizes by a common integer H . For fixed harmonic depth, quota overshoot, diagonals, and rounding are $O_u(H)$ whereas the displayed costs are $\Theta(H^2)$. Send first H , then the harmonic depth, and finally the rational approximation to their limits. Together with Lemma 5.7, this proves

$$V(u) \geq C_{\text{mix}}(u) \quad (\varphi + 2 \leq u \leq z_*). \quad (150)$$

The construction above applies directly in the open interval. At $u = \varphi + 2$ and $u = z_*$ the same statement follows by approaching from the interior, so the endpoints are included.

5.4 The large-upper-limit lower bound by cap-free transfer

Lemma 5.8 (Bounded cap-free transfer). *Let $u \geq 1$, and let $V_\infty(P)$ denote the deterministic size-asymptotic value at the cap-free endpoint, restricted to $0 \leq p_j \leq P$. Then*

$$V(u) \geq V_\infty(u - 1). \quad (151)$$

In particular,

$$V(u) \geq R_{\text{obl}} \quad \left(u \geq 1 + \frac{1}{r} = 2.752620747896442 \dots \right). \quad (152)$$

Proof. Given a finite-cap algorithm, construct a cap-free algorithm on instances with $p_j \leq u - 1$ by running a virtual copy of the finite-cap algorithm. Copy ordinary tests and processing actions. When the virtual algorithm runs an untouched job raw for u time, physically test and immediately process it in $1 + p_j \leq u$ time, hide the answer from the virtual copy, and wait for the virtual clock. Every physical completion is no later than its virtual completion. Moreover, $\min\{u, 1 + p_j\} = 1 + p_j$, so finite-cap and cap-free optima coincide. Thus we obtain the following game-value comparison. For every finite-cap algorithm $\mathcal{A}$, the simulation produces a cap-free algorithm $\mathcal{B}$ and, pointwise on every instance with $p_j \leq u - 1$,

$$\text{ALG}_{\mathcal{A}} \geq \text{ALG}_{\mathcal{B}}, \quad \text{OPT}_u = \text{OPT}_\infty.$$

Consequently

$$\text{CR}_\infty(\mathcal{A}) \geq \text{CR}_\infty(\mathcal{B}; p \leq u - 1) \geq V_\infty(u - 1).$$

Taking the infimum over $\mathcal{A}$ proves Equation (151).

The harmonic cap-free family attaining R_{obl} has limiting largest processing time

$$1 + \log(1 + y_*) = 1 + \frac{1}{2} \log z_* = \frac{z_* - 1}{2} = \frac{1}{r}, \quad y_* = \sqrt{z_*} - 1. \quad (153)$$

Approximation from below also gives the endpoint, proving Equation (152). $\square$

5.5 Completion of the curve and consistency at the joins

The lower-bound side of Theorem 5.1 is now complete: Proposition 5.3 supplies the first four branches, Equation (150) the fifth, and Equation (152) the last. The raw algorithm, Lemmas 5.4 and 5.5, supplies the matching upper bounds through $\varphi + 2$. The remaining upper bounds are proved in Section 5.6.

For clarity, all five joins are exact:

$$\begin{aligned}\rho_I(u_\phi) &= u_\phi, \\ \rho_I(u_0) &= 1 + \frac{1}{\sqrt{s_0}} = 1.682327803828019\dots, \\ 1 + \frac{1}{\sqrt{(\varphi+2)-1}} &= \varphi = C_{\text{mix}}(\varphi+2), \\ C_{\text{mix}}(z_*) &= 1 + r = R_{\text{obl}}.\end{aligned}$$

At $u = 1$, both initial branches equal one. Representative values of the new mixed branch are

u	$V(u)$	m	t
3.800000	1.598851588	0.539725635	1.138984621
4.000000	1.583283633	0.428863572	1.330517832
4.071679486	1.579204806	0.381330539	1.412236296
4.300000	1.571692026	0.198387083	1.734651379

Remark 5.9 (Asymptotic scope). All statements above concern the fixed- u size-asymptotic ratio. Below the plateau, the upper bounds have additive error $O_u(n)$. On $u \in [z_*, \infty]$ the same algorithm has one uniform $O(n)$ error. We do not claim a finite-instance endpoint inequality with one additive constant.

The next subsection, Section 5.6, proves the matching middle- and high- u upper bounds and thereby completes the proof of Theorem 5.1.

5.6 Matching upper bound for the common-upper model

We now prove the upper bounds on the two ranges on which capped outcomes matter. Besides completing the upper-bound side of the common-upper curve, the proof records explicitly why an outcome on the optional cap cannot be treated as an ordinary deferred outcome. Throughout this subsection,

$$C = 1 + c, \quad \varphi = \frac{1 + \sqrt{5}}{2},$$

and $z_* > 1$ is the solution of $z_* - 3 = \log z_*$. We also set

$$r = \frac{2}{z_* - 1}.$$

For $c \in [r, 1/\varphi]$, let $m = m(c) \in [0, 1/\varphi]$ be determined by

$$\operatorname{artanh} m - m = 1 - \frac{1}{c} + \frac{1}{2} \log \left(1 + \frac{2}{c} \right), \quad u = 1 + \frac{2}{c} - m. \quad (154)$$

The right-hand side of the first equation is strictly increasing in c , as is the left-hand side in m . Consequently $m(c)$ is increasing and $u(c)$ is decreasing. The endpoints are

$$(c, m, u) = \left(\frac{1}{\varphi}, \frac{1}{\varphi}, \varphi + 2 \right) \quad \text{and} \quad (c, m, u) = (r, 0, z_*).$$

Theorem 5.10. *For every $u \in [\varphi + 2, \infty]$, the formally specified algorithm `ADAPTIVETHRESHOLD` from algorithm 1 satisfies*

$$\text{ALG} \leq \begin{cases} (1 + c) \text{OPT} + O_u(n), & \varphi + 2 \leq u \leq z_*, \quad c \text{ given by (154)}, \\ (1 + r) \text{OPT} + O(n), & u \geq z_*. \end{cases}$$

The constant in the last $O(n)$ is absolute and uniform over the extended interval $u \in [z_, \infty]$. In particular these are upper bounds on the deterministic size-asymptotic competitive ratio.*

Proof.

At $u = \infty$ this is Theorem 4.3, with the same algorithm and the same four-endpoint bank. For the finite-cap calculations below, fix $u < \infty$.

The algorithm. Use `ADAPTIVETHRESHOLD`(J, c) from algorithm 1; on the high interval use $c = r$. For the analysis, immediately before a test let x be the number of jobs not yet tested, N the number of earlier strictly positive outcomes, and D the number of those outcomes deferred to the final tail. Thus

$$y = \frac{D - CN}{x} \leq 0 \tag{155}$$

and the threshold is

$$a_c(y) = \begin{cases} 1, & y \leq -1, \\ 1 + \frac{1}{2} \log\left(\frac{c+2}{c-2y}\right), & -1 \leq y \leq 0. \end{cases} \tag{156}$$

This is the same threshold as Equation (43). On the mixed interval $c \geq r$, and hence

$$a_c(y) \leq a_c(0) \leq a_r(0) = \frac{1}{r} < u - 1.$$

The same inequality holds on the high interval, where we use $c = r$. Thus every outcome in the capped region $[u - 1, u]$ is deferred.

The state y never increases. Indeed, the numerator in (155) changes by 0, $-C$, or $-c$ after a zero, a positive immediate outcome, or a positive deferred outcome, respectively, while x decreases. This monotonicity will allow us to allocate every pair cost when its relevant endpoint becomes known.

Exact pair objective and the five endpoints. Write

$$\bar{p}_i = \min\{p_i, u - 1\}.$$

The clairvoyant optional optimum is SPT on jobs of effective lengths $1 + \bar{p}_i = \min\{1 + p_i, u\}$. For two jobs tested in the order $i < j$, let

$$e_{ij} = \begin{cases} (p_i - p_j)_+, & i \text{ is processed immediately,} \\ 1 + (p_j - p_i)_+, & i \text{ is deferred and } j \text{ is immediate,} \\ 1, & i \text{ and } j \text{ are deferred.} \end{cases} \tag{157}$$

The first line applies whenever i is immediate, regardless of the action taken on j . Direct completion-time accounting gives

$$\text{ALG} = \sum_i (1 + p_i) + \sum_{i < j} (1 + \min\{p_i, p_j\} + e_{ij}), \tag{158}$$

$$\text{OPT} = \sum_i (1 + \bar{p}_i) + \sum_{i < j} (1 + \min\{\bar{p}_i, \bar{p}_j\}). \quad (159)$$

It is therefore natural to introduce

$$\Gamma_n = \sum_{i < j} (e_{ij} + \min\{p_i, p_j\} - C \min\{\bar{p}_i, \bar{p}_j\}). \quad (160)$$

Equations (158)–(160) yield the exact identity

$$\text{ALG} - C \text{OPT} = \Gamma_n - c \binom{n}{2} + \sum_i [(1 + p_i) - C(1 + \bar{p}_i)]. \quad (161)$$

The final sum is nonpositive. Below the cap its i -th summand is $-c(1 + p_i)$; on the cap it is at most $1 - cu \leq 0$. Thus it is enough to control Γ_n .

Fix a symbolic immediate/deferred path. Coordinatewise convexity in (160) reduces a worst outcome vector to the five formal endpoint types

type	outcome	action
Z	0	immediate
E	0^+	immediate
M	$a_c(y)$	immediate
Q	$a_c(y)^+$	deferred
U	u	deferred and capped.

(162)

For an immediate coordinate the two positive endpoints are 0^+ and the live threshold; the genuine zero is kept separate because it does not enter N . A deferred coordinate below $u - 1$ moves to the live threshold or to $u - 1$. On $[u - 1, u]$, the offline contribution is flat and the algorithmic contribution is nondecreasing, so the latter endpoint moves to u . The symbolic categories, and therefore all later threshold states, remain unchanged during each coordinate move. This is the only endpoint reduction used below.

A local pair allocation. Separate the earlier E -outcomes from the substantive outcomes M, Q, U . Let

$$P = \#\{\text{earlier } M, Q, U\}, \quad E = \#\{\text{earlier } E\}, \quad D = \#\{\text{earlier } Q, U\},$$

and let K be the number of earlier U -outcomes. Among the ordinary substantive outcomes, let I, d be the numbers of M, Q -outcomes and let A_I, A_D be the sums of their processing values. Thus $P = I + d + K$ and $D = d + K$.

At a current ordinary endpoint, with $a = a_c(y)$, allocate the relaxed local charges

$$\ell_Z = \ell_E = D, \quad \ell_M = D + a(x + D - CP), \quad \ell_Q = D + a(D - CP). \quad (163)$$

For a capped endpoint the exact charge is instead

$$\begin{aligned} \ell_U &= D + A_D + uK - C(A_I + A_D + (u - 1)K) \\ &= d - CA_I - cA_D + Kh_U, \quad h_U = 1 + u - C(u - 1). \end{aligned} \quad (164)$$

These formulae follow by checking the finitely many ordered endpoint pairs. In particular, an earlier immediate outcome already allocated its online processing delay when it appeared, so a current cap contributes only $-Cp$ against that outcome. Against an earlier ordinary

deferred outcome it contributes $p - Cp = -cp$, and against an earlier cap it contributes h_U . The same check shows

$$\Gamma_n \leq \sum_i \ell_i; \quad (165)$$

the difference is exactly the sum of the nonnegative own-processing terms allocated by the M -endpoints. Formula (164), rather than a substitution $p = u$ into ℓ_Q , is essential before caps have been put into a prefix.

Threshold monotonicity gives $A_I \geq aI$ and $A_D \geq ad$. Comparing (164) with the hypothetical Q -charge at the same state gives the exact identity

$$\begin{aligned} \ell_U - \ell_Q &= C(aI - A_I) + c(ad - A_D) + K[1 - c(u - 1 - a)] \\ &\leq K[1 - c(u - 1 - a)]. \end{aligned} \quad (166)$$

The reusable four-endpoint bank. The next potential is the parameterized form of the cap-free bank from Section 4.2. The states Z, E, M, Q are the same four ordinary endpoints; only their target coefficient changes from r to c . At $c = r$ the formulas below coincide term for term with the obligatory certificate. Keeping the parameter visible lets us add the finite-cap reserve without proving a second ordinary endpoint theorem. Define

$$\eta = \frac{D - CP}{x}, \quad b = \frac{E}{x}, \quad y = \eta - Cb. \quad (167)$$

For $-1 \leq y \leq 0$, put

$$f_c(y) = \frac{c+2}{4} + \frac{c-2y}{4} \left(\log \frac{c-2y}{c+2} - 1 \right), \quad (168)$$

$$h_c(y) = a_c(y)(1+y),$$

$$G_c(y, b) = f_c(y) + bh_c(y) + \frac{C}{2}b^2. \quad (169)$$

For $y \leq -1$, set instead

$$G_c(y, b) = g_c(\eta), \quad g_c(\eta) = \frac{(1+\eta)_+^2}{2C}. \quad (170)$$

The two definitions are C^1 across $y = -1$. Indeed, feasibility implies $0 \leq b \leq 1/C$ there, and both sides have value $Cb^2/2$ and derivatives

$$(G_y, G_b) = (b, Cb).$$

They are also nonnegative: $f_c(-1) = 0$, $f'_c = a_c - 1 \geq 0$, and $h_c, b, g_c \geq 0$.

Use the base potential

$$W_{\text{base}}(x, P, E, D) = xD + x^2 G_c(y, b). \quad (171)$$

In the raw state (x, P, E, D, K) , the endpoint directions are

	x	P	E	D	K
v_Z	-1	0	0	0	0
v_E	-1	0	1	0	0
v_M	-1	1	0	0	0
v_Q	-1	1	0	1	0
v_U	-1	1	0	1	1.

(172)

Let $J_T = \ell_T + \nabla W_{\text{base}} \cdot v_T$, where for the moment $T \in \{Z, E, M, Q\}$. In Region I, direct differentiation and the identities

$$\begin{aligned} -2f_c + (y - C)f'_c + a_c(1 + y) &= 0, \\ 1 - 2f_c + (y - c)f'_c + a_c y &= 0 \end{aligned} \quad (173)$$

give

$$\begin{aligned} \frac{J_Z}{x} &= (-2f_c + yf'_c) + b(-h_c + yh'_c), \\ \frac{J_E}{x} &= b[-h_c + (y - C)h'_c + C], \\ \frac{J_M}{x} &= b[-h_c + (y - C)h'_c + a_c C], \\ \frac{J_Q}{x} &= b[-h_c + (y - c)h'_c + a_c C]. \end{aligned} \quad (174)$$

All four quantities are nonpositive. For the first one,

$$-2f_c + yf'_c = a_c(c - y) - C \leq 0; \quad (175)$$

the right-hand side is zero at $y = -1$ and is nonincreasing because $a'_c(c - y) - a_c \leq 1 - a_c \leq 0$. Also $h_c, h'_c \geq 0$. For the remaining three drifts, observe that

$$h_c + (c - y)h'_c = a_c C + (c - y)(1 + y)a'_c \geq a_c C. \quad (176)$$

The Q -bracket in (174) is therefore nonpositive; the M -bracket is the Q -bracket minus h'_c , and the E -bracket is the M -bracket minus $C(a_c - 1)$.

In Region II, the same calculation gives

$$\begin{aligned} J_Z/x &= J_E/x = -2g_c + \eta g'_c, \\ J_M/x &= -2g_c + (\eta - C)g'_c + 1 + \eta, \\ J_Q/x &= 1 - 2g_c + (\eta - c)g'_c + \eta. \end{aligned} \quad (177)$$

For $-1 \leq \eta \leq 0$, substitution of $g'_c = (1 + \eta)/C$ makes these values $-(1 + \eta)/C$, $-(1 + \eta)/C$, and 0, respectively. For $\eta \leq -1$ their signs are immediate from $g_c = g'_c = 0$. Hence the base bank pays every ordinary endpoint.

The cap reserve. Define

$$Z(c) = 2 + \frac{1}{c} + \frac{1}{2} \log \left(1 + \frac{2}{c} \right), \quad \delta = Z(c) - u. \quad (178)$$

On the mixed curve, (154) says precisely that

$$\delta = \operatorname{artanh} m. \quad (179)$$

For $0 \leq t \leq m$, set

$$\beta(t) = ct[\delta - \operatorname{artanh} t], \quad \mathcal{V}(t) = \int_t^m \beta(s) ds, \quad (180)$$

and extend $\mathcal{V}$ by zero for $t \geq m$. Since $\beta(m) = 0$, this extension is C^1 . With

$$q = \frac{K}{x}, \quad \mu = \frac{q}{1 + q}, \quad \mathcal{H}(q) = (1 + q)^2 \mathcal{V}(\mu), \quad (181)$$

define the full potential

$$W = xD + x^2(G_c(y, b) + \mathcal{H}(q)). \quad (182)$$

It is nonnegative.

For an ordinary endpoint, K is fixed while x decreases, so the reserve contributes, after division by x ,

$$L_0(q) = -2\mathcal{H}(q) + q\mathcal{H}'(q). \quad (183)$$

A cap also increments K , and its contribution is $L_0 + \mathcal{H}'$. On the active range $\mu \leq m$, differentiation of (181) gives

$$\mathcal{H}' = 2(1+q)\mathcal{V} - \beta(\mu), \quad L_0 = -2(1+q)\mathcal{V} - q\beta(\mu) \leq 0. \quad (184)$$

Thus the reserve can only improve every ordinary drift.

For a cap, feasibility $D \leq P$ and $P \geq K$ gives

$$y = \frac{D - C(P + E)}{x} \leq -\frac{cK}{x} = -cq. \quad (185)$$

While $\mu \leq m$, we have $q \leq m/(1-m) \leq \varphi$ and hence $cq \leq 1$. Monotonicity of a_c , together with (166), yields

$$\frac{\ell_U - \ell_Q}{x} \leq cq \left[\delta - \frac{1}{2} \log(1+2q) \right] =: B(q). \quad (186)$$

The change of variables in (181) satisfies

$$\operatorname{artanh} \mu = \frac{1}{2} \log(1+2q), \quad (1+q)\beta(\mu) = B(q). \quad (187)$$

Consequently the cap residual is cancelled exactly:

$$L_0 + \mathcal{H}' + B(q) = 0 \quad (\mu \leq m). \quad (188)$$

When $\mu \geq m$, the reserve vanishes. At $q_* = m/(1-m)$, the bracket in (166), evaluated at the largest possible threshold $a_c(-cq_*)$, is zero by (179) and (187). For $q \geq q_*$, (185) only decreases the threshold, so that bracket is nonpositive. The fifth Hamiltonian inequality therefore holds for every state and for arbitrary interleavings of capped and ordinary outcomes.

Initial calibration. Initially $y = b = q = 0$, and

$$f_c(0) = \frac{1}{2} - \frac{c}{4} \log\left(1 + \frac{2}{c}\right). \quad (189)$$

Elementary integration in (180) gives

$$\mathcal{V}(0) = \frac{c}{2}(\delta - m). \quad (190)$$

Indeed,

$$\int_0^m t \operatorname{artanh} t \, dt = \frac{(m^2 - 1) \operatorname{artanh} m + m}{2}.$$

Using (154) and (179), we obtain the exact calibration

$$f_c(0) + \mathcal{H}(0) = f_c(0) + \mathcal{V}(0) = \frac{c}{2}. \quad (191)$$

Thus $W_{\text{initial}} = cn^2/2$.

Finite jobs, and uniformity on the plateau. We spell out the regularity needed to pass from the fluid Hamiltonian to unit job transitions. It is important here to restrict attention to feasible states. In Region I, $y \geq -1$ implies

$$-xy = C(P + E) - D = cP + (P - D) + CE \leq x. \quad (192)$$

All terms in the middle expression are nonnegative. Hence $P/x, D/x, q \leq 1/c$ and $b \leq 1/C$, so every normalized derivative of the Region-I perspective in (171) is bounded on the feasible cone. In Region II the raw base bank is simply

$$xD + \frac{(x + D - CP)_+^2}{2C}, \quad (193)$$

which has bounded one-sided Hessians. The two pieces are C^1 at $y = -1$, as verified above, and the positive-part square is C^1 at $\eta = -1$.

For the reserve $R(x, K) = x^2 \mathcal{H}(K/x)$, on its active cone $0 \leq q \leq q_*$ one has bounded $\mathcal{H}''$, and

$$R_{xx} = 2\mathcal{H} - 2q\mathcal{H}' + q^2\mathcal{H}'', \quad R_{xK} = \mathcal{H}' - q\mathcal{H}'', \quad R_{KK} = \mathcal{H}''. \quad (194)$$

For $q \geq q_*$, $R = 0$; moreover $\mathcal{H}(q_*) = \mathcal{H}'(q_*) = 0$. Thus the join is C^1 with bounded one-sided Hessians. At $x = 0$, set $R = 0$: for $K > 0$ it is already locally zero, while at $(x, K) = (0, 0)$ two-homogeneity gives zero gradient. Altogether W is $C^{1,1}$ on the feasible state cone. On the mixed interval its regularity constant may depend on the fixed value of u . When $c = r$ and $m = 0$, however, the reserve is identically zero and the base bank depends only on the absolute constant r ; its regularity constant is therefore uniform over all $u \geq z_*$. This includes the last transition from $x = 1$ to $x = 0$.

It follows that a unit transition has the Taylor remainder

$$\ell_i + W_{i+1} - W_i \leq O_u(1). \quad (195)$$

The endpoint $E = 0^+$ is interpreted one-sidedly. For a fixed symbolic word, replace all E -coordinates by positive values ϵ . Its state trajectory is unchanged, and the exact finite pair objective, being a finite sum of affine and minimum terms, converges as $\epsilon \downarrow 0$. Equivalently, retain an $O(\epsilon n^2)$ error and take this limit for each fixed n . Therefore the formal E -Hamiltonian transfers to genuine outcome vectors without an asymptotic loss.

Summing (195), using $W \geq 0$, and applying (191) gives

$$\sum_i \ell_i \leq \frac{c}{2} n^2 + O_u(n). \quad (196)$$

Together with (161) and (165), this yields

$$\text{ALG} - C \text{OPT} \leq -c \binom{n}{2} + \frac{c}{2} n^2 + O_u(n) = O_u(n). \quad (197)$$

At $u = z_*$, one has $c = r$, $m = 0$, and $Z(r) = z_*$. The cap reserve is identically zero, while (191) remains valid. If u is increased above z_* , the only adverse fifth-endpoint bracket $1 - r(u - 1 - a_r(y))$ decreases, and the capped diagonal term in (161) also decreases.

Here is the promised uniform finite calculation. The base bank and its feasible cone contain no u , so there is one constant κ , depending only on r , for which every ordinary or capped transition with $u \geq z_*$ satisfies

$$\ell_i + W_{i+1} - W_i \leq \kappa.$$

Because $W_0 = rn^2/2$ and $W_n = 0$, telescoping gives

$$\sum_i \ell_i \leq \frac{r}{2}n^2 + \kappa n. \quad (198)$$

In the diagonal term of Equation (161), a value below the cap contributes $-r(1 + p_i)$. A capped value contributes at most $1 - ru \leq 0$. With the convention $\bar{p}_i = p_i$ at $u = \infty$, the first formula applies there and the capped endpoint is absent. Hence the diagonal sum is nonpositive uniformly on $[z_*, \infty]$. Combining Equations (161), (165) and (198) now yields

$$\text{ALG} - (1 + r) \text{OPT} \leq -r \binom{n}{2} + \frac{r}{2}n^2 + \kappa n = \left(\kappa + \frac{r}{2}\right)n. \quad (199)$$

Thus, with $B = \kappa + r/2$,

$$\text{ALG} \leq (1 + r) \text{OPT} + Bn \quad \text{for every } u \in [z_*, \infty], \quad (200)$$

and B is independent of u . The estimate is genuinely uniform rather than the result of sending a parameter-dependent remainder to infinity. This proves Theorem 5.10. Finally, because $u > 1$ on the present range, every effective offline length is at least one and $\text{OPT} \geq n(n + 1)/2$. The additive term in (197) vanishes in the size-asymptotic ratio. $\square$

6 Randomized common upper bounds: the exact four-piece curve

We return to the finite common-upper model and allow private randomization against an oblivious adversary. The cap u is fixed throughout this section, but the algorithm is not told the processing-time multiset. This section proves Theorem 2.3. The proof has three parts. First we analyze two policies for an announced distribution: run everything raw, or test everything using a maximum-density stationary threshold. A survival-function argument shows that their worst distribution is binary. We then learn the better policy from a sublinear pilot, and finally use random labellings and Yao's principle for the matching lower bounds.

6.1 Offline and stationary fluid coefficients

Assume initially that $u > 1$ and write

$$s = u - 1. \quad (201)$$

Let D be a finite distribution on $[0, u]$, let P, P' be independent with law D , and use the survival function

$$H(t) = \Pr(P \geq t), \quad H(t) = 0 \quad (t > u). \quad (202)$$

The convention at atoms does not affect any integral below. Since a clairvoyant job has effective length $1 + \min\{s, P\}$, the layer-cake identity and Lemma 3.1 give the leading offline coefficient

$$\Omega_u(D) := \frac{1}{2} \mathbb{E} \min\{q_u(P), q_u(P')\} = \frac{1}{2} \left(1 + \int_0^s H(t)^2 dt\right), \quad (203)$$

$$\text{OPT}_u(M) = n^2 \Omega_u(D_M) + O_u(n) \quad (204)$$

for every empirical distribution D_M .

We next recall the maximum-density stationary policy in a form suited to the cap. For a processing-time prefix of mass a and first moment m , its completion density is $a/(1+m)$. Choose a maximum-density prefix, include all zeros, and put

$$\tau = \frac{1+m}{a}. \quad (205)$$

The one-point addition/removal test puts every value below τ in the prefix and every value above τ outside it; an atom at equality may be split. Equivalently,

$$\mathbb{E}(\tau - P)^+ = 1, \quad \int_0^\tau H(t) dt = \tau - 1. \quad (206)$$

The equation has a unique solution $\tau \geq 1$.

The stationary policy tests all jobs in a uniform random order, processes a positive prefix outcome immediately after its test, and after the last test processes the residual jobs in SPT order. The following formula is the finite-cap analogue of Lemma 7.1.

Lemma 6.1 (Stationary survival formula). *The doubled leading coefficient of the stationary policy is*

$$\mathcal{T}_u(D) = \begin{cases} \tau + \int_\tau^u H(t)^2 dt, & \tau \leq u, \\ \tau, & \tau > u. \end{cases} \quad (207)$$

Consequently the better of the raw and stationary policies has leading ratio

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s H(t)^2 dt}. \quad (208)$$

Proof. Let $h = 1 - a$ be the residual mass and let K_L be the minimum kernel of two residual draws, with their subprobability masses left unnormalized. The random-order pair calculation gives the stationary coefficient

$$(1+m) \left(1 - \frac{a}{2}\right) + \frac{K_L}{2}. \quad (209)$$

If $\tau \leq u$, the threshold property gives

$$K_L = \tau h^2 + \int_\tau^u H(t)^2 dt, \quad (1+m) \left(1 - \frac{a}{2}\right) = \frac{\tau}{2}(1 - h^2).$$

Their sum is one half of the first line of (207). If $\tau > u$, the whole distribution is in the prefix and (209) equals $\tau/2$. Raw execution has leading coefficient $u/2$, and division by (203) proves (208). $\square$

6.2 Flattening the survival function

We now maximize (208) over arbitrary D . Suppose first that $1 \leq \tau \leq u$ and put

$$y = \frac{\tau - 1}{\tau}. \quad (210)$$

By (206), y is the average of H on $[0, \tau]$. Since H is nonincreasing, its average on every initial subinterval is at least y . Thus Cauchy–Schwarz gives

$$\int_0^a H(t)^2 dt \geq ay^2 \quad (0 \leq a \leq \tau), \quad (211)$$

and $H(t) \leq y$ almost everywhere for $t > \tau$.

Replace H on $(0, \tau)$ by the constant y and leave its tail unchanged. This remains a nonincreasing survival function, preserves (206) and $\mathcal{T}_u(D)$, and by (211) can only decrease the offline coefficient. Hence it can only increase the ratio. We may therefore assume

$$H(t) = y \quad (0 < t < \tau). \quad (212)$$

If $u - 1 \leq \tau \leq u$, the offline integral lies entirely in the flat part, and $H(t)^2 \leq y^2$ on the remaining tail. It follows that

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s H^2} \leq B_u(\tau) := \frac{\tau + (u - \tau)y^2}{1 + (u - 1)y^2}. \quad (213)$$

The numerator displayed on the right is at most u , so the raw minimum is inactive.

Now suppose $\tau < u - 1$. Put

$$L = u - 1 - \tau, \quad E = \int_{\tau}^{u-1} H(t)^2 dt, \quad G = \int_{u-1}^u H(t)^2 dt. \quad (214)$$

Monotonicity gives

$$0 \leq E \leq Ly^2, \quad G \leq E/L. \quad (215)$$

Consequently

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s H^2} \leq \frac{\tau + \frac{u - \tau}{u - 1 - \tau} E}{1 + \tau y^2 + E}. \quad (216)$$

For fixed τ this is linear-fractional in E , so its derivative has a constant sign. Its maximum on $[0, Ly^2]$ occurs at an endpoint. The two endpoint values are

$$A(\tau) := \frac{\tau}{1 + \tau y^2} = \frac{\tau^2}{\tau^2 - \tau + 1}, \quad (217)$$

$$B_u(\tau) = \frac{\tau + (u - \tau)y^2}{1 + (u - 1)y^2}. \quad (218)$$

They are attained by rectangular survival functions: respectively a binary distribution on $\{0, \tau\}$ and one on $\{0, u\}$.

It remains to cover $\tau > u$. Equation (206) and the zero extension of H give $\int_0^u H = \tau - 1 \geq u - 1$. The average of a nonincreasing function on $[0, u - 1]$ is no smaller than its average on $[0, u]$, and hence

$$\int_0^{u-1} H(t)^2 dt \geq (u - 1) \left(\frac{\tau - 1}{u} \right)^2 \geq \frac{(u - 1)^3}{u^2}. \quad (219)$$

Here the raw policy is chosen, and its ratio is at most $B_u(u)$. We have proved the reduction

$$\sup_D \frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s H(t)^2 dt} \leq \max \left\{ \sup_{1 \leq \tau \leq u-1} A(\tau), \sup_{1 \leq \tau \leq u} B_u(\tau) \right\}, \quad (220)$$

where the first supremum is omitted for $u < 2$. In fact equality holds in (220), because the rectangular survival functions described above attain both families.

6.3 The two scalar maximizations

The first family satisfies

$$A'(\tau) = \frac{\tau(2-\tau)}{(\tau^2 - \tau + 1)^2}. \quad (221)$$

Thus its maximum is attained at $\tau = \min\{2, u-1\}$ and equals $4/3$ as soon as $u \geq 3$.

For the second family, substitution of (210) gives

$$B_u(\tau) = \frac{\tau^2 u + 2\tau^2 - 2\tau u - \tau + u}{\tau^2 u - 2\tau u + 2\tau + u - 1}. \quad (222)$$

The sign of its derivative is the sign of

$$g_u(\tau) = (2\tau - 1)^2 - u(\tau - 1)^2. \quad (223)$$

For $u \leq 4$ this is positive on $[1, \infty)$. For $u > 4$, its unique root above one is

$$\tau_*(u) = \frac{u-2+\sqrt{u}}{u-4} = \frac{\sqrt{u}-1}{\sqrt{u}-2}, \quad (224)$$

and B_u increases before this point and decreases afterwards. The identity $\tau_*(u) = u$ is equivalent to

$$u^3 - 6u^2 + 5u - 1 = 0. \quad (225)$$

The cubic is strictly increasing beyond its last critical point and changes sign there exactly once; its unique root above one is u_r from (5). Hence

$$\max_{1 \leq \tau \leq u} B_u(\tau) = B_u(u) = \frac{u^3}{u^3 - 2u^2 + 3u - 1} = \frac{u^3}{u^2 + (u-1)^3}, \quad u \leq u_r, \quad (226)$$

$$= B_u(\tau_*(u)) = 1 + \frac{1}{2(\sqrt{u}-1)}, \quad u \geq u_r. \quad (227)$$

There is no missing branch from A . On $2 \leq u \leq 3$, clearing the positive denominators in $B_u(u) - A(u-1)$ and putting $v = u-2$ leaves

$$v^4 + 3v^3 + 3v^2 + 3v + 3 > 0.$$

On $3 \leq u \leq u_r$, one has $B_u(u) > 4/3$. On $u_r \leq u \leq 25/4$, (227) is at least $4/3$, with equality exactly at $u = 25/4$; afterwards $A(2) = 4/3$ is larger. Combining these observations with (220) proves the announced upper bound

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s H(t)^2 dt} \leq R_{\text{rand}}(u) \quad (228)$$

for every finite distribution D .

The junction statements now follow directly. Equations (225) to (227) give continuity at u_r , and the square-root branch equals $4/3$ at $u = 25/4$. Differentiating (226) shows that its only interior maximum above one occurs at $u = (3 + \sqrt{3})/2$; substitution gives (9).

6.4 Learning the announced policy

The policy used for the upper bound must not know D . The next lemma removes this announcement at lower-order cost.

Lemma 6.2 (Sublinear-pilot transfer). *For every fixed $u > 1$, there are randomized nonanticipating policies $\mathcal{A}_{n,u}$ such that every fixed labelled input with empirical distribution D satisfies*

$$\mathbb{E} \text{ALG}_{\mathcal{A}_{n,u}} \leq \frac{n^2}{2} \min\{u, \mathcal{T}_u(D)\} + O_u(n^{11/6}). \quad (229)$$

Proof. Let $k = \lfloor n^{2/3} \rfloor$, $d = \lfloor n^{1/3} \rfloor$, and $\eta = u/d$. Keep zero as a separate category and round each positive value upward to one of the d grid endpoints. The algorithm privately permutes the labels, tests the first k , and forms the sample histogram. On that histogram it compares the raw template with every stationary grid-prefix template. It applies a minimizing template to the remaining labels; in the stationary case, prefix outcomes are processed immediately and the residual tail in SPT order. Sampled positive jobs are inserted in the corresponding known part of the schedule.

Let $\hat{D}$ and D^g be the sample and population grid histograms. Hypergeometric variance and Cauchy–Schwarz over the $d + 1$ categories give

$$\mathbb{E} \|\hat{D} - D^g\|_1 \leq \sqrt{\frac{d+1}{k}} = O(n^{-1/6}). \quad (230)$$

For a fixed prefix, the stationary coefficient is a quadratic polynomial in the category masses, with coefficients bounded in terms of u . It is therefore $O_u(1)$ -Lipschitz in total variation, uniformly over all grid prefixes. The empirical-minimization sandwich loses at most twice the right-hand side of (230). Zero-preserving upward rounding changes every normalized work and pair kernel by $O_u(1/d)$.

Finally, placing the sample before the main random-order template affects at most $O(nk + k^2)$ bounded operation–job pairs. The already tested sample has the same bound when raw is selected. After normalization by n^2 , the total error is

$$O_u \left(\sqrt{d/k} + 1/d + k/n \right) = O_u(n^{-1/6}).$$

This proves (229). The initial private permutation makes the law identical for every fixed public labelling. $\square$

Together with (204) and (228), Lemma 6.2 yields

$$\mathbb{E} \text{ALG}_{\mathcal{A}_{n,u}} \leq R_{\text{rand}}(u) \text{OPT}_u + O_u(n^{11/6}), \quad (231)$$

which is (7) with $\varepsilon_u(n) = O_u(n^{-1/6})$. For $0 < u \leq 1$, raw execution is exactly clairvoyantly optimal, so no learning is required.

6.5 Binary oblivious lower bounds

It remains to show that the two endpoint policies used above cannot be improved in the worst case by a more adaptive algorithm. We first record a binary random-labelling envelope.

Lemma 6.3 (Binary cap envelope). *Fix $u > 1$, $x \in [0, 1]$, and put $z = 1 - x$. Let M_n have $xn + O(1)$ jobs of value u and $zn + O(1)$ zero jobs. For every randomized nonanticipating algorithm, even one told M_n , some fixed labelling satisfies*

$$\frac{\mathbb{E} \text{ALG}}{n^2} \geq \begin{cases} \frac{u}{2} - o_u(1), & x \geq (u-1)/u, \\ \frac{1+x+ux^2}{2} - o_u(1), & x \leq (u-1)/u. \end{cases} \quad (232)$$

Every labelling has

$$\frac{\text{OPT}_u}{n^2} = \frac{1 + (u-1)x^2}{2} + O_u(1/n). \quad (233)$$

Proof. First fix a deterministic policy and place the occurrences uniformly on the public labels. List labels in the order in which the policy first touches them, by either a test or a raw execution, and expose analytically also the hidden value of a raw-touched occurrence. The map from placements to this occurrence word is a bijection: equal word prefixes give equal observable histories, hence the same next label and touch kind, and induction recovers the whole placement. The word is therefore a uniform permutation, and the test/raw choice before its next entry is predictable from the preceding word.

Take $\delta_n = n^{-1/4}$. Uniform-prefix hypergeometric concentration, followed by maximal Azuma–Hoeffding for the predictably selected tests, gives one event of probability $1 - o(1)$ on which every first-touch prefix ending with at least $\delta_n n$ untouched labels and containing t tests satisfies

$$|N_0(t) - zt| \leq \Delta_n, \quad \Delta_n = O(\delta_n^{-1} \sqrt{n \log n}) = o(n). \quad (234)$$

Here $N_0(t)$ is the number of tested zeros. The event is simultaneous, so we do not condition on the policy’s adaptive stopping time.

Edit the suffix by running raw every still-untouched job that the original schedule would later test, deleting its later known-processing operation. Only $\delta_n n + O(1)$ jobs change, so the edit changes total completion cost by $o_u(n^2)$. Let q be the final tested fraction of the edited schedule. On (234), every prefix of its completion curve is dominated, up to $o(1)$ vertical error, by the divisible items

item	completion capacity	work per completion
zero-test module	zq	$1/z$
tested cap processing	xq	u
raw jobs	$1 - q$	u

When $z = 0$, the zero-test module is simply omitted. Indeed, t tests complete at most $zt + \Delta_n$ zeros and expose at most $xt + \Delta_n$ cap jobs; raw and known cap processing both use u work per completion. The ordinary divisible-knapsack exchange therefore upper-bounds completed mass at every work value. Integrating the remaining mass gives a lower bound on total completion time.

If $z \leq 1/u$, every displayed item costs at least u per completion, so the normalized area is at least $u/2 - o_u(1)$. If $z \geq 1/u$, the zero-test module comes first. For fixed q its remaining-mass area is

$$F_z(q) = q - \frac{zq^2}{2} + \frac{u}{2}(1 - zq)^2, \quad F'_z(q) = (1 - zq)(1 - uz). \quad (235)$$

It is minimized at $q = 1$, giving $F_z(1) = (1 + x + ux^2)/2$. The suffix edit, vertical repair, and bad event all contribute $o_u(n^2)$.

The argument holds for each deterministic policy. For a randomized policy, fix its seed, average the result over seeds and the finite uniform placement space, and select a placement whose seed-expected cost is at least the joint average. This placement is fixed before the seed and is therefore an oblivious input. Finally, the effective lengths of the two job types are 1 and u , so (233) follows from Lemma 3.1. $\square$

We now select the binary masses. If $1 < u \leq u_r$, take

$$x = \frac{u-1}{u}. \quad (236)$$

This is the boundary $z = 1/u$ in Lemma 6.3, and its online and offline coefficients have ratio

$$\frac{u}{1 + (u-1)((u-1)/u)^2} = \frac{u^3}{u^2 + (u-1)^3}. \quad (237)$$

If $u_r \leq u \leq 25/4$, take

$$x = \frac{1}{\sqrt{u}-1}. \quad (238)$$

The definition of u_r is precisely the point from which $x \leq (u-1)/u$. Since $(u-1)x^2 = 1+2x$, the second line of (232) divided by (233) becomes

$$1 + \frac{x}{2} = 1 + \frac{1}{2(\sqrt{u}-1)}. \quad (239)$$

Irrational masses are approximated by empirical rational frequencies; all rounding and diagonal terms are $O_u(n)$.

For the last branch, the cap distribution is replaced by a fixed bounded one.

Lemma 6.4 (Balanced zero–two lower bound). *For every $u \geq 25/4$ and every randomized algorithm, arbitrarily large even n admit a fixed labelling with $n/2$ jobs of value zero and $n/2$ jobs of value two such that*

$$\mathbb{E} \text{ALG} \geq n^2 - o_u(n^2), \quad \text{OPT}_u = \frac{3}{4}n^2 + O(n). \quad (240)$$

Proof. Use the same trace bijection, predictable-urn event, and suffix edit as in Lemma 6.3. For a realized tested fraction q , the three fluid items now have capacities and costs

item	completion capacity	work per completion
zero-test module	$q/2$	2
tested two processing	$q/2$	2
raw jobs	$1 - q$	u

Every item costs at least two per completion. Thus the completion envelope lies below $\min\{1, t/2\}$ independently of q , whose remaining-mass area is one. This proves the online estimate after the $o_u(n^2)$ repairs and the finite Yao selection. Since $u > 3$, the two effective lengths are one and three; their equally weighted minimum kernel gives offline coefficient $3/4$. $\square$

Equations (237) and (239), together with Lemma 6.4, match (231) in every range. For $0 < u \leq 1$, raw execution is both an online and a clairvoyant optimum. This completes the proof of the exact curve in Theorem 2.3.

7 Randomized obligatory testing: the sharp ratio is $4/3$

We now keep obligatory tests but change the adversary. It fixes the labelled vector $p \in [0, \infty)^n$ before the algorithm draws its random bits. The multiset is not announced and no upper bound on the processing times is assumed. This section proves Theorem 2.4.

7.1 The normalized offline objective

For a fixed vector p , put

$$K(p) = \frac{1}{n^2} \sum_{i,j} \min\{p_i, p_j\}, \quad O(p) = \frac{1 + K(p)}{2}, \quad (241)$$

$$D(p) = \frac{n + \sum_i p_i}{2}. \quad (242)$$

The pair identity Equation (20) is equivalently

$$\text{OPT}(p) = n^2 O(p) + D(p). \quad (243)$$

The linear correction $D(p)$ will be kept rather than hidden: this makes the analysis valid for unbounded processing times.

7.2 Stationary prefixes

A set E is a *processing-time prefix* if every value in E is at most every value outside it. We require that E contain every zero job; otherwise a supposedly deferred zero completes at its test and the following identity is false. Write $L = [n] \setminus E$ and

$$e = |E|, \quad a = \frac{e}{n}, \quad h = 1 - a, \quad m = \frac{1}{n} \sum_{i \in E} p_i, \quad K_X = \frac{1}{n^2} \sum_{i,j \in X} \min\{p_i, p_j\}. \quad (244)$$

Every early-late pair has minimum equal to its early value, so

$$K(p) = K_E + 2hm + K_L. \quad (245)$$

Define

$$P(p, E) = (1 + m) \left(1 - \frac{a}{2}\right) + \frac{K_L}{2}. \quad (246)$$

The stationary policy for E tests all jobs in a uniformly random order, processes a positive early job immediately after its test, and defers every late job. After the final test it processes the late jobs in SPT order.

Lemma 7.1 (Exact stationary cost). *For every processing-time prefix E containing all zero jobs,*

$$\mathbb{E} \text{ALG}_{\text{stat}}(p, E) = n^2 P(p, E) + \frac{e + \sum_i p_i}{2}. \quad (247)$$

Proof. Let

$$W = n + \sum_{i \in E} p_i, \quad L_E = e + \sum_{i \in E} p_i.$$

The discovery phase consists of n blocks: an early block has length $1 + p_i$ and completes its job, while a late block is its unit test and completes no job. For an early job, each other

block precedes it with probability $1/2$. Its expected completion time is $W/2 + (1 + p_i)/2$, and the total early contribution is

$$\frac{eW + L_E}{2}. \quad (248)$$

Every late job sees the common offset W . Its tail contributes

$$(n - e)W + \sum_{i \in L} p_i + \sum_{\substack{i < j \\ i, j \in L}} \min\{p_i, p_j\}. \quad (249)$$

Adding Equations (248) and (249) and collecting diagonal terms gives Equation (247). $\square$

For a nonempty prefix define its completion density by

$$\rho(E) = \frac{a}{1 + m}. \quad (250)$$

Choose a maximum-density prefix and put

$$\theta = \frac{1 + m}{a} = \frac{1}{\rho(E)}. \quad (251)$$

All zeros belong to every maximizer: adding a zero increases the numerator without increasing the denominator.

Lemma 7.2 (Density threshold). *There is a maximum-density prefix for which every early value is at most θ and every late value is at least θ . Values equal to θ may be assigned to either side.*

Proof. Removing boundary mass δ at value x changes the density to $(a - \delta)/(1 + m - x\delta)$. Cross multiplication shows that this is larger exactly when $x > \theta$, so no selected boundary value can exceed θ . The symmetric addition argument excludes an unselected value below θ . Equality preserves the density. $\square$

The next identity is the algebraic source of the constant $4/3$.

Lemma 7.3 (Exact and robust certificates). *For every ordered prefix split with $a > 0$ and $\theta = (1 + m)/a$,*

$$2\theta(4O - 3P) = (\theta - 2)^2 + 4(\theta K_E - m^2) + \theta(K_L - \theta h^2). \quad (252)$$

Consequently:

- (i) *if early values are at most θ and late values are at least θ , then $P \leq 4O/3$;*
- (ii) *if early values are at most $\theta + s$ and late values are at least $\theta - s$, where $s \geq 0$, then*

$$P \leq \frac{4}{3}O + \frac{2}{3}s. \quad (253)$$

Proof. Expansion using $m = a\theta - 1$, $h = 1 - a$, and Equation (245) gives Equation (252). Under exact separation, $xy \leq \theta \min\{x, y\}$ for early x, y , hence $m^2 \leq \theta K_E$; for late x, y , one has $\min\{x, y\} \geq \theta$, hence $K_L \geq \theta h^2$. This proves (i).

Under the relaxed inequalities,

$$\theta K_E - m^2 \geq -sK_E, \quad K_L - \theta h^2 \geq -sh^2.$$

Moreover $K_E \leq am \leq a^2\theta$. Substitution into Equation (252) yields

$$P \leq \frac{4}{3}O + \frac{2}{3}sa^2 + \frac{1}{6}sh^2 \leq \frac{4}{3}O + \frac{2}{3}s,$$

because $(2/3)a^2 + (1/6)(1 - a)^2 \leq 2/3$ on $[0, 1]$. $\square$

Combining the exact certificate with Lemma 7.1 and Equation (243) already gives a finite statement useful below.

Corollary 7.4 (Announced multiset). *If the multiset is announced, the stationary policy for a maximum-density prefix satisfies*

$$\mathbb{E} \text{ALG}_{\text{stat}}(p, E) \leq \frac{4}{3} \text{OPT}(p).$$

Indeed, the correction in Equation (247) is at most $D(p)$. Equality in the quadratic coefficient of Equation (252) forces $\theta = 2$, zero mass $1/2$, and all positive mass at two. This predicts the lower-bound instance before any probabilistic argument is used.

We shall also use the following tail-safe estimate. Direct expansion gives

$$P - O = \frac{h + am - K_E}{2}. \quad (254)$$

If all early values are at most $R \geq 1$, then $m \leq aR$ and therefore

$$P(p, E) \leq O(p) + \frac{R}{2}. \quad (255)$$

7.3 Learning the prefix from a sublinear sample

Fix the universal cutoff

$$B = 32. \quad (256)$$

For an integer d , let $\eta = B/d$. Keep zero as a separate category, partition $(0, B]$ into the consecutive bins $((b-1)\eta, b\eta]$, and add one overflow category for values above B . A finite positive value is rounded upward to the endpoint $q_b = b\eta$.

Let S be a uniformly random subset of k labels. If μ_b and $\hat{\mu}_b$ are the population and sample fractions of category b , put

$$\Delta = \sum_b |\hat{\mu}_b - \mu_b|. \quad (257)$$

Lemma 7.5 (Histogram error). *For $1 \leq k < n$,*

$$\mathbb{E}\Delta \leq \sqrt{\frac{d+2}{k}}.$$

Proof. The count in category b is hypergeometric and its normalized variance is at most μ_b/k . Thus $\mathbb{E}|\hat{\mu}_b - \mu_b| \leq \sqrt{\mu_b/k}$. Summation and Cauchy-Schwarz give the claim. $\square$

For a finite category prefix J , define

$$\hat{a}(J) = \sum_{b \in J} \hat{\mu}_b, \quad \hat{m}(J) = \sum_{b \in J} q_b \hat{\mu}_b, \quad \hat{\rho}(J) = \frac{\hat{a}(J)}{1 + \hat{m}(J)}.$$

Let $\hat{\rho}$ be the maximum and $\hat{\theta} = 1/\hat{\rho}$. The learned early set is not an arbitrary maximizing category prefix. It is its *threshold closure*

$$\hat{J} = \{\text{finite categories } b : q_b \leq \hat{\theta}\}. \quad (258)$$

Lemma 7.6 (Threshold closure). *The prefix Equation (258) has the same maximum sample density. Every category in it has endpoint at most $\hat{\theta}$, and every finite category outside it has endpoint greater than $\hat{\theta}$, including categories with zero sample count.*

Proof. The removal/addition calculation in Lemma 7.2 applies to every positive-mass sample category. Empty categories do not change the sample statistics, and adding mass at equality preserves the identity $1 + \hat{m} = \hat{a}\hat{\theta}$. Closing the prefix therefore preserves optimality and makes the two pointwise assertions tautological. $\square$

For large n , set

$$k = \lfloor n^{3/4} \rfloor, \quad d = \lfloor n^{1/4} \rfloor, \quad \eta = B/d. \quad (259)$$

The algorithm draws a uniform permutation and first tests its first k jobs, leaving positive sample jobs pending. It computes the closed sample prefix. If

$$\hat{\rho} < \frac{2}{B}, \quad (260)$$

it enters *fallback mode*: test every remaining job and then process all positive jobs in SPT order. Otherwise it enters *learned mode*: process positive sampled early jobs in SPT order; test the remaining jobs in their random order, processing early outcomes immediately; and finish the deferred tail in SPT order. Overflow is never early.

7.4 Good samples, fallback, and bad-event excess

Put

$$\gamma = \frac{1}{B(B+1)}, \quad G = \{\Delta \leq \gamma\}. \quad (261)$$

Markov's inequality and Lemma 7.5 give

$$\mathbb{P}(G^c) \leq B(B+1) \sqrt{\frac{d+2}{k}}. \quad (262)$$

Assume G and learned mode. Let E be the population set selected by the closed category prefix, and let $a, m, \theta = (1+m)/a$ be its true statistics. Since $\hat{\theta} \leq B/2$ and $\hat{a} = (1+\hat{m})/\hat{\theta} \geq 2/B$, we have $a \geq 1/B$. For this fixed category set,

$$|a - \hat{a}| \leq \Delta, \quad |m - \hat{m}| \leq B\Delta + \eta.$$

Consequently

$$|\theta - \hat{\theta}| \leq \frac{3}{2}B^2\Delta + B\eta. \quad (263)$$

Every early actual value is at most $\hat{\theta}$, while every late value is greater than $\hat{\theta} - \eta$. Thus the robust certificate applies with

$$s = \frac{3}{2}B^2\Delta + B\eta + \eta. \quad (264)$$

The sample-first order is not exactly the fully stationary comparator, but the difference is only lower order. Conditional on the unordered sample, write e_S, e_R for the early counts and

$$W_S = k + \sum_{i \in E \cap S} p_i, \quad W_R = n - k + \sum_{i \in E \setminus S} p_i.$$

Comparing the two independent random internal orders with one fully random order, the early-cost difference is

$$\frac{e_R W_S - e_S W_R}{2} \leq \frac{1}{2}nk(1 + B/2).$$

Postponing positive sampled early jobs until all sample tests finish adds at most $k^2(1+B/2)$. The late tail is identical because every zero is early. Therefore, on a good learned sample,

$$\mathbb{E}[\text{ALG} \mid S] \leq \frac{4}{3} \text{OPT} + n^2 \left[B^2 \Delta + \frac{2}{3}(B+1)\eta \right] + (1+B/2)(nk/2 + k^2). \quad (265)$$

On a good fallback sample, comparison of sample and population densities shows that the true maximum-density reciprocal satisfies $\theta_* \geq B/4 = 8$. The exact threshold inequalities imply

$$K(p) \geq \frac{(\theta_* - 1)^2}{\theta_*}, \quad O(p) \geq \frac{\theta_* - 1 + 1/\theta_*}{2} \geq \frac{57}{16}. \quad (266)$$

For every test order, test-all-then-SPT satisfies

$$\text{ALG}_{\text{fb}} \leq \text{OPT} + \frac{n(n-1)}{2}. \quad (267)$$

Indeed, if there are z zeros, their total displacement from the first z SPT positions is at most $zn - z(z+1)/2$, while delaying the positive SPT tail adds $(n-z)(n-z-1)/2$; the sum is $n(n-1)/2$. Combining Equations (266) and (267) gives $\text{ALG}_{\text{fb}} < 4 \text{OPT} / 3$.

Because processing times are unbounded, a bad-event estimate must control excess over $4 \text{OPT} / 3$, not raw cost. In learned mode every early value is at most $B/2$, so Equation (255) bounds the stationary excess by $(B/4)n^2$; in fallback mode Equation (267) bounds it by $n^2/2$. Averaging Equation (265) and these bad-event bounds, and using Equation (262), yields the explicit uniform estimate

$$\mathbb{E} \text{ALG} \leq \frac{4}{3} \text{OPT} + n^2 \left[9472 \sqrt{\frac{d+2}{k}} + \frac{704}{d} \right] + 17(nk/2 + k^2). \quad (268)$$

For $n \geq 12^4$, the floors in Equation (259) give $\sqrt{(d+2)/k} \leq 2n^{-1/4}$, $1/d \leq 2n^{-1/4}$, and $k \leq n^{3/4}$. The right-hand side of Equation (268) is at most

$$\frac{4}{3} \text{OPT} + 20378n^{7/4}. \quad (269)$$

For smaller n , test-all-then-SPT is covered by the same deliberately loose constant. Finally $\text{OPT} \geq n^2/2$ proves the multiplicative form in Equation (10).

7.5 The balanced binary lower bound

Let $n = 2m$ and assign m zeros and m twos uniformly to the labels. Fix a deterministic nonanticipating algorithm. In its adaptive order of new tests, let $X_r = 1$ for a zero and $X_r = -1$ for a two, and $S_k = \sum_{r \leq k} X_r$. Conditional on every history, the untested labels remain a uniform subset, so (S_k) is a uniform balanced sign bridge even though the test order is adaptive. Put

$$\Delta_0 = \frac{1}{2} \max_{0 \leq k \leq n} |S_k|. \quad (270)$$

Lemma 7.7 (Bridge discrepancy).

$$\mathbb{E} \Delta_0 \leq \frac{\sqrt{2} n}{\sqrt{n-1}}.$$

Proof. For $0 \leq k \leq m$, $M_k = S_k/(n-k)$ is a martingale. Doob's L^2 inequality and the hypergeometric variance $\mathbb{E}S_m^2 = n^2/[4(n-1)]$ give

$$\mathbb{E} \max_{k \leq m} |S_k|^2 \leq \frac{4n^2}{n-1}.$$

Time reversal gives the same bound for the second half, and Cauchy–Schwarz then yields $\mathbb{E} \max_k |S_k| \leq 2\sqrt{2}n/\sqrt{n-1}$. $\square$

At physical time t , suppose the algorithm has completed k tests and x processing operations of two-jobs. At most $k/2 + \Delta_0 + x$ jobs are complete, while $k + 2x \leq t$. Pathwise,

$$\#\text{completed by } t \leq t/2 + \Delta_0.$$

The area under the unfinished-job curve therefore gives

$$\text{ALG} \geq \int_0^{2(n-\Delta_0)} (n - \Delta_0 - t/2) dt = (n - \Delta_0)^2. \quad (271)$$

By Lemma 7.7, $\mathbb{E} \text{ALG} \geq n^2 - O(n^{3/2})$. Every input in the distribution has

$$\text{OPT} = 3m^2 + 2m = \frac{3}{4}n^2 + n.$$

Conditioning a randomized algorithm on its private seed and averaging over the finite placement distribution shows that one fixed labelling attains at least this average. That labelling is chosen before the private seed and is therefore an oblivious input. Together with Equation (269), this completes the proof of Theorem 2.4.

8 Blind execution and instance-optimal two-phase policies

We now turn to $\text{BE}(L)$. Running an untouched job blindly consumes its actual processing time and completes it; testing consumes one unit, reveals the value, and permits the processing operation to be delayed. The adversary fixes an arbitrary labelled vector in $[0, L]^n$ before the private seed is drawn. This section proves Theorem 2.5.

8.1 The distributional benchmark

Let D be a probability distribution on $[0, L]$ and let $P \sim D$. Write

$$\mu = \mathbb{E}P. \quad (272)$$

Blind execution completes one unit of job mass using expected work μ . The best possible test module is described by

$$\rho(D) = \sup_{0 \leq g \leq 1} \frac{\mathbb{E}g(P)}{1 + \mathbb{E}[Pg(P)]}, \quad \tau(D) = \frac{1}{\rho(D)}. \quad (273)$$

The function g selects outcomes that will be processed immediately after a test. For every such selection rule,

$$\tau \mathbb{E}g(P) - \mathbb{E}[Pg(P)] = \mathbb{E}[(\tau - P)g(P)] \leq \mathbb{E}(\tau - P)^+. \quad (274)$$

The maximizing rule selects all values below τ , no values above it, and an arbitrary fraction of a boundary atom. At equality,

$$\mathbb{E}(\tau - P)^+ = 1. \quad (275)$$

This equation has a unique solution: its left side is continuous and, after becoming positive, strictly increasing. If

$$a = \Pr(P < \tau), \quad \ell = \mathbb{E}[P \mathbf{1}_{P < \tau}],$$

then, with the boundary convention just described,

$$a\tau - \ell = 1, \quad \tau = \frac{1 + \ell}{a}. \quad (276)$$

Thus a full unit of tests together with all selected low outcomes is a divisible module of completion mass a , work $1 + \ell$, and work per completion τ .

If $\tau \geq \mu$, blind work has at least the completion density of any test module, and pure blind execution is optimal at the fluid scale. Suppose now that $\tau < \mu$. After testing a fraction q , the canonical schedule is

$$\begin{aligned} \underbrace{\text{tests with immediate } p < \tau}_{\text{discovery}} &\longrightarrow \underbrace{\text{known } \tau < p < \mu \text{ in SPT}}_{\text{medium}} \\ &\longrightarrow \underbrace{\text{untouched jobs blind}}_{\text{blind block}} \longrightarrow \underbrace{\text{known } p > \mu \text{ in SPT}}_{\text{high tail}}. \end{aligned} \quad (277)$$

For the explicit cost, first assume no atom lies at τ or μ , and put

$$\begin{aligned} \mathcal{L} &= \{p < \tau\}, & \mathcal{M} &= \{\tau < p < \mu\}, & \mathcal{H} &= \{p > \mu\}, \\ a &= \Pr(P \in \mathcal{L}), & \ell &= \mathbb{E}[P \mathbf{1}_{\mathcal{L}}], & m &= \mathbb{E}[P \mathbf{1}_{\mathcal{M}}], \\ d &= \Pr(P \in \mathcal{H}), & K_M &= \mathbb{E}[\min(P, Q) \mathbf{1}_{P, Q \in \mathcal{M}}], & K_H &= \mathbb{E}[\min(P, Q) \mathbf{1}_{P, Q \in \mathcal{H}}], \end{aligned}$$

where P, Q are independent with law D . For $r = 1 - q$, remaining-mass accounting gives

$$\begin{aligned} F_D(q) &= (1 + \ell) \left(q - \frac{aq^2}{2} \right) + qm(r + qd) + \frac{q^2 K_M}{2} \\ &\quad + \mu \left(r q d + \frac{r^2}{2} \right) + \frac{q^2 K_H}{2}. \end{aligned} \quad (278)$$

Atoms at τ are split between the low module and residual stock; atoms at μ may be placed on either side of the blind block. The same formula then holds for the resulting virtual classes.

Each term in Equation (278) has a direct area interpretation. During the test module, work $(1 + \ell)dq$ reduces unfinished mass from 1 to $1 - aq$. The medium work qm delays the untouched mass r and high mass qd , while K_M is the internal SPT pair term. The blind block delays all high jobs and itself loses mass linearly. Finally K_H is the high-tail pair term. Expansion gives the quadratic

$$F_D(q) = \frac{\mu}{2} + Aq + Bq^2, \quad (279)$$

where

$$A = 1 + \ell + m + \mu(d - 1) = 1 - \mathbb{E}(\mu - P)^+, \quad (280)$$

$$B = -\frac{a(1+\ell)}{2} + m(d-1) + \frac{K_M}{2} + \mu\left(\frac{1}{2} - d\right) + \frac{K_H}{2}. \quad (281)$$

The envelope proof below implies $B \geq 0$. In the short-test regime $\tau < \mu$, monotonicity in Equation (275) gives $A < 0$, and the optimal tested fraction is therefore

$$q^* = \begin{cases} 1, & B = 0, \\ \min\{1, -A/(2B)\}, & B > 0. \end{cases} \quad (282)$$

We define

$$\Phi(D) = \min_{0 \leq q \leq 1} F_D(q), \quad (283)$$

with $\Phi(D) = \mu/2$ in the regime $\tau \geq \mu$.

8.2 The fixed-fraction completion envelope

We first prove that Equation (277) is globally optimal in a fluid relaxation. Let D have finite support $0 = p_0 < p_1 < \dots < p_s \leq L$, with masses D_i . Fix a final tested fraction q . At an arbitrary schedule prefix, let

$$\begin{aligned} t &= \text{tested mass}, & b &= \text{blindly completed mass}, \\ c_i &= \text{processed tested mass of class } i > 0. \end{aligned}$$

The completed mass and idealized work are

$$y = D_0 t + b + \sum_{i>0} c_i, \quad x = t + \mu b + \sum_{i>0} p_i c_i, \quad (284)$$

subject to

$$0 \leq t \leq q, \quad 0 \leq b \leq 1 - q, \quad 0 \leq c_i \leq D_i t. \quad (285)$$

Define $H_{D,q}(x)$ as the largest completed mass satisfying these constraints with used work at most x . Every fluid schedule that ultimately tests mass q obeys this envelope, hence

$$\text{cost} \geq \int_0^\infty (1 - H_{D,q}(x)) dx. \quad (286)$$

Lemma 8.1 (Fixed- q envelope). *For every $q \in [0, 1]$, the right side of Equation (286) equals $F_D(q)$ and is attained by the four-block fluid policy. If $\tau \geq \mu$, it is at least $\mu/2$ for every q , with equality at $q = 0$.*

Proof. Split a boundary atom at τ into selected and residual virtual classes. Let $\mathcal{L}$ be the selected positive classes and put

$$a = D_0 + \sum_{i \in \mathcal{L}} D_i, \quad w = 1 + \sum_{i \in \mathcal{L}} p_i D_i, \quad w/a = \tau.$$

For the low part of any feasible prefix, write

$$y_L = D_0 t + \sum_{i \in \mathcal{L}} c_i, \quad x_L = t + \sum_{i \in \mathcal{L}} p_i c_i.$$

The density optimality in Equation (274) gives $y_L/x_L \leq a/w$. With $\lambda = y_L/a$, the revelation constraints imply $\lambda \leq t \leq q$, while density gives $\lambda w \leq x_L$. We may therefore replace the entire low prefix by λ full test modules, preserving completions and weakly reducing work.

Relax each residual capacity from $c_i \leq D_i t$ to $c_i \leq D_i q$. The result is an ordinary divisible knapsack with items

item	completion capacity	work per completion
full test module	aq	τ
residual tested class i	qD_i^R	p_i
blind module	$1 - q$	μ

The relaxation is pointwise tight. While greedy is partway through the test item, run the corresponding fraction of tests and all selected low outcomes. After all q test modules have run, every residual tested capacity is truly available. Greedy therefore realizes the original envelope at every work budget.

When $\tau < \mu$, the increasing work-per-completion order is exactly Equation (277); integrating it gives Equation (278). When $\tau \geq \mu$, the blind item has no worse density than the best test module and every excluded class has $p_i \geq \tau \geq \mu$. One all-blind item of capacity one pointwise dominates every $q > 0$ solution. Finally, feasibility is affine in q , so $H_{D,q}(x)$ is concave in q for each x . Hence its area complement $F_D(q)$ is convex, proving $B \geq 0$. $\square$

8.3 Adaptive first touches are a uniform urn

The fluid proof fixed q , but an announced finite policy may choose whom to touch next and when to stop testing as arbitrary functions of its history. The next finite lemma is the deferred-decisions statement needed to control such a policy.

Distinguish occurrences even when their processing times coincide. Let J be the public labels, Ω the occurrence tokens, and $v : \Omega \rightarrow [0, L]$ their values. A placement is a bijection $\sigma : J \rightarrow \Omega$. Fix a deterministic proper policy, equivalently fix the private seed of a randomized policy. Its first-touch labels $\ell_1(\sigma), \dots, \ell_n(\sigma)$ form a permutation of J . Define

$$R_{\mathcal{A}}(\sigma)_k = \sigma(\ell_k(\sigma)). \quad (287)$$

Lemma 8.2 (Adaptive-reveal bijection). *The map $\sigma \mapsto R_{\mathcal{A}}(\sigma)$ is a bijection from placements to permutations of Ω . Moreover, whether first touch k is a test or a blind execution is determined by the preceding revealed values $v(R_1), \dots, v(R_{k-1})$.*

Proof. Suppose $R_{\mathcal{A}}(\sigma) = R_{\mathcal{A}}(\sigma')$. A lockstep induction on k shows that the two executions have the same observable history before touch k . They have seen the same preceding values, so the deterministic policy performs the same intervening known-job operations, chooses the same next new label, and chooses the same test/blind kind. This remains true after blind execution because its realized duration is recorded in the public history. Hence $\ell_k(\sigma) = \ell_k(\sigma')$ for all k . Now Equation (287) gives $\sigma(\ell_k) = \sigma'(\ell_k)$ for every k ; since the ℓ_k enumerate J , the placements are equal. The reveal map is injective between finite sets of the same cardinality $n!$, hence bijective. The same induction under only a common revealed prefix proves predictability of the touch kind. $\square$

Consequently a uniform placement is carried to a uniform random permutation of occurrence tokens, and the test indicator is predictable. No conditional suffix compiler is needed. Duplicate values cause no problem because the bijection acts on occurrences.

For finite support, standard prefix concentration and a predictable martingale give the following simultaneous event. Stop while at least δn labels remain untouched, and put

$$e_n = \sqrt{8n \log n}, \quad \Gamma_n = (1 + 1/\delta)e_n. \quad (288)$$

Except with probability $o(1)$, for every earlier touch prefix and every class i ,

$$|\#\text{tested class } i - D_{n,i}\#\text{tests}| \leq \Gamma_n, \quad (289)$$

$$|\text{blind work} - \mu_n \# \text{blind jobs}| \leq L\Gamma_n. \quad (290)$$

Indeed, Hoeffding's inequality for without-replacement prefixes controls the remaining urn proportions. Multiplying the centered next-draw indicator by the predictable test selector gives a martingale with bounded increments; maximal Azuma–Hoeffding controls all stopping prefixes at once. The same argument with the predictable blind selector and bounded values proves Equation (290). Importantly, we never condition on the eventual tested fraction.

8.4 The announced-multiset lower bound

Fix a policy seed. Stop after $\lfloor (1 - \delta)n \rfloor$ first touches and let $Q_{n,\delta}$ be the number of tests completed by then, divided by n . Edit the suffix: every still-untouched job that would later be tested is instead executed blindly at its first original operation, and its later processing record is deleted. At most $3\delta n + O(1)$ bounded operations are moved or deleted. Pairwise delay accounting changes total cost by at most

$$O((L + 1)\delta n^2). \quad (291)$$

The edited schedule performs exactly $Q_{n,\delta}n$ tests.

Work on the single event Equations (289) and (290) and only now set $q = Q_{n,\delta}$. Normalize mass and work by n , and let $\gamma_n = \Gamma_n/n$. At any operation boundary, let t, b, z, c_i denote tested mass, blindly completed mass, zero mass completed by tests, and processed known positive mass of class i . The event gives

$$z \leq D_{n,0}t + \gamma_n, \quad c_i \leq D_{n,i}t + \gamma_n.$$

Set $c'_i = (c_i - \gamma_n)^+$. Then

$$c'_i \leq D_{n,i}t, \quad t \leq q, \quad b \leq 1 - q.$$

If x is the actual normalized prefix work, then

$$t + \mu_n b + \sum_i p_i c'_i \leq x + L(\gamma_n + \delta), \quad (292)$$

$$z + b + \sum_i c_i \leq D_{n,0}t + b + \sum_i c'_i + (s + 1)\gamma_n. \quad (293)$$

By the definition of the empirical envelope,

$$\text{completed fraction by } x \leq H_{D_n,q}(x + L(\gamma_n + \delta)) + (s + 1)\gamma_n. \quad (294)$$

Horizontal shifting loses at most the shift itself because $0 \leq 1 - H \leq 1$; the vertical slack is multiplied by a horizon at most $L + 1$. After restoring Equation (291), the fixed- q envelope lemma therefore gives, pathwise on the good event,

$$\frac{C_A}{n^2} \geq F_{D_n}(Q_{n,\delta}) - O_D(\gamma_n + \delta + 1/n) \geq \Phi(D_n) - O_D(\gamma_n + \delta + 1/n). \quad (295)$$

The long-test regime uses the $\mu_n/2$ dominance part of Lemma 8.1. Averaging first over the uniform placement and then over private seeds proves the same lower bound for every randomized announced policy.

The random-labelling and announced minimax values coincide. A uniform labelling is a valid finite Yao distribution, giving one inequality. For the other, privately relabel the physical jobs before running an optimal deterministic urn policy; against every fixed assignment, the policy sees a uniform assignment. Thus the lower bound above applies to the announced oblivious minimax problem, not merely to one randomized input model.

8.5 A growing grid and arbitrary bounded multisets

The number of distinct values may grow with n . Keep zero as a separate cell, partition $(0, L]$ into

$$K_n = \lfloor n^{1/6} \rfloor \quad (296)$$

bins, and round every positive value upward; let $h_n = L/K_n$ be the mesh. Zero preservation matters because a tested zero completes at its test.

Coupling each value with its rounded value changes total normalized processing work by at most h_n . Running corresponding operation words therefore changes normalized completion area by at most h_n . In particular,

$$|\Phi(D) - \Phi(D^{\text{grid}})| \leq h_n. \quad (297)$$

One way to see this without assuming a stable numerical threshold is to transport a policy through the coupling: attach a private auxiliary original mark to every rounded observation and make all decisions using that mark. The zero-preserving coupling makes the completion events coincide.

Apply the predictable urn event to all $K_n + 1$ cells and take $\delta_n = n^{-1/6}$. Repairing every class loses at most $(K_n + 1)\Gamma_n/n$ completion mass. The rounded analogue of Equation (294) is

$$\text{completed fraction by } x \leq H_q^{\text{grid}}(x + \zeta_n) + \zeta_n, \quad (298)$$

where

$$\zeta_n = O_L\left(\frac{(K_n + 1)\Gamma_n}{n} + h_n + \delta_n\right) = O_L(n^{-1/6}\sqrt{\log(n + 2)}). \quad (299)$$

Integrating exactly as above and using nonnegativity on the bad event yields the announced lower bound

$$\frac{\text{VAL}_{\text{ann}}(M)}{n^2} \geq \Phi(D_M) - O_L(n^{-1/6}\sqrt{\log(n + 2)}). \quad (300)$$

For the matching upper bound, compute an optimal canonical template for the rounded empirical multiset, choose $\lfloor q^*n \rfloor$ uniformly random labels to test, and execute the four blocks. Pairwise delay accounting expresses its expectation as bounded one-occurrence and two-distinct-occurrence kernels. The without-replacement pair coefficients differ from their product-law values by $O(1/n)$ uniformly in the tested count. Hence

$$\mathbb{E}C_n(\pi) = n^2\Psi_{D_M^{\text{grid}}}(\pi) + O_L(n) \quad (301)$$

uniformly over canonical templates π . Together with Equation (297), this proves

$$\left| \frac{\text{VAL}_{\text{ann}}(M)}{n^2} - \Phi(D_M) \right| \leq O_L(n^{-1/6}\sqrt{\log(n + 2)}). \quad (302)$$

8.6 Removing the announcement

The universal policy first draws a uniform random permutation and blindly executes

$$k_n = \lceil n^{1/2} \rceil \quad (303)$$

pilot jobs. Their observed durations give a grid histogram $\hat{D}_n$. The policy minimizes the canonical quadratic over every grid threshold, blind-block insertion position, and $q \in [0, 1]$, then runs the selected four-block template on the remaining jobs in the unused random order.

Let D_n and D_n^R be the full and remaining grid histograms. Sampling without replacement gives

$$\mathbb{E}\|\hat{D}_n - D_n\|_1 \leq \sqrt{\frac{K_n + 1}{k_n}}, \quad \mathbb{E}\|\hat{D}_n - D_n^R\|_1 \leq \frac{n}{n - k_n} \sqrt{\frac{K_n + 1}{k_n}}. \quad (304)$$

For a fixed canonical template, formula Equation (278) consists only of bounded one- and two-draw expectations. Therefore, uniformly in the grid and template,

$$|\Psi_D(\pi) - \Psi_E(\pi)| \leq C_L \|D - E\|_1, \quad (305)$$

for example with $C_L = 20(L + 1)$. If $\hat{\pi}$ minimizes the sampled objective and π_R minimizes the remaining objective, the empirical optimization sandwich is

$$\begin{aligned} \Psi_{D_n^R}(\hat{\pi}) &\leq \Psi_{\hat{D}_n}(\hat{\pi}) + C_L \|\hat{D}_n - D_n^R\|_1 \\ &\leq \Psi_{\hat{D}_n}(\pi_R) + C_L \|\hat{D}_n - D_n^R\|_1 \\ &\leq \Phi(D_n^R) + 2C_L \|\hat{D}_n - D_n^R\|_1. \end{aligned} \quad (306)$$

Conditional on the pilot, the remaining order is uniform and the learned template is fixed, so Equation (301) applies. The pilot itself has work at most Lk_n and delays at most n jobs, adding at most $Lnk_n = O_L(n^{3/2})$. Combining Equations (297) and (304) with Equations (305) and (306) gives

$$\frac{\mathbb{E}C_{\mathcal{A}^*}(p)}{n^2} \leq \Phi(D_M) + O_L\left(n^{-1/6} \sqrt{\log(n + 2)}\right) \quad (307)$$

uniformly over every fixed labelled vector $p \in [0, L]^n$.

For the converse, reveal M to an arbitrary policy for free and draw a uniform labelling. The announced lower bound Equation (300) applies seedwise. Some fixed labelling in the support has cost at least the average, which proves Equation (13). This completes the proof of Theorem 2.5.

8.7 Why the medium block is necessary

Let

$$\Pr(P = 0) = \frac{1}{5}, \quad \Pr(P = 9) = \frac{1}{5}, \quad \Pr(P = 16) = \frac{3}{5}.$$

Then $\mu = 57/5$ and $\tau = 5$. Thus tested zeros are immediate, known nine-jobs precede the blind block, and known sixteen-jobs follow it. Formula Equation (278) becomes

$$F_D(q) = \frac{57}{10} - \frac{44}{25}q + \frac{11}{10}q^2.$$

Its minimizer is $q^* = 4/5$ and its value is 4.996. If all tested nonzero jobs are instead deferred until after the blind block, the best value is 5.04. The gap is a positive constant times n^2 : the medium SPT block is not cosmetic, and a policy of the simpler form “threshold, then blind, then all tested residuals” is not instance optimal.

9 Machine-checked verification

The accompanying Lean 4 development [dMU21] uses mathlib [The20] and is pinned to version 4.30.0. Its root module imports the complete deterministic, randomized-obligatory, and blind-execution developments. The source contains no `sorry`, `admit`, or project-defined axiom. This section records the precise proof boundary; Appendix C gives the detailed dictionary for the deterministic exact curve. The randomized finite common-upper curve of Section 6 is currently analytic only and is not included in the machine-checked claims below.

9.1 Deterministic exact ratios

The theorem `SchedulingPaper.exact_ratio_main` packages both directions of the six-piece finite-cap curve and the obligatory endpoint. Its upper fields contain literal transcript-only strategies, termination proofs, and $O(n)$ certificates; its lower fields quantify over arbitrary deterministic strategy families and return genuine fixed inputs after the formal replay theorem. The formal curve includes all five joins, continuity, the plateau, and the global maximum.

9.2 Randomized obligatory testing

The module `SchedulingPaper.RandomizedFourThirds` exports the operational upper theorem

```
theorem unknownMultiset_operational_expectedCost_le_20378 ... :  
  uniformAverage (physicalSampledRunCost ...) <=  
    4 / 3 * finiteObligatoryOPT p + 20378 * R ^ 7
```

At the concrete parameter choice $n \leq R^4$, this is Equation (269). The proof below this interface includes the maximum-density optimizer, exact and robust $4/3$ certificates, threshold closure for sample-empty bins, finite hypergeometric concentration, the cutoff-32 fallback, sample-first accounting, bad-event excess, and termination of the sampled transcript strategy.

The same module exports `oblivious_iid_binary_lower_online_actualOPT`. It accepts an arbitrary finite family of public-transcript strategies, assumes only that the supplied fuel completes every fair binary input, and returns a fixed binary input whose average cost has coefficient $4n/(3n+5)$. The formal development identifies its denominator with the literal clairvoyant obligatory optimum. Since $4n/(3n+5) \rightarrow 4/3$, the upper and lower exports give the machine-checked sharp asymptotic value.

9.3 Blind-execution instance optimality

This formalization uses an operation semantics in which completion of a blind job records its realized duration. The trace-bijection and observed-trace modules prove the occurrence-token reveal bijection of Lemma 8.2, its lockstep causality theorem, the predictable test selector, and the exact selected-count and blind-work identities.

The deterministic analytic layer checks the full-module replacement, fractional-knapsack envelope, long-test dominance, revelation repair, horizontal integral shift, formula Equation (278), its quadratic normal form and vertex minimizer, and the three-point witness. The finite layer checks all-grid repair, bounded one- and two-job kernels, histogram stability, the canonical physical operation word, a blind-pilot compiler, and the announced adaptive lower bound.

The end-to-end theorem is exported as `boundedUniform_blindPilot_concrete_rate` in `RandomizedOptionalUnknownRates`. In mathematical notation, for $m = \lfloor n^{1/16} \rfloor \geq 2$, every $p \in [0, L]^n$, and every complete announced policy π , it proves

$$\frac{\mathbb{E}C_{\text{learned}}(p)}{n^2} \leq \frac{\mathbb{E}C_{\pi}(p)}{n^2} + \frac{7830(L+1)^2}{m}. \quad (308)$$

The learned algorithm depends only on n, L and its private permutations, not on the multiset. Lean separately proves $\lfloor n^{1/16} \rfloor \rightarrow \infty$ and hence that the displayed error tends to zero. It also discharges the zero-mean case for the same algorithm. This is a slower rate than the exponential-tail analysis in Equation (307), but it proves the same uniform $o_L(n^2)$ theorem using finite second-moment and checkpoint bounds.

The formal statements use finite uniform averages rather than measure-valued random seeds. This is sufficient for the advertised algorithms and Yao distributions, all of which use finite permutations, finite pilot choices, and finite tie-breaking coins. Classical real analysis and noncomputable exact minimizers are used in certificates; no floating-point program is extracted.

10 Discussion and open questions

The two common-upper curves identify the leading quadratic constants of the deterministic and oblivious-randomized games for every finite u ; the deterministic analysis also includes $u = \infty$. Their scope is deliberately precise: one machine, all release dates zero, unit tests, nonpreemptive operations, total completion time, and one common upper limit when the cap is finite. The deterministic upper proof loses at most a constant on each of the n discrete transitions. This loss is asymptotically negligible, but it is not a fixed additive constant over the whole instance. On $u \in [z_*, \infty]$ its per-transition constant is uniform in u .

For finite u , the randomized curve is learned with an $O_u(n^{11/6})$ remainder. This exponent is only a convenient consequence of the pilot/grid parameters; sharper empirical-process estimates should reduce it without changing the four leading branches. The randomized proof uses the oblivious assumption essentially. Its random-labelling lower bound does not apply to an adversary that observes the private choices, so the adaptive-adversary curve remains open.

The randomized obligatory theorem changes both the algorithmic power and the adversary: the input is fixed obliviously, while the algorithm may privately permute labels. Its $4/3$ upper bound is uniform over unbounded processing times and unknown multisets. The additive $O(n^{7/4})$ term is unlikely to be optimal. The announced stationary theorem suggests that sharper empirical-process estimates, avoiding the coarse histogram and sample-first schedule, could reduce it substantially without changing the leading constant.

The blind-execution theorem has a different benchmark. It is instance-optimal up to $o(n^2)$ relative to the announced oblivious minimax value, but it is not a bounded competitive-ratio theorem relative to the clairvoyant SPT optimum in Equation (18). This distinction is unavoidable for the present statement: on sparse instances the clairvoyant cost can be subquadratic and the permitted error may dominate it. Natural next questions are whether a finer, instance-dependent error can recover a meaningful competitive guarantee on those sparse inputs, and whether the fixed bounded-support assumption can be replaced by a moment condition.

Conjecture 10.1 (Endpoint additive competitiveness). There exist a deterministic online algorithm and a constant $B < \infty$ such that every instance at the cap-free endpoint satisfies

$$\text{ALG}(I) \leq R_\star \text{OPT}(I) + B.$$

The present Adaptive-Threshold algorithm is the natural first candidate. In Equations (51) and (52) the exact local contributions contain favorable negative terms that are intentionally discarded before the bank-account calculation. A discrete potential that retains those terms might prove Conjecture 10.1; alternatively, a finite-instance obstruction might show that a different policy is needed. The current lower bound does not distinguish these possibilities. In particular, our result neither proves nor claims the stronger ordinary inequality with $B = 0$.

The analogous endpoint-additive question is open for each nontrivial finite-cap deterministic branch as well: our claims concern CR_∞ and the infimum of admissible additive coefficients. Also open are adaptive-adversary randomization, heterogeneous upper limits, nonuniform test times in the exact-ratio regime, weights, and more than one machine. The deterministic common-upper curve shows why a direct extension is unlikely to have one universal threshold: already in the uniform case the optimal policy changes form at five analytically distinct junctions.

The lower construction also suggests a structural principle for future lower bounds: hard instances reveal many decreasing positive levels, with continuum increment $dp = ds/(1+s)$, and postpone zero jobs to the end. Our argument proves that this pattern is sufficient. It remains open whether every asymptotically extremal construction must have this form.

11 Conclusion

Uniform tests produce three different but related models. For deterministic common-upper execution, the exact answer is the six-piece curve Equation (4); its cap-free endpoint is $R_\star = 1.570574096649395\dots$ and its global maximum is $u_\diamond = 1.866760399173862\dots$. The proof combines binary stopping games, a harmonic adversary, and adaptive bank potentials.

Against an oblivious adversary, randomization gives the exact four-piece finite-cap curve Equation (6). Comparing raw execution with a maximum-density stationary threshold, and flattening the survival function, reduces its upper bound to two scalar binary families. A sublinear pilot learns the right family without knowing the multiset, while binary random labellings give all matching lower branches. At the obligatory endpoint, a uniform unbounded-support argument learns the stationary threshold without any boundedness assumption, and the balanced half-zero, half-two instance proves that $4/3$ remains sharp.

When untested jobs instead run for their actual duration, the right answer is instance dependent. Tests, known residual classes, and blind work become divisible completion modules; their fractional-knapsack order yields the four-block policy and the quadratic benchmark $\Phi(D)$. The adaptive-reveal bijection and a simultaneous urn bound show that neither adaptive stopping nor prior knowledge of the multiset improves its leading n^2 coefficient. Together, these results identify completion density and pairwise delay as the common organizing principles behind deterministic competitive ratios, randomized learning, and instance-optimal testing.

Acknowledgements

An initial improvement over the then-current deterministic lower bound for obligatory testing was obtained with the Bolzano open-source multi-agent research system [GHK⁺26].

That preliminary result motivated the present project. Most of the subsequent conjecture generation, proof development, counterexample search, exposition, and Lean formalization was carried out with substantial assistance from OpenAI’s GPT-5.6 through interactive Codex sessions. In particular, the randomized common-upper curve was developed in a separate model run and then integrated with the instance-optimal framework. The author selected the research directions, reviewed and revised the arguments, and takes responsibility for the claims in this paper.

A Finite rational checkpoint for the cap-free lower bound

The lower bound at $u = \infty$ takes two limits: first the scaling $H \rightarrow \infty$, and then the number K of positive levels tends to infinity. This appendix is not needed for that proof. It gives one exact finite instance showing that the leading-term calculation already predicts the true ratio accurately.

Take

$$K = 1000, \quad \alpha = \frac{9}{8}, \quad \gamma = \frac{679}{1000}, \quad \xi = \frac{K}{\alpha} = \frac{8000}{9}, \quad H = 9000.$$

There are H jobs at each positive level and ξH zero jobs. All multiplicities are integral, and all processing times defined by Equation (31) are rational.

Using exact rational arithmetic in Equations (30) and (34) to (36) gives the leading-coefficient comparison

$$\frac{Q_K}{D_K} = 1.5705287442294085 \dots > 1.5705.$$

The displayed decimal is only for orientation; all comparisons below are between integers or rational numbers.

For the offline schedule, retaining the terms that are linear in H gives

$$\text{OPT}_H = H^2 D_K + H L_{\text{off}}, \quad (309)$$

where

$$\begin{aligned} L_{\text{off}} &= \frac{1}{2} \left(\xi + \sum_{i=0}^{K-1} (1 + p_i) \right) \\ &= \frac{1}{2} \left(\xi + 2K + \frac{K\gamma}{\xi} + \sum_{s=1}^{K-1} \frac{K-s}{\xi+s} \right) = 1656.440800628232 \dots \end{aligned} \quad (310)$$

Again, the middle expression is the exact rational value used in the comparison.

The discrete shortest-prefix calculation behind Equation (28) also retains favorable diagonal terms that vanish after division by H^2 . Before dropping them it gives

$$\text{ALG}_H \geq H^2 Q_K + \frac{H}{2} \left(\xi + \sum_{i=0}^{K-1} p_i + \sum_{i=0}^{K-1} s_i \right) \geq H^2 Q_K.$$

Here the s_i ’s are the nonnegative completion fractions from the lower proof. Exact rational arithmetic verifies

$$2000H^2 Q_K > 3141(H^2 D_K + H L_{\text{off}}). \quad (311)$$

Consequently, the fixed instance obtained by the replay argument in Section 4.1 satisfies

$$\frac{\text{ALG}_H}{\text{OPT}_H} \geq \frac{H^2 Q_K}{H^2 D_K + H L_{\text{off}}} = 1.5705286254871234 \dots > \frac{3141}{2000} = 1.5705.$$

Thus the checkpoint includes the finite- H correction; it is not merely a comparison of the two leading coefficients.

B Independent certificate by bubbling caps to a prefix

For completeness, we give a second proof of the quadratic estimate. This argument is also a useful audit of the fifth-endpoint calculation: it never inserts $p = u$ into an ordinary deferred formula before the caps have actually been moved.

Work first in normalized fluid mass. Immediately before an adjacent pair XU , let E_0 be the preceding E -mass, P the preceding substantive mass, D its deferred submass, x the remaining mass, and w the mass of U -endpoints strictly after the pair. Put

$$y = \frac{D - C(P + E_0)}{x}, \quad \eta = \frac{D - CP}{x} = y + C \frac{E_0}{x} \leq 0. \quad (312)$$

Direct differentiation of the exact pair objective (160), including the mutual pair and all future caps, gives the two nontrivial commutators

$$UM - MU = 1 + \left(1 + \eta - \frac{Cw}{x}\right) (y - c) a'_c(y), \quad (313)$$

$$UQ - QU = \left(\eta - \frac{cw}{x}\right) (y - c) a'_c(y). \quad (314)$$

Here $UX - XU$ means “new objective after moving U left” minus the old objective. The second quantity is nonnegative. For the first, if $B = 1 + \eta - Cw/x \leq 0$, its second term is nonnegative. If $B > 0$, then $B \leq 1$, and on $-1 < y \leq 0$

$$\frac{(c - y)B}{c - 2y} \leq 1.$$

Thus (313) is nonnegative as well. Below $y = -1$, the threshold is flat. Direct evaluation also gives

$$UZ - ZU = UE - EU = 1. \quad (315)$$

The suffix terms involving w are crucial: they make every adjacent exchange valid even while other caps remain to the right. Repeated swaps therefore put all caps into one prefix.

After that prefix there is no future cap. Let e, b, d be the preceding E -mass, substantive mass, and deferred substantive mass, and define

$$y = \frac{d - C(e + b)}{x}, \quad v = \frac{e}{x}, \quad h = d - Cb, \quad k = x + d - Cb.$$

The four ordinary commutators are

$$\begin{aligned} ME - EM &= a_c + a'_c(C - y) \frac{k}{x}, & QE - EQ &= 1 - a'_c(C - y)(-y - Cv), \\ MZ - ZM &= a_c - a'_c y \frac{k}{x}, & QZ - ZQ &= 1 - a'_c y \frac{h}{x}. \end{aligned} \quad (316)$$

They are nonnegative. Indeed, in the varying region $h/x = y + Cv \in [y, 0]$ and $k/x \in [0, 1]$. With $t = -y \in [0, 1]$,

$$a'_c(C - y)t = \frac{t(C + t)}{c + 2t} \leq 1, \quad a'_c t^2 = \frac{t^2}{c + 2t} \leq 1; \quad (317)$$

the first inequality is equivalent to $(t - 1)(t + c) \leq 0$. In the flat region $a'_c = 0$. Hence Z, E can be moved to the tail, and a worst word has the form

$$U^\mu \{M, Q\}^* \{Z, E\}^*. \quad (318)$$

On the middle block use the simpler potential

$$\widehat{W}(x, N, D) = xD + x^2 f_c \left(\frac{D - CN}{x} \right).$$

The identity

$$2f_c - yf'_c = \max\{0, -Cf'_c, -Cf'_c + a_c(1 + y), 1 - cf'_c + a_c y\} \quad (319)$$

is a direct rewriting of (173); it pays the entire ordinary continuation. If the cap prefix has mass μ , then after it $x = 1 - \mu$, $N = D = \mu$, and $y = -c\mu/(1 - \mu)$. Its internal cap-pair reward plus its worst continuation is therefore

$$F_{c,u}(\mu) = \frac{2 - c(u - 1)}{2} \mu^2 + \mu(1 - \mu) + (1 - \mu)^2 f_c \left(-\frac{c\mu}{1 - \mu} \right). \quad (320)$$

For $0 < \mu < 1/C$, differentiation gives

$$F'_{c,u}(\mu) = c\mu [Z(c) - u - \operatorname{artanh} \mu]. \quad (321)$$

On the mixed curve the unique maximizer in this range is therefore $\mu = m$, and

$$F_{c,u}(m) = f_c(0) + \frac{c}{2} (\operatorname{artanh} m - m) = \frac{c}{2}. \quad (322)$$

Here $m \leq 1/C$, since $c, m \leq 1/\varphi$. On the flat part $\mu \geq 1/C$,

$$F_{c,u}(\mu) = \mu - \frac{c(u - 1)}{2} \mu^2. \quad (323)$$

Its derivative at $1/C$ is

$$\frac{c(1 + m) - 1}{C} \leq 0,$$

and only decreases thereafter. Thus $F_{c,u}(\mu) \leq c/2$ on the whole unit interval. At $c = r, m = 0, u = z_*$, the same calculation is maximized at $\mu = 0$, and increasing u only lowers (320).

To make this independent exchange proof finite, fix $\varepsilon > 0$ and first replace the last $\varepsilon n + O(1)$ outcomes by genuine zeros. Earlier states and actions are unchanged, and only $O(\varepsilon n^2 + n)$ pair terms are affected. All cap and ordinary exchanges in the retained prefix then occur with $x \geq \varepsilon$; their displayed fluid commutators have uniform Taylor remainders there. More precisely, after assigning mass $1/n$ to every job, a normalized adjacent exchange has remainder $O_{u,\varepsilon}(n^{-3})$; there are at most n^2 exchanges. At the join $y = -1$, split an exchange at the join and use the two one-sided formulae, whose values agree. The artificial zeros already form the terminal block, so no unbubbled suffix cap invalidates (316). The normalized one-job remainder in the continuation-potential calculation is $O_{u,\varepsilon}(n^{-2})$. For fixed ε , first let $n \rightarrow \infty$, and then let $\varepsilon \downarrow 0$. Equations (319)–(322) give

$$\Gamma_n \leq c \binom{n}{2} + o(n^2),$$

which is an independent size-asymptotic certificate for the two upper bounds above.

C Machine-checked deterministic theorem and its correspondence

This appendix records the public theorem exported by the Lean 4 development [dMU21] associated with this manuscript and explains how its formal predicates correspond to the deterministic statements in Theorems 2.1 and 2.2. The development uses the mathlib library [The20] and is pinned to Lean and mathlib version 4.30.0. Its root module imports `SchedulingPaper.MainTheorems`, whose final theorem is reproduced below. Apart from the typographic rendering of Unicode symbols, this is a verbatim excerpt.

```
theorem exact_ratio_main
  : (∀ {u : ℝ}, 0 < u →
    FiniteExactRatioConclusion u (exactCurve u)) ∧
    ObligatoryExactRatioConclusion RStar := by
  constructor
  · intro u hu
    exact finite_exact_ratio hu
  · exact obligatory_exact_ratio
```

The two conclusion records occurring in the theorem are:

```
structure FiniteExactRatioConclusion (u ratio : ℝ) : Prop where
  lower : LowerBound.FixedSizeLowerBound (.finite u) ratio
  upper :
    ∃ strategy : ∀ n, Online.Strategy n,
      UpperBound.CompletesByAnalysisFuel (.finite u) strategy ∧
      SizeAsymptoticUpper
        (UpperBound.strategyCost (.finite u) strategy)
        (UpperBound.fixedOfflineCost (.finite u)) ratio

structure ObligatoryExactRatioConclusion (ratio : ℝ) : Prop where
  lower : LowerBound.FixedSizeLowerBound .infinite ratio
  upper :
    ∃ strategy : ∀ n, Online.Strategy n,
      UpperBound.CompletesByAnalysisFuel .infinite strategy ∧
      SizeAsymptoticUpper
        (UpperBound.strategyCost .infinite strategy)
        (UpperBound.fixedOfflineCost .infinite) ratio
```

C.1 Objects on the two sides

The following table gives the direct dictionary between the formal objects and the notation of the manuscript.

Lean object	Manuscript object	Meaning
<code>Cap.finite u</code>	finite common upper limit u	processing times lie in $[0, u]$; testing and raw execution are available
<code>Cap.infinite</code>	obligatory testing	processing times are nonnegative and raw execution is unavailable
<code>exactCurve u</code>	$V(u)$	the six branches in Equation (4), including all five verified joins
<code>RStar</code>	$R_\star$	$1 + 2/(z_\star - 1)$, where $z_\star > 1$ is the unique solution of $z_\star - 3 = \log z_\star$
<code>fixedOfflineCost</code>	OPT	the clairvoyant shortest-first cost of the effective lengths $q_u(p)$
<code>strategyCost</code>	ALG	the total completion cost of the operational online transcript

A formal strategy on n jobs is a deterministic function from the public transcript to its next action. A test publishes the revealed processing time; raw execution publishes no hidden value. Thus the strategy has exactly the information available to an online algorithm in the manuscript. The formal action type uses the standard no-idling normalization appropriate when all jobs are released at time zero. Processing times and comparisons are exact real numbers.

C.2 Expansion of the lower and upper fields

For a cap and a candidate value R , `FixedSizeLowerBound` expands to the following statement. For every size-indexed family of deterministic strategies $\mathcal{B}_n$, every $\varepsilon > 0$, and every N , there are an $n \geq N$, a genuine fixed admissible n -job input I , and a finite settled execution such that either the execution leaves a job incomplete or

$$(R - \varepsilon) \text{OPT}(I) \leq \text{ALG}_{\mathcal{B}_n}(I). \quad (324)$$

An incomplete settled run represents infinite cost. In particular, for algorithms that complete all inputs, (324) is the usual lower-bound sequence on arbitrarily large fixed instances. The lower constructions are first described by adaptive revelation oracles, but the formal replay theorem freezes every such interaction into the fixed input quantified here; this is the machine-checked counterpart of Lemma 3.3.

The predicate `SizeAsymptoticUpper` is the epsilon formulation

$$\forall \varepsilon > 0 \exists N \forall n \geq N \forall I, \quad \text{ALG}_{\mathcal{A}_n}(I) \leq (R + \varepsilon) \text{OPT}(I). \quad (325)$$

The accompanying completion field proves that every job is done after the common analysis fuel $2n + 1$, rather than merely assigning a cost to a partial trace. Consequently, Equations (324) and (325) are the two matching operational statements that characterize the optimal value in Equation (23). The Lean theorem packages these two directions; it does not introduce a separate real-valued infimum or `limsup` object.

Before projecting to `SizeAsymptoticUpper`, the checked upper proof constructs a stronger `FixedUpperCertificate`: for the chosen cap it contains a constant $C \geq 0$ such that, for every n and every fixed input,

$$\text{ALG}_{\mathcal{A}_n}(I) \leq R \text{OPT}(I) + Cn. \quad (326)$$

The quadratic lower bounds Equations (21) and (22) then imply (325). For $u \geq z_\star$, the separately verified plateau certificate uses the same strategy and the same C , independent of u , including the obligatory endpoint. This uniformity is stronger than the quantifier order displayed by `exact_ratio_main`, which only exposes one strategy after a cap has been fixed.

C.3 Correspondence with the two deterministic theorems

The first conjunct of `exact_ratio_main` quantifies over every finite $u > 0$. Substituting $R = \text{exactCurve } u = V(u)$, its lower field gives the fixed-instance lower direction of Theorem 2.2, while its upper field supplies a completing strategy and the matching upper direction. The definition of `exactCurve` is literally the six-way case split in Equation (4); the Lean development also proves its joins, continuity, plateau, and global maximum at u_∞ .

The second conjunct substitutes $R = \text{RStar} = R_\star$ in the obligatory model. Its lower field is Theorem 2.1(i), and its completing upper strategy together with (326) is Theorem 2.1(ii). The common formal infrastructure also proves that the offline benchmark is the shortest-first schedule of effective lengths and that the transcript cost is exactly its suffix-weighted pair-accounting expression.

C.4 Proof boundary

The root Lean module imports the complete lower- and upper-bound assemblies. The public theorem has no operational interface hypotheses, and the development contains no `sorry`, `admit`, or project-defined `axiom`. It does use `mathlib`’s classical real analysis and noncomputable choices of exact real parameters. Thus it certifies deterministic mathematical strategy terms, not an extracted floating-point implementation.

The theorem recorded here is specifically the size-asymptotic result. It does not assert the pointwise inequality $\text{ALG} \leq R \text{OPT}$ without a remainder, nor admissibility of the endpoint coefficient with one fixed additive constant. The formal library separately derives additive admissibility of every coefficient $R + \varepsilon$ from (326). The machine-checked mixed-regime upper bound follows the cap-reserve potential proof; the bubbling argument in Appendix B remains an independent manuscript cross-check.

References

- [AE21] Susanne Albers and Alexander Eckl. Explorable uncertainty in scheduling with non-uniform testing times. In Christos Kaklamanis and Asaf Levin, editors, *Approximation and Online Algorithms: 18th International Workshop, WAOA 2020, Revised Selected Papers*, volume 12806 of *Lecture Notes in Computer Science*, pages 127–142. Springer, 2021. doi:10.1007/978-3-030-80879-2_9.
- [BS26] Felix Buld and Andreas S. Schulz. Scheduling with testing: Competitive algorithms for minimizing the total weighted completion time in the adversarial model, 2026. arXiv:2606.25166.
- [DDN⁺22] Fanny Dufossé, Christoph Dürr, Noël Nadal, Denis Trystram, and Óscar C. Vázquez. Scheduling with a processing time oracle. *Applied Mathematical Modelling*, 104:701–720, 2022. doi:10.1016/j.apm.2021.12.020.
- [DEL24] Konstantinos Dogeas, Thomas Erlebach, and Ya-Chun Liang. Scheduling with obligatory tests. In Timothy Chan, Johannes Fischer, John Iacono, and Grzegorz Herman, editors, *32nd Annual European Symposium on Algorithms (ESA 2024)*, volume 308 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 48:1–48:14, Dagstuhl, Germany, 2024. Schloss Dagstuhl – Leibniz-Zentrum für Informatik. doi:10.4230/LIPIcs.ESA.2024.48.

- [DEMM20] Christoph Dürr, Thomas Erlebach, Nicole Megow, and Julie Meißner. An adversarial model for scheduling with testing. *Algorithmica*, 82(12):3630–3675, 2020. doi:10.1007/s00453-020-00742-2.
- [DKL⁺23] Christoph Damerius, Peter Kling, Minming Li, Chenyang Xu, and Ruilong Zhang. Scheduling with a limited testing budget: Tight results for the offline and oblivious settings. In *31st Annual European Symposium on Algorithms (ESA 2023)*, volume 274 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 38:1–38:15. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2023. doi:10.4230/LIPIcs.ESA.2023.38.
- [dMU21] Leonardo de Moura and Sebastian Ullrich. The Lean 4 theorem prover and programming language. In *Automated Deduction – CADE 28*, volume 12699 of *Lecture Notes in Computer Science*, pages 625–635. Springer, 2021. doi:10.1007/978-3-030-79876-5_37.
- [GCH24] Mingyang Gong, Zhi-Zhong Chen, and Kuniteru Hayashi. Approximation algorithms for multiprocessor scheduling with testing to minimize the total job completion time. *Algorithmica*, 86(5):1400–1427, 2024. doi:10.1007/s00453-023-01198-w.
- [GHK⁺26] Jan Grebík, Pavel Hubáček, Martin Koutecký, Matěj Kripner, Václav Rozhoň, Robert Šámal, and Adrián Zámečník. Bolzano: Case studies in LLM-assisted mathematical research, 2026. URL: <https://arxiv.org/abs/2604.16989>, arXiv:2604.16989.
- [Kah91] Simon Kahan. A model for data in motion. In *Proceedings of the 23rd Annual ACM Symposium on Theory of Computing (STOC 1991)*, pages 265–277. Association for Computing Machinery, 1991. doi:10.1145/103418.103449.
- [KLL⁺26] Bob Krekelberg, Alison Hsiang-Hsuan Liu, Fu-Hong Liu, Prudence W. H. Wong, and Xiao-Ou Zhang. The power of amortization on minimizing total completion time with explorable uncertainty, 2026. Version 2, revised 8 May 2026. arXiv:2503.13357.
- [LL26] Kao-Chuan Liang and Ya-Chun Liang. The completion-threshold framework for obligatory-test scheduling on multiple machines, 2026. Version 2, revised 14 July 2026. arXiv:2606.02029.
- [LLWZ23] Alison Hsiang-Hsuan Liu, Fu-Hong Liu, Prudence W. H. Wong, and Xiao-Ou Zhang. The power of amortization on scheduling with explorable uncertainty. In Jaroslav Byrka and Andreas Wiese, editors, *Approximation and Online Algorithms: 21st International Workshop, WAOA 2023*, volume 14297 of *Lecture Notes in Computer Science*, pages 90–103. Springer, 2023. doi:10.1007/978-3-031-49815-2_7.
- [LMS19] Retsef Levi, Thomas L. Magnanti, and Yaron Shaposhnik. Scheduling with testing. *Management Science*, 65(2):776–793, 2019. doi:10.1287/mnsc.2017.2973.
- [LMS24] Retsef Levi, Thomas L. Magnanti, and Yaron Shaposhnik. Scheduling with testing of heterogeneous jobs. *Management Science*, 70(5):2934–2953, 2024. doi:10.1287/mnsc.2023.4833.

- [MPT94] Rajeev Motwani, Steven J. Phillips, and Eric Torng. Nonclairvoyant scheduling. *Theoretical Computer Science*, 130(1):17–47, 1994. doi:10.1016/0304-3975(94)90151-1.
- [The20] The mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020)*, pages 367–381. Association for Computing Machinery, 2020. doi:10.1145/3372885.3373824.