

Scheduling with Uniform Tests: Information and Optimization

Václav Rozhoň

August 2026

Abstract

A machine faces many jobs whose lengths are hidden. Spending one unit of time to inspect a job may reveal a short job that should be finished now, or it may reveal nothing useful while every other job waits. When should the machine keep looking, and when should it start working?

We study this question for one machine and total completion time. We consider four natural rules: every job must be tested; a unit-time revealing optimization can replace a raw time u by a shorter revealed length; an untested job can be run blindly for its true hidden length; or a unit-time optimization can shorten it without revealing the new length. Together these models separate the two possible benefits of spending one unit before execution: learning which job to do next and making the job faster.

We give exact asymptotic answers for these models, both as competitive ratios and as instance-optimal guarantees. The resulting algorithms are often quite simple, which may make them useful in practice.

Contents

1	Introduction	3
1.1	Warmup: every job has to be tested	3
1.2	One algorithm for every input	5
1.3	Generalizations of the model	6
1.4	Information without speedup: blind execution	6
1.5	Blind optimization	7
1.6	Revealing optimization	9
1.7	Techniques	11
1.8	Related work	11
1.9	Roadmap	12
2	Models and common preliminaries	13
2.1	Information and adversaries	14
2.2	Offline optima and pair accounting	14
2.3	Maximum-density modules and fluid completion envelopes	17
2.4	Competitive-ratio conventions	19
2.5	Test-order adversaries	20
2.6	Probabilistic tools	20
2.7	Notation and fluid-to-finite transfer	23

3	Deterministic revealing optimization: the exact curve	25
3.1	Policy guarantees	27
3.1.1	Low and intermediate caps	27
3.1.2	High caps	31
3.2	Branchwise lower bounds	38
3.2.1	The binary stopping game and its lower envelope	38
3.2.2	The cap-free harmonic core	40
3.2.3	The unrestricted mixed lower bound	46
3.2.4	The large-upper-limit lower bound by cap-free transfer	48
3.3	Assembly of the curve	49
3.3.1	The shape of the claimed curve	49
3.3.2	Completion of the curve and consistency at the joins	49
3.4	The obligatory endpoint	50
4	Revealing-optimization instance optimality	50
4.1	The fixed-fraction fluid benchmark	51
4.2	Adaptive policies and the announced lower bound	52
4.3	Removing the announcement	53
5	Randomized revealing optimization: the exact four-piece curve	55
5.1	Offline and stationary fluid coefficients	55
5.2	Flattening the survival function	57
5.3	The two scalar maximizations	58
5.4	Worst-case upper bound from instance optimality	59
5.5	Binary oblivious lower bounds	59
5.6	The randomized obligatory endpoint	61
5.6.1	Normalization of the offline pair identity	62
5.6.2	Stationary prefixes	62
5.6.3	Learning the prefix from a sublinear sample	63
5.6.4	Good samples, fallback, and bad-event excess	64
5.6.5	Transfer of the zero-two lower bound	66
6	Obligatory-testing instance optimality	66
6.1	The maximum-density benchmark	68
6.2	The announced upper bound	69
6.3	No adaptive announced policy has a better leading cost	69
6.4	Removing the announcement	70
7	Blind execution: instance-optimal four-block policies	71
7.1	The distributional benchmark	72
7.2	The fixed-fraction completion envelope	74
7.3	Adaptive first touches are a uniform urn	74
7.4	The announced lower bound	75
7.5	A growing grid and arbitrary bounded multisets	76
7.6	Removing the announcement	76
7.7	Why the medium block is necessary	77

8	Blind optimization: speedup without revelation	77
8.1	The deterministic worst-case curve	79
8.2	Proof of randomized instance optimality	82
8.3	The exact worst-case curve	84
	Acknowledgements	85
A	Machine-checked verification	86
B	Finite rational checkpoint for the cap-free lower bound	88
C	Independent certificate by bubbling caps to a prefix	89
D	Machine-checked deterministic theorem and its correspondence	91
D.1	Objects on the two sides	92
D.2	Expansion of the lower and upper fields	92
D.3	Correspondence with the two deterministic theorems	93
D.4	Proof boundary	93

1 Introduction

Suppose that we have many jobs waiting for one machine. We want to minimize the sum of the times at which they finish. The length of each job is written on a folded piece of paper. If we knew all the lengths, the right schedule would be clear: we would simply run the jobs from shortest to longest. But the papers are folded, and opening one of them takes one unit of machine time. Should we open another paper, or should we start working?

Opening a paper may reveal a tiny job that we can finish immediately. It may also reveal a huge job that is best left for the end. On the other hand, looking has a cost. If one hundred jobs are still unfinished, one unit spent opening a paper adds one hundred units to the sum of completion times. The machine must therefore balance two uses of its time: learning which jobs are short and completing jobs that are already understood.

Variants of this question have appeared in both stochastic and adversarial scheduling with testing [LMS19, DEMM20, DEL24]. This work improves earlier results and studies new models. We start with the cleanest version, in which every paper has to be opened. We then allow a job to be run without opening its paper. Depending on what this means, we obtain three further variants. The same algorithmic ideas—random sampling, thresholding, and shortest-first execution—give exact asymptotic answers in all four models.

1.1 Warmup: every job has to be tested

Consider the following problem. There are n jobs. Testing job i takes one unit of time and reveals a number $p_i \geq 0$. The job then needs another p_i units of processing; if $p_i = 0$, it completes with the test. For now every job has to be tested: testing is *obligatory*. The formal model is given in Section 2.

We compare an online scheduler with the shortest-first schedule that knows all values p_i in advance. The *competitive ratio* is the online cost divided by this offline optimum. We care about its asymptotic value. Since every job must be tested, even the optimum has cost $\Omega(n^2)$, so we ignore additive terms of size $o(n^2)$.

The obligatory-testing model for total completion time was introduced by Dogeas, Erlebach, and Liang [DEL24]. Their work and a later lower bound of Liang and Liang [LL26]

left a gap between the best deterministic upper and lower bounds. They also left the randomized case open. We close both gaps.

Theorem 1.1 (Obligatory testing, informal version of Theorems 3.13 and 5.5). *In the obligatory-testing model, the best deterministic asymptotic competitive ratio is $R_{\text{OT}}^{\text{det}} = 1.570574\dots$. Against an adversary that fixes the input before the algorithm draws its random bits, the best randomized ratio is $R_{\text{OT}}^{\text{rand}} = 4/3$. The two optimal policies are sketched in algorithms 1 and 2.*

Here and below, *shortest first* means processing revealed jobs in nondecreasing order of p_i . The following pseudocode gives a little more detail about the two policies. The formulas are not meant to be memorable.

Algorithm 1 Deterministic threshold

- 1: Put $c = 0.570574\dots$ and $R = 1 + c = R_{\text{OT}}^{\text{det}}$.
- 2: Before test $i+1$, where $0 \leq i < n$, let c_i be the number of positive jobs processed immediately and d_i the number deferred.
- 3: Put

$$y_i = \frac{d_i - R(c_i + d_i)}{n - i}.$$

- 4: Use threshold $T_i = 1$ if $y_i \leq -1$; otherwise use

$$T_i = 1 + \frac{1}{2} \ln \frac{c + 2}{c - 2y_i}.$$

- 5: Test the next job. Process it now if $p \leq T_i$; otherwise defer it.
 - 6: After all tests, finish deferred jobs shortest first.
-

Algorithm 2 Learned randomized threshold $\mathcal{A}_n^*$

- 1: Set $B_n = O(n^{1/20})$ and test $O(n^{3/4})$ random jobs to estimate the job-length distribution $X \sim \hat{D}$.
- 2: For each candidate threshold $0 \leq t \leq B_n$, put

$$a(t) = \Pr(X \leq t), \quad m(t) = \mathbb{E}[X \mathbf{1}_{X \leq t}],$$

and define

$$\hat{\tau} = \min_{\substack{0 \leq t \leq B_n \\ a(t) > 0}} \frac{1 + m(t)}{a(t)}.$$

- 3: If $\hat{\tau} > B_n$, test every remaining job and then finish all revealed jobs shortest first.
 - 4: Otherwise process the sampled jobs with $p \leq \hat{\tau}$. Keep testing jobs, processing them immediately when $p \leq \hat{\tau}$ and deferring them otherwise.
 - 5: After all tests, finish deferred jobs shortest first.
-

Both algorithms follow the same principle: process short jobs as soon as possible and leave long jobs for the end. The difficult part is choosing the threshold. A deterministic algorithm has no representative view of the unseen jobs. Its decisions are therefore more complicated: in fact, its threshold must change with the history. The exact update is given in steps 3 and 4 of algorithm 1. The matching lower bound in Section 3.2.2 uses an adversary that reveals jobs with decreasing positive values of p_i while hiding many zero jobs until the end.

Randomization gives the algorithm a global view of the input. It first tests a sublinear random sample and estimates the distribution of job lengths. Because the labels are privately shuffled, the remaining empirical distribution stays close to the sampled one throughout the run. The algorithm therefore faces essentially the same distribution at every step and can use one fixed threshold.

The choice of that threshold has a simple interpretation. For a candidate t , a full round of tests together with the immediate processing of outcomes at most t uses normalized work $1 + m(t)$ and completes mass $a(t)$. Thus

$$\frac{1 + m(t)}{a(t)}$$

is the work per immediate completion of this testing module; the unit term includes the tests of *all* jobs, including those deferred to the tail. The algorithm minimizes this quantity and uses its optimum $\hat{\tau}$ as the actual threshold. Equivalently, adding an outcome of length p improves the module exactly when $p < \hat{\tau}$.

The matching randomized lower bound is correspondingly simple: half of the jobs have length zero and half have length two, and their labels are shuffled. Until the shuffle is uncovered, one unit of work completes only about half a job. Counting the area under the number of unfinished jobs gives the lower bound $4/3$. The formal argument is in Section 5.6.

1.2 One algorithm for every input

Let us return to algorithm 2. The ratio $4/3$ describes only the hardest input. It says much less on an easy one. Suppose, for example, that all processing times are very large compared with the unit test—a plausible regime in many applications. Then the straightforward strategy that first tests everything and subsequently runs the jobs shortest first is nearly optimal. The worst-case guarantee of algorithm 2 still tells us only that its cost is at most $1.33 \dots$ times the optimum.

In fact, the same algorithm satisfies a much stronger per-input guarantee: it is *instance-optimal* on every class with a fixed upper bound on job lengths.

Theorem 1.2 (Obligatory-testing instance optimality, informal version of Theorem 6.1). *Fix $L < \infty$. The algorithm $\mathcal{A}^*$ in algorithm 2 has the following property: for every size- n multiset $M \subseteq [0, L]$ and every online algorithm $\mathcal{A}'$,*

$$\text{cost}(\mathcal{A}^*) \leq \text{cost}(\mathcal{A}') + o_L(n^2),$$

where both costs are measured on M and include the algorithms' randomness. The competing algorithm $\mathcal{A}'$ may even be designed specifically for M and told the entire input multiset in advance, though not its assignment to the public labels.

Here and in the instance-optimal statements below, cost on a multiset uses the order-oblivious convention of Remark 2.1. A private shuffle makes the original labeling irrelevant, so the comparison is made separately for every multiset M .

Both assumptions in the theorem are necessary. Without a fixed bound L there is no randomized instance-optimal algorithm, and without randomization there is no deterministic one; see Theorem 6.2.

Two points about the theorem are worth emphasizing. First, the comparison is made separately on every multiset. We must therefore compete not only with one fixed online

algorithm, but also with algorithms optimized for this particular M . Allowing $\mathcal{A}'$ to know M makes this strongest possible online benchmark explicit.

Second, the existence of some instance-optimal algorithm is perhaps not surprising. One could imagine a meta-algorithm that samples the input to estimate M and then runs an optimal policy for that estimate. The substantive part of the theorem is that no complicated meta-algorithm is needed. The optimal leading-order policy for a known input is itself a simple threshold policy, so $\mathcal{A}^*$ has essentially the same form as algorithm 2: sample the input distribution, choose the maximum-density threshold, and use it for the rest of the schedule.

1.3 Generalizations of the model

We next weaken the obligatory model in two directions. First, we may allow a job to run without being tested. Second, the unit preliminary action may have two different benefits. It may reveal the true running time p_i before the job is run. It may also be an optimization procedure that replaces a common raw running time u by a shorter time $p_i \leq u$. For example, a generic piece of code solves every task within time u , but an AI agent can spend one unit of time adapting the code to a particular input and make it faster.

These two benefits are independent. An optimization may reveal its new running time p_i , giving the scheduler both a faster job and information about where the job belongs. Or it may leave p_i hidden until the optimized job finishes.

These choices give the following four models.

model	without the unit action	with the unit action	where discussed
obligatory testing	cannot be run	reveals p_i and unlocks the job	Intro 1.1; Sections 3.4, 5.6 and 6
blind execution	runs for its hidden length p_i	reveals p_i before execution	Intro 1.4; Section 7
blind optimization	runs for raw time u	replaces u by a still-hidden $p_i \leq u$	Intro 1.5; Section 8
revealing optimization	runs for raw time u	reveals p_i and replaces u by p_i	Intro 1.6; Sections 3 to 5

The last three rows separate the two benefits of the unit preliminary action. Blind execution studies information without speedup and is related to the stochastic model of Levi, Magnanti, and Shaposhnik [LMS19]. Blind optimization studies speedup without advance information and is related to the model of Damerius et al. [DKL⁺23]. Revealing optimization, introduced by Dürr et al. [DEMM20], contains both benefits.

The three optional models have somewhat different optimal policies and input-specific benchmarks, so we state their results separately. Their randomized algorithms nevertheless share one design principle: privately shuffle the jobs, use a sublinear sample to estimate the input distribution, and optimize the rest of the schedule using that information.

1.4 Information without speedup: blind execution

Suppose an untested job can simply be run until it finishes. Its running time is its true hidden length p_i . Testing does not make the job shorter; it only tells us p_i before we decide when to run it. We call this *blind execution*.

Without an upper bound on the job lengths, no randomized algorithm has a finite competitive ratio, even if we ignore an additive $o(n^2)$ term; see Theorem 7.2 for a simple

counterexample. For every fixed bound L , however, sampling again gives one algorithm that is optimal on every bounded multiset.

Theorem 1.3 (Blind-execution instance optimality, informal version of Theorem 7.1). *Fix $L < \infty$. There is one randomized algorithm $\mathcal{A}^*$ such that, for every size- n multiset $M \subseteq [0, L]$ and every online algorithm $\mathcal{A}'$,*

$$\text{cost}(\mathcal{A}^*) \leq \text{cost}(\mathcal{A}') + o_L(n^2).$$

To describe the algorithm, let $X \sim \hat{D}$ denote the estimated job-length distribution and put $\hat{\mu} = \mathbb{E}X$. As in algorithm 2, define

$$a(t) = \Pr(X \leq t), \quad m(t) = \mathbb{E}[X \mathbf{1}_{X \leq t}], \quad \hat{\tau} = \min_{\substack{t \geq 0 \\ a(t) > 0}} \frac{1 + m(t)}{a(t)}.$$

The optimal policy is more subtle than algorithm 2, because running a job blindly can be better than testing it. Sometimes, after the initial sample, the best choice is to run every remaining job blindly. Even when further testing is useful, it need not be useful all the way to the end of the input.

For a simple illustration, suppose that jobs have only two possible lengths. The obligatory-testing policy would process short outcomes immediately and defer long ones. Now consider the last untouched job. If it is short, we want to run it next. If it is long, it can still be run next, as the first job in the arbitrarily ordered long tail. Thus in either case the useful action is to run the job; testing it first only adds one unit of work. This is why the value of testing can disappear before every job has been tested.

In general, the policy tests only a fraction q of the jobs and uses the threshold $\hat{\tau}$ from algorithm 2 to decide which tested jobs to process immediately. After testing stops, deferred jobs with $\hat{\tau} \leq p < \hat{\mu}$ are processed before the untouched jobs: their known lengths are smaller than the expected duration $\hat{\mu}$ of a blind execution. The untouched jobs are then run blindly, and tested jobs with $p \geq \hat{\mu}$ form the final tail. The optimal fraction q minimizes the explicit quadratic $F_{\text{BE}, \hat{D}}$ from Equation (321).

Algorithm 3 Instance-optimal blind execution

- 1: Shuffle the jobs and run $o(n)$ of them blindly; use their observed lengths to estimate $\hat{D}$ and $\hat{\mu}$.
 - 2: Compute $\hat{\tau} = \min_{t \geq 0, a(t) > 0} (1 + m(t))/a(t)$.
 - 3: If $\hat{\tau} \geq \hat{\mu}$, put $q = 0$. Otherwise choose $q \in \arg \min_{0 \leq z \leq 1} F_{\text{BE}, \hat{D}}(z)$.
 - 4: Test a random q -fraction of the remaining jobs. Whenever a test reveals $p < \hat{\tau}$, process that job immediately; defer all other outcomes.
 - 5: After the tests stop, process the deferred jobs with $\hat{\tau} \leq p < \hat{\mu}$, shortest first.
 - 6: Run every untouched job blindly.
 - 7: Finish the remaining tested jobs shortest first.
-

1.5 Blind optimization

The two remaining models start from the same operation. Every job takes a known raw time u , and one unit of optimization reduces its running time to $p_i \leq u$. They differ only in whether optimization reveals p_i . We start with the case in which it does not. Before execution, all optimized jobs still look the same.

This is the simplest case. Randomization is necessary for instance optimality: a binary adversarial-labeling argument rules out every deterministic policy; see Theorem 8.1. The deterministic worst-case problem nevertheless has an exact and very simple answer. Up to $u = 2$, run every job raw. After $u = 2$, optimize and run every job immediately in an arbitrary fixed order. A hidden-stopping adversary proves that adaptively mixing the two actions cannot improve this choice.

Theorem 1.4 (Deterministic blind-optimization curve, informal version of Theorem 8.3). *The exact deterministic asymptotic competitive ratio is*

$$R_{\text{BO}}^{\text{det}}(u) = \begin{cases} 1, & 0 < u \leq 1, \\ u, & 1 \leq u \leq 2, \\ \frac{\sqrt{4u^3 - 4u + 1} - 1}{2(u - 1)}, & u \geq 2. \end{cases} \quad (1)$$

It grows as $\sqrt{u} + o(1)$.

With randomization, we can sample a few jobs to estimate the only number that matters: the mean μ of the optimized lengths. The policy in algorithm 4 then compares the two average times per job, u and $1 + \mu$, and uses the better action for all remaining jobs.

Theorem 1.5 (Blind-optimization instance optimality, informal version of Theorem 8.5). *Consider the blind-optimization model with some $u < \infty$. There is one randomized algorithm $\mathcal{A}^*$ such that, for every size- n multiset $M \subseteq [0, u]$ and every online algorithm $\mathcal{A}'$,*

$$\text{cost}(\mathcal{A}^*) \leq \text{cost}(\mathcal{A}') + o_u(n^2).$$

Algorithm 4 Instance-optimal blind optimization

- 1: Shuffle the jobs, optimize and run $\lceil n^{2/3} \rceil$ of them, and let $\hat{\mu}$ be the mean observed length.
 - 2: Treat $\hat{\mu}$ as the true mean of the input.
 - 3: If $1 + \hat{\mu} < u$, optimize and then run every remaining job.
 - 4: Otherwise run every remaining job raw for time u .
-




Figure 1: The exact deterministic and randomized ratios for blind optimization.

More precisely, if μ_M is the mean optimized length in M , the best possible leading value is

$$\frac{n^2}{2} \min\{u, 1 + \mu_M\}.$$

Knowing the instance-optimal algorithm does not yet give its competitive ratio: we must still find the worst input. The resulting ratio is unbounded and grows on the order of $\sqrt{u}$.

Theorem 1.6 (Blind-optimization curve, informal version of Theorem 8.6). *The exact randomized asymptotic competitive ratio $R_{\text{BO}}^{\text{rand}}(u)$ is 1 for $u \leq 1$ and is given by the explicit two-branch curve in Figure 1 for $u > 1$. It grows as $\sqrt{u}/2 + 1/2 + o(1)$ as $u \rightarrow \infty$.*

1.6 Revealing optimization

Now assume that optimization also reveals p_i . Thus the unit action buys both information and speed. We call this *revealing optimization*.

The two extreme regimes are easy to understand. If $u \leq 1$, the test alone costs at least as much as raw execution, so we should never test. If u gets very large, raw execution is unattractive, so the problem approaches obligatory testing. The interesting question is what happens between these regimes.

This model was introduced by Dürr et al. [DEMM20]. Their results showed that intermediate values of u are the difficult ones. We determine the exact answer for every u , for both deterministic and randomized algorithms.

Theorem 1.7 (Revealing-optimization curves, informal version of Theorems 3.1 and 5.1). *For every $u > 0$, the exact deterministic and randomized asymptotic competitive ratios $R_{\text{RO}}^{\text{det}}(u)$ and $R_{\text{RO}}^{\text{rand}}(u)$ are the six-piece and four-piece curves in Figures 2 and 3. The deterministic curve reaches $1.87\dots$ and then falls to the obligatory value $1.57\dots$. The randomized curve reaches $1.63\dots$ and eventually becomes $4/3$.*

Write the five deterministic transition points as

$$\begin{aligned} u_1^{\text{det}} &= 1, & u_2^{\text{det}} &= 1.866760\dots, & u_3^{\text{det}} &= 3.147899\dots, \\ u_4^{\text{det}} &= \varphi + 2 = 3.618033\dots, & u_5^{\text{det}} &= 4.505241\dots \end{aligned}$$

They delimit the six branches from left to right.




Figure 2: The guarantees of the three deterministic policies and their exact lower envelope. The black outline is the optimal ratio. The arrow on RAW marks its off-scale linear continuation; the teal horizontal line is the φ guarantee of zero-prefix FORCEDPREFIXUTE.

The deterministic curve is the lower envelope of three explicit policy families. We describe their shape here and defer the exact parameters to Section 3.

The first algorithm, $\text{RAW}(u)$, simply fixes an arbitrary job order and executes every job untested for the public raw time u . Its ratio is 1 for $u \leq u_1^{\text{det}}$ and u for $u > u_1^{\text{det}}$, and it supplies the lower envelope up to u_2^{det} ; see Figure 2.

The second algorithm, FORCEDPREFIXUTE in algorithm 5, tests every job and normally processes only outcomes below a fixed threshold. It makes one deliberate exception: every outcome in an initial $b(u)$ -fraction of the test order is processed immediately. Without this

forced prefix, deferring early long outcomes can create a tail that is delayed by almost all later tests; the prefix balances this tail cost against the opposite risk of processing a long job before shorter unseen jobs.

The exact parameter $b(u)$ is defined in Equation (76); it is nonnegative on $[u_2^{\det}, u_3^{\det}]$ and vanishes at $u_3^{\det}$. For $u \geq u_4^{\det}$, the same zero-prefix policy remains φ -competitive by Lemma 3.3; this continuation is shown in Figure 2 to make the lower envelope visible.

Algorithm 5 FORCEDPREFIXUTE(J, u)

- 1: Fix a test order $j_1, \dots, j_n$ and set $Q \leftarrow \emptyset$.
 - 2: Set $\theta_u \leftarrow \min\{1, u - 1\}$. If $u_2^{\det} \leq u \leq u_3^{\det}$, set $b \leftarrow b(u)$; otherwise set $b \leftarrow 0$.
 - 3: For $i = 1, \dots, n$, test j_i and observe p_{j_i} .
 - 4: If $i \leq \lfloor bn \rfloor$ or $p_{j_i} \leq \theta_u$, process j_i immediately; otherwise add it to Q .
 - 5: Process the jobs of Q in nondecreasing revealed processing time.
-

Finally, ADAPTIVETHRESHOLD is the same history-dependent policy as algorithm 1. For finite u , only its parameter c changes: on $u_4^{\det} \leq u \leq u_5^{\det}$ it is chosen by the optimization in Equation (74), after which it equals the obligatory-testing value $0.570574\dots$. The three policies form the deterministic lower envelope in Figure 2: RAW wins first, FORCEDPREFIXUTE in the middle, and ADAPTIVETHRESHOLD for large u .

For randomized algorithms, private sampling again gives instance optimality with a policy similar in shape to algorithm 3. Unlike blind execution, every tested processing time is at most the raw time u , so all tested jobs belong before the untouched raw block.

Theorem 1.8 (Revealing-optimization instance optimality, informal version of Theorem 4.1). *Fix $u < \infty$. There is one randomized algorithm $\mathcal{A}^*$ such that, for every size- n multiset $M \subseteq [0, u]$ and every online algorithm $\mathcal{A}'$,*

$$\text{cost}(\mathcal{A}^*) \leq \text{cost}(\mathcal{A}') + o_u(n^2).$$

For the randomized curve, write

$$u_1^{\text{rand}} = 1, \quad u_2^{\text{rand}} = 5.048917\dots, \quad u_3^{\text{rand}} = \frac{25}{4} = 6.25.$$

These three points delimit its four branches.

Algorithm 6 Instance-optimal revealing-optimization policy

- 1: Shuffle the jobs, test $\lceil \sqrt{n} \rceil$ of them, and use the observed lengths to estimate the input distribution.
 - 2: Compute the best threshold and the best tested fraction q .
 - 3: Test a random q -fraction; finish outcomes below the threshold at once.
 - 4: Finish all other tested jobs, shortest first.
 - 5: Run every untouched job raw for time u .
-

The pilot sample estimates the input distribution. If that estimate were exact, the best leading-order online schedule would be determined by two numbers: a threshold and the fraction q of jobs to test. Short outcomes are completed during testing, longer tested outcomes form a shortest-first tail, and the untouched jobs form the final raw block. The




Figure 3: The exact randomized revealing-optimization ratio as a function of the raw execution time u .

algorithm optimizes these two parameters on the sample and applies them after a private shuffle. The sample and its estimation error cost only $o_u(n^2)$.

For each multiset, this policy attains the best online leading cost. The competitive ratio is obtained by maximizing the ratio between that cost and the clairvoyant optimum over inputs. The worst input can be reduced to two job lengths, leaving a one-variable optimization.

The curves in Figures 2 and 3 are not monotone. This is not a contradiction. For very small u , testing is useless, while as $u \rightarrow \infty$ the model converges to the obligatory-testing setting of Section 3.4. Intermediate values are harder because raw execution gives the clairvoyant scheduler an additional useful option.

The exact formulas and matching constructions are in Sections 3 and 5.

1.7 Techniques

The proofs repeatedly use three pictures.

The area of unfinished jobs. During one unit of machine work, the objective increases by the number of jobs that are still unfinished. Thus the sum of completion times is the area under the curve “number of unfinished jobs versus time.” This picture is often easier to reason about than individual completion times.

Jobs completed per unit of work. A round of tests together with the short jobs completed during that round can be viewed as one large action. It spends some amount of time and completes some fraction of the jobs. Higher-density actions—those completing more jobs per unit of work—should come first. Sorting these actions produces the thresholds, the ordered blocks, and the one-variable quadratics in the instance-optimal results.

A private shuffle. The universal algorithms start by shuffling the labels and inspecting a small prefix. The same idea is needed in the lower bound, where the policy may adaptively decide which label to touch next. If equal job lengths are temporarily distinguished as separate tokens, the order in which any deterministic policy first reveals these tokens is itself just a permutation of them. The revealed sequence is therefore sampling without replacement. One concentration event can then control every possible stopping point of the policy.

The deterministic proofs use a complementary idea. Their lower bounds make the algorithm choose between completing the current jobs and waiting for shorter future jobs. Their upper bounds maintain a threshold and charge each wrongly ordered pair to a small potential. The exact constants appear when the two possible losses are equal.

1.8 Related work

The closest papers have already appeared next to the models they introduced. We give a broader comparison here.

Scheduling with tests belongs to optimization with *explorable uncertainty*, where uncertain data can be queried at a cost; the query viewpoint goes back at least to Kahan [Kah91]. Levi, Magnanti, and Shaposhnik study a stochastic scheduling model and derive prefix-testing policies followed by an order based on expected processing time per unit weight [LMS19, LMS24]. Our four-block blind-execution policy has a similar shape, but our inputs are adversarial, need not converge to a distribution, and are not announced to the universal algorithm.

Dürr et al. [DEMM20] introduced the adversarial revealing-optimization model and obtained deterministic bounds, including the $1.87\dots$ guarantee for binary values. We close the deterministic gap for arbitrary values $p_i \in [0, u]$ and give the exact randomized curve against an oblivious adversary. Related work allows nonuniform test times [AE21, LLWZ23, KLL⁺26], multiple machines [GCH24], or weights [BS26].

Dogeas, Erlebach, and Liang [DEL24] introduced obligatory testing for total completion time. For uniform tests they proved the SIDLE upper bound $1.58\dots$ and the deterministic lower bound $\sqrt{2}$. Liang and Liang [LL26] later raised the lower bound to $3/2$. Our matching value is $1.57\dots$, and our randomized result gives the oblivious value $4/3$. More strongly, its stationary threshold is asymptotically optimal separately for every bounded input.

The processing-time oracle of Dufossé et al. [DDN⁺22] is close to blind execution, but allows only two possible hidden durations. If no tests are available and processing times are revealed only at completion, one obtains the classical nonclairvoyant model of Motwani, Phillips, and Torng [MPT94]. These semantics differ from raw execution in revealing optimization, which always lasts the declared time u .

1.9 Roadmap

Section 2 formally defines the four models, their information structures and adversaries, the corresponding offline optima, and our asymptotic competitive-ratio convention.

Section 3 then proves the exact deterministic revealing-optimization curve. Its cap-free harmonic core and parameterized endpoint bank are shared by the mixed branch and the plateau; only after the finite curve is complete does Section 3.4 derive the deterministic obligatory ratio by a uniform endpoint transfer.

Section 4 identifies the optimal leading cost separately for every bounded revealing input and learns the associated threshold policy from a sublinear sample. Section 5 maximizes this benchmark against the clairvoyant optimum and proves the exact four-piece randomized curve. Its final subsection, Section 5.6, transfers the lower bound to the unbounded obligatory endpoint and supplies the cutoff-and-fallback argument needed for a uniform $4/3$ upper bound. Section 6 then strengthens that endpoint algorithm to a single policy that is asymptotically optimal on every bounded obligatory input.

Section 7 turns to blind execution. It derives the announced-input benchmark, proves that arbitrary adaptive policies cannot beat it, and removes advance knowledge of the input by sampling; it also explains why no finite worst-case ratio is possible without a bound on job lengths. Section 8 studies speedup without revelation, proves the deterministic hidden-stopping lower bound and matching two-policy curve, proves the randomized instance-optimal theorem, and maximizes its mean-based benchmark to obtain the exact randomized worst-case curve.

The appendices document and independently audit the more delicate parts of the argument. Appendix A states precisely which results are machine checked and summarizes the corresponding Lean 4 development. Appendix B evaluates a large but finite rational instance of the cap-free lower bound, while Appendix C gives an independent cap-bubbling certificate for the deterministic upper bound. Finally, Appendix D records the exported Lean theorem and the detailed dictionary between its objects and the deterministic statements in the paper.

2 Models and common preliminaries

This section makes the four models from the introduction precise and records the common facts used throughout the paper. There are n jobs, all available at time zero, and one machine. Operations are nonpreemptive, and at most one operation can run at a time. Job j has a hidden value $p_j \geq 0$. The four models differ in what can be done before this value is known.

Obligatory testing OT.

The values p_j are arbitrary finite nonnegative numbers. Every job must first be tested for one unit. The test reveals p_j , after which its processing operation of length p_j may be performed immediately or delayed.

Blind execution BE(L).

Here $p_j \in [0, L]$. A job may be run blindly for its actual duration p_j ; the value becomes known only when that execution finishes. Alternatively, the job may be tested for one unit, revealing p_j , and its processing operation may then be delayed. Testing provides information but does not shorten the job.

Blind optimization BO(u).

Here $p_j \in [0, u]$. A job may be run raw for the known time u . Alternatively, one unit of optimization changes its running time from u to p_j without revealing p_j ; the optimized job is subsequently run blindly, possibly after a delay. Its value becomes observable only when that execution finishes. Raw execution reveals no hidden value.

Revealing optimization RO(u).

Again $p_j \in [0, u]$, and raw execution takes the known time u . The other option is a unit test that reveals p_j , followed, possibly after a delay, by processing for p_j . Thus the preliminary action provides both the speedup and its size.

The different letters L and u are intentional. In blind execution, L is only a known bound used by the uniform instance-optimal theorem. In the last two models, u is itself an available running time and hence part of every scheduling decision. The number n and the relevant parameter L or u are known to the algorithm.

The completion time C_j is the time at which job j finishes. If a test reveals $p_j = 0$, the job completes at the end of that test. A blind zero job in BE(L) completes as soon as it is selected, whereas an optimized zero job in BO(u) first pays its unit optimization cost. The objective is

$$\sum_{j=1}^n C_j.$$

Obligatory testing can be viewed as the limit of revealing optimization in which raw execution becomes unavailable. We use this only as intuition: the obligatory theorems and their algorithms are proved directly, without interchanging the limits $u \rightarrow \infty$ and $n \rightarrow \infty$.

We write $\text{cost}(\mathcal{A}, I)$ for the objective value of an online algorithm $\mathcal{A}$ on an instance I , and $\text{cost}(\text{OPT}, I)$ for the value of an offline optimum that knows all processing times. In calculations we use the shorter abbreviations

$$\text{ALG}_{\mathcal{A}}(I) = \text{cost}(\mathcal{A}, I), \quad \text{OPT}(I) = \text{cost}(\text{OPT}, I),$$

and omit algorithms or instances when they are clear. Thus OPT denotes an algorithm inside $\text{cost}(\text{OPT}, I)$ and its cost only in the abbreviated notation $\text{OPT}(I)$.

2.1 Information and adversaries

The deterministic curve results allow the usual adversarial input and use adaptive revelation only as a lower-bound device that is frozen by Lemma 2.10. The randomized results use an *oblivious* adversary: the labeled processing-time vector is fixed before the algorithm draws its private random seed. Unless explicitly stated otherwise, the algorithm is not told the multiset of values. An *announced* policy is additionally given that multiset but not its assignment to labels. Announcing the multiset can only help and is therefore a valid strengthening in lower bounds.

All randomized expectations are over private algorithmic randomness. A uniform random relabeling is private randomness, not a distributional assumption on the adversarial multiset.

Remark 2.1 (Order-oblivious convention). For a multiset M of n occurrences and a randomized policy $\mathcal{A}$, write

$$\text{cost}^{\text{oo}}(\mathcal{A}, M) := \max_{\sigma \in \mathfrak{S}_n} \mathbb{E} \text{cost}(\mathcal{A}, \sigma(M)).$$

This is the standard order-oblivious view of instance optimality: the input is a multiset and its order carries no instance information [ABC17, NRTT24]. In the present models the convention is especially harmless. Given any policy $\mathcal{A}$, let $\mathcal{A}^{\text{sym}}$ first draw a uniform permutation Π of the public labels and then simulate $\mathcal{A}$ through that relabeling. For every fixed labeling σ ,

$$\mathbb{E} \text{cost}(\mathcal{A}^{\text{sym}}, \sigma(M)) = \mathbb{E}_{\Pi} \mathbb{E} \text{cost}(\mathcal{A}, \Pi\sigma(M)),$$

which is independent of σ and at most $\text{cost}^{\text{oo}}(\mathcal{A}, M)$. Hence randomized policies may be symmetrized without loss, after which only the multiset matters. When the introduction compares costs “on M ,” it uses this convention. An announced policy knows M , but not its assignment to the public labels.

For lower bounds we use a uniform distribution over fixed labelings as a finite Yao distribution and then select one labeling whose cost is at least the average, as formalized in Lemma 2.17 below.

2.2 Offline optima and pair accounting

The offline optimum knows all values p_j . It therefore chooses the shorter available way to complete each job and orders the resulting job blocks shortest first. For later use, define

$$\lambda_u(p) := \min\{u, 1 + p\} \quad (0 < u < \infty), \quad (2)$$

and associate with job j the model-dependent *effective length*

$$\lambda_j := \begin{cases} 1 + p_j, & \text{OT}, \\ p_j, & \text{BE}(L), \\ \lambda_u(p_j), & \text{BO}(u) \text{ or RO}(u). \end{cases} \quad (3)$$

Lemma 2.2 (Effective lengths and the offline pair identity). *Fix any one of the four models and let $\lambda_{(1)} \leq \dots \leq \lambda_{(n)}$ be its effective lengths from Equation (3). A clairvoyant optimum completes the jobs in this order and has value*

$$\text{OPT} = \sum_{k=1}^n \sum_{i=1}^k \lambda_{(i)}, \quad (4)$$

$$= \sum_i \lambda_i + \sum_{i < j} \min\{\lambda_i, \lambda_j\}, \quad (5)$$

$$= \frac{1}{2} \left(\sum_i \lambda_i + \sum_{i,j} \min\{\lambda_i, \lambda_j\} \right). \quad (6)$$

In OT it tests every job. In BE(L) it runs every job blindly. In BO(u) and RO(u) it uses raw mode when $u \leq 1 + p_j$ and the unit preliminary action followed by processing otherwise.

Proof. Let $C_{[1]} \leq \dots \leq C_{[n]}$ be the ordered completion times of any feasible schedule. Completing job j consumes at least λ_j units of work belonging to that job. Hence, by time $C_{[k]}$, the machine has performed the required work of some k jobs, and

$$C_{[k]} \geq \sum_{i=1}^k \lambda_{(i)}.$$

Choose the mode specified in the lemma, keep each resulting block contiguous, and execute the blocks in nondecreasing order of λ_j . This schedule is feasible in the relevant model and attains equality for every k . Summing over k and then collecting the contribution of each unordered pair gives the first two formulas. Splitting the double sum into its diagonal and the two orientations of every unordered pair gives the symmetric form. $\square$

For any finite input object I whose occurrences have processing times $p_1, \dots, p_n$, write

$$D[I] := \frac{1}{n} \sum_{i=1}^n \delta_{p_i}. \quad (7)$$

Thus $D[M]$ and $D[p]$ are instances of the same empirical-law operator; labels matter to the online interaction, but not to $D[I]$.

For a probability distribution D and a nonnegative effective-length map ℓ , let P, P' be independent with law D and define

$$\Omega_\ell(D) := \frac{1}{2} \mathbb{E} \min\{\ell(P), \ell(P')\}. \quad (8)$$

We use the model-specific abbreviations

$$\begin{aligned} \Omega_{\text{OT}}(D) &:= \frac{1}{2} \mathbb{E} \min\{1 + P, 1 + P'\}, & \Omega_{\text{BE}}(D) &:= \frac{1}{2} \mathbb{E} \min\{P, P'\}, \\ \Omega_{\text{BO},u}(D) = \Omega_{\text{RO},u}(D) &:= \frac{1}{2} \mathbb{E} \min\{\lambda_u(P), \lambda_u(P')\}. \end{aligned} \quad (9)$$

The last coefficient is common to blind and revealing optimization; their online benchmarks differ because only a revealing test exposes p_j before execution.

Corollary 2.3 (Empirical effective-length identity). *Let Λ, Λ' be independent uniform draws from the multiset of effective lengths $\{\lambda_1, \dots, \lambda_n\}$. Then*

$$\text{OPT} = \frac{n^2}{2} \mathbb{E} \min\{\Lambda, \Lambda'\} + \frac{n}{2} \mathbb{E} \Lambda. \quad (10)$$

For a multiset M and the effective-length map ℓ of the chosen model,

$$\text{OPT}(M) = n^2 \Omega_\ell(D[M]) + \frac{n}{2} \int \ell(p) dD[M](p). \quad (11)$$

In particular, in BO(u) or RO(u),

$$\text{OPT}_u(M) = n^2 \Omega_{\text{RO},u}(D[M]) + \frac{n}{2} \int \lambda_u(p) dD[M](p). \quad (12)$$

Proof. Interpret the two sums in Equation (6) as expectations under independent uniform draws from the occurrences. Substituting $\lambda_j = \ell(p_j)$ gives Equation (11); the capped specialization uses $\ell = \lambda_u$. $\square$

The revealing-optimization and blind-optimization models have the same offline effective lengths. Their difference is entirely online: a test reveals p_j , whereas optimization does not. Blind execution is different even offline, because a clairvoyant scheduler can run a job for p_j without paying the unit test. Thus

$$\text{OPT}_{\text{BE}}(M) = n^2 \Omega_{\text{BE}}(D[M]) + \frac{n}{2} \int p dD[M](p) = \sum_i p_i + \sum_{i < j} \min\{p_i, p_j\}. \quad (13)$$

This is not the principal benchmark in Theorem 7.1. The clairvoyant value may be $o(n^2)$, whereas that theorem compares policies at the n^2 scale with the optimal announced-multiset online value $\Phi_{\text{BE}}(D[M])$.

For obligatory testing, Equations (4) and (5) specialize to

$$\text{OPT}_{\text{OT}} = \sum_{k=1}^n \left(k + \sum_{i=1}^k p_{(i)} \right), \quad (14)$$

$$= \sum_i (1 + p_i) + \sum_{i < j} (1 + \min\{p_i, p_j\}). \quad (15)$$

In particular,

$$\text{OPT}_{\text{OT}} \geq \frac{n(n+1)}{2}. \quad (16)$$

For $\text{BO}(u)$ and $\text{RO}(u)$, $\lambda_j \geq \min\{u, 1\}$ gives

$$\text{OPT}_u \geq \min\{u, 1\} \frac{n(n+1)}{2}. \quad (17)$$

There is deliberately no analogous positive lower bound for blind execution: an input consisting only of zero jobs has clairvoyant cost zero.

We also isolate the random-order identity used by several revealing arguments. For $p = (p_1, \dots, p_n)$, a set $E \subseteq [n]$ is a *processing-time prefix* if every value in E is at most every value outside it. Suppose that E contains all zero jobs, put $T = [n] \setminus E$, and use the normalized quantities

$$e = |E|, \quad a = \frac{e}{n}, \quad h = 1 - a, \quad \ell_E = \frac{1}{n} \sum_{i \in E} p_i, \quad K_X = \frac{1}{n^2} \sum_{i, j \in X} \min\{p_i, p_j\}. \quad (18)$$

The prefix property gives

$$K(D[p]) = K_E + 2h\ell_E + K_T, \quad K(D) := \iint \min\{x, y\} dD(x)dD(y). \quad (19)$$

Define its stationary leading coefficient by

$$F_{\text{OT}, D[p]}(E) = (1 + \ell_E) \left(1 - \frac{a}{2} \right) + \frac{K_T}{2}. \quad (20)$$

Lemma 2.4 (Stationary prefix pair identity). *Test all jobs in a uniformly random order, process a positive E -job immediately after its test, defer every T -job, and finally process T in SPT order. The resulting stationary policy satisfies*

$$\mathbb{E} \text{ALG}_{\text{stat}}(p, E) = n^2 F_{\text{OT}, D[p]}(E) + \frac{e + \sum_i p_i}{2}. \quad (21)$$

Proof. Let $W = n + \sum_{i \in E} p_i$ and $W_E = e + \sum_{i \in E} p_i$. The discovery phase has one block per job: an E -block has length $1 + p_i$ and completes its job, whereas a T -block is its unit test and completes no job. For an E -job, every other block precedes it with probability $1/2$, so all E -jobs contribute

$$\frac{eW + W_E}{2}. \quad (22)$$

Every T -job sees the common offset W , and the final SPT phase contributes

$$(n - e)W + \sum_{i \in T} p_i + \sum_{\substack{i < j \\ i, j \in T}} \min\{p_i, p_j\}. \quad (23)$$

Adding the two displays and collecting the diagonal terms gives Equation (21). $\square$

2.3 Maximum-density modules and fluid completion envelopes

Several randomized and instance-optimal arguments use the same fluid primitive. Testing one unit mass of jobs automatically completes its zero outcomes and may be followed by immediate processing of selected positive outcomes. We isolate that primitive, its optimal density, and the resulting fractional-knapsack envelope here.

Let D be a probability distribution on $[0, \infty)$ with finite mean and let $P \sim D$. For a measurable selector $g(p) \in [0, 1]$ on the positive values, define its completed mass and work by

$$a_D(g) := D(\{0\}) + \mathbb{E}[g(P)\mathbf{1}_{P>0}], \quad w_D(g) := 1 + \mathbb{E}[Pg(P)\mathbf{1}_{P>0}]. \quad (24)$$

The zero mass is included automatically because its test completes it.

Lemma 2.5 (Maximum-density discovery module). *There is a unique $\tau_D \geq 1$ satisfying*

$$\mathbb{E}(\tau_D - P)^+ = 1. \quad (25)$$

Choose g_ to select every positive $p < \tau_D$, reject every $p > \tau_D$, and select an arbitrary fraction of an atom at $p = \tau_D$. Put*

$$a_* = a_D(g_*), \quad w_* = w_D(g_*), \quad d\nu(p) = (1 - g_*(p))\mathbf{1}_{p>0} dD(p), \quad (26)$$

with $\nu(\{0\}) = 0$. Then

$$\frac{a_*}{w_*} = \max_g \frac{a_D(g)}{w_D(g)} = \frac{1}{\tau_D}, \quad \tau_D a_D(g) \leq w_D(g) \quad \text{for every } g. \quad (27)$$

Moreover, ν has mass $1 - a_$ and is supported on $[\tau_D, \infty)$.*

Proof. The function $h(t) = \mathbb{E}(t - P)^+$ is continuous and nondecreasing, satisfies $h(0) = 0$, and tends to infinity because $h(t) \geq t - \mathbb{E}P$. Once positive it is strictly increasing, so $h(t) = 1$ has a unique solution; $h(t) \leq t$ gives $\tau_D \geq 1$.

For an arbitrary selector g ,

$$\tau_D a_D(g) - \mathbb{E}[Pg(P)\mathbf{1}_{P>0}] = \tau_D D(\{0\}) + \mathbb{E}[(\tau_D - P)g(P)\mathbf{1}_{P>0}] \leq \mathbb{E}(\tau_D - P)^+ = 1.$$

This is the dual inequality in Equation (27). The threshold selector g_* attains equality because it keeps exactly the terms with $\tau_D - p > 0$; mass at $p = \tau_D$ does not affect the equality. Hence $\tau_D a_* = w_*$ and g_* has maximum density $1/\tau_D$. Its unselected mass is $1 - a_*$ and can occur only at values at least τ_D . $\square$

Corollary 2.6 (Empirical maximum-density prefix). *For an empirical distribution $D[p]$ from Equation (7), the maximum of*

$$\frac{|E|/n}{1 + n^{-1} \sum_{i \in E} p_i} \quad (28)$$

over processing-time prefixes E containing all zero jobs is $1/\tau_{D[p]}$. It is attained by taking every occurrence below $\tau_{D[p]}$, no occurrence above it, and any subset of occurrences equal to $\tau_{D[p]}$. Thus every selected value is at most $\tau_{D[p]}$ and every unselected value is at least $\tau_{D[p]}$.

Proof. Apply Lemma 2.5 to $D[p]$. At the boundary $p_i = \tau_{D[p]}$, adding or removing an occurrence preserves the ratio $1/\tau_{D[p]}$, so the fractional boundary selector may be replaced by any subset of the tied occurrences. $\square$

We will also use the continuous analogue of shortest-processing-time pair accounting. Let $\mathcal{M} = \sum_k c_k \delta_{v_k}$ be a finitely supported probability measure on $[0, \infty)$. Its divisible completion envelope is

$$K_{\mathcal{M}}(x) := \max \left\{ \sum_k z_k : 0 \leq z_k \leq c_k, \sum_k v_k z_k \leq x \right\}. \quad (29)$$

Lemma 2.7 (Divisible SPT area). *The envelope $K_{\mathcal{M}}$ processes the items in nondecreasing order of v_k , and*

$$\int_0^\infty (1 - K_{\mathcal{M}}(x)) dx = \frac{1}{2} \iint \min\{r, s\} d\mathcal{M}(r) d\mathcal{M}(s). \quad (30)$$

Proof. The ordering assertion is the usual fractional-knapsack exchange. In a block of completion mass c_k and work per completion v_k , unfinished mass decreases linearly. After sorting the blocks by v_k , its area contribution is

$$v_k c_k \left(1 - \sum_{h < k} c_h - \frac{c_k}{2} \right).$$

Summing these terms and collecting each unordered pair gives Equation (30). $\square$

Assume from now on that D has finite support and write $d_p = D(\{p\})$. Fix a final tested fraction $q \in [0, 1]$ and an independent completion mode that uses $v \geq 0$ work per completed job. At a fluid schedule prefix, let t be tested mass, b mass completed by the independent mode, and c_p processed tested mass of positive class p . Define

$$\mathcal{C}_{D,q}^v(x) := \max \left\{ d_0 t + b + \sum_{p>0} c_p : \begin{array}{ll} 0 \leq t \leq q, & 0 \leq b \leq 1 - q, \\ 0 \leq c_p \leq d_p t, & \\ t + vb + \sum_{p>0} p c_p \leq x \end{array} \right\}. \quad (31)$$

Lemma 2.8 (Test-module completion envelope). *Let (a_*, τ_D, ν) be supplied by Lemma 2.5. Then*

$$\mathcal{C}_{D,q}^v = K_{\mathcal{M}_{D,q,v}}, \quad \mathcal{M}_{D,q,v} := q a_* \delta_{\tau_D} + q \nu + (1 - q) \delta_v. \quad (32)$$

Consequently,

$$\int_0^\infty (1 - \mathcal{C}_{D,q}^v(x)) dx = \frac{1}{2} \iint \min\{r, s\} d\mathcal{M}_{D,q,v}(r) d\mathcal{M}_{D,q,v}(s). \quad (33)$$

The three terms in $\mathcal{M}_{D,q,v}$ are respectively the maximum-density test module, the residual tested classes, and the independent completion mode.

Proof. Split every positive support class virtually into its g_* -selected and residual parts. For a feasible point in Equation (31), let y_L, x_L be the completion and work contributed by the tests, tested zeros, and processed selected outcomes. Applying Equation (27) to the processed fractions within the first t units of tests gives

$$\tau_D y_L \leq x_L, \quad y_L \leq a_* t.$$

With $\lambda = y_L/a_*$, replace this part by λ full test modules. The replacement preserves completed mass, weakly decreases work, and uses $0 \leq \lambda \leq t \leq q$. Residual class p has capacity at most $q\nu(\{p\})$, while the independent mode has capacity $1 - q$. Thus every feasible point maps to the divisible knapsack in Equation (32).

Conversely, order the knapsack items by work per completion, placing the test module before residual items at a tie. The independent item can be run whenever its position in this order requires. A partial test-module item is realized by the corresponding fraction of tests and selected outcomes. Once all q test modules have run, every residual capacity is available; this is early enough because Lemma 2.5 gives $\text{supp } \nu \subseteq [\tau_D, \infty)$. Hence the greedy curve is feasible at every work budget, proving the envelope identity. The area formula follows from Lemma 2.7. $\square$

Lemma 2.9 (Zero-preserving transport). *Suppose distributions D, D' admit a coupling (P, P') such that $P = 0$ if and only if $P' = 0$ and $|P - P'| \leq h$ almost surely. Transporting the same fluid policy through this coupling preserves all completion events and changes accumulated processing work by at most h . Consequently, each model-specific fluid value Φ defined below satisfies*

$$|\Phi(D) - \Phi(D')| \leq h. \quad (34)$$

Proof. Couple each infinitesimal job occurrence and transport decisions using its coupled mark, exposing that mark only when the model exposes the corresponding processing time. Zero preservation makes test completions coincide. At every prefix, the total mass that has received processing is at most one, so replacing its processing times changes cumulative work by at most h . One unfinished-mass curve is therefore contained in a horizontal h -shift of the other, and its area changes by at most h . Taking the infimum over policies gives one inequality; exchanging D, D' gives the other. $\square$

2.4 Competitive-ratio conventions

Every asymptotic statement in this paper allows a uniform additive $o(n^2)$ term. Formally, a policy family $\mathcal{A}$ has asymptotic competitive ratio at most R if there is a deterministic sequence $\varepsilon(n) \rightarrow 0$ such that every n -job instance satisfies

$$\mathbb{E} \text{ALG}_{\mathcal{A}}(I) \leq R \text{OPT}(I) + \varepsilon(n)n^2. \quad (35)$$

For a deterministic policy the expectation is omitted. The optimal ratio in a model is the infimum of such R over all admissible policy families.

For obligatory testing and for either model with fixed raw time $u > 0$, Equations (16) and (17) show that OPT is bounded below by a positive constant times n^2 . In these models the definition above is equivalent to the usual worst-case ratio after taking $n \rightarrow \infty$. This equivalence fails for blind execution, where OPT can be $o(n^2)$. Accordingly, Theorem 7.1 makes the absolute statement $\mathbb{E} \text{ALG} = n^2 \Phi_{\text{BE}}(D[M]) + o_L(n^2)$ and does not claim a bounded ratio to the clairvoyant optimum on every sparse input.

2.5 Test-order adversaries

The lower bound will assign processing times according to the order in which tests finish. This is only a proof device.

Lemma 2.10 (Replay). *Fix a deterministic online algorithm. Suppose all untested jobs have the same known parameters, and an adversary assigns processing times deterministically from the transcript and test-completion order. The resulting interaction is reproduced by a fixed input instance.*

Proof. Run the interaction once and record the value revealed for every tested job label. Assign those values to the corresponding labels in advance. An induction over operation completions shows that the deterministic algorithm chooses the same tests, processing operations, and idle intervals on the fixed instance. If the algorithm fails to test or complete some job forever, its competitive ratio is already infinite; otherwise every relevant label is recorded. $\square$

2.6 Probabilistic tools

We collect here the concentration estimates used in the randomized sections. All of them are finite-space statements. We use without further mention the union bound, Markov's inequality $\mathbb{P}(Y \geq t) \leq \mathbb{E}Y/t$ for $Y \geq 0$, and the tail identity $\mathbb{E}Y = \int_0^\infty \mathbb{P}(Y \geq t) dt$.

The next deferred-decisions lemma turns adaptive label choices into sampling without replacement. Occurrence tokens distinguish equal processing times.

Lemma 2.11 (Adaptive first-touch permutation). *Let J be n public labels, let Ξ be n occurrence tokens, and let a placement $\sigma : J \rightarrow \Xi$ be a bijection. Fix a deterministic nonanticipating policy that eventually first-touches every label. Write $\ell_1(\sigma), \dots, \ell_n(\sigma)$ for its labels in order of first touch and define*

$$R_A(\sigma)_k = \sigma(\ell_k(\sigma)). \quad (36)$$

Then $\sigma \mapsto R_A(\sigma)$ is a bijection from placements to permutations of Ξ . The action selected at first touch k is a function of the preceding tokens and is therefore predictable with respect to the analytically exposed token order. Consequently, a uniform placement induces a uniform random permutation of the occurrences.

Proof. Suppose two placements induce the same occurrence order. A lockstep induction shows that the runs have the same observable history before each first touch: the preceding tokens and actions are the same, and any observation made by the model is determined by them. The deterministic policy therefore performs the same intervening operations, chooses the same new label, and selects the same action. Thus all first-touch labels agree. Equation (36) then makes the placements equal on every label. The map is injective between two sets of size $n!$, hence bijective. The same induction stopped before touch k proves predictability. Exposing a token value that the policy itself does not see only enriches the analytical filtration and does not affect the conclusion. $\square$

Lemma 2.12 (Chernoff bounds). *Let $X = \sum_{i=1}^m X_i$, where the X_i are independent Bernoulli random variables, and write $\mu = \mathbb{E}X$. For every $\delta > 0$,*

$$\mathbb{P}(X \geq (1 + \delta)\mu) \leq \left(\frac{e^\delta}{(1 + \delta)^{1+\delta}} \right)^\mu \leq \exp\left(-\frac{\delta^2 \mu}{2 + \delta} \right). \quad (37)$$

For $0 \leq \delta \leq 1$,

$$\mathbb{P}(X \leq (1 - \delta)\mu) \leq \exp\left(-\frac{\delta^2\mu}{2}\right). \quad (38)$$

These are the standard exponential-moment bounds of Chernoff [Che52].

Lemma 2.13 (Sampling without replacement). *Let $x_1, \dots, x_N \in [0, 1]$, let $X_1, \dots, X_N$ be a uniformly random permutation of these numbers, and put $\bar{x} = N^{-1} \sum_i x_i$. For $1 \leq k \leq N$ and $t \geq 0$,*

$$\mathbb{P}\left(\left|\sum_{i=1}^k X_i - k\bar{x}\right| \geq t\right) \leq 2 \exp\left(-\frac{2t^2}{k}\right). \quad (39)$$

Moreover, if $\hat{x}_k = k^{-1} \sum_{i=1}^k X_i$ and $\sigma^2 = N^{-1} \sum_i (x_i - \bar{x})^2$, then, for $N > 1$,

$$\text{Var}(\hat{x}_k) = \frac{N - k}{k(N - 1)} \sigma^2 \leq \frac{1}{4k}. \quad (40)$$

If the population is partitioned into d classes with population histogram p and sample histogram $\hat{p}$, the same variance formula and Cauchy–Schwarz give

$$\mathbb{E}\|\hat{p} - p\|_1 \leq \sqrt{\frac{d}{k}}. \quad (41)$$

Finally, if $H_{k,j}$ counts class j among the first k entries, then

$$\mathbb{P}\left(\max_{0 \leq k \leq N} \max_{1 \leq j \leq d} |H_{k,j} - kp_j| \geq t\right) \leq 2d(N + 1) \exp\left(-\frac{2t^2}{N}\right), \quad (42)$$

$$\mathbb{E} \max_{0 \leq k \leq N} \max_{1 \leq j \leq d} |H_{k,j} - kp_j| \leq C\sqrt{N \ln(2d(N + 1))} \quad (43)$$

for a universal constant C .

Proof. The estimate Equation (39) is Hoeffding’s sampling-without-replacement inequality [Hoe63, Section 6]. The variance identity is the elementary finite-population correction. For a class of mass p_j , it gives $\mathbb{E}|\hat{p}_j - p_j| \leq \sqrt{p_j/k}$; summing and applying Cauchy–Schwarz proves Equation (41). Apply Equation (39) to every binary class indicator and union-bound over k and j to obtain Equation (42). Integrating that tail bound proves Equation (43). $\square$

Lemma 2.14 (Martingale maximal inequalities). *Let $(M_k, \mathcal{F}_k)_{k=0}^N$ be a real martingale with $M_0 = 0$. If it is square integrable, then Doob’s inequality gives*

$$\mathbb{E} \max_{0 \leq k \leq N} |M_k|^2 \leq 4\mathbb{E}|M_N|^2. \quad (44)$$

If instead $|M_k - M_{k-1}| \leq b_k$ almost surely for deterministic b_k , then maximal Azuma–Hoeffding gives, for every $t \geq 0$,

$$\mathbb{P}\left(\max_{0 \leq k \leq N} |M_k| \geq t\right) \leq 2 \exp\left(-\frac{t^2}{2 \sum_{k=1}^N b_k^2}\right). \quad (45)$$

We use the classical forms due to Doob [Doo53] and Azuma [Azu67]; the maximal version of the latter follows by applying the exponential-supermartingale proof at the first crossing time.

Lemma 2.15 (Predictable sampling from a random permutation). *Let $X_1, \dots, X_N$ be a uniform random permutation of fixed vectors $x_1, \dots, x_N \in [0, B]^d$, and let $\bar{x} = N^{-1} \sum_i x_i$. If $A_k \in [0, 1]$ is measurable before X_k is exposed, then for $\delta, \eta \in (0, 1)$, with probability at least $1 - \eta$, simultaneously for every coordinate ℓ and every $m \leq (1 - \delta)N$,*

$$\left| \sum_{k \leq m} A_k(X_{k,\ell} - \bar{x}_\ell) \right| \leq (1 + \delta^{-1})E_{N,d,B,\eta}, \quad E_{N,d,B,\eta} := B \sqrt{2N \ln \left(\frac{4d(N+1)}{\eta} \right)}. \quad (46)$$

The statement remains valid after conditioning on independent private randomness used to choose the predictable sequence.

Proof. For each coordinate, apply Equation (39) after scaling by B , and union-bound over coordinates and prefix lengths. Except with probability at most $\eta/2$, every centered prefix sum has absolute value at most $E = E_{N,d,B,\eta}$. Before draw $k \leq (1 - \delta)N$, the mean $\tilde{x}_{k-1,\ell}$ of the remaining urn therefore satisfies

$$|\tilde{x}_{k-1,\ell} - \bar{x}_\ell| \leq \frac{E}{\delta N}.$$

For each ℓ ,

$$M_{m,\ell} = \sum_{k \leq m} A_k(X_{k,\ell} - \tilde{x}_{k-1,\ell})$$

is a martingale with increments bounded by B . The maximal Azuma–Hoeffding inequality Equation (45), followed by a union bound over ℓ , puts all $|M_{m,\ell}|$ below E except with probability at most $\eta/2$. The predictable drift is at most E/δ , which proves Equation (46). Conditioning and then averaging proves the final assertion. $\square$

Lemma 2.16 (Lipschitz empirical minimization). *Let Π be a common family of templates and let $F_D(\pi)$ be their objectives for distributions D . Suppose that, for some metric d ,*

$$\sup_{\pi \in \Pi} |F_D(\pi) - F_E(\pi)| \leq Ld(D, E). \quad (47)$$

If $\Phi(D) = \min_{\pi \in \Pi} F_D(\pi)$, then

$$|\Phi(D) - \Phi(E)| \leq Ld(D, E). \quad (48)$$

Moreover, if π_E minimizes F_E , then

$$F_D(\pi_E) \leq \Phi(D) + 2Ld(D, E). \quad (49)$$

Proof. Evaluating the E -objective at a minimizer for D , and conversely, proves Equation (48). For the second claim, insert $F_E(\pi_E) = \Phi(E)$ between $F_D(\pi_E)$ and $\Phi(D)$ and apply Equations (47) and (48). $\square$

Lemma 2.17 (Finite Yao selection). *Let ν be a finite-support distribution on inputs, and let $g(A, I)$ be an integrable real payoff. If*

$$\mathbb{E}_{I \sim \nu} g(A, I) \geq L \quad (50)$$

for every deterministic strategy A , then for every randomized strategy $\mathcal{A}$ there is a fixed I in the support of ν such that

$$\mathbb{E}_{A \sim \mathcal{A}} g(A, I) \geq L. \quad (51)$$

Proof. Condition on the random seed of $\mathcal{A}$ and apply Equation (50). Averaging over the seed shows that the ν -average of $\mathbb{E}_{A \sim \mathcal{A}} g(A, I)$ is at least L , so at least one input in the finite support attains this average. This is the one-sided finite form of Yao’s principle [Yao77] used in our lower bounds. $\square$

2.7 Notation and fluid-to-finite transfer

We use one notation hierarchy in all four models. An uppercase Roman D denotes a probability law, and $D[I]$ is the empirical law Equation (7) of a finite input object I . Calligraphic uppercase letters denote curves, sets, or policies. For a model M , an input law D , and a fluid policy template π , we write

$$\mathcal{C}_{M,D,\pi}(x) = \text{maximum completed mass by work } x, \quad F_{M,D}(\pi) = \int_0^\infty (1 - \mathcal{C}_{M,D,\pi}(x)) dx. \quad (52)$$

Uppercase Greek letters are leading n^2 -coefficients:

$$\Omega_M(D) = \text{offline fluid cost}, \quad \Phi_M(D) = \inf_{\pi} F_{M,D}(\pi) = \text{instance-optimal online value}. \quad (53)$$

Model parameters such as u are appended to the model subscript when needed. Finally, R_M^{det} and R_M^{rand} denote worst-case ratios. We reserve R for a candidate ratio and $r = R - 1$ for its excess. Lowercase letters denote local scalars; in particular a is completion mass, w is work, q is a tested fraction, μ is a mean, κ is completion density, and $\tau = 1/\kappa$ is work per completion. Specialized formulas may suppress a template or a fixed model parameter, but retain these semantic roles.

The following reusable results isolate the finite-size argument shared by the instance-optimal sections. They deliberately separate the model-specific task—constructing a fluid envelope and repairing a finite prefix into a feasible fluid point—from the generic completion-area and learning steps.

Lemma 2.18 (Uniform first-touch sampling event). *Fix values $p_1, \dots, p_n \in [0, L]$, partition their occurrence tokens into $s + 1$ cells with empirical masses $d_0, \dots, d_s$, and put $\mu = n^{-1} \sum_i p_i$. Place the tokens uniformly on the public labels, fix a deterministic policy, and let X_k be the value encountered at first touch k . If A_k is the predictable indicator that this touch uses the revealing mode, then for every $\delta, \eta \in (0, 1)$, with probability at least $1 - \eta$, simultaneously for every $m \leq (1 - \delta)n$,*

$$\left| \sum_{k \leq m} A_k \mathbf{1}_{X_k \in j} - d_j \sum_{k \leq m} A_k \right| \leq \Gamma_{\text{cell}} \quad (0 \leq j \leq s), \quad (54)$$

$$\left| \sum_{k \leq m} (1 - A_k) X_k - \mu \sum_{k \leq m} (1 - A_k) \right| \leq \Gamma_{\text{work}}, \quad (55)$$

where

$$\Gamma_{\text{cell}} = (1 + \delta^{-1}) E_{n,s+1,1,\eta/2}, \\ \Gamma_{\text{work}} = (1 + \delta^{-1}) E_{n,1,L,\eta/2}.$$

The statement remains true after conditioning on an independent policy seed.

Proof. By Lemma 2.11, the first-touch values form a uniform permutation and A_k is predictable. Apply Lemma 2.15 to the cell-indicator vector with selector A_k and to the scalar value with selector $1 - A_k$, assigning failure probability $\eta/2$ to each application. $\square$

Lemma 2.19 (Bounded suffix edit). *Consider two completing no-idling schedules on the same n jobs, each of total work at most Wn . Suppose one is obtained from the other by changing the operations of at most h jobs, altering at most ch operations of individual*

length at most W , while preserving the relative order and completion events of all unaffected jobs. Then

$$|\text{cost}(\mathcal{A}) - \text{cost}(\tilde{\mathcal{A}})| \leq (c+1)Wn. \quad (56)$$

Proof. Write completion cost as operation length times the number of unfinished jobs. Altered operations contribute at most $chWn$. At every unaffected operation, the unfinished counts in the two schedules differ only on the h affected jobs; summing this discrepancy over total work at most Wn adds at most hWn . $\square$

Theorem 2.20 (Fluid-to-finite completion transfer). *Fix a law D and a family of templates Π . Suppose every envelope $\mathcal{C}_{D,\pi}$ is nondecreasing, takes values in $[0, 1]$, and equals one for $x \geq H$, where $H < \infty$ is common to $\pi \in \Pi$. Put*

$$F_D(\pi) = \int_0^\infty (1 - \mathcal{C}_{D,\pi}(x)) dx, \quad \Phi(D) = \inf_{\pi \in \Pi} F_D(\pi), \quad (57)$$

and assume $0 \leq \Phi(D) \leq B$.

Let $\mathcal{A}$ be a random finite schedule on n jobs. Suppose that on an event G there are an edited completing schedule $\tilde{\mathcal{A}}$ and a possibly random template $\pi \in \Pi$ such that

$$\text{cost}(\mathcal{A}) \geq \text{cost}(\tilde{\mathcal{A}}) - \varepsilon n^2, \quad (58)$$

$$\frac{N_{\tilde{\mathcal{A}}}(nx)}{n} \leq \mathcal{C}_{D,\pi}(x + \alpha) + \beta \quad (0 \leq x \leq H), \quad (59)$$

where $N_{\tilde{\mathcal{A}}}(t)$ is the number of jobs completed by time t . If $\mathbb{P}(G^c) \leq \eta$, then

$$\frac{\mathbb{E} \text{cost}(\mathcal{A})}{n^2} \geq \Phi(D) - \varepsilon - \alpha - H\beta - \eta B. \quad (60)$$

The same conclusion holds seedwise under a uniform finite placement; in that case, for each randomized policy, finite Yao selection produces one fixed oblivious labeling satisfying the bound.

Proof. Total completion time is the area under the unfinished-job curve. On G , Equation (59) and the fact that the envelope is one after H give

$$\begin{aligned} \frac{\text{cost}(\tilde{\mathcal{A}})}{n^2} &\geq \int_0^H (1 - \mathcal{C}_{D,\pi}(x + \alpha) - \beta) dx \\ &\geq F_D(\pi) - \alpha - H\beta \geq \Phi(D) - \alpha - H\beta. \end{aligned}$$

Use Equation (58), average over G , and use nonnegativity on G^c . The lost factor is at most $\eta\Phi(D) \leq \eta B$, proving Equation (60). The final assertion is Lemma 2.17 applied after conditioning on the policy seed. $\square$

Lemma 2.21 (Cell-capacity repair). *Let D have support $0 = p_0 < p_1 < \dots < p_s$ with masses d_j , and consider the common envelope $\mathcal{C}_{D,q}^v$ from Equation (31). At a finite schedule prefix, let t, b, z be normalized tested mass, independent-mode mass, and zero completions, and let c_j be normalized processed mass of positive class j . Suppose*

$$t \leq q, \quad b \leq 1 - q, \quad z \leq d_0 t + \gamma, \quad c_j \leq d_j t + \gamma, \quad (61)$$

and, with $c'_j = (c_j - \gamma)^+$,

$$t + vb + \sum_{j=1}^s p_j c'_j \leq x + \alpha. \quad (62)$$

Then

$$z + b + \sum_{j=1}^s c_j \leq \mathcal{C}_{D,q}^v(x + \alpha) + (s + 1)\gamma. \quad (63)$$

Proof. The repaired variables t, b, c'_j satisfy every constraint in Equation (31) at work budget $x + \alpha$. Replacing z by $d_0 t$ and each c_j by c'_j loses at most $(s + 1)\gamma$ completed mass. $\square$

Lemma 2.22 (Pilot-learning transfer). *Let Π be a common template family, suppose $\Phi(D) = \inf_{\pi \in \Pi} F_D(\pi)$, and assume*

$$\sup_{\pi \in \Pi} |F_D(\pi) - F_E(\pi)| \leq L_0 d(D, E). \quad (64)$$

Let $\hat{D}$ be a pilot estimate, D^{rem} the law of the remaining jobs, and let $\hat{\pi}$ minimize $F_{\hat{D}}$. If the complete learned schedule, including its pilot, satisfies conditionally

$$\mathbb{E}[\text{cost}(\mathcal{A}) \mid \hat{D}, D^{\text{rem}}] \leq n^2 F_{D^{\text{rem}}}(\hat{\pi}) + \varepsilon n^2, \quad (65)$$

then

$$\frac{\mathbb{E} \text{cost}(\mathcal{A})}{n^2} \leq \Phi(D) + 2L_0 \mathbb{E}d(\hat{D}, D^{\text{rem}}) + L_0 \mathbb{E}d(D^{\text{rem}}, D) + \varepsilon. \quad (66)$$

Proof. By Lemma 2.16, $F_{D^{\text{rem}}}(\hat{\pi}) \leq \Phi(D^{\text{rem}}) + 2L_0 d(\hat{D}, D^{\text{rem}})$. The value part of the same lemma gives $\Phi(D^{\text{rem}}) \leq \Phi(D) + L_0 d(D^{\text{rem}}, D)$. Substitute both estimates in Equation (65) and average. $\square$

3 Deterministic revealing optimization: the exact curve

The goal of this section is to prove Theorem 3.1, the rigorous deterministic part of the informal revealing-optimization curve theorem Theorem 1.7. We now allow raw execution for a finite common time u . All jobs have this same known upper limit, and a test reveals $p_j \in [0, u]$. By Equation (2) and Lemma 2.2, the offline effective length is $\lambda_u(p_j) = \min\{u, 1 + p_j\}$, and the offline optimum schedules these effective lengths in nondecreasing order. We write $R_{\text{RO}}^{\text{det}}(u)$ for the optimal deterministic size-asymptotic ratio at this fixed common upper limit.

The answer has six branches separated by five transition points. Put

$$\varphi = \frac{1 + \sqrt{5}}{2}, \quad u_1^{\text{det}} = 1, \quad u_2^{\text{det}} = \frac{1 + \sqrt{3 + 2\sqrt{5}}}{2} = 1.866760399173862 \dots \quad (67)$$

Let $s_0 > 1$ be the root of

$$s_0^3 = (s_0 + 1)^2, \quad u_3^{\text{det}} = 1 + s_0 = 3.147899035704787 \dots \quad (68)$$

The fourth transition point is

$$u_4^{\text{det}} = \varphi + 2 = 3.618033988749895 \dots \quad (69)$$

For $s = u - 1$, put

$$T_s(z) = sz^2 + (s^3 + s^2 + 1)z - (s^2 + s + 1)(s + 2), \quad (70)$$

and let $R_{\text{quad}}(u)$ be the positive root of

$$sR^2 + (s^3 + s^2 + 1)R - (s^2 + s + 1)(s + 2) = 0. \quad (71)$$

Denote the left-hand side by $T_s(R)$. Equivalently,

$$R_{\text{quad}}(u) = \frac{-1 - u + 2u^2 - u^3 + \sqrt{u^6 - 4u^5 + 10u^4 - 6u^3 - 3u^2 + 6u - 3}}{2(u - 1)}. \quad (72)$$

At the other end, let $u_5^{\text{det}} > 1$ solve $u_5^{\text{det}} - 3 = \ln u_5^{\text{det}}$ and define

$$\begin{aligned} r &= \frac{2}{u_5^{\text{det}} - 1}, & R_{\text{OT}}^{\text{det}} &= 1 + r = 1.570574096649395\dots, \\ u_5^{\text{det}} &= 4.505241495792883\dots \end{aligned} \quad (73)$$

For $u_4^{\text{det}} \leq u \leq u_5^{\text{det}}$, define $R_{\text{mix}}(u) = 1 + c$ by the unique solution (c, m) of

$$\text{artanh } m - m = 1 - \frac{1}{c} + \frac{1}{2} \ln \left(1 + \frac{2}{c} \right), \quad u = 1 + \frac{2}{c} - m, \quad (74)$$

with $r \leq c \leq 1/\varphi$ and $0 \leq m \leq 1/\varphi$.

Theorem 3.1 (Exact revealing-optimization curve). *For every fixed $0 < u < \infty$, the optimal deterministic size-asymptotic ratio in the revealing-optimization model is*

$$R_{\text{RO}}^{\text{det}}(u) = \begin{cases} 1, & 0 < u \leq u_1^{\text{det}}, \\ u, & u_1^{\text{det}} \leq u \leq u_2^{\text{det}}, \\ R_{\text{quad}}(u), & u_2^{\text{det}} \leq u \leq u_3^{\text{det}}, \\ 1 + \frac{1}{\sqrt{u-1}}, & u_3^{\text{det}} \leq u \leq u_4^{\text{det}}, \\ R_{\text{mix}}(u), & u_4^{\text{det}} \leq u \leq u_5^{\text{det}}, \\ R_{\text{OT}}^{\text{det}}, & u \geq u_5^{\text{det}}. \end{cases} \quad (75)$$

For every branch there is an explicit deterministic algorithm satisfying

$$\text{cost}(\mathcal{A}, I) \leq R_{\text{RO}}^{\text{det}}(u) \text{cost}(\text{OPT}, I) + O_u(n).$$

On $u \geq u_5^{\text{det}}$ the remainder is one absolute $O(n)$ term, uniform over all finite u . Conversely, for every deterministic algorithm $\mathcal{A}$ and every $\varepsilon > 0$, there are arbitrarily large fixed instances I such that

$$\text{cost}(\mathcal{A}, I) \geq (R_{\text{RO}}^{\text{det}}(u) - \varepsilon) \text{cost}(\text{OPT}, I).$$

For $u \geq u_5^{\text{det}}$, these lower-bound instances may be chosen cap-free: $p_j < u - 1$ for every job. Thus both witnesses on the plateau are uniform in the sense needed to pass separately to the obligatory endpoint.

The proof combines two kinds of lower bounds. Binary hidden-stopping instances give the first four branches, while a continuous capped version of the harmonic construction gives the mixed branch and the plateau. Three explicit policies match them: raw execution, a forced-prefix uniform threshold, and a history-dependent adaptive threshold. The upper-bound witness can be read from the following policy map. The algorithms themselves are specified formally in algorithms 7 to 9.

range	policy	parameter choice
$0 < u \leq u_2^{\det}$	RAW	none
$u_2^{\det} < u \leq u_3^{\det}$	FORCEDPREFIXUTE	$b = b(u)$ from (76)
$u_3^{\det} < u \leq u_4^{\det}$	FORCEDPREFIXUTE	$b = 0$
$u_4^{\det} < u < u_5^{\det}$	ADAPTIVETHRESHOLD	$c = c(u)$ from (74)
$u_5^{\det} \leq u < \infty$	ADAPTIVETHRESHOLD	$c = r$

3.1 Policy guarantees

3.1.1 Low and intermediate caps

The first policy is formalized in algorithm 7.

Algorithm 7 RAW(J, u)

Input: jobs J and a finite common upper limit u .

- 1: Fix an arbitrary order $j_1, \dots, j_n$.
 - 2: **For** $i = 1, \dots, n$, execute j_i untested for u time.
-

For $u \leq u_2^{\det}$, use RAW. It is optimal when $u \leq u_1^{\det}$; when $u > 1$, every effective length $\lambda_u(p_j)$ from Equation (2) is at least one, so the ratio is at most u .

For the next interval, let $R = R_{\text{quad}}(u)$, $s = u - 1$, and define

$$K = s^2 + s + 1, \quad D = K + sR, \quad b = \frac{K - s^2R}{D}, \quad \chi = \frac{1 - b}{R}, \quad \alpha = \frac{K}{D}. \quad (76)$$

The defining equation for R gives

$$0 \leq b < \chi < \alpha < 1, \quad \chi = \frac{s(s+1)}{D}, \quad b = (s+1)\alpha - s; \quad (77)$$

For completeness, these inequalities follow without numerical root isolation. Since T_s is increasing on positive arguments,

$$T_s\left(\frac{K}{s^2}\right) = \frac{K}{s^3}(-s^3 + s^2 + 2s + 1) \geq 0 \quad (s \leq s_0), \quad (78)$$

and hence $b = (K - s^2R)/D \geq 0$, with equality only at $s = s_0$. Moreover,

$$\alpha - \chi = \frac{1}{D}, \quad \chi - b = \frac{s^2R - 1}{D} > 0. \quad (79)$$

For the last sign,

$$9T_s(5/3) = f(s) := 6s^3 - 12s^2 - 2s - 3 < 0. \quad (80)$$

throughout the present interval. Indeed, on $[4/5, 1]$ the polynomial is decreasing and $f(4/5) = -1151/125$; on $[1, s_0]$ it has one interior minimum, while $f(1) = -11$ and $f(s_0) < f(13/6) = -95/36$. Here $s_0 < 13/6$ follows by substituting $13/6$ in $s^3 - (s+1)^2$. Thus $R > 5/3$, while $s > 4/5$, and $s^2R > 16/15$. Finally $K < D$ gives $\alpha < 1$. In particular, $b = 0$ occurs only at $u = u_3^{\det}$. The remaining two elementary branches use the single procedure in algorithm 8.

Algorithm 8 FORCEDPREFIXUTE(J, u, b)

- Input:** jobs J , finite $u > 1$, and $b \in [0, 1]$.
- 1: Fix a test order $j_1, \dots, j_n$; set $k \leftarrow \lfloor bn \rfloor$, $\theta_u \leftarrow \min\{1, u - 1\}$, and $Q \leftarrow \emptyset$.
 - 2: **For** $i = 1, \dots, n$ **do**:
 - 3: Test j_i for one unit and observe p_{j_i} .
 - 4: **If** $i \leq k$ or $p_{j_i} \leq \theta_u$, process j_i immediately.
 - 5: **Otherwise**, add j_i to Q .
 - 6: Process the jobs of Q in nondecreasing revealed processing time.
-

On $u_2^{\det} \leq u \leq u_3^{\det}$, use FORCEDPREFIXUTE with b from Equation (76). Rounding bn changes the cost by $O_u(n)$.

Here is the complete binary calculation. Adjacent exchanges place as many long jobs as possible in the forced prefix and put all suffix-long jobs before suffix zeros. If x is the total long mass, then the prefix-long mass is $a = \min\{b, x\}$ and the deferred-long mass is $d = (x - b)_+$. After division by n^2 , the leading excess $G(x) = A - RO$ is

$$G(x) = \begin{cases} \frac{1-R}{2} + u \left(x - \frac{x^2}{2} \right) - \frac{Rs}{2} x^2, & 0 \leq x \leq b, \\ \frac{1-R}{2} + u \left(b - \frac{b^2}{2} \right) + (1-b)(x-b) + \frac{s}{2}(x-b)^2 - \frac{Rs}{2} x^2, & b \leq x \leq 1. \end{cases} \quad (81)$$

The second quadratic is strictly concave, has its stationary point at $x = \alpha$, and satisfies $G(\alpha) = 0$; after substituting Equation (76), the latter identity is exactly Equation (71). Since $b < \alpha$, its right derivative at b is positive. The left derivative satisfies $G'_-(b) = G'_+(b) + s(1-b) > 0$, and the first quadratic is concave, so it is increasing on $[0, b]$. Thus $G(x) \leq 0$ for every $x \in [0, 1]$, including the limiting case $b = 0$. This proves the binary upper bound without a hidden local-versus-global step.

For $u_2^{\det} \leq u \leq 2$, arbitrary values reduce to binary values by independent Bernoulli extremalization, in the following order. First raise *every* suffix-deferred value to u . This cannot lower ALG, while its offline effective length was already u . Only after this deterministic step, replace a prefix value p by u with probability p/u and by zero otherwise, and replace a suffix-immediate value $p \leq u - 1$ by u with probability p and by zero otherwise. Notice that the u realization of a suffix-immediate job becomes deferred, so status is not preserved. Nevertheless the online check is pairwise. For jobs i before j , the leading pair contribution is

$$g_{ij} = \begin{cases} 1 + p_i, & i \text{ is immediate,} \\ 2 + p_j, & i \text{ is deferred and } j \text{ is immediate,} \\ 2 + \min\{p_i, p_j\}, & i, j \text{ are deferred.} \end{cases} \quad (82)$$

A rounded prefix job stays immediate and preserves its expected value. If i is suffix-immediate, its rounded contribution is one when it becomes zero and at least two when it becomes u , hence its expectation is at least $1 + p_i$. If i is deferred and j is suffix-immediate, then i has been raised to u and the expected pair contribution is $2 + up_j \geq 2 + p_j$; a deferred-deferred pair can only increase. Thus expected ALG does not decrease, up to its $O_u(n)$ diagonal terms.

The expected effective lengths are, respectively,

$$1 + \frac{u-1}{u}p \leq \lambda_u(p), \quad 1 + (u-1)p \leq 1 + p, \quad (83)$$

and u for a deferred value. Since the minimum is jointly concave, $\mathbb{E} \min(\lambda'_i, \lambda'_j) \leq \min(\lambda_i, \lambda_j)$. Thus some binary realization has competitive excess at least that of the original instance.

It remains to justify arbitrary values for $2 \leq u \leq u_3^{\det}$. The following quadratic lemma records the entire endpoint calculation.

Lemma 3.2 (UTE endpoint game). *For $2 \leq u \leq u_3^{\det}$, UTE with the parameters Equation (76) satisfies*

$$\text{ALG} \leq R_{\text{quad}}(u) \text{OPT} + O_u(n). \quad (84)$$

Proof. Fix the symbolic prefix/immediate/deferred decisions. Convexity in one coordinate reduces a prefix job to $p \in \{0, u\}$, a suffix-immediate job to $p \in \{0, 1\}$, and a suffix-deferred job to $p \in \{1^+, u\}$. For the last claim, on $1 < p < u - 1$ the deferred-deferred term has coefficient $1 - R < 0$ multiplying a concave minimum; on $[u - 1, u]$, OPT is flat and ALG is nondecreasing. Adjacent exchanges give the canonical endpoint order.

Normalize counts by n . Let a be prefix- u mass, d suffix- u mass, t suffix- 1^+ mass, and m suffix-immediate-one mass. Their constraints are

$$0 \leq a \leq b, \quad d, t, m \geq 0, \quad d + t + m \leq 1 - b. \quad (85)$$

Pairwise accounting using Equation (82) gives

$$A = \frac{1}{2} + (s + 1) \left(a - \frac{a^2}{2} \right) + (1 - b)(d + t + m) - \frac{m^2}{2} + \frac{s}{2} d^2, \quad (86)$$

$$O = \frac{1}{2} + \frac{(a + d + t + m)^2}{2} + \frac{s - 1}{2} (a + d)^2. \quad (87)$$

Moving m to t does not change O and increases A , so $m = 0$. Put $x = a + d$ and write $F(x) = A - RO$ after optimizing the other variables. The derivative of the gap in t is $1 - b - R(x + t)$, so its maximizing value is $(\chi - x)_+$. At fixed x, t , moving mass from d to a leaves O unchanged and changes A at rate $s(1 - x) + b - a \geq 0$. Hence

$$t = (\chi - x)_+, \quad a = \min\{b, x\}, \quad d = (x - b)_+. \quad (88)$$

For $x \geq \chi$, the gap $A - RO$ is the binary concave quadratic already discussed; it is tangent to zero at $x = \alpha$. For $x < \chi$, put

$$C_0 = -\frac{R - 1}{2} + \frac{(1 - b)^2}{2R}, \quad \gamma = s - R(s - 1). \quad (89)$$

Substitution gives

$$A - RO = C_0 + a(s + b - sx) - \frac{a^2}{2} + \frac{\gamma x^2}{2}. \quad (90)$$

One has $\gamma > 0$ on $1 \leq s \leq s_0$. Indeed, $T_s(2) = s(s^2 - s + 1) > 0$, so for $s \leq 2$ the equation $T_s(R) = 0$ gives $R < 2$, and hence $\gamma > 2 - s \geq 0$. For $2 \leq s \leq s_0 < 13/6$,

$$16T_s(7/4) = 12s^3 - 20s^2 + s - 4 > 0; \quad (91)$$

the polynomial is 14 at $s = 2$ and has positive derivative thereafter. Thus $R < 7/4$ and $\gamma > 7/4 - 3s/4 > 1/8$.

On $0 \leq x \leq b$, one has $a = x$, and

$$A - RO = C_0 + (s + b)x - \frac{s + 1 + R(s - 1)}{2} x^2. \quad (92)$$

Its derivative is decreasing, and the derivative at b multiplied by D is

$$R(1 + s^2 + Rs^2(s - 1)) > 0, \quad (93)$$

so the maximum is at b . On $b \leq x \leq \chi$, Equation (90) is convex and its maximum is at b or χ . The endpoint χ lies on the binary branch. Finally,

$$F(b) - F(\chi) = \frac{\chi - b}{2} [2sb - \gamma(b + \chi)], \quad (94)$$

and direct use of Equation (76) gives

$$D[\gamma(b + \chi) - 2sb] = -s + (1 + s - s^3)R + s^2(s - 1)R^2 > 0. \quad (95)$$

For a rational sign certificate, T_s is increasing on positive arguments and

$$9T_s(5/3) = 6s^3 - 12s^2 - 2s - 3 < 0, \quad (96)$$

so $R > 5/3$. If the right-hand side of Equation (95) is denoted by $q_s(R)$, then

$$3 \partial_R q_s(5/3) = 7s^3 - 10s^2 + 3s + 3 > 0. \quad (97)$$

The last polynomial is increasing for $s \geq 1$, so q_s is increasing for $R \geq 5/3$. Finally,

$$9q_s(5/3) = 10s^3 - 25s^2 + 6s + 15 > 0. \quad (98)$$

For the last sign, write $s = 3/2 + z$; the expression becomes $3/2 - 3z/2 + 20z^2 + 10z^3$. It is at least $3/2$ for $-1/2 \leq z \leq 0$, and for $z \geq 0$ it is larger than $3/2 - 9/320$. These inequalities prove that every endpoint gap is nonpositive. Diagonals and prefix rounding contribute $O_u(n)$. $\square$

At $u = u_3^{\det}$, $b = 0$. The same zero-prefix algorithm remains exact through $u = u_4^{\det}$ and continues with a φ guarantee afterwards. We state the continuation because it makes the three-policy lower envelope in Figure 2 literal rather than schematic.

Lemma 3.3 (Zero-prefix extension). *For every $u \geq u_3^{\det}$, $\text{FORCEDPREFIXUTE}(J, u, 0)$ from algorithm 8 is*

$$\max \left\{ \varphi, 1 + \frac{1}{\sqrt{u - 1}} \right\} \quad (99)$$

competitive up to $O_u(n)$. Thus its guarantee is $1 + 1/\sqrt{u - 1}$ on $[u_3^{\det}, u_4^{\det}]$ and φ on $[u_4^{\det}, \infty)$.

Proof. The same coordinatewise reduction leaves capped-deferred mass d , boundary-deferred mass t , immediate-one mass m , and zeros. With $s = u - 1$, these masses satisfy $d, t, m \geq 0$ and $d + t + m \leq 1$, and the leading costs are

$$A = \frac{1}{2} + d + t + m - \frac{m^2}{2} + \frac{sd^2}{2}, \quad O = \frac{1}{2} + \frac{(d + t + m)^2}{2} + \frac{(s - 1)d^2}{2}. \quad (100)$$

Move m to t and put $y = d + t$. For fixed y , the ratio is fractional-linear in d^2 , so its maximum is on one of two faces:

$$\frac{1 + 2y}{1 + y^2} \leq \varphi \quad (d = 0), \quad 1 + \frac{2y}{1 + sy^2} \leq 1 + \frac{1}{\sqrt{s}} \quad (d = y). \quad (101)$$

The first one-variable function is maximized at $y = 1/\varphi$ and the second at $y = 1/\sqrt{s}$, which gives the two displayed constants. The second value dominates the first exactly when $u \leq u_4^{\det}$. Binary instances attain the second value. $\square$

3.1.2 High caps

We now prove the upper bounds on the two ranges on which capped outcomes matter. Besides completing the upper-bound side of the revealing-optimization curve, the proof records explicitly why an outcome at the raw cap cannot be treated as an ordinary deferred outcome. Throughout this subsection, $R = 1 + c$, and we retain φ , $u_5^{\det}$, and r from Equations (67) and (73).

For every $c > 0$ define the history-dependent threshold

$$a_c(y) = \begin{cases} 1, & y \leq -1, \\ 1 + \frac{1}{2} \ln\left(\frac{c+2}{c-2y}\right), & -1 \leq y \leq 0. \end{cases} \quad (102)$$

The single algorithm used throughout the last two branches is the following. It does not use raw execution and therefore does not depend on u .

Algorithm 9 ADAPTIVETHRESHOLD(J, c)

Input: jobs J and parameter $c > 0$.

- 1: Fix an arbitrary test order $j_1, \dots, j_n$.
 - 2: Set $N \leftarrow 0$, $N_{\text{def}} \leftarrow 0$, and $Q \leftarrow \emptyset$.
 - 3: **For** $i = 1, \dots, n$ **do**:
 - 4: Set $x \leftarrow n - i + 1$ and $y \leftarrow (N_{\text{def}} - (1 + c)N)/x$.
 - 5: Set $a \leftarrow a_c(y)$ according to Equation (102).
 - 6: Test j_i for one unit and observe p_{j_i} .
 - 7: **If** $p_{j_i} \leq a$, process j_i immediately.
 - 8: **Otherwise**, add j_i to Q and set $N_{\text{def}} \leftarrow N_{\text{def}} + 1$.
 - 9: **If** $p_{j_i} > 0$, set $N \leftarrow N + 1$.
 - 10: Process the jobs of Q in nondecreasing order of their revealed processing times.
-

For $c \in [r, 1/\varphi]$, let $m = m(c) \in [0, 1/\varphi]$ and $u = u(c)$ be determined by the common parametrization Equation (74). The right-hand side of the first equation is strictly increasing in c , as is the left-hand side in m . Consequently $m(c)$ is increasing and $u(c)$ is decreasing. The endpoints are

$$(c, m, u) = \left(\frac{1}{\varphi}, \frac{1}{\varphi}, u_4^{\det}\right) \quad \text{and} \quad (c, m, u) = (r, 0, u_5^{\det}).$$

Theorem 3.4 (Middle and high revealing upper bounds). *For every finite $u \in [u_4^{\det}, \infty)$, the formally specified algorithm ADAPTIVETHRESHOLD from algorithm 9 satisfies*

$$\text{ALG} \leq \begin{cases} (1 + c) \text{OPT} + O_u(n), & u_4^{\det} \leq u \leq u_5^{\det}, \quad c \text{ given by (74)}, \\ (1 + r) \text{OPT} + O(n), & u \geq u_5^{\det}. \end{cases}$$

The constant in the last $O(n)$ is absolute and uniform over $u \in [u_5^{\det}, \infty)$. In particular these are upper bounds on the deterministic size-asymptotic competitive ratio.

Proof. Fix a finite cap u in the stated range.

The algorithm. Use ADAPTIVETHRESHOLD(J, c) from algorithm 9; on the high interval use $c = r$. For the analysis, immediately before a test let x be the number of jobs not yet

tested, N the number of earlier strictly positive outcomes, and N_{def} the number of those outcomes deferred to the final tail. Thus

$$y = \frac{N_{\text{def}} - RN}{x} \leq 0 \quad (103)$$

and the threshold is given by Equation (102). On the mixed interval $c \geq r$, and hence

$$a_c(y) \leq a_c(0) \leq a_r(0) = \frac{1}{r} < u - 1.$$

The same inequality holds on the high interval, where we use $c = r$. Thus every outcome in the capped region $[u - 1, u]$ is deferred.

The state y never increases. Indeed, the numerator in (103) changes by 0, $-R$, or $-c$ after a zero, a positive immediate outcome, or a positive deferred outcome, respectively, while x decreases. This monotonicity will allow us to allocate every pair cost when its relevant endpoint becomes known.

Exact pair objective and the five endpoints. Write

$$\bar{p}_i = \min\{p_i, u - 1\}.$$

By Equation (2) and Lemma 2.2, the clairvoyant revealing optimum is SPT on jobs of effective lengths $1 + \bar{p}_i = \min\{1 + p_i, u\}$. For two jobs tested in the order $i < j$, let

$$e_{ij} = \begin{cases} (p_i - p_j)_+, & i \text{ is processed immediately,} \\ 1 + (p_j - p_i)_+, & i \text{ is deferred and } j \text{ is immediate,} \\ 1, & i \text{ and } j \text{ are deferred.} \end{cases} \quad (104)$$

The first line applies whenever i is immediate, regardless of the action taken on j . Pairwise completion-time accounting for the online schedule gives

$$\text{ALG} = \sum_i (1 + p_i) + \sum_{i < j} (1 + \min\{p_i, p_j\} + e_{ij}), \quad (105)$$

whereas the common offline pair identity Equation (5) gives

$$\text{OPT} = \sum_i (1 + \bar{p}_i) + \sum_{i < j} (1 + \min\{\bar{p}_i, \bar{p}_j\}). \quad (106)$$

It is therefore natural to introduce

$$\Gamma_n = \sum_{i < j} (e_{ij} + \min\{p_i, p_j\} - R \min\{\bar{p}_i, \bar{p}_j\}). \quad (107)$$

Equations (105)–(107) yield the exact identity

$$\text{ALG} - R \text{OPT} = \Gamma_n - c \binom{n}{2} + \sum_i [(1 + p_i) - R(1 + \bar{p}_i)]. \quad (108)$$

The final sum is nonpositive. Below the cap its i -th summand is $-c(1 + p_i)$; on the cap it is at most $1 - cu \leq 0$. Thus it is enough to control Γ_n .

Fix a symbolic immediate/deferred path. Coordinatewise convexity in (107) reduces a worst outcome vector to the five formal endpoint types

type	outcome	action
Z	0	immediate
E	0^+	immediate
M	$a_c(y)$	immediate
Q	$a_c(y)^+$	deferred
U	u	deferred and capped.

(109)

For an immediate coordinate the two positive endpoints are 0^+ and the live threshold; the genuine zero is kept separate because it does not enter N . A deferred coordinate below $u - 1$ moves to the live threshold or to $u - 1$. On $[u - 1, u]$, the offline contribution is flat and the algorithmic contribution is nondecreasing, so the latter endpoint moves to u . The symbolic categories, and therefore all later threshold states, remain unchanged during each coordinate move. This is the only endpoint reduction used below.

A local pair allocation. Separate the earlier E -outcomes from the substantive outcomes M, Q, U . Let

$$P = \#\{\text{earlier } M, Q, U\}, \quad E = \#\{\text{earlier } E\}, \quad N_{\text{def}} = \#\{\text{earlier } Q, U\},$$

and let K be the number of earlier U -outcomes. Among the ordinary substantive outcomes, let I, d be the numbers of M, Q -outcomes and let A_I, A_D be the sums of their processing values. Thus $P = I + d + K$ and $N_{\text{def}} = d + K$.

At a current ordinary endpoint, with $a = a_c(y)$, allocate the relaxed local charges

$$\ell_Z = \ell_E = N_{\text{def}}, \quad \ell_M = N_{\text{def}} + a(x + N_{\text{def}} - RP), \quad \ell_Q = N_{\text{def}} + a(N_{\text{def}} - RP). \quad (110)$$

For a capped endpoint the exact charge is instead

$$\begin{aligned} \ell_U &= N_{\text{def}} + A_D + uK - R(A_I + A_D + (u - 1)K) \\ &= d - RA_I - cA_D + Kh_U, \quad h_U = 1 + u - R(u - 1). \end{aligned} \quad (111)$$

These formulae follow by checking the finitely many ordered endpoint pairs. In particular, an earlier immediate outcome already allocated its online processing delay when it appeared, so a current cap contributes only $-Rp$ against that outcome. Against an earlier ordinary deferred outcome it contributes $p - Rp = -cp$, and against an earlier cap it contributes h_U . The same check shows

$$\Gamma_n \leq \sum_i \ell_i; \quad (112)$$

the difference is exactly the sum of the nonnegative own-processing terms allocated by the M -endpoints. Formula (111), rather than a substitution $p = u$ into ℓ_Q , is essential before caps have been put into a prefix.

Threshold monotonicity gives $A_I \geq aI$ and $A_D \geq ad$. Comparing (111) with the hypothetical Q -charge at the same state gives the exact identity

$$\begin{aligned} \ell_U - \ell_Q &= R(aI - A_I) + c(ad - A_D) + K[1 - c(u - 1 - a)] \\ &\leq K[1 - c(u - 1 - a)]. \end{aligned} \quad (113)$$

The reusable four-endpoint bank. The states Z, E, M, Q are the four ordinary, noncap endpoints. The potential below proves their inequalities simultaneously for every target parameter c on the mixed interval. The fifth endpoint U is handled only through the extra reserve. At $c = r$ that reserve vanishes, leaving a cap-free bank that will also support the obligatory endpoint without a second proof. Define

$$\eta = \frac{N_{\text{def}} - RP}{x}, \quad b = \frac{E}{x}, \quad y = \eta - Rb. \quad (114)$$

For $-1 \leq y \leq 0$, put

$$f_c(y) = \frac{c+2}{4} + \frac{c-2y}{4} \left(\ln \frac{c-2y}{c+2} - 1 \right), \quad (115)$$

$$h_c(y) = a_c(y)(1+y),$$

$$G_c(y, b) = f_c(y) + bh_c(y) + \frac{R}{2}b^2. \quad (116)$$

For $y \leq -1$, set instead

$$G_c(y, b) = g_c(\eta), \quad g_c(\eta) = \frac{(1+\eta)_+^2}{2R}. \quad (117)$$

The two definitions are C^1 across $y = -1$. Indeed, feasibility implies $0 \leq b \leq 1/R$ there, and both sides have value $Rb^2/2$ and derivatives

$$(G_y, G_b) = (b, Rb).$$

They are also nonnegative: $f_c(-1) = 0$, $f'_c = a_c - 1 \geq 0$, and $h_c, b, g_c \geq 0$.

Use the base potential

$$W_{\text{base}}(x, P, E, N_{\text{def}}) = xN_{\text{def}} + x^2G_c(y, b). \quad (118)$$

In the raw state $(x, P, E, N_{\text{def}}, K)$, the endpoint directions are

	x	P	E	N_{def}	K
v_Z	-1	0	0	0	0
v_E	-1	0	1	0	0
v_M	-1	1	0	0	0
v_Q	-1	1	0	1	0
v_U	-1	1	0	1	1.

(119)

Let $J_T = \ell_T + \nabla W_{\text{base}} \cdot v_T$, where for the moment $T \in \{Z, E, M, Q\}$. In Region I, direct differentiation and the identities

$$\begin{aligned} -2f_c + (y-R)f'_c + a_c(1+y) &= 0, \\ 1 - 2f_c + (y-c)f'_c + a_c y &= 0 \end{aligned} \quad (120)$$

give

$$\begin{aligned} \frac{J_Z}{x} &= (-2f_c + yf'_c) + b(-h_c + yh'_c), \\ \frac{J_E}{x} &= b[-h_c + (y-R)h'_c + R], \\ \frac{J_M}{x} &= b[-h_c + (y-R)h'_c + a_c R], \end{aligned}$$

$$\frac{J_Q}{x} = b[-h_c + (y - c)h'_c + a_c R]. \quad (121)$$

All four quantities are nonpositive. For the first one,

$$-2f_c + yf'_c = a_c(c - y) - R \leq 0; \quad (122)$$

the right-hand side is zero at $y = -1$ and is nonincreasing because $a'_c(c - y) - a_c \leq 1 - a_c \leq 0$. Also $h_c, h'_c \geq 0$. For the remaining three drifts, observe that

$$h_c + (c - y)h'_c = a_c R + (c - y)(1 + y)a'_c \geq a_c R. \quad (123)$$

The Q -bracket in (121) is therefore nonpositive; the M -bracket is the Q -bracket minus h'_c , and the E -bracket is the M -bracket minus $R(a_c - 1)$.

In Region II, the same calculation gives

$$\begin{aligned} J_Z/x = J_E/x &= -2g_c + \eta g'_c, \\ J_M/x &= -2g_c + (\eta - R)g'_c + 1 + \eta, \\ J_Q/x &= 1 - 2g_c + (\eta - c)g'_c + \eta. \end{aligned} \quad (124)$$

For $-1 \leq \eta \leq 0$, substitution of $g'_c = (1 + \eta)/R$ makes these values $-(1 + \eta)/R$, $-(1 + \eta)/R$, and 0, respectively. For $\eta \leq -1$ their signs are immediate from $g_c = g'_c = 0$. Hence the base bank pays every ordinary endpoint.

The cap reserve. Define

$$Z(c) = 2 + \frac{1}{c} + \frac{1}{2} \ln \left(1 + \frac{2}{c} \right), \quad \delta = Z(c) - u. \quad (125)$$

On the mixed curve, (74) says precisely that

$$\delta = \operatorname{artanh} m. \quad (126)$$

For $0 \leq t \leq m$, set

$$\beta(t) = ct[\delta - \operatorname{artanh} t], \quad \mathcal{V}(t) = \int_t^m \beta(s) ds, \quad (127)$$

and extend $\mathcal{V}$ by zero for $t \geq m$. Since $\beta(m) = 0$, this extension is C^1 . With

$$q = \frac{K}{x}, \quad \mu = \frac{q}{1 + q}, \quad \mathcal{H}(q) = (1 + q)^2 \mathcal{V}(\mu), \quad (128)$$

define the full potential

$$W = xN_{\text{def}} + x^2(G_c(y, b) + \mathcal{H}(q)). \quad (129)$$

It is nonnegative.

For an ordinary endpoint, K is fixed while x decreases, so the reserve contributes, after division by x ,

$$L_0(q) = -2\mathcal{H}(q) + q\mathcal{H}'(q). \quad (130)$$

A cap also increments K , and its contribution is $L_0 + \mathcal{H}'$. On the active range $\mu \leq m$, differentiation of (128) gives

$$\mathcal{H}' = 2(1 + q)\mathcal{V} - \beta(\mu), \quad L_0 = -2(1 + q)\mathcal{V} - q\beta(\mu) \leq 0. \quad (131)$$

Thus the reserve can only improve every ordinary drift.

For a cap, feasibility $N_{\text{def}} \leq P$ and $P \geq K$ gives

$$y = \frac{N_{\text{def}} - R(P + E)}{x} \leq -\frac{cK}{x} = -cq. \quad (132)$$

While $\mu \leq m$, we have $q \leq m/(1 - m) \leq \varphi$ and hence $cq \leq 1$. Monotonicity of a_c , together with (113), yields

$$\frac{\ell_U - \ell_Q}{x} \leq cq \left[\delta - \frac{1}{2} \ln(1 + 2q) \right] =: B(q). \quad (133)$$

The change of variables in (128) satisfies

$$\text{artanh } \mu = \frac{1}{2} \ln(1 + 2q), \quad (1 + q)\beta(\mu) = B(q). \quad (134)$$

Consequently the cap residual is cancelled exactly:

$$L_0 + \mathcal{H}' + B(q) = 0 \quad (\mu \leq m). \quad (135)$$

When $\mu \geq m$, the reserve vanishes. At $q_* = m/(1 - m)$, the bracket in (113), evaluated at the largest possible threshold $a_c(-cq_*)$, is zero by (126) and (134). For $q \geq q_*$, (132) only decreases the threshold, so that bracket is nonpositive. The fifth Hamiltonian inequality therefore holds for every state and for arbitrary interleavings of capped and ordinary outcomes.

Initial calibration. Initially $y = b = q = 0$, and

$$f_c(0) = \frac{1}{2} - \frac{c}{4} \ln\left(1 + \frac{2}{c}\right). \quad (136)$$

Elementary integration in (127) gives

$$\mathcal{V}(0) = \frac{c}{2}(\delta - m). \quad (137)$$

Indeed,

$$\int_0^m t \text{artanh } t \, dt = \frac{(m^2 - 1) \text{artanh } m + m}{2}.$$

Using (74) and (126), we obtain the exact calibration

$$f_c(0) + \mathcal{H}(0) = f_c(0) + \mathcal{V}(0) = \frac{c}{2}. \quad (138)$$

Thus $W_{\text{initial}} = cn^2/2$.

Finite jobs, and uniformity on the plateau. We spell out the regularity needed to pass from the fluid Hamiltonian to unit job transitions. It is important here to restrict attention to feasible states. In Region I, $y \geq -1$ implies

$$-xy = R(P + E) - N_{\text{def}} = cP + (P - N_{\text{def}}) + RE \leq x. \quad (139)$$

All terms in the middle expression are nonnegative. Hence $P/x, N_{\text{def}}/x, q \leq 1/c$ and $b \leq 1/R$, so every normalized derivative of the Region-I perspective in (118) is bounded on the feasible cone. In Region II the raw base bank is simply

$$xN_{\text{def}} + \frac{(x + N_{\text{def}} - RP)_+^2}{2R}, \quad (140)$$

which has bounded one-sided Hessians. The two pieces are C^1 at $y = -1$, as verified above, and the positive-part square is C^1 at $\eta = -1$.

For the reserve $\mathcal{R}(x, K) = x^2\mathcal{H}(K/x)$, on its active cone $0 \leq q \leq q_*$ one has bounded $\mathcal{H}''$, and

$$\mathcal{R}_{xx} = 2\mathcal{H} - 2q\mathcal{H}' + q^2\mathcal{H}'', \quad \mathcal{R}_{xK} = \mathcal{H}' - q\mathcal{H}'', \quad \mathcal{R}_{KK} = \mathcal{H}''. \quad (141)$$

For $q \geq q_*$, $\mathcal{R} = 0$; moreover $\mathcal{H}(q_*) = \mathcal{H}'(q_*) = 0$. Thus the join is C^1 with bounded one-sided Hessians. At $x = 0$, set $\mathcal{R} = 0$: for $K > 0$ it is already locally zero, while at $(x, K) = (0, 0)$ two-homogeneity gives zero gradient. Altogether W is $C^{1,1}$ on the feasible state cone. On the mixed interval its regularity constant may depend on the fixed value of u . When $c = r$ and $m = 0$, however, the reserve is identically zero and the base bank depends only on the absolute constant r ; its regularity constant is therefore uniform over all $u \geq u_5^{\text{det}}$. This includes the last transition from $x = 1$ to $x = 0$.

It follows that a unit transition has the Taylor remainder

$$\ell_i + W_{i+1} - W_i \leq O_u(1). \quad (142)$$

The endpoint $E = 0^+$ is interpreted one-sidedly. For a fixed symbolic word, replace all E -coordinates by positive values ε . Its state trajectory is unchanged, and the exact finite pair objective, being a finite sum of affine and minimum terms, converges as $\varepsilon \downarrow 0$. Equivalently, retain an $O(\varepsilon n^2)$ error and take this limit for each fixed n . Therefore the formal E -Hamiltonian transfers to genuine outcome vectors without an asymptotic loss.

Summing (142), using $W \geq 0$, and applying (138) gives

$$\sum_i \ell_i \leq \frac{c}{2}n^2 + O_u(n). \quad (143)$$

Together with (108) and (112), this yields

$$\text{ALG} - \text{ROPT} \leq -c \binom{n}{2} + \frac{c}{2}n^2 + O_u(n) = O_u(n). \quad (144)$$

At $u = u_5^{\text{det}}$, one has $c = r$, $m = 0$, and $Z(r) = u_5^{\text{det}}$. The cap reserve is identically zero, while (138) remains valid. If u is increased above u_5^{det} , the only adverse fifth-endpoint bracket $1 - r(u - 1 - a_r(y))$ decreases, and the capped diagonal term in (108) also decreases.

Here is the promised uniform finite calculation. The base bank and its feasible cone contain no u , so there is one constant C_{step} , depending only on r , for which every ordinary or capped transition with $u \geq u_5^{\text{det}}$ satisfies

$$\ell_i + W_{i+1} - W_i \leq C_{\text{step}}.$$

Because $W_0 = rn^2/2$ and $W_n = 0$, telescoping gives

$$\sum_i \ell_i \leq \frac{r}{2}n^2 + C_{\text{step}}n. \quad (145)$$

In the diagonal term of Equation (108), a value below the cap contributes $-r(1 + p_i)$. A capped value contributes at most $1 - ru \leq 0$. Hence the diagonal sum is nonpositive uniformly on $[u_5^{\det}, \infty)$. Combining Equations (108), (112) and (145) now yields

$$\text{ALG} - (1 + r) \text{OPT} \leq -r \binom{n}{2} + \frac{r}{2} n^2 + C_{\text{step}} n = \left(C_{\text{step}} + \frac{r}{2} \right) n. \quad (146)$$

Thus, with $B = C_{\text{step}} + r/2$,

$$\text{ALG} \leq (1 + r) \text{OPT} + Bn \quad \text{for every finite } u \in [u_5^{\det}, \infty), \quad (147)$$

and B is independent of u . The estimate is genuinely uniform rather than the result of sending a parameter-dependent remainder to infinity. This proves Theorem 3.4. Finally, because $u > 1$ on the present range, every effective offline length is at least one and $\text{OPT} \geq n(n + 1)/2$. The additive term in (144) vanishes in the size-asymptotic ratio. $\square$

3.2 Branchwise lower bounds

3.2.1 The binary stopping game and its lower envelope

The first four branches already occur on binary instances $p_j \in \{0, u\}$. The stopping point must be hidden from the algorithm; announcing it in advance gives a strictly weaker lower bound.

Lemma 3.5 (Raw-safe hidden stopping). *Fix $u > 1$, a target $R \leq u$, and a stopping fraction $\alpha \in (0, 1)$. Against a deterministic algorithm, there is a legal binary adversary whose leading excess at its first crossing is*

$$(u - R)(1 - \sigma^2) + \sigma^2 f_{u,R}(y, b), \quad (148)$$

where $0 \leq b \leq y \leq 1$ and

$$0 \leq y - \alpha < \frac{1}{n - v}. \quad (149)$$

Moreover,

$$f_{u,R}(y, b) = 1 - R + 2y + (u - 1)(1 - R)y^2 + b^2 + 2b(u - 1 - uy). \quad (150)$$

Here $\sigma \in [0, 1]$ is the fraction of jobs not already executed raw. Consequently, if

$$\min_{0 \leq b \leq \alpha} f_{u,R}(\alpha, b) \geq 0,$$

then the adversary gives ratio at least $R - O_u(1/n)$.

Proof. Start in a long phase. A fresh job that is tested receives $p = u$, while a fresh job run raw is assigned the hidden value zero. If v jobs have been run raw and L tested-long jobs have been seen, stop at the first touch for which

$$L \geq \alpha(n - v). \quad (151)$$

All untouched jobs are then assigned value zero. Only for the purpose of lower-bounding the remaining cost, we grant the algorithm this future information and allow an optimal continuation; in the actual interaction, values are still revealed only by tests. If the line is never crossed, every job was run raw: if even one job had been tested, then after the last

first touch the tested fraction would be one. On the all-zero hidden instance the ratio is $u \geq R$.

At a crossing, let e of the L long jobs already be complete, let $d = L - e$ be pending, and let $x = n - v - e - d$ jobs remain untouched. The following favorable canonical schedule lower-bounds the algorithm: run the v raw jobs; test and immediately process the e completed long jobs; perform the d noncompleting long tests; test the x zero jobs; and finally process the d pending long jobs. Indeed, if a noncompleting unit test is immediately before an available length- u completion while h jobs remain, the two orders contribute $h + hu$ and $hu + (h - 1)$, respectively; moving the completion left saves one. Raw completions are available from time zero, and identical long jobs may be relabeled so that the e completed ones are tested first. After the switch, a zero test-completion has length one and therefore precedes a pending operation of length u . These exchanges respect precedence.

The exact costs of this relaxed schedule and of OPT are

$$A_{\text{ex}} = u \left(vn - \frac{v(v-1)}{2} \right) + (1+u) \left(e(n-v) - \frac{e(e-1)}{2} \right) + d(n-v-e) + x(d+x) - \frac{x(x-1)}{2} + u \frac{d(d+1)}{2}, \quad (152)$$

$$O_{\text{ex}} = \frac{n(n+1)}{2} + (u-1) \frac{(e+d)(e+d+1)}{2}. \quad (153)$$

Put $\nu = v/n$, $\delta = (v + e + d)/n$, and $\lambda = e/n$. Twice the quadratic coefficients in Equations (152) and (153) are

$$\begin{aligned} \mathcal{A} &= 1 + 2\delta(1 - u\nu) + (u-1)\delta^2 + 2\nu(\nu + u - 2) \\ &\quad + \lambda^2 + 2\lambda(\nu + u - 1 - u\delta), \\ \mathcal{O} &= 1 + (u-1)(\delta - \nu)^2. \end{aligned}$$

Now put $\sigma = 1 - \nu$, $y = (e + d)/(n - v)$, and $b = e/(n - v)$. Direct expansion gives exactly Equation (148) after normalizing by $n^2/2$. The first-crossing overshoot in Equation (151) is less than one: a test increases $e + d - \alpha(n - v)$ by one and a raw first touch increases it by α . This proves Equation (149), and in particular $\sigma(y - \alpha) < 1/n$. There is one small domain issue: the actual $b \leq y$ can exceed α . Put $\bar{b} = \min\{b, \alpha\}$. Then $0 \leq b - \bar{b} \leq y - \alpha$. Since $f_{u,R}$ is Lipschitz on the unit square,

$$\sigma^2 f_{u,R}(y, b) \geq \sigma^2 f_{u,R}(\alpha, \bar{b}) - O_u(1/n). \quad (154)$$

Thus the stated minimum over $0 \leq \bar{b} \leq \alpha$ suffices uniformly even when $n - v$ is small. The discarded diagonal terms are $O_u(n)$ in the unnormalized costs. Finally, the adaptive transcript defines a fixed instance by replay: every tested answer was fixed when revealed, and every raw job's zero value stayed hidden. This proves the lemma. $\square$

Minimizing Equation (150) over the algorithm's completion choice gives

$$b_*(y) = \max\{0, uy - (u-1)\}. \quad (155)$$

The upper constraint $b \leq y$ never binds, because $uy - (u-1) \leq y$ for $y \leq 1$. There are therefore two elementary quadratic branches. On the positive branch, put

$$A_u = u^2 - u + 1, \quad D_{u,R} = A_u + (u-1)R. \quad (156)$$

Then

$$\min_b f_{u,R}(y, b) = 1 - R - (u - 1)^2 + 2A_u y - D_{u,R} y^2. \quad (157)$$

Its maximizer is $\alpha = A_u/D_{u,R}$, and tangency to zero is

$$(1 - R - (u - 1)^2)D_{u,R} + A_u^2 = 0. \quad (158)$$

Expanding Equation (158) gives Equation (71). On the zero- b branch,

$$\min_b f_{u,R}(y, b) = 1 - R + 2y + (u - 1)(1 - R)y^2, \quad (159)$$

and tangency gives

$$R = 1 + \frac{1}{\sqrt{u - 1}}, \quad \alpha = \frac{1}{\sqrt{u - 1}}. \quad (160)$$

The latter point belongs to the zero- b branch iff $(u - 1)^3 \geq u^2$, with equality at $u = u_3^{\det}$.

For $R = u$, the maximum in Equation (157) is nonnegative exactly while

$$[u(u - 1)]^2 - u(u - 1) - 1 \leq 0, \quad (161)$$

whose positive equality is $u = u_2^{\det}$. We have proved the following binary lower envelope.

Proposition 3.6 (Binary lower curve). *Every deterministic algorithm has size-asymptotic ratio at least*

$$B(u) = \begin{cases} 1, & 0 < u \leq 1, \\ u, & u_1^{\det} \leq u \leq u_2^{\det}, \\ R_{\text{quad}}(u), & u_2^{\det} \leq u \leq u_3^{\det}, \\ 1 + \frac{1}{\sqrt{u - 1}}, & u \geq u_3^{\det}. \end{cases} \quad (162)$$

The lower bound uses only $p_j \in \{0, u\}$.

Proof. For $u_1^{\det} < u \leq u_2^{\det}$, use $R = u$ and the maximizing point of Equation (157). For $u_2^{\det} \leq u \leq u_3^{\det}$, use the positive root $R_{\text{quad}}(u)$ and $\alpha = A_u/D_{u,R_{\text{quad}}(u)}$; the tangency identity makes the maximum zero. The minimizing completion mass is positive in the interior and becomes zero at $u = u_3^{\det}$. For $u \geq u_3^{\det}$, use the zero- b tangency point from Equation (160). The branch conditions are precisely the two inequalities checked above. Finally, $R_{\text{RO}}^{\det}(u) \geq 1$ is trivial for $u \leq 1$. In each nontrivial case Lemma 3.5 supplies fixed binary instances by replay. $\square$

3.2.2 The cap-free harmonic core

This subsection develops the cap-free construction used by the last two branches of the revealing curve. Its notation is local to this subsection. We first record its extremal specialization and then extract the scaled form reused by the mixed branch. The construction has many equally large groups of positive jobs, revealed from longest to shortest, followed by one large group of zero jobs. The only design choice is the gap between two consecutive positive processing times. Choosing these gaps harmonically will make all decisions of the online algorithm disappear from the leading term.

Theorem 3.7 (Extremal harmonic core). *For every deterministic online algorithm $\mathcal{B}$ and every $\varepsilon > 0$, there are arbitrarily large fixed instances I with unit tests and rational processing times such that*

$$\text{ALG}_{\mathcal{B}}(I) \geq (R_{\text{OT}}^{\det} - \varepsilon) \text{OPT}(I).$$

The revelation experiment. Fix an integer K , a rational number $\xi > 0$, and a rational slack $\gamma \in (0, 1)$; one may keep $\gamma = 1/2$ throughout. For a large scaling integer H , the instance will contain H jobs of each of K positive types $0, 1, \dots, K-1$ and ξH zero jobs. We choose H to clear all denominators.

The adversary assigns processing times according to the order in which tests are started. The first H tested jobs receive processing time p_0 , the next H receive p_1 , and so on; after the K positive blocks, every remaining job receives processing time zero. Thus

$$p_0 > p_1 > \dots > p_{K-1} > 1,$$

so the algorithm sees the positive types from longest to shortest and learns about the zero jobs only at the end.

This adaptive description is only a convenient way to construct the input. After running the interaction against a fixed deterministic algorithm, assign to every job label the value revealed when that label was tested. Replaying the algorithm on this fixed assignment produces exactly the same transcript. If the algorithm never tests or completes some job, its cost is infinite and the lower bound is immediate; otherwise every job label is recorded. Hence it is enough to lower-bound the cost in the revelation experiment.

Why each phase can be relaxed to shortest processing time. Let T_ℓ be the start of the first test in positive block ℓ , and let T_K be the start of the first test in the zero block. Phase ℓ contains the completions in $(T_\ell, T_{\ell+1}]$; a completion exactly at T_ℓ belongs to the preceding phase. Each T_ℓ is the boundary between two nonpreemptive operations. Consequently, a previously tested job that is unfinished at T_ℓ has not started processing and still needs its entire processing time.

We repeatedly use the following elementary observation. Suppose jobs with remaining work $v_1, \dots, v_m$ complete after some time T . If their completion order is $q_1, \dots, q_m$, then the g th relative completion time is at least $\sum_{h \leq g} v_{q_h}$. Summing these prefix inequalities shows that their total relative completion time is minimized by arranging the v_i in nondecreasing order. This is merely a lower bound; the online algorithm need not actually use that order.

For normalized group masses $q_1, \dots, q_m$ and remaining lengths $v_1 \leq \dots \leq v_m$, this shortest-first bound has leading coefficient

$$\mathcal{S}((q_g, v_g)_{g=1}^m) = \sum_{g=1}^m \left(\frac{v_g q_g^2}{2} + q_g \sum_{h < g} v_h q_h \right). \quad (163)$$

More precisely, the corresponding contribution for groups of size $q_g H$ is $H^2 \mathcal{S} + O(H)$. The quadratic term is simply the familiar sum $v_g(1 + \dots + q_g H)$ inside one group; the cross term is the waiting caused by all shorter groups.

We ensure that

$$p_0 - p_{K-1} < 1. \quad (164)$$

In phase ℓ , an unfinished job of an older type $j < \ell$ therefore has remaining length p_j , whereas a type- ℓ job completed in its own phase needs both its test and processing, of total length $1 + p_\ell$. Their order in the shortest-first relaxation is

$$p_{\ell-1} < p_{\ell-2} < \dots < p_0 < 1 + p_\ell.$$

At T_K , an untested zero job has remaining length 1, so the relaxed tail starts with all zero jobs and then uses the positive types from shortest to longest. These are the only two places where Equation (164) is used.

The finite accounting identity. We now record the online algorithm's possible behavior. Normalize all job counts by H . Let

- $s_j \in [0, 1]$ be the mass of type- j jobs completed in their own phase;
- $d_{j,\ell} \geq 0$ be the mass of type- j jobs completed in the later phase $\ell > j$; and
-

$$X_{j,\ell} = s_j + \sum_{h=j+1}^{\ell-1} d_{j,h} \leq 1 \quad (\ell > j)$$

be the mass of type j completed before phase ℓ . Thus $X_{j,\ell+1}$ is the mass completed by the end of phase ℓ .

At time T_ℓ , the machine has already performed the tests of the ℓH jobs in earlier positive blocks and has processed every earlier job that has completed. Thus

$$\frac{T_\ell}{H} \geq \ell + \sum_{j < \ell} p_j X_{j,\ell}. \quad (165)$$

For each phase, charge its completions their common start time from Equation (165), and charge their relative completion times using Equation (163). Do the same at T_K for the remaining positive jobs and the ξH zero jobs. Expanding and collecting equal variables gives

$$\begin{aligned} \frac{\text{ALG}}{H^2} &\geq C_K + \sum_{j=0}^{K-1} \left(\frac{s_j^2}{2} - \theta_j s_j \right) \\ &\quad + \sum_{j < \ell} d_{j,\ell} \left[\ell - j - \theta_j - \sum_{j < k < \ell} (p_j - p_k) X_{k,\ell+1} \right] - O(1/H), \end{aligned} \quad (166)$$

where K, ξ, γ are fixed while $H \rightarrow \infty$, and

$$\theta_j = 1 - \xi(p_j - 1) - \sum_{k > j} (p_j - p_k - 1), \quad (167)$$

$$C_K = \frac{\xi^2}{2} + 2\xi K + K^2 + P_K, \quad P_K = \sum_{j=0}^{K-1} \left(j + \frac{1}{2} \right) p_j. \quad (168)$$

For completeness, here is how to audit Equation (166) without guessing its origin. In phase ℓ , the masses in the shortest-first relaxation are

$$d_{\ell-1,\ell}, d_{\ell-2,\ell}, \dots, d_{0,\ell}, s_\ell$$

with respective lengths

$$p_{\ell-1}, p_{\ell-2}, \dots, p_0, 1 + p_\ell.$$

Multiply their total mass by the right-hand side of Equation (165) and add the functional Equation (163). In the final phase use masses

$$\xi, 1 - X_{K-1,K}, \dots, 1 - X_{0,K}$$

and lengths $1, p_{K-1}, \dots, p_0$. The constant terms sum to C_K . The terms containing only s_j are $s_j^2/2 - \theta_j s_j$; the coefficient of each $d_{j,\ell}$ is precisely the bracket in Equation (166). Thus the displayed identity is just the phasewise shortest-first bound with the phase start times retained. No optimality assumption about the online schedule is hidden in it.

The meaning of the expression is useful. Completing an s_j fraction of a type in its own phase creates a quadratic completion cost but saves later waiting cost, producing $s_j^2/2 - \theta_j s_j$. A completion postponed to a later positive phase is represented by $d_{j,\ell}$ and its bracket. We next choose the processing times so that every such bracket is nonnegative and all θ_j are equal.

The harmonic choice and its telescoping cancellation. Set

$$p_{K-1} = 1 + \frac{\gamma}{\xi}, \quad p_i = p_{i+1} + \frac{1}{\xi + K - 1 - i} \quad (0 \leq i < K - 1). \quad (169)$$

In other words, measured upward from the shortest type, the consecutive gaps are

$$\frac{1}{\xi + 1}, \frac{1}{\xi + 2}, \dots, \frac{1}{\xi + K - 1}.$$

This is the key design choice. If $m = K - 1 - j$, then

$$p_j - 1 = \frac{\gamma}{\xi} + \sum_{h=1}^m \frac{1}{\xi + h},$$

$$\sum_{k>j} (p_j - p_k) = \sum_{h=1}^m \frac{h}{\xi + h}.$$

Adding the first line multiplied by ξ to the second line makes each denominator cancel:

$$\xi(p_j - 1) + \sum_{k>j} (p_j - p_k) = \gamma + \sum_{h=1}^m \frac{\xi + h}{\xi + h} = \gamma + m. \quad (170)$$

Substitution in Equation (167) gives the same value for every type,

$$\theta_j = 1 - \gamma. \quad (171)$$

This is why the increments in Equation (169) are harmonic: the factor ξ coming from the zero block and the multiplicity h coming from the h shorter positive types add up to the denominator $\xi + h$.

The delayed-completion terms are now harmless as well. Since $X_{k,\ell+1} \leq 1$, their brackets are at least

$$\begin{aligned} \ell - j - (1 - \gamma) - \sum_{j < k < \ell} (p_j - p_k) \\ = \gamma + \sum_{j < k < \ell} (1 - (p_j - p_k)) > 0, \end{aligned}$$

where the final inequality follows from Equation (164). We may therefore drop every term containing $d_{j,\ell}$. The remaining minimization is not a multivariable optimization problem: complete one square for each j ,

$$\frac{s_j^2}{2} - (1 - \gamma)s_j = \frac{1}{2}(s_j - (1 - \gamma))^2 - \frac{1}{2}(1 - \gamma)^2.$$

Because $1 - \gamma \in (0, 1)$, this minimum is feasible for $s_j \in [0, 1]$. Hence every online algorithm obeys

$$\liminf_{H \rightarrow \infty} \frac{\text{ALG}}{H^2} \geq Q_K := C_K - \frac{K}{2}(1 - \gamma)^2. \quad (172)$$

It remains to compute the offline benchmark. Knowing all processing times, the optimum runs the zero jobs first and the positive types from shortest to longest. Equivalently, the pairwise contribution of two jobs is $1 + \min\{p_i, p_j\}$. Its leading coefficient is

$$D_K = \frac{\xi^2}{2} + \xi K + \frac{K^2}{2} + P_K, \quad \lim_{H \rightarrow \infty} \frac{\text{OPT}}{H^2} = D_K. \quad (173)$$

The four terms respectively account for zero-zero pairs, zero-positive pairs, the unit-test part of positive-positive pairs, and the processing part of positive-positive pairs. Thus the finite- K lower-bound certificate is simply Q_K/D_K .

Passing to a smooth curve. Write

$$\alpha = \frac{K}{\xi}$$

for the ratio of positive mass to zero mass, and keep α fixed while $K \rightarrow \infty$. We restrict to $0 < \alpha < e - 1$. Then

$$p_0 - p_{K-1} = \sum_{h=1}^{K-1} \frac{1}{\xi + h} < \ln\left(1 + \frac{K}{\xi}\right) = \ln(1 + \alpha) < 1,$$

which verifies Equation (164).

The processing contribution P_K has the exact form

$$P_K = \frac{K^2}{2} \left(1 + \frac{\gamma}{\xi}\right) + \frac{1}{2} \sum_{h=1}^{K-1} \frac{(K-h)^2}{\xi + h}. \quad (174)$$

Indeed, the common baseline $1 + \gamma/\xi$ contributes the first term, and swapping the order of summation shows that the increment $1/(\xi + h)$ receives total weight $(K-h)^2/2$. After division by ξ^2 , the sum becomes a Riemann integral:

$$\begin{aligned} \frac{P_K}{\xi^2} &\rightarrow I(\alpha) = \frac{\alpha^2}{2} + \frac{1}{2} \int_0^\alpha \frac{(\alpha - t)^2}{1 + t} dt \\ &= \int_0^\alpha (\alpha - t)(1 + \ln(1 + t)) dt \\ &= \frac{1}{2}(1 + \alpha)^2 \ln(1 + \alpha) - \frac{\alpha}{2} - \frac{\alpha^2}{4}. \end{aligned} \quad (175)$$

The second integral form follows either by integration by parts or by summing the limiting processing-time curve $1 + \ln(1 + t)$ against the remaining positive mass $\alpha - t$. The subtraction in Equation (172) is only $O(K)$, whereas $C_K, D_K = \Theta(K^2)$. Since $C_K - D_K = \xi K + K^2/2$, we obtain

$$R(\alpha) := \lim_{K \rightarrow \infty} \frac{Q_K}{D_K} = 1 + \frac{\alpha + \alpha^2/2}{\frac{1}{2} + \frac{\alpha}{2} + \frac{\alpha^2}{4} + \frac{1}{2}(1 + \alpha)^2 \ln(1 + \alpha)}. \quad (176)$$

This last one-variable maximization is where the constant in the theorem comes from. Put $z = (1 + \alpha)^2$. Then

$$R(z) = 1 + \frac{2(z - 1)}{1 + z + z \ln z}. \quad (177)$$

The derivative of the fraction has the sign of

$$3 - z + \ln z.$$

For $z > 1$ this expression is strictly decreasing, is positive at $z = 1$, and tends to $-\infty$. Hence it has one zero $u_5^{\det}$, characterized by

$$u_5^{\det} - 3 = \ln u_5^{\det}.$$

The derivative is positive before $u_5^{\det}$ and negative afterward, so this is the global maximum. At the root,

$$1 + u_5^{\det} + u_5^{\det} \ln u_5^{\det} = (u_5^{\det} - 1)^2,$$

and therefore

$$R(u_5^{\det}) = 1 + \frac{2}{u_5^{\det} - 1} = R_{\text{OT}}^{\det}.$$

Moreover, $\sqrt{u_5^{\det}} - 1 = 1.12255\dots < e - 1$, so the maximizer lies in the range where Equation (164) holds.

Proof of Theorem 3.7. Choose a rational α sufficiently close to $\alpha_\star = \sqrt{u_5^{\det}} - 1$, and choose any rational $\gamma \in (0, 1)$. For a sufficiently large K , set $\xi = K/\alpha$ and use Equation (169); then $Q_K/D_K > R_{\text{OT}}^{\det} - \varepsilon/2$. All parameters are rational. Let H tend to infinity through multiples of their common denominators. Equations (172) and (173) make the actual ratio exceed $R_{\text{OT}}^{\det} - \varepsilon$ for all sufficiently large H . Finally, the replay argument at the beginning of the proof converts the revelation experiment into a fixed instance for the given deterministic algorithm. $\square$

Lemma 3.8 (Scaled harmonic core). *Fix a total normalized mass $s > 0$ and a parameter $t \in [1, e)$. Against every deterministic obligatory-testing algorithm, there are scales $M_\nu \rightarrow \infty$ and finite rational multilevel instances with $sM_\nu + o(M_\nu)$ jobs such that*

$$\frac{\text{OPT}}{M_\nu^2} \longrightarrow O_0 = \frac{s^2}{4}(1 + t^{-2} + 2 \ln t), \quad \liminf_{\nu \rightarrow \infty} \frac{\text{ALG}}{M_\nu^2} \geq A_0^{\text{lb}} := O_0 + \frac{s^2}{2}(1 - t^{-2}). \quad (178)$$

Their normalized total positive processing mass converges to

$$\frac{1}{M_\nu} \sum_{j: p_j > 0} p_j \longrightarrow L = s \ln t, \quad (179)$$

and their largest processing time converges to $1 + \ln t < 2$. The adaptive revelation is converted by replay into a fixed instance.

Proof. The case $t = 1$ is obtained by continuity from instances with vanishing positive mass, so suppose $1 < t < e$. In the construction above set $\alpha = t - 1 = K/\xi$. Before rescaling, the total normalized mass is $K + \xi = \xi t$. Give every unit of normalized mass the common factor $s/(\xi t)$.

The phase accounting, harmonic cancellation, and completion of squares in Equations (166), (170) and (172) are homogeneous of degree two in the job masses. The Riemann-sum limit Equation (175) substituted into Equation (173) therefore gives

$$O_0 = \frac{s^2}{4}(1 + t^{-2} + 2 \ln t).$$

Likewise $C_K - D_K = \xi K + K^2/2$, while the $O(K)$ square-completion loss in Equation (172) disappears after division by the quadratic scale. After the same rescaling this gives the guaranteed lower benchmark

$$A_0^{\text{lb}} - O_0 = \frac{s^2}{2}(1 - t^{-2}).$$

Finally, summing the limiting processing-time curve $1 + \ln(1 + x)$ over the positive mass gives $L = s \ln t$, and Equation (169) gives $p_0 \rightarrow 1 + \ln t$. Rational approximation, clearing denominators, and the replay argument used in the theorem produce a diagonal sequence of arbitrarily large fixed rational instances with the stated limit and liminf bounds. $\square$

3.2.3 The unrestricted mixed lower bound

For $u_4^{\det} < u < u_5^{\det}$, neither the binary family nor the pure cap-free family is tight. The correct adversary puts a capped block before a scaled harmonic core. Two lemmas ensure that raw execution and early processing of caps cannot escape it.

The cap-free construction needed below is exactly the scaled harmonic core from Lemma 3.8; no second phase-accounting proof is needed here. We now prepend capped jobs to that reusable core.

Lemma 3.9 (Cap deferral). *Prepend capped mass m with value $p = u$ to a later cap-free block of mass $s = 1 - m$ and total processing mass L . Suppose every later value satisfies $1 + p \leq u$ and*

$$s(u - 1) - L - m \geq 0. \quad (180)$$

Then processing any capped job before all later jobs are complete cannot decrease the leading online cost.

Proof. A cap completed after its test block but before a later job of value p has mutual cost $1 + u$; leaving it pending until after that job gives $2 + p$. The exchange saves $u - 1 - p \geq 0$. If cap mass q is instead completed inside the cap-test block, it pairs with all later jobs add $q[s(u - 1) - L]$. Its largest possible saving against not-yet-tested caps is $qm - q^2/2$. Relative to leaving every cap pending, the change is therefore at least

$$q[s(u - 1) - L - m] + \frac{q^2}{2} \geq 0. \quad (181)$$

The finite diagonal discrepancy is lower order. $\square$

After cap deferral, pairwise accounting gives the offline coefficient O and the guaranteed online lower benchmark A^{lb} :

$$A^{\text{lb}} = A_0^{\text{lb}} + m(1 - m) + m(1 + L) + \frac{u}{2}m^2, \quad (182)$$

$$O = O_0 + m(1 - m + L) + \frac{u}{2}m^2. \quad (183)$$

Indeed, a cap-later pair contributes $2 + p$ online and $1 + p$ offline, whereas a cap-cap pair contributes $2 + u$ online and u offline; collecting these pairs yields the two cross terms and the cap squares above. In particular, $A^{\text{lb}} - O = (A_0^{\text{lb}} - O_0) + m$.

Lemma 3.10 (Raw-safe quota and replay). *Fix $m \in (0, 1)$ and a later cap-free block satisfying the hypotheses of Lemma 3.9. Against every deterministic revealing algorithm there are fixed instances whose asymptotic excess is at least the cap-free hybrid benchmark $A^{\text{lb}} - RO$ from Equations (182) and (183) for every target $R \leq u$.*

Proof. In the initial phase, a tested fresh job receives $p = u$, while a job run raw is assigned hidden value zero. If v raw jobs and ℓ tested caps have been touched, stop at the first crossing

$$\frac{\ell}{n - v} \geq m. \quad (184)$$

With $S = n - v$, the overshoot is less than one, so $\ell = mS + O(1)$. If no crossing occurs, no test can have occurred: after all first touches, any positive ℓ would make the ratio in Equation (184) equal one. Thus every job was run raw and the all-zero hidden instance has ratio $u \geq R$.

At a crossing, scale the prescribed later instance to the remaining $S - \ell$ first touches, absorbing the one-job cap overshoot and integer rounding into its zero group. A later test reveals the next value in this fixed type stream; a later raw action consumes the next value but keeps it hidden from the revealing algorithm.

Move every initial raw-zero completion to the start. It has the same length u as cap processing, and crossing a unit test only lowers completion cost. With

$$Z_v = vn - \frac{v(v-1)}{2}, \quad (185)$$

this prefix contributes uZ_v online and Z_v offline. Delete it. The remaining S labels consist, up to one-job rounding, of capped fraction m and the prescribed later cap-free block.

If the revealing algorithm later runs a fresh assigned value p raw, a physical cap-free simulation tests and immediately processes that job in time $1+p \leq u$, suppresses the answer from the virtual revealing algorithm, and idles until its virtual u -block ends. Inductively every physical completion is no later than its virtual revealing completion. Since the finite-cap and cap-free offline lengths are both $1+p$ on the later block,

$$\text{ALG}_{\text{RO}} - R \text{OPT}_{\text{RO}} \geq (u - R)Z_v + (\text{ALG}_{\text{hyb}} - R \text{OPT}_{\text{hyb}}) - O_u(n). \quad (186)$$

The first term is nonnegative. Finally, record every assigned value on its job label. Replaying this fixed vector produces the same transcript because the values learned only by the physical simulation were hidden from the virtual algorithm. This proves legality. $\square$

Apply Lemma 3.8 with $s = 1 - m$. Substitution into Equations (182) and (183) gives the guaranteed two-parameter lower ratio

$$\mathcal{R}_u(m, t) = 1 + \frac{m + \frac{(1-m)^2}{2}(1-t^{-2})}{\frac{(1-m)^2}{4}(1+t^{-2} + 2\ln t) + m(1-m)(1+\ln t) + \frac{u}{2}m^2}. \quad (187)$$

At an interior stationary point with value $1 + c$, differentiation and the zero-gap equation give

$$\ln t = u - 2 - \frac{1}{c}, \quad t^2 = \frac{1-m}{1+m} \left(1 + \frac{2}{c}\right). \quad (188)$$

Eliminating t gives precisely Equation (74). Conversely, those equations make $\mathcal{R}_u(m, t) = 1 + c$.

The branch runs from

$$(u, m, t) = \left(u_4^{\det}, \frac{1}{\varphi}, 1\right) \quad \text{to} \quad (u, m, t) = (u_5^{\det}, 0, \sqrt{u_5^{\det}}). \quad (189)$$

It satisfies $u \geq u_4^{\det}$, $\ln t < 1$, and $m \leq 1/\varphi$. Therefore

$$(1-m)(u-1-\ln t) > (1-m)\varphi \geq \frac{1}{\varphi} \geq m, \quad (190)$$

so Equation (180) holds; also $1+p < 3 < u$ for every later positive value in a sufficiently fine finite construction. The functions

$$g(c) = 1 - \frac{1}{c} + \frac{1}{2} \ln \left(1 + \frac{2}{c}\right), \quad h(m) = \text{artanh } m - m \quad (191)$$

have $g'(c) = 2/[c^2(c+2)] > 0$ and $h'(m) = m^2/(1-m^2) > 0$. Hence $m(c) = h^{-1}(g(c))$ is strictly increasing, while

$$u(c) = 1 + \frac{2}{c} - m(c) \quad (192)$$

is strictly decreasing. The identities $g(r) = h(0) = 0$ and $g(1/\varphi) = h(1/\varphi)$ give the two endpoints in Equation (189); consequently every $u \in [u_4^{\det}, u_5^{\det}]$ has exactly one parameter pair.

To pass to fixed finite instances, first approximate (m, t) rationally while retaining the strict cap inequalities, then choose a sufficiently fine harmonic family, and finally multiply all rational block sizes by a common integer H . For fixed harmonic depth, quota overshoot, diagonals, and rounding are $O_u(H)$ whereas the displayed costs are $\Theta(H^2)$. Send first H , then the harmonic depth, and finally the rational approximation to their limits. Together with Lemma 3.10, this proves

$$R_{\text{RO}}^{\det}(u) \geq R_{\text{mix}}(u) \quad (u_4^{\det} \leq u \leq u_5^{\det}). \quad (193)$$

The construction above applies directly in the open interval. At $u = u_4^{\det}$ and $u = u_5^{\det}$ the same statement follows by approaching from the interior, so the endpoints are included.

3.2.4 The large-upper-limit lower bound by cap-free transfer

Lemma 3.11 (Bounded cap-free transfer). *Let $u \geq 1$, and let $L_{\text{OT}}(P)$ denote the deterministic size-asymptotic value in the cap-free limit, restricted to $0 \leq p_j \leq P$. Then*

$$R_{\text{RO}}^{\det}(u) \geq L_{\text{OT}}(u-1). \quad (194)$$

In particular,

$$R_{\text{RO}}^{\det}(u) \geq R_{\text{OT}}^{\det} \left(u \geq 1 + \frac{1}{r} = 2.752620747896442 \dots \right). \quad (195)$$

Proof. Given a finite-cap algorithm, construct a cap-free algorithm on instances with $p_j \leq u-1$ by running a virtual copy of the finite-cap algorithm. Copy ordinary tests and processing actions. When the virtual algorithm runs an untouched job raw for u time, physically test and immediately process it in $1+p_j \leq u$ time, hide the answer from the virtual copy, and wait for the virtual clock. Every physical completion is no later than its virtual completion. Moreover, $\min\{u, 1+p_j\} = 1+p_j$, so finite-cap and cap-free optima coincide. Thus we obtain the following game-value comparison. For every finite-cap algorithm $\mathcal{A}$, the simulation produces a cap-free algorithm $\mathcal{B}$ and, pointwise on every instance with $p_j \leq u-1$,

$$\text{ALG}_{\mathcal{A}} \geq \text{ALG}_{\mathcal{B}}, \quad \text{OPT}_u = \text{OPT}_{\infty}.$$

Consequently

$$\text{CR}_{\infty}(\mathcal{A}) \geq \text{CR}_{\infty}(\mathcal{B}; p \leq u-1) \geq L_{\text{OT}}(u-1).$$

Taking the infimum over $\mathcal{A}$ proves Equation (194).

The harmonic cap-free family attaining $R_{\text{OT}}^{\det}$ has limiting largest processing time

$$1 + \ln(1 + y_*) = 1 + \frac{1}{2} \ln u_5^{\det} = \frac{u_5^{\det} - 1}{2} = \frac{1}{r}, \quad y_* = \sqrt{u_5^{\det}} - 1. \quad (196)$$

Approximation from below also gives the endpoint, proving Equation (195). $\square$

3.3 Assembly of the curve

3.3.1 The shape of the claimed curve

The global-maximum assertion follows directly from the displayed formulas. Only the algebraic branch needs a check. Implicit differentiation of $T_s(R_{\text{quad}}(u)) = 0$, with $s = u - 1$, gives

$$R'_{\text{quad}}(u) = -\frac{R^2 + (3s^2 + 2s)R - 3(s+1)^2}{2sR + s^3 + s^2 + 1}. \quad (197)$$

Using $T_s(R) = 0$, its numerator factors as

$$\frac{2s^3 + s^2 - 1}{s} \left(R - \frac{2s^3 + 3s^2 - 2}{2s^3 + s^2 - 1} \right).$$

On the relevant interval $s \geq u_2^{\text{det}} - 1 > 2/3$, the prefactor is positive, while the fraction in parentheses is at most $5/3$ because $4s^3 - 4s^2 + 1 > 0$. The sign certificate (80) gives $R > 5/3$. Hence $R'_{\text{quad}}(u) < 0$. The reciprocal branch and the mixed parametrization are also decreasing, so $R_{\text{RO}}^{\text{det}}(u)$ attains its global maximum only at $u = u_2^{\text{det}}$.

3.3.2 Completion of the curve and consistency at the joins

The branchwise lower bounds are: Proposition 3.6 supplies the first four branches, Equation (193) the fifth, and Equation (195) the last. The raw algorithm, Lemmas 3.2 and 3.3, supplies the matching upper bounds through u_4^{det} , while Section 3.1.2 supplies the last two. Together these statements prove Theorem 3.1.

For clarity, all five joins are exact:

$$\begin{aligned} R_{\text{quad}}(u_2^{\text{det}}) &= u_2^{\text{det}}, \\ R_{\text{quad}}(u_3^{\text{det}}) &= 1 + \frac{1}{\sqrt{s_0}} = 1.682327803828019\dots, \\ 1 + \frac{1}{\sqrt{u_4^{\text{det}} - 1}} &= \varphi = R_{\text{mix}}(u_4^{\text{det}}), \\ R_{\text{mix}}(u_5^{\text{det}}) &= 1 + r = R_{\text{OT}}^{\text{det}}. \end{aligned}$$

At $u = u_1^{\text{det}}$, both initial branches equal one. Representative values of the new mixed branch are

u	$R_{\text{RO}}^{\text{det}}(u)$	m	t
3.800000	1.598851588	0.539725635	1.138984621
4.000000	1.583283633	0.428863572	1.330517832
4.071679486	1.579204806	0.381330539	1.412236296
4.300000	1.571692026	0.198387083	1.734651379

Remark 3.12 (Asymptotic scope). All statements above concern the fixed- u size-asymptotic ratio. Below the plateau, the upper bounds have additive error $O_u(n)$. On $u \in [u_5^{\text{det}}, \infty)$ the same algorithm has one uniform $O(n)$ error. The unbounded obligatory endpoint is not included in the statement of the curve; it is derived after the finite- u proof in Section 3.4.

The next subsection, Section 3.1.2, proves the matching middle- and high- u upper bounds and thereby completes the proof of Theorem 3.1.

3.4 The obligatory endpoint

The obligatory model is the unbounded endpoint of revealing optimization, but the passage to that endpoint is not a formal substitution $u = \infty$: both the algorithm and the additive error must be uniform in u . The finite curve has just established exactly those two properties, so the endpoint now follows by a short transfer.

Theorem 3.13 (Deterministic obligatory testing). *Let $u_5^{\det} > 1$ be the unique solution of $u_5^{\det} - 3 = \ln u_5^{\det}$, and set*

$$R_{\text{OT}}^{\det} = 1 + \frac{2}{u_5^{\det} - 1} = 1.570574096649395 \dots$$

For every deterministic online algorithm $\mathcal{B}$ and every $\varepsilon > 0$, there are arbitrarily large obligatory-testing instances I such that

$$\text{cost}(\mathcal{B}, I) \geq (R_{\text{OT}}^{\det} - \varepsilon) \text{cost}(\text{OPT}, I).$$

Conversely, there are a deterministic algorithm $\mathcal{A}$ and an absolute constant B such that every n -job obligatory instance satisfies

$$\text{cost}(\mathcal{A}, I) \leq R_{\text{OT}}^{\det} \text{cost}(\text{OPT}, I) + Bn.$$

Consequently the optimal deterministic size-asymptotic competitive ratio is exactly $R_{\text{OT}}^{\det}$.

Proof. For the upper bound, use `ADAPTIVETHRESHOLD` with $c = r$. This algorithm never runs a job raw and is independent of u . Given a finite obligatory instance, choose any finite

$$u > \max\{u_5^{\det}, 1 + \max_j p_j\}.$$

Its execution is identical in the revealing model with cap u , and every offline effective length is $\min\{u, 1 + p_j\} = 1 + p_j$. Thus the revealing and obligatory optima coincide. The constant B in Equation (147) is independent of u , so the same inequality holds for the obligatory instance.

For the lower bound, regard an arbitrary obligatory algorithm $\mathcal{B}$ as a revealing algorithm at the fixed cap $u = u_5^{\det}$ that simply never uses raw execution. The plateau lower bound in Theorem 3.1 may be witnessed by cap-free instances with $p_j < u_5^{\det} - 1$. On those instances the online transcript is unchanged and $\min\{u_5^{\det}, 1 + p_j\} = 1 + p_j$, so again the two offline optima coincide. The finite-cap lower inequality therefore gives the claimed obligatory lower bound for arbitrarily large instances. $\square$

4 Revealing-optimization instance optimality

Fix $u < \infty$. An untouched job may be run raw for the public time u , or tested for one unit; a test reveals $p_i \in [0, u]$, after which the processing operation of length p_i may be delayed. The goal of this section is to prove Theorem 4.1, the rigorous version of the informal revealing-optimization theorem Theorem 1.8.

For a distribution D on $[0, u]$, we define an explicit value $\Phi_{\text{RO},u}(D)$ in Equation (206). It is obtained by choosing one threshold and minimizing a quadratic over the tested fraction q . For a multiset M , use the empirical distribution $D[M]$ from Equation (7).

Theorem 4.1 (Revealing-optimization instance optimality). *For every fixed $u > 0$ there exist randomized nonanticipating policies $\mathcal{A}_{n,u}^*$, depending only on n, u , and a deterministic sequence $\varepsilon_u(n) \rightarrow 0$ such that every size- n multiset $M \subseteq [0, u]$ has the following properties. For every fixed labeling σ ,*

$$\mathbb{E} \text{cost}(\mathcal{A}_{n,u}^*, \sigma(M)) \leq n^2 \Phi_{\text{RO},u}(D[M]) + \varepsilon_u(n)n^2. \quad (198)$$

Conversely, for every randomized nonanticipating policy $\mathcal{A}$, even an announced policy, some fixed oblivious labeling σ satisfies

$$\mathbb{E} \text{cost}(\mathcal{A}, \sigma(M)) \geq n^2 \Phi_{\text{RO},u}(D[M]) - \varepsilon_u(n)n^2. \quad (199)$$

One may take

$$\varepsilon_u(n) = O_u\left(n^{-1/6} \sqrt{\ln(n+2)}\right). \quad (200)$$

The proof has three layers. First, for a fixed final tested fraction q , a completion envelope reduces every fluid schedule to an ordinary fractional knapsack. Second, an adaptive first-touch trace turns every deterministic policy under a uniform placement into predictable sampling without replacement. One event then controls every touch prefix, so the realized stopping fraction can be substituted pathwise. Finally, a growing grid and a sublinear sample make the result uniform over arbitrary bounded multisets without announcing the input.

4.1 The fixed-fraction fluid benchmark

Let D initially have finite support, let $P \sim D$, and put $\mu = \mathbb{E}P$. A test automatically completes a zero outcome. Apply Lemma 2.5 and choose its threshold selector g_* . Write

$$a = a_D(g_*), \quad \ell = \mathbb{E}[Pg_*(P)\mathbf{1}_{P>0}], \quad \tau_D = \frac{1+\ell}{a}. \quad (201)$$

Write the positive support as $p_1, \dots, p_s$, and let $\nu_i = \nu(\{p_i\})$ be the residual masses from Lemma 2.5. Thus

$$\sum_i \nu_i = 1 - a, \quad p_i \geq \tau_D \text{ whenever } \nu_i > 0. \quad (202)$$

For a fixed final tested fraction $q \in [0, 1]$, define the model envelope

$$\mathcal{C}_{\text{RO},D,q} := \mathcal{C}_{D,q}^u, \quad (203)$$

where the right-hand side is the common envelope Equation (31) with independent-mode work $v = u$; the independent mode is raw execution.

Lemma 4.2 (Fixed- q completion envelope). *The envelope $\mathcal{C}_{\text{RO},D,q}$ is the divisible-knapsack curve with items*

item	completion capacity	work per completion
maximum-density test module	aq	τ_D
residual tested class i	$q\nu_i$	p_i
raw module	$1 - q$	u

If $\tau_D < u$, the greedy order is the test module, the residual classes in nondecreasing p_i , and the raw module. If $\tau_D \geq u$, the all-raw curve pointwise dominates every choice with $q > 0$.

Proof. Apply Lemma 2.8 with $v = u$. It gives exactly the three displayed kinds of items. If $\tau_D < u$, their work-per-completion values obey $\tau_D \leq p_i \leq u$. If $\tau_D \geq u$, every test and residual item costs at least u per completion, while the $q = 0$ envelope consists of one raw item of capacity one and cost u . $\square$

Put

$$\ell_\nu = \sum_i p_i \nu_i = \mu - \ell, \quad K_\nu = \sum_{i,j} \min\{p_i, p_j\} \nu_i \nu_j. \quad (204)$$

The common area identity Equation (33), specialized through Lemma 4.2, gives

$$F_{\text{RO},u,D}(q) = (1 + \ell) \left(q - \frac{aq^2}{2} \right) + q\ell_\nu(1 - q) + \frac{q^2 K_\nu}{2} + \frac{u(1 - q)^2}{2}. \quad (205)$$

Indeed, the four terms are respectively the test-module area, residual work delaying the raw block, internal residual SPT pairs, and the homogeneous raw block. Define

$$\Phi_{\text{RO},u}(D) = \begin{cases} \min_{0 \leq q \leq 1} F_{\text{RO},u,D}(q), & \tau_D < u, \\ u/2, & \tau_D \geq u. \end{cases} \quad (206)$$

This is an explicit one-dimensional optimization. In fact

$$F_{\text{RO},u,D}(q) = \frac{u}{2} + Aq + Bq^2, \quad A = 1 + \mu - u, \quad B = -\frac{a(1 + \ell)}{2} - \ell_\nu + \frac{K_\nu}{2} + \frac{u}{2}. \quad (207)$$

For each work budget, convexly combining feasible points for q_1, q_2 gives a feasible point for the corresponding convex combination of the two fractions. Hence $\mathcal{C}_{\text{RO},D,q}(x)$ is concave in q , its remaining-mass area is convex, and $B \geq 0$. Moreover,

$$\tau_D < u \iff \mathbb{E}(u - P) > 1 \iff A < 0.$$

Thus the minimizer is

$$q^* = 1 \quad (B = 0), \quad q^* = \min\{1, -A/(2B)\} \quad (B > 0). \quad (208)$$

We have proved pathwise at the fluid level that a policy with realized final tested fraction q has area at least $F_{\text{RO},u,D}(q) \geq \Phi_{\text{RO},u}(D)$.

4.2 Adaptive policies and the announced lower bound

We now transfer the fluid envelope to a finite policy which may choose labels and its stopping time adaptively. Equal-valued jobs are first distinguished as occurrence tokens, and we fix the private seed of a completing policy. For the present finite-support law, take zero and the $K = s$ positive atoms as the cells, with masses d_j . Apply Lemma 2.18 with failure probability n^{-2} , using its notation X_k, A_k , and let $\Gamma_n = \Gamma_{\text{cell}}$. With that probability, simultaneously for every cell j and every touch prefix $m \leq (1 - \delta)n$,

$$\left| \sum_{k \leq m} A_k \mathbf{1}_{X_k \in j} - d_j \sum_{k \leq m} A_k \right| \leq \Gamma_n. \quad (209)$$

Stop the first-touch sequence when only $\delta n + O(1)$ labels remain. In the suffix, replace every future test of an untouched job by raw execution at its first original operation and

delete its later known processing. At most $\delta n + O(1)$ jobs and three operations per affected job change. Thus Lemma 2.19, with $W = u + 1$, gives

$$C_{\text{original}} \geq C_{\text{edited}} - O_u(\delta n^2 + n). \quad (210)$$

The edited schedule has a fixed final tested fraction q , but we do not condition on its value.

On the single event Equation (209), consider any operation prefix and normalize counts and work by n . Let t, b, z, c_j denote tested mass, raw mass, tested-zero completions, and processed known mass from positive cell j , and put $\gamma_n = \Gamma_n/n$. The urn event gives the capacity hypotheses of Lemma 2.21; raw work is exactly ub , so its work hypothesis holds with $\alpha = 0$. Therefore

$$\text{completed fraction at work } x \leq C_{\text{RO},D,q}(x) + (K + 1)\gamma_n. \quad (211)$$

The completion horizon is at most $H = 1 + u$, and $0 \leq \Phi_{\text{RO},u}(D) \leq u/2$. Thus Theorem 2.20, with the suffix edit as $\varepsilon = O_u(\delta + 1/n)$ and $\beta = (K + 1)\gamma_n$, gives

$$\frac{C_{\text{original}}}{n^2} \geq \Phi_{\text{RO},u}(D) - O_{u,K}(\gamma_n + \delta + 1/n). \quad (212)$$

For an arbitrary multiset, take

$$K_n = \lfloor n^{1/6} \rfloor, \quad h_n = u/K_n, \quad \delta_n = n^{-1/6}, \quad (213)$$

keep zero separate, and round every positive value upward. If two distributions admit a zero-preserving coupling with pointwise displacement at most h , the transport lemma Lemma 2.9 gives

$$|\Phi_{\text{RO},u}(D) - \Phi_{\text{RO},u}(D')| \leq h. \quad (214)$$

For the finite repair, raising a processed value to its bin endpoint adds at most h_n to normalized prefix work, while Lemma 2.21 loses at most $(K_n + 1)\gamma_n$ completed mass. With the parameters above,

$$(K_n + 1)\frac{\Gamma_n}{n} = O\left(n^{-1/6}\sqrt{\ln(n+2)}\right). \quad (215)$$

Apply Theorem 2.20 directly to the rounded grid, with $\alpha = h_n$, $\beta = (K_n + 1)\gamma_n$, $H = 1 + u$, and $B = u/2$. Together with the suffix edit and Equations (214) and (215), it proves that every announced policy has uniform-placement cost at least

$$n^2\Phi_{\text{RO},u}(D[M]) - O_u\left(n^{11/6}\sqrt{\ln(n+2)}\right). \quad (216)$$

The final assertion of Theorem 2.20 selects a fixed oblivious labeling before the policy seed. This proves Equation (199).

4.3 Removing the announcement

It remains to attain the benchmark without an announcement. We first record the finite cost of one canonical template: select N_{test} labels uniformly, test them in random order while immediately processing outcomes in a fixed low prefix, then process the tested residual stock in SPT order and finally run the other $n - N_{\text{test}}$ jobs raw. If $q = N_{\text{test}}/n$, direct pair accounting gives, uniformly over the empirical grid histogram and the chosen prefix,

$$\mathbb{E}C_n(\pi) = n^2F_{\text{RO},u,D}(\pi) + O_u(n), \quad (217)$$

where the template π records the low prefix and q , and $F_{\text{RO},u,D}(\pi)$ is the right-hand side of Equation (205). To verify this, expand the testing phase into one-job terms and relative-order kernels for pairs of tested jobs; expand the residual SPT block into its work against raw jobs and its minimum kernel; and add the exact raw-block cost $u(n - N_{\text{test}})(n - N_{\text{test}} + 1)/2$. For distinct occurrences the finite coefficients

$$\frac{N_{\text{test}}(N_{\text{test}} - 1)}{n(n - 1)}, \quad \frac{N_{\text{test}}(n - N_{\text{test}})}{n(n - 1)}, \quad \frac{(n - N_{\text{test}})(n - N_{\text{test}} - 1)}{n(n - 1)}$$

differ from $q^2, q(1 - q), (1 - q)^2$ by $O(1/n)$. Every kernel is bounded in terms of u , so fewer than n^2 pair terms create only $O_u(n)$ error.

Use the grid from Equation (213) and take a private random permutation. Test its first

$$k_n = \lceil n^{1/2} \rceil \quad (218)$$

labels, keep positive sample jobs pending, and form their rounded histogram $\hat{D}_n$. Among all grid prefixes containing zero and all $q \in [0, 1]$, minimize the quadratic Equation (205). Apply the selected template to the remaining labels in their unused random order, then finish the positive sample jobs.

Let D_n and D_n^{rem} be the full and remaining rounded histograms. The sampling-without-replacement histogram estimate Equation (41) gives

$$\mathbb{E} \|\hat{D}_n - D_n\|_1 \leq \sqrt{\frac{K_n + 1}{k_n}}, \quad \|D_n^{\text{rem}} - D_n\|_1 \leq \frac{2k_n}{n - k_n}. \quad (219)$$

For a fixed template, the quantities a, ℓ, ℓ_ν in Equation (205) are bounded one-draw expectations and K_ν is a bounded two-draw expectation. Inserting an intermediate product measure shows

$$|F_{\text{RO},u,D}(\pi) - F_{\text{RO},u,E}(\pi)| \leq 4(u + 1) \|D - E\|_1, \quad (220)$$

uniformly over the grid and template. The same bound holds for the minima. The maximum-density and completion-envelope results Lemmas 2.5 and 4.2 show that the minimum over this common template family is exactly $\Phi_{\text{RO},u}(D)$.

Condition on the sample. The learned template is fixed and the unused labels are uniformly ordered, so Equation (217) applies. Including the sample and its final tail, whose cost is $O_u(nk_n)$, gives the conditional hypothesis of Lemma 2.22 with $\varepsilon = O_u(k_n/n + 1/n)$. Apply that lemma with total-variation distance, $L_0 = 4(u + 1)$, and Equation (219). Finally use Equation (214). Since

$$h_n = O_u(n^{-1/6}), \quad \sqrt{(K_n + 1)/k_n} = O(n^{-1/6}), \quad k_n/n = O(n^{-1/2}),$$

the resulting bound is

$$\mathbb{E} \text{cost}(\mathcal{A}_{n,u}^*, \sigma(M)) \leq n^2 \Phi_{\text{RO},u}(D[M]) + O_u(n^{11/6}) \quad (221)$$

for every fixed labeling. Together with the announced lower bound Equation (216), this proves Theorem 4.1. The larger logarithmic lower-bound error gives the common sequence in Equation (200).

5 Randomized revealing optimization: the exact four-piece curve

The previous section identified the optimal leading online cost for every fixed input in the revealing-optimization model. We now compare that cost with the clairvoyant optimum and maximize over all inputs. The cap u is fixed, but the algorithm is not told the processing-time multiset. This section proves Theorem 5.1, the rigorous randomized half of the informal curve theorem Theorem 1.7. Its three transition points are

$$u_1^{\text{rand}} = 1, \quad u_2^{\text{rand}} = 5.048917339522306\dots, \quad u_3^{\text{rand}} = \frac{25}{4}. \quad (222)$$

Here $u_2^{\text{rand}} > 1$ is the unique root of $u^3 - 6u^2 + 5u - 1 = 0$ above one.

Theorem 5.1 (Randomized revealing-optimization curve). *For every fixed $u > 0$, the optimal randomized size-asymptotic competitive ratio against an oblivious adversary is*

$$R_{\text{RO}}^{\text{rand}}(u) = \begin{cases} 1, & 0 < u \leq u_1^{\text{rand}}, \\ \frac{u^3}{u^2 + (u-1)^3}, & u_1^{\text{rand}} \leq u \leq u_2^{\text{rand}}, \\ 1 + \frac{1}{2(\sqrt{u}-1)}, & u_2^{\text{rand}} \leq u \leq u_3^{\text{rand}}, \\ \frac{4}{3}, & u \geq u_3^{\text{rand}}. \end{cases} \quad (223)$$

More precisely, there are randomized nonanticipating algorithms $\mathcal{A}_{n,u}$ and numbers $\varepsilon_u(n) \rightarrow 0$ such that every fixed labeled vector $p \in [0, u]^n$ satisfies

$$\mathbb{E} \text{cost}(\mathcal{A}_{n,u}, p) \leq R_{\text{RO}}^{\text{rand}}(u) \text{cost}(\text{OPT}_u, p) + \varepsilon_u(n)n^2. \quad (224)$$

Conversely, for every randomized online algorithm $\mathcal{A}$ and every $\varepsilon > 0$, there are arbitrarily large n and a fixed vector $p \in [0, u]^n$ for which

$$\mathbb{E} \text{cost}(\mathcal{A}, p) \geq (R_{\text{RO}}^{\text{rand}}(u) - \varepsilon) \text{cost}(\text{OPT}_u, p). \quad (225)$$

The curve is continuous and has global maximum

$$\max_{u>0} R_{\text{RO}}^{\text{rand}}(u) = \frac{27 + 6\sqrt{3}}{23} = 1.625752384583185\dots, \quad (226)$$

attained at $u = (3 + \sqrt{3})/2$.

The preceding instance-optimal theorem identifies the best policy on each input. Here we compare that value with the clairvoyant minimum kernel and maximize over all empirical distributions. A survival-function flattening argument reduces the outer optimization to binary distributions. Two scalar maximizations give the four branches above. A sublinear sample removes advance knowledge of the input, and binary random labelings give matching Yao lower bounds.

5.1 Offline and stationary fluid coefficients

Assume initially that $u > 1$ and write

$$s = u - 1. \quad (227)$$

Let D be a finite distribution on $[0, u]$, let P, P' be independent with law D , and use the survival function

$$S_D(t) = \mathbb{P}(P > t), \quad S_D(t) = 0 \quad (t > u). \quad (228)$$

The convention at atoms does not affect any integral below. Since a clairvoyant job has effective length $1 + \min\{s, P\}$, the layer-cake identity and Corollary 2.3 give

$$\Omega_{\text{RO},u}(D) = \frac{1}{2} \left(1 + \int_0^s S_D(t)^2 dt \right), \quad (229)$$

$$\text{OPT}_u(M) = n^2 \Omega_{\text{RO},u}(D[M]) + \frac{n}{2} \int \lambda_u(p) dD[M](p) \quad (230)$$

for every empirical distribution $D[M]$.

We next specialize the maximum-density module Lemma 2.5 to a form suited to the cap. If its threshold prefix has mass a and first moment ℓ , put

$$\tau_D = \frac{1 + \ell}{a}. \quad (231)$$

The common threshold equation and the layer-cake identity give

$$\mathbb{E}(\tau_D - P)^+ = 1, \quad \int_0^{\tau_D} S_D(t) dt = \tau_D - 1. \quad (232)$$

The stationary policy tests all jobs in a uniform random order, processes a positive prefix outcome immediately after its test, and after the last test processes the residual jobs in SPT order. The following formula is the stationary pair identity specialized to a finite cap.

Lemma 5.2 (Stationary survival formula). *The doubled leading coefficient of the stationary policy is*

$$\mathcal{T}_u(D) = \begin{cases} \tau_D + \int_{\tau_D}^u S_D(t)^2 dt, & \tau_D \leq u, \\ \tau_D, & \tau_D > u. \end{cases} \quad (233)$$

Consequently the better of the raw and stationary policies has leading ratio

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s S_D(t)^2 dt}. \quad (234)$$

Proof. Let $h = 1 - a$ be the residual mass and let K_T be the minimum kernel of two residual draws, with their subprobability masses left unnormalized. The stationary prefix identity Lemma 2.4, applied first to empirical laws and then by continuity, gives the leading coefficient

$$(1 + \ell) \left(1 - \frac{a}{2} \right) + \frac{K_T}{2}. \quad (235)$$

If $\tau_D \leq u$, the threshold property gives

$$K_T = \tau_D h^2 + \int_{\tau_D}^u S_D(t)^2 dt, \quad (1 + \ell) \left(1 - \frac{a}{2} \right) = \frac{\tau_D}{2} (1 - h^2).$$

Their sum is one half of the first line of (233). If $\tau_D > u$, the whole distribution is in the prefix and (235) equals $\tau_D/2$. Raw execution has leading coefficient $u/2$, and division by (229) proves (234). $\square$

This is where the instance-optimal theorem enters the curve calculation. In Equation (205), $q = 0$ is the raw policy and $q = 1$ is precisely the stationary maximum-density policy. Hence

$$2\Phi_{\text{RO},u}(D) \leq \min\{u, \mathcal{T}_u(D)\}, \quad (236)$$

and Equation (234) is an upper bound on the instance-specific ratio $\Phi_{\text{RO},u}(D)/\Omega_{\text{RO},u}(D)$. Interior values of q can strictly improve individual inputs; the point of the next two subsections is that the two endpoint policies already give the sharp value after maximizing over all D . The binary lower bounds at the end apply to every policy, not only to these endpoints.

5.2 Flattening the survival function

We now maximize (234) over arbitrary D . Suppose first that $1 \leq \tau_D \leq u$ and put

$$y = \frac{\tau_D - 1}{\tau_D}. \quad (237)$$

By (232), y is the average of S_D on $[0, \tau_D]$. Since S_D is nonincreasing, its average on every initial subinterval is at least y . Thus Cauchy–Schwarz gives

$$\int_0^a S_D(t)^2 dt \geq ay^2 \quad (0 \leq a \leq \tau_D), \quad (238)$$

and $S_D(t) \leq y$ almost everywhere for $t > \tau_D$.

Replace S_D on $(0, \tau_D)$ by the constant y and leave its tail unchanged. This remains a nonincreasing survival function, preserves (232) and $\mathcal{T}_u(D)$, and by (238) can only decrease the offline coefficient. Hence it can only increase the ratio. We may therefore assume

$$S_D(t) = y \quad (0 < t < \tau_D). \quad (239)$$

If $u - 1 \leq \tau_D \leq u$, the offline integral lies entirely in the flat part, and $S_D(t)^2 \leq y^2$ on the remaining tail. It follows that

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s S_D(t)^2 dt} \leq B_u(\tau_D) := \frac{\tau_D + (u - \tau_D)y^2}{1 + (u - 1)y^2}. \quad (240)$$

The numerator displayed on the right is at most u , so the raw minimum is inactive.

Now suppose $\tau_D < u - 1$. Put

$$L = u - 1 - \tau_D, \quad E = \int_{\tau_D}^{u-1} S_D(t)^2 dt, \quad G = \int_{u-1}^u S_D(t)^2 dt. \quad (241)$$

Monotonicity gives

$$0 \leq E \leq Ly^2, \quad G \leq E/L. \quad (242)$$

Consequently

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s S_D(t)^2 dt} \leq \frac{\tau_D + \frac{u - \tau_D}{u - 1 - \tau_D} E}{1 + \tau_D y^2 + E}. \quad (243)$$

For fixed τ_D this is linear-fractional in E , so its derivative has a constant sign. Its maximum on $[0, Ly^2]$ occurs at an endpoint. The two endpoint values are

$$A(\tau_D) := \frac{\tau_D}{1 + \tau_D y^2} = \frac{\tau_D^2}{\tau_D^2 - \tau_D + 1}, \quad (244)$$

$$B_u(\tau_D) = \frac{\tau_D + (u - \tau_D)y^2}{1 + (u - 1)y^2}. \quad (245)$$

They are attained by rectangular survival functions: respectively a binary distribution on $\{0, \tau_D\}$ and one on $\{0, u\}$.

It remains to cover $\tau_D > u$. Equation (232) and the zero extension of S_D give $\int_0^u S_D(t) dt = \tau_D - 1 \geq u - 1$. The average of a nonincreasing function on $[0, u - 1]$ is no smaller than its average on $[0, u]$, and hence

$$\int_0^{u-1} S_D(t)^2 dt \geq (u - 1) \left(\frac{\tau_D - 1}{u} \right)^2 \geq \frac{(u - 1)^3}{u^2}. \quad (246)$$

Here the raw policy is chosen, and its ratio is at most $B_u(u)$. We have proved the reduction

$$\sup_D \frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s S_D(t)^2 dt} \leq \max \left\{ \sup_{1 \leq \tau \leq u-1} A(\tau), \sup_{1 \leq \tau \leq u} B_u(\tau) \right\}, \quad (247)$$

where the first supremum is omitted for $u < 2$. In fact equality holds in (247), because the rectangular survival functions described above attain both families.

5.3 The two scalar maximizations

The first family satisfies

$$A'(\tau) = \frac{\tau(2 - \tau)}{(\tau^2 - \tau + 1)^2}. \quad (248)$$

Thus its maximum is attained at $\tau = \min\{2, u - 1\}$ and equals $4/3$ as soon as $u \geq 3$.

For the second family, substitution of (237) gives

$$B_u(\tau) = \frac{\tau^2 u + 2\tau^2 - 2\tau u - \tau + u}{\tau^2 u - 2\tau u + 2\tau + u - 1}. \quad (249)$$

The sign of its derivative is the sign of

$$g_u(\tau) = (2\tau - 1)^2 - u(\tau - 1)^2. \quad (250)$$

For $u \leq 4$ this is positive on $[1, \infty)$. For $u > 4$, its unique root above one is

$$\tau_*(u) = \frac{u - 2 + \sqrt{u}}{u - 4} = \frac{\sqrt{u} - 1}{\sqrt{u} - 2}, \quad (251)$$

and B_u increases before this point and decreases afterwards. The identity $\tau_*(u) = u$ is equivalent to

$$u^3 - 6u^2 + 5u - 1 = 0. \quad (252)$$

The cubic is strictly increasing beyond its last critical point and changes sign there exactly once; its unique root above one is u_2^{rand} from (222). Hence

$$\max_{1 \leq \tau \leq u} B_u(\tau) = B_u(u) = \frac{u^3}{u^3 - 2u^2 + 3u - 1} = \frac{u^3}{u^2 + (u - 1)^3}, \quad u \leq u_2^{\text{rand}}, \quad (253)$$

$$= B_u(\tau_*(u)) = 1 + \frac{1}{2(\sqrt{u} - 1)}, \quad u \geq u_2^{\text{rand}}. \quad (254)$$

There is no missing branch from A . On $2 \leq u \leq 3$, clearing the positive denominators in $B_u(u) - A(u - 1)$ and putting $v = u - 2$ leaves

$$v^4 + 3v^3 + 3v^2 + 3v + 3 > 0.$$

On $3 \leq u \leq u_2^{\text{rand}}$, one has $B_u(u) > 4/3$. On $u_2^{\text{rand}} \leq u \leq u_3^{\text{rand}}$, (254) is at least $4/3$, with equality exactly at $u = u_3^{\text{rand}}$; afterwards $A(2) = 4/3$ is larger. Combining these observations with (247) proves the announced upper bound

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s S_D(t)^2 dt} \leq R_{\text{RO}}^{\text{rand}}(u) \quad (255)$$

for every finite distribution D .

The junction statements now follow directly. Equations (252) to (254) give continuity at u_2^{rand} , and the square-root branch equals $4/3$ at $u = u_3^{\text{rand}}$. Differentiating (253) shows that its only interior maximum above one occurs at $u = (3 + \sqrt{3})/2$; substitution gives (226).

5.4 Worst-case upper bound from instance optimality

No second learning argument is needed. Apply the universal unannounced policy of Theorem 4.1 to $D = D[M]$. The endpoint comparison Equation (236) and the scalar bound Equation (255) give

$$\Phi_{\text{RO},u}(D[M]) \leq R_{\text{RO}}^{\text{rand}}(u) \Omega_{\text{RO},u}(D[M]).$$

The exact empirical identity Equation (230) has a nonnegative linear correction. Therefore the instance-optimal upper bound immediately yields

$$\mathbb{E} \text{ALG}_{\mathcal{A}_{n,u}} \leq R_{\text{RO}}^{\text{rand}}(u) \text{OPT}_u + o_u(n^2), \quad (256)$$

which is (224). For $0 < u \leq 1$, raw execution is exactly clairvoyantly optimal, so no learning is required.

5.5 Binary oblivious lower bounds

It remains to show that the two endpoint policies used above cannot be improved in the worst case by a more adaptive algorithm. We first record a binary random-labeling envelope.

Lemma 5.3 (Binary cap envelope). *Fix $u > 1$, $x \in [0, 1]$, and put $z = 1 - x$. Let M_n have $xn + O(1)$ jobs of value u and $zn + O(1)$ zero jobs. For every randomized nonanticipating algorithm, even an announced one, some fixed labeling satisfies*

$$\frac{\mathbb{E} \text{ALG}}{n^2} \geq \begin{cases} \frac{u}{2} - o_u(1), & x \geq (u-1)/u, \\ \frac{1+x+ux^2}{2} - o_u(1), & x \leq (u-1)/u. \end{cases} \quad (257)$$

Every labeling has

$$\frac{\text{OPT}_u}{n^2} = \frac{1 + (u-1)x^2}{2} + O_u(1/n). \quad (258)$$

Proof. First fix a deterministic policy. If it does not complete all jobs, the claim is immediate. Otherwise place the occurrences uniformly on the public labels, distinguishing equal-valued occurrences as tokens. By Lemma 2.11, the analytically exposed occurrence word is a uniform permutation, and the test/raw choice before its next entry is predictable.

Take $\delta_n = n^{-1/4}$. The predictable-permutation bound Lemma 2.15, applied to the zero indicator and the test selector, gives one event of probability $1 - o(1)$ on which every first-touch prefix ending with at least $\delta_n n$ untouched labels and containing t tests satisfies

$$|N_0(t) - zt| \leq \Delta_n, \quad \Delta_n = O(\delta_n^{-1} \sqrt{n \ln n}) = o(n). \quad (259)$$

Here $N_0(t)$ is the number of tested zeros. The event is simultaneous, so we do not condition on the policy's adaptive stopping time.

Edit the suffix by running raw every still-untouched job that the original schedule would later test, deleting its later known-processing operation. Only $\delta_n n + O(1)$ jobs change, so the edit changes total completion cost by $o_u(n^2)$. Let q be the final tested fraction of the edited schedule. On (259), every prefix of its completion curve is dominated, up to $o(1)$ vertical error, by the divisible items

item	completion capacity	work per completion
zero-test module	zq	$1/z$
tested cap processing	xq	u
raw jobs	$1 - q$	u

When $z = 0$, the zero-test module is simply omitted. Indeed, t tests complete at most $zt + \Delta_n$ zeros and expose at most $xt + \Delta_n$ cap jobs; raw and known cap processing both use u work per completion. The ordinary divisible-knapsack exchange therefore upper-bounds completed mass at every work value. Integrating the remaining mass gives a lower bound on total completion time.

If $z \leq 1/u$, every displayed item costs at least u per completion, so the normalized area is at least $u/2 - o_u(1)$. If $z \geq 1/u$, the zero-test module comes first. For fixed q its remaining-mass area is

$$F_z(q) = q - \frac{zq^2}{2} + \frac{u}{2}(1 - zq)^2, \quad F'_z(q) = (1 - zq)(1 - uz). \quad (260)$$

It is minimized at $q = 1$, giving $F_z(1) = (1 + x + ux^2)/2$. The suffix edit, vertical repair, and bad event all contribute $o_u(n^2)$.

The argument holds for each deterministic policy. For a randomized policy, fix its seed, average the result over seeds and the finite uniform placement space, and select a placement whose seed-expected cost is at least the joint average. This placement is fixed before the seed and is therefore an oblivious input. Finally, the effective lengths of the two job types are 1 and u , so (258) follows from Lemma 2.2. $\square$

We now select the binary masses. If $u_1^{\text{rand}} < u \leq u_2^{\text{rand}}$, take

$$x = \frac{u - 1}{u}. \quad (261)$$

This is the boundary $z = 1/u$ in Lemma 5.3, and its online and offline coefficients have ratio

$$\frac{u}{1 + (u - 1)((u - 1)/u)^2} = \frac{u^3}{u^2 + (u - 1)^3}. \quad (262)$$

If $u_2^{\text{rand}} \leq u \leq u_3^{\text{rand}}$, take

$$x = \frac{1}{\sqrt{u} - 1}. \quad (263)$$

The definition of u_2^{rand} is precisely the point from which $x \leq (u - 1)/u$. Since $(u - 1)x^2 = 1 + 2x$, the second line of (257) divided by (258) becomes

$$1 + \frac{x}{2} = 1 + \frac{1}{2(\sqrt{u} - 1)}. \quad (264)$$

Irrational masses are approximated by empirical rational frequencies; all rounding and diagonal terms are $O_u(n)$.

For the last branch, the cap distribution is replaced by a fixed bounded one.

Lemma 5.4 (Balanced zero–two lower bound). *For every $u \geq u_3^{\text{rand}}$ and every randomized algorithm, arbitrarily large even n admit a fixed labeling with $n/2$ jobs of value zero and $n/2$ jobs of value two such that*

$$\mathbb{E} \text{ALG} \geq n^2 - o_u(n^2), \quad \text{OPT}_u = \frac{3}{4}n^2 + O(n). \quad (265)$$

Proof. Use the same trace bijection, predictable-urn event, and suffix edit as in Lemma 5.3. For a realized tested fraction q , the three fluid items now have capacities and costs

item	completion capacity	work per completion
zero-test module	$q/2$	2
tested two processing	$q/2$	2
raw jobs	$1 - q$	u

Every item costs at least two per completion. Thus the completion envelope lies below $\min\{1, t/2\}$ independently of q , whose remaining-mass area is one. This proves the online estimate after the $o_u(n^2)$ repairs and finite Yao selection Lemma 2.17. Since $u > 3$, the two effective lengths are one and three; their equally weighted minimum kernel gives offline coefficient $3/4$. $\square$

Equations (262) and (264), together with Lemma 5.4, match (256) in every range. For $0 < u \leq 1$, raw execution is both an online and a clairvoyant optimum. This completes the proof of the exact curve in Theorem 5.1.

5.6 The randomized obligatory endpoint

We now pass from the finite-cap randomized curve to the unbounded obligatory endpoint. The adversary fixes the labeled vector $p \in [0, \infty)^n$ before the algorithm draws its random bits. The multiset is not announced, and no upper bound on the processing times is assumed. The finite-cap zero–two lower bound transfers directly. The upper bound needs one additional uniformization: a cutoff and fallback protect the finite-cap stationary policy from an arbitrarily large value missed by its sample. Letting the cutoff grow slowly also makes the same policy instance-optimal on every bounded class. This subsection proves Theorem 5.5, the rigorous randomized part of the informal obligatory-testing result Theorem 1.1 from the introduction.

Theorem 5.5 (Randomized obligatory testing). *There are a universal constant C and a randomized nonanticipating algorithm $\mathcal{A}_n^*$, depending only on n , such that every fixed vector $p \in [0, \infty)^n$ satisfies*

$$\mathbb{E} \text{cost}(\mathcal{A}_n^*, p) \leq \frac{4}{3} \text{cost}(\text{OPT}, p) + Cn^{19/10} \leq \left(\frac{4}{3} + 2Cn^{-1/10} \right) \text{cost}(\text{OPT}, p). \quad (266)$$

A fixed-cutoff companion $\mathcal{A}_n^{(32)}$, using $B = 32$ and the more conservative fallback criterion $\hat{\kappa} < 2/B$, satisfies the sharper explicit estimate

$$\mathbb{E} \text{cost}(\mathcal{A}_n^{(32)}, p) \leq \frac{4}{3} \text{cost}(\text{OPT}, p) + 20378n^{7/4}. \quad (267)$$

Conversely, for every randomized algorithm $\mathcal{A}$ and every $\varepsilon > 0$, arbitrarily large even n admit a fixed labeling of $n/2$ zero jobs and $n/2$ jobs of processing time two such that

$$\mathbb{E} \text{cost}(\mathcal{A}, p) \geq \left(\frac{4}{3} - \varepsilon \right) \text{cost}(\text{OPT}, p), \quad \text{cost}(\text{OPT}, p) = \frac{3}{4}n^2 + n. \quad (268)$$

Hence the exact randomized size-asymptotic competitive ratio against an oblivious adversary is $R_{\text{OT}}^{\text{rand}} = 4/3$.

For an announced multiset, the algorithmic core is the processing-time prefix with maximum completions per unit discovery work. Testing jobs in a uniformly random order gives an exact pair identity and a $4/3$ algebraic certificate. The universal algorithm learns an approximately separated prefix from a sublinear sample; a cutoff and a fallback schedule protect it from unbounded values missed by the sample. The matching lower bound is inherited from the zero–two endpoint of the finite-cap curve. We first prove the uniform upper bound and then give the short transfer.

5.6.1 Normalization of the offline pair identity

For a fixed vector p , use $D[p]$ and its minimum kernel $K(D[p])$ from Equation (19). The specialization in Equation (9) and the exact linear correction are

$$\Omega_{\text{OT}}(D[p]) = \frac{1 + K(D[p])}{2}, \quad \Delta_{\text{OT}}(p) = \frac{n + \sum_i p_i}{2}. \quad (269)$$

The empirical effective-length identity Equation (11), with $\ell(p) = 1 + p$, gives

$$\text{OPT}(p) = n^2 \Omega_{\text{OT}}(D[p]) + \Delta_{\text{OT}}(p). \quad (270)$$

The linear correction $\Delta_{\text{OT}}(p)$ will be kept rather than hidden: this makes the analysis valid for unbounded processing times.

5.6.2 Stationary prefixes

Use the processing-time-prefix notation Equations (18) and (19) and the coefficient $F_{\text{OT}, D[p]}(E)$ from Equation (20). We require that E contain every zero job; otherwise a supposedly deferred zero completes at its test. The stationary policy and its exact cost are supplied by the common pair identity Lemma 2.4.

For a nonempty prefix define its completion density by

$$\kappa(E) = \frac{a}{1 + \ell_E}. \quad (271)$$

Choose a maximum-density prefix and put

$$\tau = \frac{1 + \ell_E}{a} = \frac{1}{\kappa(E)}. \quad (272)$$

All zeros belong to every maximizer: adding a zero increases the numerator without increasing the denominator. The empirical prefix corollary Corollary 2.6 lets us choose a maximizer with

$$\tau = \tau_{D[p]}, \quad p_i \leq \tau \ (i \in E), \quad p_i \geq \tau \ (i \notin E), \quad (273)$$

where tied occurrences may be assigned to either side.

The next identity is the algebraic source of the constant $4/3$.

Lemma 5.6 (Exact and robust certificates). *For every ordered prefix split with $a > 0$ and $\tau = (1 + \ell_E)/a$,*

$$2\tau(4\Omega_{\text{OT}}(D[p]) - 3F_{\text{OT}, D[p]}(E)) = (\tau - 2)^2 + 4(\tau K_E - \ell_E^2) + \tau(K_T - \tau h^2). \quad (274)$$

Consequently:

- (i) if early values are at most τ and late values are at least τ , then $F_{\text{OT},D[p]}(E) \leq 4\Omega_{\text{OT}}(D[p])/3$;
- (ii) if early values are at most $\tau + s$ and late values are at least $\tau - s$, where $s \geq 0$, then

$$F_{\text{OT},D[p]}(E) \leq \frac{4}{3}\Omega_{\text{OT}}(D[p]) + \frac{2}{3}s. \quad (275)$$

Proof. Expansion using $\ell_E = a\tau - 1$, $h = 1 - a$, and Equation (19) gives Equation (274). Under exact separation, $xy \leq \tau \min\{x, y\}$ for early x, y , hence $\ell_E^2 \leq \tau K_E$; for late x, y , one has $\min\{x, y\} \geq \tau$, hence $K_T \geq \tau h^2$. This proves (i).

Under the relaxed inequalities,

$$\tau K_E - \ell_E^2 \geq -sK_E, \quad K_T - \tau h^2 \geq -sh^2.$$

Moreover $K_E \leq a\ell_E \leq a^2\tau$. Substitution into Equation (274) yields

$$F_{\text{OT},D[p]}(E) \leq \frac{4}{3}\Omega_{\text{OT}}(D[p]) + \frac{2}{3}sa^2 + \frac{1}{6}sh^2 \leq \frac{4}{3}\Omega_{\text{OT}}(D[p]) + \frac{2}{3}s,$$

because $(2/3)a^2 + (1/6)(1-a)^2 \leq 2/3$ on $[0, 1]$. $\square$

Combining the exact certificate with Lemma 2.4 and Equation (270) already gives a finite statement useful below.

Corollary 5.7 (Announced multiset). *If the multiset is announced, the stationary policy for a maximum-density prefix satisfies*

$$\mathbb{E} \text{ALG}_{\text{stat}}(p, E) \leq \frac{4}{3} \text{OPT}(p).$$

Indeed, the correction in Equation (21) is at most $\Delta_{\text{OT}}(p)$. Equality in the quadratic coefficient of Equation (274) forces $\tau = 2$, zero mass $1/2$, and all positive mass at two. This predicts the lower-bound instance before any probabilistic argument is used.

We also use the following tail-safe estimate. Direct expansion gives

$$F_{\text{OT},D[p]}(E) - \Omega_{\text{OT}}(D[p]) = \frac{h + a\ell_E - K_E}{2}. \quad (276)$$

If all early values are at most $L \geq 1$, then $\ell_E \leq aL$ and therefore

$$F_{\text{OT},D[p]}(E) \leq \Omega_{\text{OT}}(D[p]) + \frac{L}{2}. \quad (277)$$

5.6.3 Learning the prefix from a sublinear sample

For the main algorithm, set the slowly growing cutoff

$$B = B_n = 32 + n^{1/20}. \quad (278)$$

For an integer d , let $\eta = B/d$. Keep zero as a separate category, partition $(0, B]$ into the consecutive bins $((b-1)\eta, b\eta]$, and add one overflow category for values above B . A finite positive value is rounded upward to the endpoint $q_b = b\eta$.

Let S be a uniformly random subset of k labels. If μ_b and $\hat{\mu}_b$ are the population and sample fractions of category b , put

$$\Delta = \sum_b |\hat{\mu}_b - \mu_b|. \quad (279)$$

Lemma 5.8 (Histogram error). *For $1 \leq k < n$,*

$$\mathbb{E}\Delta \leq \sqrt{\frac{d+2}{k}}.$$

Proof. This is Equation (41), with the zero bin and the overflow bin included among the $d+2$ categories. $\square$

For a finite category prefix J , define

$$\hat{a}(J) = \sum_{b \in J} \hat{\mu}_b, \quad \hat{\ell}(J) = \sum_{b \in J} q_b \hat{\mu}_b, \quad \hat{\kappa}(J) = \frac{\hat{a}(J)}{1 + \hat{\ell}(J)}.$$

Let $\hat{\kappa}$ be the maximum and $\hat{\tau} = 1/\hat{\kappa}$. The learned early set is not an arbitrary maximizing category prefix. It is its *threshold closure*

$$\hat{J} = \{\text{finite categories } b : q_b \leq \hat{\tau}\}. \quad (280)$$

Lemma 5.9 (Threshold closure). *The prefix Equation (280) has the same maximum sample density. Every category in it has endpoint at most $\hat{\tau}$, and every finite category outside it has endpoint greater than $\hat{\tau}$, including categories with zero sample count.*

Proof. The removal/addition calculation in Corollary 2.6 applies to every positive-mass sample category. Empty categories do not change the sample statistics, and adding mass at equality preserves the identity $1 + \hat{\ell} = \hat{a}\hat{\tau}$. Closing the prefix therefore preserves optimality and makes the two pointwise assertions tautological. $\square$

For large n , set

$$k = \lfloor n^{3/4} \rfloor, \quad d = \lfloor n^{1/4} \rfloor, \quad \eta = B/d. \quad (281)$$

The algorithm draws a uniform permutation and first tests its first k jobs, leaving positive sample jobs pending. It computes the closed sample prefix. If

$$\hat{\kappa} < \frac{1}{B}, \quad (282)$$

it enters *fallback mode*: test every remaining job and then process all positive jobs in SPT order. Otherwise it enters *learned mode*: process positive sampled early jobs in SPT order; test the remaining jobs in their random order, processing early outcomes immediately; and finish the deferred tail in SPT order. Overflow is never early.

5.6.4 Good samples, fallback, and bad-event excess

Put

$$\gamma = \frac{1}{B(B+1)}, \quad G = \{\Delta \leq \gamma\}. \quad (283)$$

Markov's inequality, recalled in Section 2.6, and Lemma 5.8 give

$$\mathbb{P}(G^c) \leq B(B+1) \sqrt{\frac{d+2}{k}}. \quad (284)$$

Assume G and learned mode. Let E be the population set selected by the closed category prefix, and let $a, \ell_E, \tau = (1 + \ell_E)/a$ be its true statistics. Since $\hat{\tau} \leq B$ and $\hat{a} = (1 + \hat{\ell})/\hat{\tau} \geq 1/B$, we have $a \geq 1/(B+1)$. For this fixed category set,

$$|a - \hat{a}| \leq \Delta, \quad |\ell_E - \hat{\ell}| \leq B\Delta + \eta.$$

Consequently

$$|\tau - \hat{\tau}| \leq 2B(B+1)\Delta + (B+1)\eta. \quad (285)$$

Every early actual value is at most $\hat{\tau}$, while every late value is greater than $\hat{\tau} - \eta$. Thus the robust certificate applies with

$$s = 2B(B+1)\Delta + (B+2)\eta. \quad (286)$$

The sample-first order is not exactly the fully stationary comparator, but the difference is only lower order. Conditional on the unordered sample, write e_S, e_T for the sample and remaining early counts and

$$W_S = k + \sum_{i \in E \cap S} p_i, \quad W_T = n - k + \sum_{i \in E \setminus S} p_i.$$

Comparing the two independent random internal orders with one fully random order, the early-cost difference is

$$\frac{e_T W_S - e_S W_T}{2} \leq \frac{1}{2} n k (1 + B).$$

Postponing positive sampled early jobs until all sample tests finish adds at most $k^2(1+B)$. The late tail is identical because every zero is early. Therefore, on a good learned sample,

$$\mathbb{E}[\text{ALG} \mid S] \leq \frac{4}{3} \text{OPT} + n^2 \left[\frac{4}{3} B(B+1)\Delta + \frac{2}{3} (B+2)\eta \right] + (1+B)(nk/2 + k^2). \quad (287)$$

On a good fallback sample, comparison of sample and population densities shows that the true maximum-density reciprocal satisfies $\tau_{D[p]} \geq B/4 \geq 8$. The exact threshold inequalities imply

$$K(D[p]) \geq \frac{(\tau_{D[p]} - 1)^2}{\tau_{D[p]}}, \quad \Omega_{\text{OT}}(D[p]) \geq \frac{\tau_{D[p]} - 1 + 1/\tau_{D[p]}}{2} \geq \frac{57}{16}. \quad (288)$$

For every test order, test-all-then-SPT satisfies

$$\text{ALG}_{\text{fb}} \leq \text{OPT} + \frac{n(n-1)}{2}. \quad (289)$$

Indeed, if there are z zeros, their total displacement from the first z SPT positions is at most $zn - z(z+1)/2$, while delaying the positive SPT tail adds $(n-z)(n-z-1)/2$; the sum is $n(n-1)/2$. Combining Equations (288) and (289) gives $\text{ALG}_{\text{fb}} < 4 \text{OPT}/3$.

Because processing times are unbounded, a bad-event estimate must control excess over $4 \text{OPT}/3$, not raw cost. In learned mode every early value is at most B , so Equation (277) bounds the stationary excess by $(B/2)n^2$; in fallback mode Equation (289) bounds it by $n^2/2$. Averaging Equation (287) and these bad-event bounds, and using Equation (284), yields the uniform estimate

$$\mathbb{E} \text{ALG} \leq \frac{4}{3} \text{OPT} + n^2 \left[\frac{B(B+1)(3B+8)}{6} \sqrt{\frac{d+2}{k}} + \frac{2B(B+2)}{3d} \right] + (1+B)(nk/2 + k^2). \quad (290)$$

For $n \geq 12^4$, the floors in Equation (281) give $\sqrt{(d+2)/k} \leq 2n^{-1/4}$, $1/d \leq 2n^{-1/4}$, and $k \leq n^{3/4}$. Since $B_n = O(n^{1/20})$, the three error terms in Equation (290) are respectively $O(n^{19/10})$, $O(n^{37/20})$, and $O(n^{9/5})$. Enlarging one universal constant covers smaller n and proves the first inequality in Equation (266). The bound $\text{OPT} \geq n^2/2$ gives its multiplicative form.

For completeness, the fixed-cutoff companion uses $B = 32$ and the more conservative fallback criterion $\hat{\kappa} < 2/B$. Repeating the same calculation with learned threshold at most $B/2$ gives

$$\mathbb{E} \text{ALG} \leq \frac{4}{3} \text{OPT} + n^2 \left[9472 \sqrt{\frac{d+2}{k}} + \frac{704}{d} \right] + 17(nk/2 + k^2).$$

The preceding floor bounds give $\mathbb{E} \text{ALG} \leq 4 \text{OPT} / 3 + 20378n^{7/4}$, with the same deliberately loose constant covering smaller n . This proves Equation (267).

5.6.5 Transfer of the zero–two lower bound

Fix the finite cap $u = u_3^{\text{rand}}$. Any randomized obligatory algorithm can be viewed as a revealing-optimization algorithm at this cap that simply never uses raw execution. Apply Lemma 5.4. Its fixed hard input has only values zero and two, so every revealing effective length is

$$\min\{u_3^{\text{rand}}, 1 + p_j\} = 1 + p_j.$$

The online transcript and the offline optimum are therefore exactly the same in the finite-cap and obligatory models. For even n , the common offline value is

$$\text{OPT} = \frac{3}{4}n^2 + n,$$

whereas the selected labeling has expected online cost $n^2 - o(n^2)$. Its ratio tends to $4/3$, which proves Equation (268). Together with Equation (266), this completes the proof of Theorem 5.5.

6 Obligatory-testing instance optimality

The goal of this section is to prove Theorem 6.1, the rigorous version of the “one algorithm for every input” statement Theorem 1.2. The algorithm is exactly the growing-cutoff policy $\mathcal{A}_n^*$ from Section 5.6. The new point is that, on every bounded input, the policy attains the best leading cost available even to an announced policy.

For a multiset $M = \{p_1, \dots, p_n\}$, we define below an explicit number $\Phi_{\text{OT}}(D[M])$, using the common empirical notation Equation (7). It is obtained by scanning the processing times in increasing order, choosing the prefix with the largest number of completions per unit of discovery work, and evaluating the stationary policy for that prefix.

Theorem 6.1 (Obligatory-testing instance optimality). *There is a single randomized nonanticipating policy $\mathcal{A}^*$, depending only on n and not on the multiset, with the following property. For every fixed $L < \infty$ there is a deterministic sequence $\varepsilon_L(n) \rightarrow 0$ such that, for every size- n multiset $M \subseteq [0, L]$ and every fixed labeling σ ,*

$$\mathbb{E} \text{cost}(\mathcal{A}^*, \sigma(M)) \leq n^2 \Phi_{\text{OT}}(D[M]) + \varepsilon_L(n)n^2. \quad (291)$$

Conversely, for every randomized policy $\mathcal{A}$, even an announced policy, some fixed oblivious labeling σ satisfies

$$\mathbb{E} \text{cost}(\mathcal{A}, \sigma(M)) \geq n^2 \Phi_{\text{OT}}(D[M]) - \varepsilon_L(n)n^2. \quad (292)$$

One may take $\varepsilon_L(n) = O_L(n^{-1/5})$. Thus $\mathcal{A}^$ is asymptotically instance-optimal, uniformly over all bounded inputs.*

Both estimates below are stated with the common order $O_L(n^{-1/5})$.

Theorem 6.2 (When instance optimality is impossible). (i) *There is no deterministic instance-optimal policy, even when all job lengths lie in $[0, 2]$. More precisely, for every deterministic policy $\mathcal{A}$, even an announced one, and every even n , there is a fixed labeling of the input with $n/2$ zero jobs and $n/2$ length-two jobs for which*

$$\text{cost}(\mathcal{A}) \geq \frac{9}{8}n^2. \quad (293)$$

On the same input, an announced randomized policy $\mathcal{A}'$ has

$$\text{cost}(\mathcal{A}') = n^2 + \frac{3}{4}n.$$

Thus $\text{cost}(\mathcal{A}) > \text{cost}(\mathcal{A}') + n^2/8 - O(n)$.

(ii) *If job lengths are not bounded by a fixed constant, there is no randomized instance-optimal policy with one uniform additive $o(n^2)$ term. More precisely, for every randomized policy $\mathcal{A}$ independent of the input and every sufficiently large n , there is an input with lengths in $[0, n^2]$, a fixed labeling, and an announced randomized policy $\mathcal{A}'$ such that*

$$\text{cost}(\mathcal{A}) \geq \text{cost}(\mathcal{A}') + \frac{1}{8}n^2. \quad (294)$$

Proof. As in the introduction, $\text{cost}(\mathcal{A})$ includes the expectation over the private randomness of $\mathcal{A}$. For part (i), write $n = 2m$. Against the fixed deterministic policy, reveal length two on the first m labels it tests and zero on the remaining m . As usual, replay freezes this interaction into one fixed oblivious labeling. If the policy does not finish, the claim is immediate. Otherwise, let h be the number of length-two jobs completed before the first zero is tested. By that time all m positive jobs have been tested, so at least $m + 2h$ work has elapsed. The first h completed positive jobs contribute at least $3h(h+1)/2$. At that point the remaining jobs consist of m unit tests of zero jobs and $m - h$ processing operations of length two. Even SPT order for this remaining work gives

$$\begin{aligned} \text{cost}(\mathcal{A}, M) &\geq \frac{3h(h+1)}{2} + (2m-h)(m+2h) + \frac{m(m+1)}{2} \\ &\quad + m(m-h) + (m-h)(m-h+1) \\ &= \frac{9}{2}m^2 + \frac{1}{2}h^2 + \frac{3}{2}m + \frac{1}{2}h \geq \frac{9}{8}n^2. \end{aligned}$$

The comparison policy $\mathcal{A}'$ tests in a uniformly random order, completes zeros at their tests, defers every length-two job, and finally processes the deferred jobs. This is the stationary policy with the zero jobs early. Its exact cost identity Equation (21) gives

$$\text{cost}(\mathcal{A}') = n^2 + \frac{3}{4}n.$$

This proves the first claim.

For part (ii), put $H = n^2$ and first run $\mathcal{A}$ on the input M^0 in which every job has length H . For a fixed private seed, let X_k be the number of completed jobs immediately before the k th test. Every processing operation contributes the same $Hn(n+1)/2$ in total, regardless of its position. The k th test delays $n - X_k$ unfinished jobs. Hence the excess over the policy $\mathcal{A}'$ that tests and processes each job immediately satisfies

$$\text{cost}(\mathcal{A}) - \text{cost}(\mathcal{A}') = \mathbb{E} \sum_{k=1}^n (k-1 - X_k). \quad (295)$$

If this sum has expectation at least $n^2/8$, we are done.

Otherwise, replace one uniformly random label by a zero and leave every other length equal to H ; call the resulting input M^1 . Conditional on the seed, the rank K at which the zero label is tested is uniform on $\{1, \dots, n\}$. Until that test, the run agrees with the all- H run, and therefore X_K length- H jobs complete before the zero. From Equation (295),

$$\mathbb{E}X_K = \frac{1}{n}\mathbb{E}\sum_{k=1}^n X_k > \frac{n-1}{2} - \frac{n}{8} = \frac{3n}{8} - \frac{1}{2}. \quad (296)$$

Forcing the zero job behind X_K length- H jobs adds HX_K to the clairvoyant SPT lower bound. On the other hand, an announced policy can test in a private random order, defer positive jobs until it finds the zero, and then finish the positive jobs. Its completion order is SPT, and moving its unit tests away from the corresponding processing operations costs at most n^2 . Thus

$$\text{cost}(\mathcal{A}) - \text{cost}(\mathcal{A}') \geq H \left(\frac{3n}{8} - \frac{1}{2} \right) - n^2.$$

For $H = n^2$ this exceeds $n^2/8$ for all sufficiently large n . Averaging over the possible zero labels shows that one fixed oblivious labeling has the same lower bound. This proves part (ii). $\square$

6.1 The maximum-density benchmark

We first define the benchmark for an arbitrary finitely supported probability distribution D on $[0, \infty)$. Apply the maximum-density module Lemma 2.5, choose its threshold selector g_* , and put

$$a_* = a_D(g_*), \quad w_* = w_D(g_*) = a_*\tau_D. \quad (297)$$

The residual measure ν from Equation (26) has mass $1 - a_*$ and support in $[\tau_D, \infty)$. For an empirical measure one may equivalently sort the occurrences and maximize Equation (28) over processing-time prefixes E containing all zeros. A tied boundary class may be placed on either side, by Corollary 2.6.

Define

$$\Phi_{\text{OT}}(D) = w_* \left(1 - \frac{a_*}{2} \right) + \frac{1}{2} \iint \min\{p, q\} d\nu(p) d\nu(q). \quad (298)$$

For $D = D[M]$ and a whole-occurrence maximizing prefix E , this is precisely $F_{\text{OT}, D[M]}(E)$ from Equation (20). The next lemma proves both that the value is independent of choices at a tied atom and that it lower-bounds every fluid policy.

Lemma 6.3 (Pointwise completion envelope). *For scaled time $s \geq 0$, let*

$$\mathcal{C}_{\text{OT}, D}(s) = \max \left\{ D(\{0\})T + \sum_{p>0} c_p : \begin{array}{l} 0 \leq T \leq 1, \quad 0 \leq c_p \leq D(\{p\})T, \\ T + \sum_{p>0} pc_p \leq s \end{array} \right\}. \quad (299)$$

Then $\mathcal{C}_{\text{OT}, D}$ is attained simultaneously at all times by the stationary maximum-density schedule: repeat infinitesimal full modules g_ until all jobs have been tested, then process the residual measure ν in SPT order. Moreover,*

$$\Phi_{\text{OT}}(D) = \int_0^\infty (1 - \mathcal{C}_{\text{OT}, D}(s)) ds. \quad (300)$$

Proof. The envelope in Equation (299) is $\mathcal{C}_{D,1}^v$ from Equation (31); the independent-mode term has zero capacity, so v is immaterial. By Lemma 2.8, its item measure is

$$a_*\delta_{\tau_D} + \nu.$$

The greedy realization first repeats the full maximum-density module and then processes ν in SPT order, proving the pointwise assertion. Since ν has mass $1 - a_*$ and is supported above τ_D , the common area formula Equation (33) expands to

$$\begin{aligned} \int_0^\infty (1 - \mathcal{C}_{\text{OT},D}(s)) ds &= \tau_D \left(\frac{a_*^2}{2} + a_*(1 - a_*) \right) + \frac{1}{2} \iint \min\{p, q\} d\nu(p) d\nu(q) \\ &= w_* \left(1 - \frac{a_*}{2} \right) + \frac{1}{2} \iint \min\{p, q\} d\nu(p) d\nu(q), \end{aligned}$$

because $w_* = a_*\tau_D$. This is Equation (298) and is unaffected by how a boundary atom is split. $\square$

6.2 The announced upper bound

Suppose the input is announced but its labeling is not. Choose a maximizing whole-occurrence prefix E , test labels in a uniformly random order, process an E -job immediately, and finish the residual jobs in SPT order. The exact identity Equation (21) and Equation (298) give

$$\mathbb{E} \text{cost}(\text{stat}, M) = n^2 \Phi_{\text{OT}}(D[M]) + \frac{1}{2} \left(|E| + \sum_i p_i \right). \quad (301)$$

In particular, if $M \subseteq [0, L]$, the remainder is at most $(1 + L)n/2$. The announced optimum is therefore at most the claimed fluid value plus $O_L(n)$.

6.3 No adaptive announced policy has a better leading cost

Assign the occurrences of M uniformly to the public labels and fix a deterministic nonanticipating policy. We may assume that it completes every job, since otherwise the lower bound is immediate. In the adaptive order in which it tests new labels, the revealed occurrences form a uniform random permutation $P_1, \dots, P_n$ of M . This is Lemma 2.11, with equal-valued occurrences kept distinct as temporary tokens. Because every first touch is a test, no suffix edit is needed and we may use the sharper all-prefix sampling bound directly.

Keep zero separate, partition $(0, L]$ into

$$K_n = \lfloor n^{1/5} \rfloor \quad (302)$$

cells, and round every positive value upward to its cell endpoint. Let D^+ be the resulting empirical law and $h_n = L/K_n$ the mesh. For each cell j , let $H_{k,j}$ count its occurrences among the first k tests and let d_j be its mass in D^+ . The uniform-prefix bound Equation (42) gives, except on an event of probability at most n^{-2} ,

$$|H_{k,j} - kd_j| \leq \Gamma_n \quad \text{for all } k, j, \quad \Gamma_n = O\left(\sqrt{n \ln(n+2)}\right). \quad (303)$$

Put $\gamma_n = \Gamma_n/n$. At an operation prefix, let $t = k/n$, let z be the normalized number of completed zero jobs, and let c_j be the completed positive mass from cell j . Equation (303)

gives the capacity hypotheses of Lemma 2.21 with $q = 1$ and $b = 0$. Rounding upward adds at most h_n to normalized processed work, so the work hypothesis holds with $\alpha = h_n$. Since $\mathcal{C}_{D^+,1}^v = \mathcal{C}_{\text{OT},D^+}$, the lemma gives, simultaneously at every prefix,

$$\frac{N_{\mathcal{A}}(nx)}{n} \leq \mathcal{C}_{\text{OT},D^+}(x + h_n) + (K_n + 1)\gamma_n. \quad (304)$$

The completion horizon is at most $H = L + 2$, and the immediate test-process policy gives $0 \leq \Phi_{\text{OT}}(D^+) \leq (L + 1)/2$. Apply Theorem 2.20 with $\varepsilon = 0$, $\alpha = h_n$, $\beta = (K_n + 1)\gamma_n$, and $\eta = n^{-2}$. Finally, zero-preserving transport Lemma 2.9 changes the fluid value by at most h_n . Because $(K_n + 1)\gamma_n = O(n^{-3/10}\sqrt{\ln(n+2)})$, every deterministic announced policy has uniform-placement expected cost at least

$$n^2\Phi_{\text{OT}}(D[M]) - O_L(n^{9/5}). \quad (305)$$

The finite-Yao conclusion built into Theorem 2.20 gives the same lower bound on one fixed oblivious placement for every randomized policy.

6.4 Removing the announcement

It remains to prove Equation (291). Use the same sampling, threshold, and fallback rules as $\mathcal{A}_n^*$ in Section 5.6. Its full parameter choice is

$$k = \lfloor n^{3/4} \rfloor, \quad d = \lfloor n^{1/4} \rfloor, \quad B_n = 32 + n^{1/20}, \quad \eta = B_n/d. \quad (306)$$

The policy privately permutes the labels, tests the first k , rounds positive values upward to the B_n -grid, and uses the threshold closure of a maximum-density sample prefix.

Fix L . For all sufficiently large n , $B_n > L$ and the grid has no overflow job. Moreover, the full sample prefix has density at least $1/(1+L+\eta) > 1/B_n$, so fallback never occurs. Let D^+ be the rounded population histogram, D^{rem} the rounded histogram of the remaining jobs, $\hat{D}$ the sample histogram, and $\Delta = \|D^+ - \hat{D}\|_1$. By Lemma 5.8 and deletion of the k sample occurrences,

$$\mathbb{E}\Delta \leq \sqrt{\frac{d+2}{k}}, \quad \|D^{\text{rem}} - D^+\|_1 \leq \frac{2k}{n-k}. \quad (307)$$

For a fixed grid prefix J , let $F_{\text{OT},D}(J)$ be the right-hand side of Equation (298) when J is used as the early set, whether or not it is optimal for D . If D, E are grid histograms supported on $[0, L + \eta]$, then

$$|F_{\text{OT},D}(J) - F_{\text{OT},E}(J)| \leq 3(L + 2)\|D - E\|_1. \quad (308)$$

Indeed, early mass changes by at most the ℓ_1 distance, early first moment by at most $(L + \eta)$ times that distance, and the residual pair kernel $\min\{p, q\}$ changes by at most twice this amount under the two product measures. The displayed loose constant covers the three terms.

By Lemma 6.3, a maximum-density prefix minimizes $F_{\text{OT},D}(J)$ over grid prefixes. Let $\hat{J}$ be the prefix learned from the sample. Rounding each processing time upward by at most η preserves zeros, so Lemma 2.9 gives

$$\Phi_{\text{OT}}(D^+) \leq \Phi_{\text{OT}}(D[M]) + \eta. \quad (309)$$

Conditional on the unordered sample, compare the actual sample-first schedule with the fully stationary policy for $\hat{J}$. The calculation in Section 5.6 bounds the extra cost by

$O_L(nk + k^2)$. The diagonal term in the exact stationary identity is $O_L(n)$, and using the rounded values can only increase the comparator's processing delays. Therefore

$$\mathbb{E}[\text{cost}(\mathcal{A}^*, M) \mid S] \leq n^2 F_{\text{OT}, D^{\text{rem}}}(\hat{J}) + O_L(nk + k^2 + n). \quad (310)$$

This is the conditional hypothesis of Lemma 2.22. Apply that lemma with metric $d(D, E) = \|D - E\|_1$, $L_0 = 3(L + 2)$, and Equation (307); then use Equation (309). Since

$$\sqrt{d/k} = O(n^{-1/4}), \quad \eta = O(n^{-1/5}), \quad k/n = O(n^{-1/4}),$$

we obtain

$$\mathbb{E} \text{cost}(\mathcal{A}^*, \sigma(M)) \leq n^2 \Phi_{\text{OT}}(D[M]) + O_L(n^{9/5}) \quad (311)$$

for every fixed labeling σ . Together with Equation (305), this proves Theorem 6.1.

Finally, the exact certificate Equation (274) says that $\Phi_{\text{OT}}(D)$ is at most $4/3$ times the leading clairvoyant SPT value for every D . Equality is attained by $D = \frac{1}{2}\delta_0 + \frac{1}{2}\delta_2$. Thus the sharp $4/3$ theorem is exactly the worst-input consequence of the stronger instance-optimal description.

7 Blind execution: instance-optimal four-block policies

We now turn to $\text{BE}(L)$. Running an untouched job blindly consumes its actual processing time and completes it; testing consumes one unit, reveals the value, and permits the processing operation to be delayed. The adversary fixes an arbitrary labeled vector in $[0, L]^n$ before the private seed is drawn. The goal of this section is to prove Theorem 7.1, the rigorous version of the informal blind-execution theorem Theorem 1.3.

Theorem 7.1 (Blind-execution instance optimality). *Fix $L < \infty$. There exist a randomized nonanticipating policy $\mathcal{A}^*$, which knows n, L but not the processing-time multiset, and a deterministic sequence $\varepsilon_L(n) \rightarrow 0$ with the following properties. For every multiset $M \subseteq [0, L]$ of size n and every fixed labeling σ ,*

$$\mathbb{E} \text{cost}(\mathcal{A}^*, \sigma(M)) \leq n^2 \Phi_{\text{BE}}(D[M]) + \varepsilon_L(n)n^2. \quad (312)$$

For every randomized policy $\mathcal{A}$, even an announced policy, some fixed oblivious labeling σ satisfies

$$\mathbb{E} \text{cost}(\mathcal{A}, \sigma(M)) \geq n^2 \Phi_{\text{BE}}(D[M]) - \varepsilon_L(n)n^2. \quad (313)$$

The benchmark Φ_{BE} is defined explicitly in Section 7.1; it is the minimum of a one-variable quadratic. One may take $\varepsilon_L(n) = O_L(n^{-1/6} \sqrt{\ln(n+2)})$.

The boundedness assumption in Theorem 7.1 cannot be removed. In fact, without it even the much weaker worst-case guarantee is impossible.

Theorem 7.2 (No finite ratio for unbounded blind execution). *There are no randomized policies $\mathcal{A}_n$, constant $R < \infty$, and sequence $\eta(n) \rightarrow 0$ such that every vector $p \in [0, \infty)^n$ satisfies*

$$\mathbb{E} \text{cost}(\mathcal{A}_n, p) \leq R \text{cost}(\text{OPT}, p) + \eta(n)n^2. \quad (314)$$

Proof. Suppose that such a guarantee exists, and run $\mathcal{A}_n$ on the all-zero instance. Its offline cost is zero, so if Z_n denotes the expected online cost, then $Z_n \leq \eta(n)n^2 = o(n^2)$. We may assume that the policy finishes every job, since otherwise its cost is infinite.

Fix its private seed and order the labels by their first touch on this zero instance. Every first touch completes its job: a blind zero takes no time, while a tested zero completes with its unit test. If the job at rank h is tested, that test is performed while $n - h + 1$ jobs are unfinished and hence contributes $n - h + 1$ to the objective. Consequently

$$Z_n \geq \mathbb{E} \sum_{h: \text{rank } h \text{ is tested}} (n - h + 1). \quad (315)$$

Now choose one label uniformly and change only its length from zero to $H = n^2$. Until this exceptional label is first touched, the policy sees exactly the zero-instance history. If its rank h is touched blindly, its nonpreemptive execution occupies the machine for H time units while $n - h + 1$ jobs are unfinished. Averaging over the exceptional label and the private seed therefore gives

$$\begin{aligned} \mathbb{E} \text{cost}(\mathcal{A}_n, p) &\geq \frac{H}{n} \mathbb{E} \sum_{h: \text{rank } h \text{ is blind}} (n - h + 1) \\ &\geq \frac{H}{n} \left(\frac{n(n+1)}{2} - Z_n \right) = H \left(\frac{n}{2} - o(n) \right). \end{aligned} \quad (316)$$

Some fixed choice of the exceptional label has at least this expected cost. For every such labeling, the clairvoyant schedule runs all zero jobs first and the long job last, so $\text{cost}(\text{OPT}, p) = H$. With $H = n^2$, Equation (316) contradicts Equation (314) for all sufficiently large n . $\square$

The proof first replaces a round of tests and its immediate short outcomes by one divisible completion module. For a fixed tested fraction q , a fractional-knapsack envelope then forces the four-block order from the introduction and gives its exact quadratic cost. To handle an arbitrary adaptive policy, we show that its first-touch order is a bijective relabeling of the occurrence permutation; predictable sampling controls all touch prefixes at once. A growing grid extends the bound to arbitrary bounded multisets, and a blind sample removes advance knowledge of the input.

7.1 The distributional benchmark

Let D be a probability distribution on $[0, L]$ and let $P \sim D$. Write

$$\mu = \mathbb{E}P. \quad (317)$$

Blind execution completes one unit of job mass using expected work μ . Apply Lemma 2.5 to D , choose its threshold selector g_* , and let ν be the resulting residual measure. The best possible test-module density and its reciprocal are

$$\kappa_D := \max_g \frac{a_D(g)}{w_D(g)} = \frac{1}{\tau_D}. \quad (318)$$

The selector processes all positive values below τ_D immediately after their tests, rejects all values above it, and may split a boundary atom. With that boundary convention, put

$$a = a_D(g_*), \quad \ell = \mathbb{E}[Pg_*(P)\mathbf{1}_{P>0}].$$

The identities in Lemma 2.5 give

$$a\tau_D - \ell = 1, \quad \tau_D = \frac{1 + \ell}{a}. \quad (319)$$

Thus a full unit of tests together with all selected low outcomes is a divisible module of completion mass a , work $1 + \ell$, and work per completion τ_D .

If $\tau_D \geq \mu$, blind work has at least the completion density of any test module, and pure blind execution is optimal at the fluid scale. Suppose now that $\tau_D < \mu$. After testing a fraction q , the canonical schedule is

$$\begin{aligned}
\underbrace{\text{tests with immediate } p < \tau_D}_{\text{discovery}} &\longrightarrow \underbrace{\text{known } \tau_D < p < \mu \text{ in SPT}}_{\text{medium}} \\
&\longrightarrow \underbrace{\text{untouched jobs blind}}_{\text{blind block}} \longrightarrow \underbrace{\text{known } p > \mu \text{ in SPT}}_{\text{high tail}}.
\end{aligned} \tag{320}$$

For the explicit cost, first assume no atom lies at τ_D or μ , and put

$$\begin{aligned}
\mathcal{L} &= \{p < \tau_D\}, & \mathcal{M} &= \{\tau_D < p < \mu\}, & \mathcal{H} &= \{p > \mu\}, \\
a &= \mathbb{P}(P \in \mathcal{L}), & \ell &= \mathbb{E}[P \mathbf{1}_{\mathcal{L}}], & \ell_M &= \mathbb{E}[P \mathbf{1}_{\mathcal{M}}], \\
d &= \mathbb{P}(P \in \mathcal{H}), & K_M &= \mathbb{E}[\min(P, P') \mathbf{1}_{P, P' \in \mathcal{M}}], & K_H &= \mathbb{E}[\min(P, P') \mathbf{1}_{P, P' \in \mathcal{H}}],
\end{aligned}$$

where P, P' are independent with law D . Put $\bar{q} = 1 - q$. Remaining-mass accounting gives

$$\begin{aligned}
F_{\text{BE}, D}(q) &= (1 + \ell) \left(q - \frac{aq^2}{2} \right) + q\ell_M(\bar{q} + qd) + \frac{q^2 K_M}{2} \\
&\quad + \mu \left(\bar{q}qd + \frac{\bar{q}^2}{2} \right) + \frac{q^2 K_H}{2}.
\end{aligned} \tag{321}$$

Atoms at τ_D are split between the low module and residual stock; atoms at μ may be placed on either side of the blind block. The same formula then holds for the resulting virtual classes.

Each term in Equation (321) has a direct area interpretation. During the test module, work $(1 + \ell)dq$ reduces unfinished mass from 1 to $1 - aq$. The medium work $q\ell_M$ delays the untouched mass $\bar{q}$ and high mass qd , while K_M is the internal SPT pair term. The blind block delays all high jobs and itself loses mass linearly. Finally K_H is the high-tail pair term. Expansion gives the quadratic

$$F_{\text{BE}, D}(q) = \frac{\mu}{2} + Aq + Bq^2, \tag{322}$$

where

$$A = 1 + \ell + \ell_M + \mu(d - 1) = 1 - \mathbb{E}(\mu - P)^+, \tag{323}$$

$$B = -\frac{a(1 + \ell)}{2} + \ell_M(d - 1) + \frac{K_M}{2} + \mu \left(\frac{1}{2} - d \right) + \frac{K_H}{2}. \tag{324}$$

The envelope proof below implies $B \geq 0$. In the short-test regime $\tau_D < \mu$, monotonicity in Equation (25) gives $A < 0$, and the optimal tested fraction is therefore

$$q^* = \begin{cases} 1, & B = 0, \\ \min\{1, -A/(2B)\}, & B > 0. \end{cases} \tag{325}$$

We define

$$\Phi_{\text{BE}}(D) = \min_{0 \leq q \leq 1} F_{\text{BE}, D}(q), \tag{326}$$

with $\Phi_{\text{BE}}(D) = \mu/2$ in the regime $\tau_D \geq \mu$.

7.2 The fixed-fraction completion envelope

We first prove that Equation (320) is globally optimal in a fluid relaxation. Let D have finite support $0 = p_0 < p_1 < \dots < p_s \leq L$, with masses d_i . Fix a final tested fraction q . At an arbitrary schedule prefix, let

$$\begin{aligned} t &= \text{tested mass,} & b &= \text{blindly completed mass,} \\ c_i &= \text{processed tested mass of class } i > 0. \end{aligned}$$

The completed mass and idealized work are

$$y = d_0 t + b + \sum_{i>0} c_i, \quad x = t + \mu b + \sum_{i>0} p_i c_i, \quad (327)$$

subject to

$$0 \leq t \leq q, \quad 0 \leq b \leq 1 - q, \quad 0 \leq c_i \leq d_i t. \quad (328)$$

Define the model envelope

$$\mathcal{C}_{\text{BE},D,q} := \mathcal{C}_{D,q}^\mu, \quad (329)$$

using the common envelope Equation (31). Every fluid schedule that ultimately tests mass q obeys this envelope, hence

$$\text{cost} \geq \int_0^\infty (1 - \mathcal{C}_{\text{BE},D,q}(x)) dx. \quad (330)$$

Lemma 7.3 (Fixed- q envelope). *If $\tau_D < \mu$, then for every $q \in [0, 1]$ the right side of Equation (330) equals $F_{\text{BE},D}(q)$ and is attained by the four-block fluid policy. If $\tau_D \geq \mu$, that right side is at least $\mu/2$ for every q , with equality at $q = 0$.*

Proof. Apply Lemma 2.8 with independent-mode work $v = \mu$. The resulting item measure is

$$qa\delta_{\tau_D} + q\nu + (1 - q)\delta_\mu.$$

When $\tau_D < \mu$, sorting these items by work per completion gives exactly the low test module, residual classes below μ , the blind module, and residual classes above μ , which is the four-block order Equation (320). Expanding the common area formula Equation (33) across these four blocks gives Equation (321).

When $\tau_D \geq \mu$, every test and residual item costs at least μ per completion. The $q = 0$ envelope, one blind item of capacity one and cost μ , therefore pointwise dominates every $q > 0$ envelope and has area $\mu/2$. Finally, feasibility in Equations (327) and (328) is affine in q , so $\mathcal{C}_{\text{BE},D,q}(x)$ is concave in q for each x . In the short-test regime its area complement $F_{\text{BE},D}(q)$ is convex, proving $B \geq 0$. $\square$

7.3 Adaptive first touches are a uniform urn

The fluid proof fixed q , but an announced finite policy may choose whom to touch next and when to stop testing as arbitrary functions of its history. The next finite lemma is the deferred-decisions statement needed to control such a policy.

For a size- n multiset whose values lie among $0 = p_0 < p_1 < \dots < p_s \leq L$, write D_n for its empirical distribution, $D_{n,i} = D_n(\{p_i\})$, and $\mu_n = \int p dD_n(p)$.

Distinguish occurrences even when their processing times coincide and fix a deterministic proper policy, equivalently the private seed of a randomized policy. Apply Lemma 2.18 to the $s + 1$ classes with failure probability n^{-2} , and put $\Gamma_n = \max\{\Gamma_{\text{cell}}, \Gamma_{\text{work}}\}$. On the

resulting event, for every touch prefix leaving at least δn labels untouched and every class i ,

$$|\# \text{tested class } i - D_{n,i} \# \text{tests}| \leq \Gamma_n, \quad (331)$$

$$|\text{blind work} - \mu_n \# \text{blind jobs}| \leq \Gamma_n. \quad (332)$$

Importantly, the common event does not condition on the eventual tested fraction.

7.4 The announced lower bound

Fix a policy seed. Stop after $\lfloor (1 - \delta)n \rfloor$ first touches and let $Q_{n,\delta}$ be the number of tests completed by then, divided by n . Edit the suffix: every still-untouched job that would later be tested is instead executed blindly at its first original operation, and its later processing record is deleted. At most $\delta n + O(1)$ jobs and three operations per affected job change. The bounded suffix-edit lemma Lemma 2.19, with $W = L + 1$, gives

$$O((L + 1)\delta n^2). \quad (333)$$

The edited schedule performs exactly $Q_{n,\delta}n$ tests.

Work on the single event Equations (331) and (332) and only now set $q = Q_{n,\delta}$. Normalize mass and work by n , and let $\gamma_n = \Gamma_n/n$. At any operation boundary, let t, b, z, c_i denote tested mass, blindly completed mass, zero mass completed by tests, and processed known positive mass of class i . The event gives

$$z \leq D_{n,0}t + \gamma_n, \quad c_i \leq D_{n,i}t + \gamma_n.$$

If x is the actual normalized prefix work, the blind-work discrepancy and the suffix edit give

$$t + \mu_n b + \sum_i p_i (c_i - \gamma_n)^+ \leq x + (L + 1)(\gamma_n + \delta). \quad (334)$$

Thus all hypotheses of Lemma 2.21 hold with $\alpha = (L + 1)(\gamma_n + \delta)$, and it gives

$$\text{completed fraction by } x \leq \mathcal{C}_{\text{BE}, D_n, q}(x + (L + 1)(\gamma_n + \delta)) + (s + 1)\gamma_n. \quad (335)$$

The envelope horizon is at most $H = L + 1$, and pure blind execution gives $0 \leq \Phi_{\text{BE}}(D_n) \leq L/2$. Apply Theorem 2.20 with the suffix-edit error from Equation (333), $\alpha = (L + 1)(\gamma_n + \delta)$, and $\beta = (s + 1)\gamma_n$. It gives

$$\frac{\text{cost}(\mathcal{A}, M)}{n^2} \geq F_{\text{BE}, D_n}(Q_{n,\delta}) - O_{L,s}(\gamma_n + \delta + 1/n) \geq \Phi_{\text{BE}}(D_n) - O_{L,s}(\gamma_n + \delta + 1/n). \quad (336)$$

The long-test regime uses the $\mu_n/2$ dominance part of Lemma 7.3; the same transfer applies. Its finite-Yao conclusion also proves the lower bound for randomized announced policies.

The random-labeling and announced minimax values coincide. The transfer lemma gives one inequality. For the other, privately relabel the physical jobs before running an optimal deterministic urn policy; against every fixed assignment, the policy sees a uniform assignment. Thus the lower bound above applies to the announced oblivious minimax problem, not merely to one randomized input model. Write $\text{VAL}_{\text{ann}}(M)$ for this common minimax value.

7.5 A growing grid and arbitrary bounded multisets

The number of distinct values may grow with n . Keep zero as a separate cell, partition $(0, L]$ into

$$K_n = \lfloor n^{1/6} \rfloor \quad (337)$$

bins, and round every positive value upward; let $h_n = L/K_n$ be the mesh. Zero preservation matters because a tested zero completes at its test. Write $D[M]^{\text{grid}}$ for the rounded empirical distribution and $\mu_M^{\text{grid}} = \int p dD[M]^{\text{grid}}(p)$ for its mean.

Coupling each value with its rounded value and applying Lemma 2.9 gives

$$|\Phi_{\text{BE}}(D) - \Phi_{\text{BE}}(D^{\text{grid}})| \leq h_n. \quad (338)$$

This avoids any assumption that the numerical optimizer or threshold is stable under rounding.

Apply Lemma 2.18 to all $K_n + 1$ cells and take $\delta_n = n^{-1/6}$. The model-specific blind-work discrepancy supplies the horizontal error; Lemma 2.21 supplies the vertical one. After rounding, both are bounded by

$$\zeta_n = O_L\left(\frac{(K_n + 1)\Gamma_n}{n} + h_n + \delta_n\right) = O_L(n^{-1/6}\sqrt{\ln(n+2)}). \quad (339)$$

Consequently the rounded completion curve is bounded by $\mathcal{C}_{\text{BE}, D[M]^{\text{grid}}, q}(x + \zeta_n) + \zeta_n$. Apply Theorem 2.20 with $H = L + 1$, $B = L/2$, the suffix-edit error, and the failure probability from the urn lemma. Together with Equation (338), this yields the announced lower bound

$$\frac{\text{VAL}_{\text{ann}}(M)}{n^2} \geq \Phi_{\text{BE}}(D[M]) - O_L(n^{-1/6}\sqrt{\ln(n+2)}). \quad (340)$$

For the matching upper bound, compute an optimal canonical template for the rounded empirical multiset, choose $\lfloor q^*n \rfloor$ uniformly random labels to test, and execute the four blocks. Pairwise delay accounting expresses its expectation as bounded one-occurrence and two-distinct-occurrence kernels. The without-replacement pair coefficients differ from their product-law values by $O(1/n)$ uniformly in the tested count. Hence if π records the virtual classes and the tested fraction,

$$\mathbb{E}C_n(\pi) = n^2 F_{\text{BE}, D[M]^{\text{grid}}}(\pi) + O_L(n) \quad (341)$$

uniformly over canonical templates π . Together with Equation (338), this proves

$$\left| \frac{\text{VAL}_{\text{ann}}(M)}{n^2} - \Phi_{\text{BE}}(D[M]) \right| \leq O_L(n^{-1/6}\sqrt{\ln(n+2)}). \quad (342)$$

7.6 Removing the announcement

The universal policy first draws a uniform random permutation and blindly executes

$$k_n = \lceil n^{1/2} \rceil \quad (343)$$

sample jobs. Their observed durations give a grid histogram $\hat{D}_n$. The policy minimizes the canonical quadratic over every grid threshold, blind-block insertion position, and $q \in [0, 1]$, then runs the selected four-block template on the remaining jobs in the unused random order.

Let D_n and D_n^{rem} be the full and remaining grid histograms. Applying Equation (41) to the grid classes gives

$$\begin{aligned}\mathbb{E}\|\hat{D}_n - D_n\|_1 &\leq \sqrt{\frac{K_n + 1}{k_n}}, & \mathbb{E}\|\hat{D}_n - D_n^{\text{rem}}\|_1 &\leq \frac{n}{n - k_n} \sqrt{\frac{K_n + 1}{k_n}}, \\ \|D_n^{\text{rem}} - D_n\|_1 &\leq \frac{2k_n}{n - k_n}.\end{aligned}\tag{344}$$

For a fixed canonical template, formula Equation (321) consists only of bounded one- and two-draw expectations. Therefore, uniformly in the grid and template,

$$|F_{\text{BE},D}(\pi) - F_{\text{BE},E}(\pi)| \leq C_L \|D - E\|_1,\tag{345}$$

for example with $C_L = 20(L+1)$. Let $\hat{\pi}$ minimize the sampled objective. Conditional on the sample, the remaining order is uniform and the learned template is fixed, so Equation (341) applies. The sample itself has work at most Lk_n and delays at most n jobs, adding at most $Lnk_n = O_L(n^{3/2})$. Hence

$$\mathbb{E}[\text{cost}(\mathcal{A}^*, p) \mid \hat{D}_n, D_n^{\text{rem}}] \leq n^2 F_{\text{BE},D_n^{\text{rem}}}(\hat{\pi}) + O_L(nk_n + n).$$

Apply Lemma 2.22 with $d(D, E) = \|D - E\|_1$, $L_0 = C_L$, and Equation (344); then use Equation (338). This gives

$$\frac{\mathbb{E} \text{cost}(\mathcal{A}^*, p)}{n^2} \leq \Phi_{\text{BE}}(D[M]) + O_L\left(n^{-1/6} \sqrt{\ln(n+2)}\right)\tag{346}$$

uniformly over every fixed labeled vector $p \in [0, L]^n$.

For the converse, reveal M to an arbitrary policy for free and draw a uniform labeling. The announced lower bound Equation (340) applies seedwise. Finite Yao selection Lemma 2.17 gives a fixed labeling with cost at least the average, which proves Equation (313). This completes the proof of Theorem 7.1.

7.7 Why the medium block is necessary

Let

$$\mathbb{P}(P = 0) = \frac{1}{5}, \quad \mathbb{P}(P = 9) = \frac{1}{5}, \quad \mathbb{P}(P = 16) = \frac{3}{5}.$$

Then $\mu = 57/5$ and $\tau_D = 5$. Thus tested zeros are immediate, known nine-jobs precede the blind block, and known sixteen-jobs follow it. Formula Equation (321) becomes

$$F_{\text{BE},D}(q) = \frac{57}{10} - \frac{44}{25}q + \frac{11}{10}q^2.$$

Its minimizer is $q^* = 4/5$ and its value is 4.996. If all tested nonzero jobs are instead deferred until after the blind block, the best value is 5.04. The gap is a positive constant times n^2 : the medium SPT block is not cosmetic, and a policy of the simpler form “threshold, then blind, then all tested residuals” is not instance optimal.

8 Blind optimization: speedup without revelation

We finally separate the two benefits that were bundled together in the revealing-optimization model. Raw execution still takes the known time u , and one unit of optimization still reduces the processing time to $p_i \in [0, u]$. The difference is that optimization does not

reveal p_i . An optimized job must be started blindly and its duration becomes known only when it finishes. We call this the *blind-optimization* model $\text{BO}(u)$.

We study deterministic worst-case policies, and randomized policies against an oblivious adversary. The deterministic optimum has a particularly simple form: run everything raw up to $u = 2$, and optimize everything after that point. Proving that no adaptive mixture improves these two policies uses the same raw-safe hidden-stopping idea as the revealing-optimization lower bound. Randomization is still essential for learning the input: a private random sample is representative under every fixed labeling, whereas an adversary can make a deterministic exploration history atypical.

This small change removes the threshold decision entirely. It also shows that the bounded randomized ratio in the revealing model is genuinely an information effect.

The section first proves that deterministic instance optimality is impossible and determines the exact deterministic worst-case curve. It then proves two closely related randomized statements. The first, Theorem 8.5, is the rigorous version of the informal instance-optimality theorem Theorem 1.5. The second, Theorem 8.6, is the rigorous version of Theorem 1.6 and maximizes that instance-specific value relative to the clairvoyant optimum. Keeping the results together avoids repeating the same blind-block reduction.

For a multiset M , use $D[M]$ from Equation (7) and put

$$\mu_M = \int p dD[M](p) = \frac{1}{n} \sum_{i=1}^n p_i$$

and define

$$\Phi_{\text{BO},u}(D[M]) = \frac{1}{2} \min\{u, 1 + \mu_M\}. \quad (347)$$

The two terms have an immediate meaning. Running every job raw uses u units per completion. Optimizing and then blindly processing every job uses $1 + \mu_M$ units per completion on average.

Randomization is necessary for attaining this benchmark on every input.

Theorem 8.1 (No deterministic instance-optimal policy). *For every unannounced deterministic nonanticipating policy $\mathcal{A}$ and every even n , there are a fixed labeled input $I \in \{0, 2\}^n$ and an announced randomized policy $\mathcal{A}'$ such that*

$$\text{cost}(\mathcal{A}, I) \geq \mathbb{E} \text{cost}(\mathcal{A}', I) + \frac{1}{8} n^2. \quad (348)$$

Consequently no deterministic policy is asymptotically instance-optimal in $\text{BO}(2)$.

Proof. Run $\mathcal{A}$ against the following deterministic revelation rule. Give a job value zero if $\mathcal{A}$ runs it raw, and value two if $\mathcal{A}$ optimizes it. Optimization reveals nothing, and the value two is seen only when the optimized job is subsequently processed. Since $\mathcal{A}$ is deterministic, recording the resulting value of every label produces a fixed labeled input with the same transcript.

Let m be the number of optimized jobs on this input. Every raw job consumes two units before completion, whereas every optimized job consumes three. Irrespective of interleaving, the completion times are therefore bounded below by the shortest-first schedule of $n - m$ blocks of length two and m blocks of length three. Hence

$$\text{cost}(\mathcal{A}, I) \geq n^2 + n + \frac{m^2 + m}{2}. \quad (349)$$

The multiset has $n - m$ zeros and m twos. If $m \leq n/2$, let $\mathcal{A}'$ privately permute the jobs and optimize and process them one at a time in that order. Every position sees the average block length $(n + 2m)/n$, so

$$\mathbb{E} \text{cost}(\mathcal{A}', I) = \frac{(n + 1)(n + 2m)}{2}. \quad (350)$$

Subtracting this from Equation (349) gives $(n - m)(n - m + 1)/2 \geq n^2/8$.

If $m > n/2$, let $\mathcal{A}'$ run every job raw. Its cost is $n(n + 1)$, and the difference in Equation (349) is $(m^2 + m)/2 > n^2/8$. This proves Equation (348). $\square$

An optimization operation reveals no information. We will repeatedly use the following normal form.

Lemma 8.2 (Blind-block normal form). *For every nonanticipating policy $\mathcal{A}$ in $\text{BO}(u)$ there is a nonanticipating policy $\mathcal{A}^{\text{nf}}$ which completes every optimized job as one contiguous block of length $1 + p_i$ and satisfies*

$$\text{cost}(\mathcal{A}^{\text{nf}}, I) \leq \text{cost}(\mathcal{A}, I) \quad (351)$$

on every input I . If $\mathcal{A}$ is deterministic, then so is $\mathcal{A}^{\text{nf}}$.

Proof. Maintain a virtual execution of $\mathcal{A}$. When the virtual policy optimizes a job, record that fact and advance the virtual clock by one unit, but perform no physical work. Immediately before the virtual policy later processes that job, perform its postponed optimization physically and then process it. Observations that occur earlier in physical time are buffered until the corresponding virtual completion. Thus the virtual policy sees exactly its original history and makes the same decisions. Moving a unit of optimization to the right leaves the optimized job and every later job no later than before, while jobs completed in between move one unit earlier. Applying this to every separated optimization proves (351). $\square$

8.1 The deterministic worst-case curve

For $u > 1$ define

$$G_{\text{BO}}(u) := \frac{\sqrt{4u^3 - 4u + 1} - 1}{2(u - 1)}. \quad (352)$$

The two deterministic policies needed below are RAW, which runs every job for time u , and OPTIMIZEALL, which fixes an arbitrary label order and completes every job as a blind block of length $1 + p_i$.

Theorem 8.3 (Exact deterministic blind-optimization curve). *The optimal deterministic size-asymptotic competitive ratio in $\text{BO}(u)$ is*

$$R_{\text{BO}}^{\text{det}}(u) = \begin{cases} 1, & 0 < u \leq 1, \\ u, & 1 \leq u \leq 2, \\ G_{\text{BO}}(u), & u \geq 2. \end{cases} \quad (353)$$

The policy RAW is optimal for $u \leq 2$, and OPTIMIZEALL is optimal for $u \geq 2$; both are optimal at $u = 2$. In particular,

$$R_{\text{BO}}^{\text{det}}(u) = \sqrt{u} + o(1) \quad (u \rightarrow \infty). \quad (354)$$

We first prove the adaptive lower bound. Its stopping rule is the direct blind-block analogue of Lemma 3.5.

Lemma 8.4 (Blind raw-safe hidden stopping). *Fix $u > 1$, a target $R \leq u$, and $\alpha \in (0, 1)$. Against every deterministic policy there is a fixed binary input with $p_j \in \{0, u\}$ on which*

$$\text{ALG} - R \text{OPT} \geq \frac{n^2}{2} \left((u - R)(1 - \sigma^2) + \sigma^2 f_{u,R}^{\text{BO}}(y) \right) - O_u(n), \quad (355)$$

where $\sigma \in [0, 1]$, $0 \leq y - \alpha < 1/(\sigma n)$ when $\sigma > 0$, and

$$f_{u,R}^{\text{BO}}(y) = 1 - R + 2uy - (u + R(u - 1))y^2. \quad (356)$$

Consequently, if $f_{u,R}^{\text{BO}}(\alpha) \geq 0$, then every deterministic policy has size-asymptotic ratio at least R .

Proof. By Lemma 8.2, it is enough to lower-bound a policy whose optimized jobs are contiguous blind blocks. Start in a long phase. Give value zero to every job completed raw and value u to every optimized job. If v raw jobs and L optimized-long jobs have been completed, stop at the first completion for which $L > 0$ and

$$L \geq \alpha(n - v). \quad (357)$$

Give value zero to all remaining jobs. The answers are deterministic functions of the transcript, so the replay induction of Lemma 2.10 freezes the interaction into one fixed labeled binary input.

If the policy fails to complete all jobs, its ratio is infinite. Otherwise, if the line is never crossed, then $L = 0$ after all jobs are completed: otherwise $n - v = L$ at the end and $L \geq \alpha L$. Thus the algorithm ran every job raw on the all-zero input and has ratio $u \geq R$.

Suppose the line is crossed. Put $x = n - v - L$. For a lower bound, discard idle time, put the already completed raw blocks before the already completed optimized-long blocks, reveal the future, and let all x remaining zero jobs be optimized. These exchanges only favor the algorithm: among the past blocks, shortest first is optimal, while no untouched future job could have completed before the crossing. The resulting exact cost and the clairvoyant optimum are

$$A_{\text{ex}} = u \left(vn - \frac{v(v-1)}{2} \right) + (u+1) \left(L(n-v) - \frac{L(L-1)}{2} \right) + \frac{x(x+1)}{2}, \quad (358)$$

$$O_{\text{ex}} = \frac{n(n+1)}{2} + (u-1) \frac{L(L+1)}{2}. \quad (359)$$

Here the offline optimum first completes the $n - L$ zero jobs as unit optimized blocks and then runs the L long jobs raw.

Set

$$\sigma = \frac{n-v}{n}, \quad y = \frac{L}{n-v}. \quad (360)$$

Twice the quadratic coefficients of Equations (358) and (359) are

$$\begin{aligned} \mathcal{A} &= u(1 - \sigma^2) + \sigma^2(1 + 2uy - uy^2), \\ \mathcal{O} &= (1 - \sigma^2) + \sigma^2(1 + (u-1)y^2). \end{aligned}$$

Subtracting $R\mathcal{O}$ from $\mathcal{A}$ gives exactly the bracket in (355). Immediately before the crossing, $L - \alpha(n - v) < 0$. An optimized completion increases this expression by one and a raw completion by α , so

$$0 \leq y - \alpha < \frac{1}{n-v} = \frac{1}{\sigma n}. \quad (361)$$

The quadratic $f_{u,R}^{\text{BO}}$ is uniformly Lipschitz on $[0, 1]$ for fixed u and $R \leq u$. Hence $\sigma^2 f_{u,R}^{\text{BO}}(y) \geq \sigma^2 f_{u,R}^{\text{BO}}(\alpha) - O_u(1/n)$. The exact diagonal terms contribute another $O_u(n)$ before normalization. If $u - R$ and $f_{u,R}^{\text{BO}}(\alpha)$ are nonnegative, this proves the claimed lower bound. $\square$

Proof of Theorem 8.3. If $u \leq 1$, raw execution is an offline-optimal block for every job, so the ratio is one. If $1 < u \leq 2$, use

$$R = u, \quad \alpha = \frac{1}{u}. \quad (362)$$

Then $f_{u,u}^{\text{BO}}(1/u) = 2 - u \geq 0$, and Lemma 8.4 gives the lower bound u . Conversely, every offline effective length is at least one, whereas RAW has cost $un(n+1)/2$, so its ratio is at most u .

Now let $u \geq 2$ and put $R = G_{\text{BO}}(u)$. The definition (352) is equivalently the positive solution of

$$(R - 1)(u + R(u - 1)) = u^2. \quad (363)$$

Moreover $R \leq u$, with equality only at $u = 2$. Choose

$$\alpha = \frac{u}{u + R(u - 1)}. \quad (364)$$

This is the maximizer of Equation (356), and (363) gives

$$f_{u,R}^{\text{BO}}(\alpha) = 1 - R + \frac{u^2}{u + R(u - 1)} = 0. \quad (365)$$

Thus Lemma 8.4 proves the lower bound R .

It remains to analyze OPTIMIZEALL. Put $a(p) = 1 + p$. For a fixed multiset, its worst label order has the $a(p_i)$ in nonincreasing order. If P, P' are independent draws from the empirical distribution, online pair accounting and the empirical offline identity Corollary 2.3 give

$$\text{cost}(\text{OPTIMIZEALL}) \leq \frac{n^2}{2} \mathbb{E} \max\{a(P), a(P')\} + \frac{n}{2} \mathbb{E} a(P), \quad (366)$$

$$\text{OPT} = \frac{n^2}{2} \mathbb{E} \min\{\lambda_u(P), \lambda_u(P')\} + \frac{n}{2} \mathbb{E} \lambda_u(P). \quad (367)$$

Write $S_D(t) = \mathbb{P}(P > t)$ for $0 \leq t \leq u$. Layer cake yields

$$\mathbb{E} \max\{a(P), a(P')\} = 1 + \int_0^u (2S_D(t) - S_D(t)^2) dt, \quad (368)$$

$$\mathbb{E} \min\{\lambda_u(P), \lambda_u(P')\} = 1 + \int_0^{u-1} S_D(t)^2 dt. \quad (369)$$

Set

$$b = \frac{1}{u-1} \int_0^{u-1} S_D(t) dt. \quad (370)$$

Cauchy–Schwarz bounds the square integral on $[0, u-1]$ from below by $(u-1)b^2$. Since S_D is nonincreasing,

$$T := \int_{u-1}^u S_D(t) dt \leq S_D(u-1) \leq b. \quad (371)$$

The function $h \mapsto 2h - h^2$ is increasing and concave on $[0, 1]$. Cauchy–Schwarz on the prefix and Jensen’s inequality on the unit tail therefore give

$$\mathbb{E} \max\{a(P), a(P')\} \leq 1 + 2ub - ub^2, \quad (372)$$

$$\mathbb{E} \min\{\lambda_u(P), \lambda_u(P')\} \geq 1 + (u - 1)b^2. \quad (373)$$

Consequently their ratio is at most

$$g_u(b) = \frac{1 + 2ub - ub^2}{1 + (u - 1)b^2}. \quad (374)$$

The derivative of g_u changes sign exactly once on $[0, 1]$. At its maximum, if $R = g_u(b)$, the stationary equation $g'_u(b) = 0$ says

$$u(1 - b) = R(u - 1)b, \quad b = \frac{u}{u + R(u - 1)}. \quad (375)$$

Substitution into $g_u(b) = R$ gives (363). That equation has the unique positive solution $G_{\text{BO}}(u)$; hence

$$\max_{0 \leq b \leq 1} g_u(b) = G_{\text{BO}}(u). \quad (376)$$

All survival-function inequalities above are tight for a binary multiset with a b -fraction of values u and the remaining values zero, labeled so that the long optimized blocks occur first. Equations (366)–(374) prove $\text{cost}(\text{OPTIMIZEALL}) \leq R \text{OPT} + O_u(n)$. This matches the lower bound and proves (353). The asymptotic formula (354) follows directly from (352). $\square$

Theorem 8.5 (Blind-optimization instance optimality). *For every fixed $u < \infty$ there exist a randomized nonanticipating policy $\mathcal{A}_{\text{BO}}$, depending only on n, u , and a sequence $\varepsilon_u(n) \rightarrow 0$ such that every size- n multiset $M \subseteq [0, u]$ and every fixed labeling σ satisfy*

$$\mathbb{E} \text{cost}(\mathcal{A}_{\text{BO}}, \sigma(M)) \leq n^2 \Phi_{\text{BO}, u}(D[M]) + \varepsilon_u(n)n^2. \quad (377)$$

Conversely, for every randomized nonanticipating policy $\mathcal{A}$, even an announced one, some fixed oblivious labeling σ satisfies

$$\mathbb{E} \text{cost}(\mathcal{A}, \sigma(M)) \geq n^2 \Phi_{\text{BO}, u}(D[M]) - \varepsilon_u(n)n^2. \quad (378)$$

Thus one policy is asymptotically optimal, at the n^2 scale, for every multiset simultaneously. Unlike the benchmarks in the preceding sections, $\Phi_{\text{BO}, u}$ depends only on the empirical mean.

The proof is simpler than in the revealing models. Because optimization reveals nothing, it can be moved next to the processing of its job. Every job is then either a raw block of known length u or an optimized blind block with average length $1 + \mu_M$. A predictable-sampling bound rules out an adaptive advantage, while a sample estimates the one number μ_M . The first subsection makes this argument formal; the second performs the outer worst-case optimization.

8.2 Proof of randomized instance optimality

In the equivalent random-labeling formulation, the labels are shuffled uniformly before the policy starts. Then the lower half simply says that no announced policy beats $n^2 \Phi_{\text{BO}, u}(D[M]) - o_u(n^2)$ in expectation.

By Lemma 8.2, we may regard every optimized job as one indivisible block of hidden length $1 + p_i$. A raw block has the known length u . Both blocks complete exactly one job.

The announced lower bound. Fix the private seed of a policy and place the announced occurrences uniformly behind the labels. We may assume that the policy completes every job. Give equal-valued occurrences distinct hidden tokens. The common first-touch lemma Lemma 2.11 makes their analytically exposed order a uniform permutation and the selector of optimized blocks predictable. A raw block does not reveal its token's value to the policy; exposing it only in the analysis enriches the filtration and is already allowed by the lemma.

Fix $\delta > 0$ and ignore the last δn first touches for the moment. The predictable-permutation estimate Lemma 2.15, applied to the processing times and the optimized-block selector, gives one event of probability $1 - o(1)$ on which every earlier prefix containing tn completed optimized blocks has total hidden processing work

$$\sum_{i \text{ optimized in the prefix}} p_i = \mu_M tn + o_\delta(n). \quad (379)$$

The event is simultaneous over all prefixes, so a stopping fraction chosen adaptively by the policy may be substituted only after the event has been fixed. Editing the last δn jobs to either mode changes total completion cost by $O_u(\delta n^2)$.

This can be made uniform rather than diagonalized. Take $\delta_n = n^{-1/4}$. The bounded selected-work estimate used in Section 7.3 has deviation

$$O_u\left((1 + \delta_n^{-1})\sqrt{n \ln(n+2)}\right) = O_u\left(n^{3/4}\sqrt{\ln(n+2)}\right)$$

and failure probability $o(1)$, simultaneously over all relevant prefixes. After division by n , both this deviation and the edited suffix contribute $O_u(n^{-1/4}\sqrt{\ln(n+2)})$ to the normalized completion-area bound.

Put $a = 1 + \mu_M$. If the final optimized fraction is q , the best possible fluid ordering puts the cheaper of the two block types first. When $a \leq u$ its area is

$$F_{a,u}(q) = a\left(q - \frac{q^2}{2}\right) + \frac{u(1-q)^2}{2}, \quad F'_{a,u}(q) = (a-u)(1-q). \quad (380)$$

It is minimized at $q = 1$, with value $a/2$. When $a \geq u$, raw blocks go first and the area is

$$F_{a,u}(q) = \frac{u(1-q^2)}{2} + \frac{aq^2}{2} = \frac{u}{2} + \frac{a-u}{2}q^2, \quad (381)$$

which is minimized at $q = 0$, with value $u/2$. An adjacent exchange proves that no interleaving has smaller area: swapping two consecutive blocks puts the shorter cost per completion first. Equations (379)–(381), with the uniform choice $\delta = \delta_n$, give

$$\frac{1}{n^2} \mathbb{E} \text{cost}(\mathcal{A}, M) \geq \frac{1}{2} \min\{u, 1 + \mu_M\} - o_u(1).$$

The estimate is uniform in the deterministic policy. Applying Lemma 2.17 and the same relabeling argument as in Section 7.4 proves (378). Together with the upper bound below, one may take $\varepsilon_u(n) = O_u(n^{-1/4}\sqrt{\ln(n+2)})$ after enlarging its constant.

One policy for all multisets. Take $k = \lceil n^{2/3} \rceil$ uniformly random sample jobs. Optimize each sample job and process it immediately, thereby observing its duration, and let $\hat{\mu}$ be their sample mean. If $1 + \hat{\mu} < u$, optimize and immediately process every remaining job in fresh random order. Otherwise run every remaining job raw.

The sample contributes at most $O_u(nk) = o_u(n^2)$ beyond the main schedule. The variance estimate Equation (40), after scaling from $[0, 1]$ to $[0, u]$, gives

$$\mathbb{E}|\hat{\mu} - \mu_M| = O_u(k^{-1/2} + k/n) = o_u(1).$$

Choosing the wrong mode can cost at most $O(n^2)|\hat{\mu} - \mu_M|$, while choosing the correct mode has leading cost $n^2\Phi_{\text{BO},u}(D[M])$. This proves (377). In particular, the universal policy needs to learn only one number, not a histogram or a threshold.

8.3 The exact worst-case curve

The clairvoyant scheduler knows p_i , uses the effective length $\lambda_u(p_i)$ from Equation (2), and orders these blocks shortest first. If P, P' are independent draws from a distribution D on $[0, u]$, its leading coefficient is $\Omega_{\text{BO},u}(D)$ from Equation (8). Consequently the instance-specific ratio from Theorem 8.5 is

$$\alpha_{\text{BO},u}(D) = \frac{\min\{u, 1 + \mathbb{E}P\}}{\mathbb{E} \min\{\lambda_u(P), \lambda_u(P')\}}. \quad (382)$$

Let $u_{\text{BO}} > 1$ denote the unique root greater than one of

$$u^3 - 2u^2 - u + 1 = 0, \quad u_{\text{BO}} = 2.2469796037 \dots \quad (383)$$

Theorem 8.6 (Exact blind-optimization curve). *The optimal randomized size-asymptotic competitive ratio in $\text{BO}(u)$ against an oblivious adversary is*

$$R_{\text{BO}}^{\text{rand}}(u) = \begin{cases} 1, & 0 < u \leq 1, \\ \frac{u^3}{u^2 + (u-1)^3}, & 1 \leq u \leq u_{\text{BO}}, \\ \frac{1}{2} \left(1 + \sqrt{\frac{u^2 + u - 1}{u - 1}} \right), & u \geq u_{\text{BO}}. \end{cases} \quad (384)$$

The two branches agree at u_{BO} . Moreover,

$$R_{\text{BO}}^{\text{rand}}(u) = \frac{\sqrt{u}}{2} + \frac{1}{2} + o(1) \quad (u \rightarrow \infty). \quad (385)$$

Proof. The claim is trivial for $u \leq 1$, because raw execution is no longer than the unit optimization operation. Assume $u > 1$ and let

$$S_D(t) = \mathbb{P}(P > t), \quad 0 \leq t \leq u.$$

The layer-cake identity applied to (8) gives

$$2\Omega_{\text{BO},u}(D) = 1 + \int_0^{u-1} S_D(t)^2 dt. \quad (386)$$

Put $S = \int_0^{u-1} S_D(t) dt$. Since S_D is nonincreasing,

$$\int_{u-1}^u S_D(t) dt \leq S_D(u-1) \leq \frac{S}{u-1}.$$

Therefore, writing $\mu = \mathbb{E}P = \int_0^u S_D(t) dt$,

$$S \geq \frac{u-1}{u} \mu. \quad (387)$$

Cauchy–Schwarz now yields

$$\int_0^{u-1} S_D(t)^2 dt \geq \frac{S^2}{u-1} \geq (u-1) \left(\frac{\mu}{u}\right)^2. \quad (388)$$

Set $b = \mu/u$. Equations (382)–(388) imply

$$\alpha_{\text{BO},u}(D) \leq f_u(b) := \frac{\min\{u, 1+ub\}}{1+(u-1)b^2}. \quad (389)$$

Equality is attained by the binary distribution

$$\mathbb{P}(P = u) = b, \quad \mathbb{P}(P = 0) = 1 - b. \quad (390)$$

Indeed, its survival function is constant b on $[0, u)$, so both inequalities in (388) are equalities.

It remains to maximize the scalar function f_u . Put

$$b_0 = \frac{u-1}{u}.$$

For $b \geq b_0$ the numerator is u and f_u decreases, so this region is maximized at b_0 . For $b \leq b_0$, differentiation gives the unique unconstrained maximizer

$$b_*(u) = \frac{u}{u-1 + \sqrt{(u-1)(u^2+u-1)}}. \quad (391)$$

The equality $b_*(u) = b_0$ is equivalent to $u^3 - 2u^2 - u + 1 = 0$. Hence b_0 is optimal up to u_{BO} and $b_*(u)$ is optimal afterwards. Substitution gives

$$f_u(b_0) = \frac{u^3}{u^2 + (u-1)^3}, \quad f_u(b_*) = \frac{1}{2} \left(1 + \sqrt{\frac{u^2+u-1}{u-1}} \right).$$

This proves (384). The uniformly relabeled binary distributions (390), together with Lemma 2.17, give the oblivious lower bounds. Together with the universal upper policy of Theorem 8.5, they prove both sides of the competitive-ratio claim. If the maximizing mass b is irrational, finite multisets use $\lfloor bn \rfloor$ copies of u ; continuity gives the same asymptotic ratio. Finally, $(u^2 + u - 1)/(u - 1) = u + 2 + 1/(u - 1)$ gives (385). $\square$

The hard distribution explains the separation. Most jobs have optimized length zero, while a carefully chosen small fraction have length u . Blind optimization is attractive on average, so the online policy optimizes the jobs. It cannot recognize a long outcome before committing to it, and an early long job delays many zeros. The clairvoyant optimum optimizes the zeros, runs the long jobs raw, and places them last. In the revealing model, by contrast, the same long outcomes can be deferred after their tests. The gap between the two curves therefore measures the scheduling value of the revealed information itself.

Acknowledgements

An initial improvement over the then-current deterministic lower bound for obligatory testing was obtained with the Bolzano open-source multi-agent research system [GHK⁺26]. That preliminary result motivated the present project. Most of the subsequent conjecture generation, proof development, counterexample search, exposition, and Lean formalization was carried out with substantial assistance from OpenAI’s GPT-5.6 through interactive Codex sessions. In particular, the randomized revealing-optimization curve was developed in a separate model run and then integrated with the instance-optimal framework. The author selected the research directions, reviewed and revised the arguments, and takes responsibility for the claims in this paper.

A Machine-checked verification

The accompanying Lean 4 development [dMU21] uses mathlib [The20] and is pinned to version 4.30.0. Its root module imports every development listed below. The following table states the proof boundary; Appendix D gives the detailed paper–Lean dictionary for the deterministic theorem.

paper result	Lean export	coverage
Deterministic revealing optimization and obligatory testing, Theorems 3.1 and 3.13	<code>SchedulingPaper.exact_ratio_main</code>	End-to-end: both bounds, fixed replayed inputs, completing strategies, linear remainders, all joins, and the obligatory endpoint.
Randomized obligatory testing, Theorem 5.5	<code>unknownMultiset_operational_expectedCost_le_20378; oblivious_iid_binary_lower_online_actualOPT</code>	End-to-end sharp worst-case value $4/3$ for the fixed-cutoff companion in Equation (267), including its finite error and a fixed binary lower-bound input. The growing-cutoff extension in Equation (266) is proved in the paper.
Obligatory-testing impossibility, Theorem 6.2	<code>ObligatoryInstance.deterministic_balanced_cost_lower; finiteSeed_exists_operational_quadratic_gap</code>	The adversarial-policy side of both parts is end-to-end. The deterministic proof includes adaptive revelation, freezing to one balanced labeled input, termination, and the sharp $9n^2/8$ cost bound. The unbounded randomized impossibility is also end-to-end for finite private seeds: literal all-long and one-zero runs are coupled up to the exceptional test, their actual completion costs are charged, and averaging selects one fixed oblivious zero label. The advertised comparison schedules enter through their checked closed-form cost bounds rather than through a second compiled transcript.
Obligatory-testing instance optimality, Theorem 6.1	<code>RandomizedAnnounced.threshold_maximizes_discovery_density; ObligatoryInstance.empiricalMaximumDensity_completion_le_line; ObligatoryInstance.empiricalPartialCompletion_le_thresholdPrefix; ObligatoryInstance.exists_empiricalObligatoryBenchmark; FluidToFinite.completion_transfer_uniform; FluidToFinite.pilot_learning_transfer</code>	The maximum-density algebra is instantiated on every concrete nonempty finite input: Lean constructs the threshold, proves its global fractional density certificate and the complete finite fractional pointwise envelope, and identifies the exact uniform cost of the announced stationary operation word, including its linear diagonal remainder. The two model-independent transfer principles are also checked. Connecting the fractional envelope to every literal online prefix, the bounded adaptive-policy lower bound, and the growing-cutoff operational learner are not yet assembled in Lean.

paper result	Lean export	coverage
Blind-execution results, Theorems 7.1 and 7.2	<code>boundedUniform_blindPilot_</code> <code>concrete_rate;</code> <code>UnboundedOperational.no_</code> <code>finite_ratio_at_quadratic_</code> <code>scale_operational</code>	End-to-end uniform $o_L(n^2)$ comparison with every complete announced policy; the checked rate is slower than the paper's rate. The unbounded impossibility is also end-to-end: the all-zero first-touch order is extracted from each literal transcript, lockstep causality preserves the exceptional label's blind rank, suffix accounting charges its actual one-long-job run, and finite-seed averaging fixes one oblivious label.
Revealing-optimization instance and randomized curve, Theorems 4.1 and 5.1	<code>RevealingOptimization.</code> <code>binaryFamilies_le_curve;</code> <code>binaryFamilies_attain_curve;</code> <code>randomizedCurve_attains_</code> <code>globalMaximum;</code> <code>RevealingOptimization.</code> <code>announcedTemplate_curve_</code> <code>upper;</code> <code>RevealingOptimization.</code> <code>RandomizedLower.adaptive_</code> <code>binary_families_attain_curve;</code> <code>RevealingOptimization.</code> <code>RandomizedYao.finiteSeed_</code> <code>binary_families_attain_curve;</code> <code>FluidToFinite.completion_</code> <code>transfer_uniform;</code> <code>FluidToFinite.pilot_learning_</code> <code>transfer</code>	The exact four-piece scalar maximization, its transition root, plateau, attainment, and global maximum are checked. The survival reduction now starts from a concrete nonempty finite input, constructs its maximum-density threshold, identifies the exact empirical survival and offline minimum-pair formulas, derives the prefix-energy and adjacent-tail inequalities, and reaches the exact curve in all three threshold regimes. The resulting announced endpoint is an explicit uniformly permuted stationary-or-raw operation word, and its finite bound retains an explicit linear diagonal remainder. Its denominator is identified exactly with the empirical clairvoyant minimum-pair leading term, including the nonnegative diagonal correction to the literal finite SPT optimum. The common completion-area and pilot-learning transfer is checked abstractly. On the lower side an exact finite Bellman induction now covers every adaptive test–raw–process decision tree and both attaining binary families. The lower tree now retains job labels, its recursive expectation is identified with the exact Bernoulli average over fixed labelled inputs, and finite Yao selects one input before an arbitrary finite private seed. The model-specific finite revealing-policy envelope and operational pilot upper bound, together with the final compiler from a literal transcript-only strategy to this labelled tree, are not yet assembled in Lean.

paper result	Lean export	coverage
Blind-optimization results, Theorems 8.1, 8.3, 8.5 and 8.6	<code>BlindOptimization.Adversary.no_deterministic_instance_optimal; deterministicCurve_operational_exact; operationalPilot_expectedCost_le_curve_normalized; pilotNormalizedError_tendsto_zero; RandomizedLower.exists_binary_mass_adaptive_curve_lower; RandomizedLower.exists_binary_mass_finite_ratio_tendsto_curve; randomizedBinaryEnvelope_attains_curve; Randomized.finite_yao_select_weighted; RandomizedCompiler.exists_binary_mass_operational_ratio_tendsto_curve</code>	The deterministic counterexample and exact deterministic curve are end-to-end, including replay, adaptive hidden stopping, completing policies, and finite linear remainders. The randomized upper is checked for literal independent pilot/main orders, with without-replacement learning error, observable threshold, exact finite offline comparison, and explicit remainder; each private seed is now a literal completing transcript-only strategy whose run cost is identified with the analyzed schedule. The distribution reduction, scalar maximum, attainment, both large-u asymptotics, fully adaptive binary decision-tree lower bound, exact finite diagonal limit, and finite weighted Yao selection (which permits the exact, possibly irrational Bernoulli mass) are checked. An arbitrary transcript-only strategy is mechanically unfolded into the corresponding labeled adaptive tree. Its structural expectation is identified with the weighted average of literal operational runs, the exact finite offline expectation (including its diagonal correction) is checked, and Yao selects one fixed labeled input against the entire private-seed family at a ratio converging to the advertised curve.

The development contains no `sorry`, `admit`, or project-defined axiom. Formal randomized statements use finite uniform averages for private seeds and explicit nonnegative finite weights for Yao input laws; these cover permutations, sample choices, tie-breaking coins, and the exact Bernoulli masses used here. Classical real analysis and noncomputable exact minimizers occur only in certificates; no floating-point program is extracted.

B Finite rational checkpoint for the cap-free lower bound

The lower bound at $u = \infty$ takes two limits: first the scaling $H \rightarrow \infty$, and then the number K of positive levels tends to infinity. This appendix is not needed for that proof. It gives one exact finite instance showing that the leading-term calculation already predicts the true ratio accurately.

Take

$$K = 1000, \quad \alpha = \frac{9}{8}, \quad \gamma = \frac{679}{1000}, \quad \xi = \frac{K}{\alpha} = \frac{8000}{9}, \quad H = 9000.$$

There are H jobs at each positive level and ξH zero jobs. All multiplicities are integral, and all processing times defined by Equation (169) are rational.

Using exact rational arithmetic in Equations (168) and (172) to (174) gives the leading-coefficient comparison

$$\frac{Q_K}{D_K} = 1.5705287442294085 \dots > 1.5705.$$

The displayed decimal is only for orientation; all comparisons below are between integers or rational numbers.

For the offline schedule, retaining the terms that are linear in H gives

$$\text{OPT}_H = H^2 D_K + H L_{\text{off}}, \quad (392)$$

where

$$\begin{aligned} L_{\text{off}} &= \frac{1}{2} \left(\xi + \sum_{i=0}^{K-1} (1 + p_i) \right) \\ &= \frac{1}{2} \left(\xi + 2K + \frac{K\gamma}{\xi} + \sum_{s=1}^{K-1} \frac{K-s}{\xi+s} \right) = 1656.440800628232 \dots \end{aligned} \quad (393)$$

Again, the middle expression is the exact rational value used in the comparison.

The discrete shortest-prefix calculation behind Equation (166) also retains favorable diagonal terms that vanish after division by H^2 . Before dropping them it gives

$$\text{ALG}_H \geq H^2 Q_K + \frac{H}{2} \left(\xi + \sum_{i=0}^{K-1} p_i + \sum_{i=0}^{K-1} s_i \right) \geq H^2 Q_K.$$

Here the s_i 's are the nonnegative completion fractions from the lower proof. Exact rational arithmetic verifies

$$2000H^2 Q_K > 3141(H^2 D_K + H L_{\text{off}}). \quad (394)$$

Consequently, the fixed instance obtained by the replay argument in Section 3.2.2 satisfies

$$\frac{\text{ALG}_H}{\text{OPT}_H} \geq \frac{H^2 Q_K}{H^2 D_K + H L_{\text{off}}} = 1.5705286254871234 \dots > \frac{3141}{2000} = 1.5705.$$

Thus the checkpoint includes the finite- H correction; it is not merely a comparison of the two leading coefficients.

C Independent certificate by bubbling caps to a prefix

For completeness, we give a second proof of the quadratic estimate. This argument is also a useful audit of the fifth-endpoint calculation: it never inserts $p = u$ into an ordinary deferred formula before the caps have actually been moved.

Work first in normalized fluid mass. Immediately before an adjacent pair XU , let E_0 be the preceding E -mass, P the preceding substantive mass, N_{def} its deferred submass, x the remaining mass, and w the mass of U -endpoints strictly after the pair. Put

$$y = \frac{N_{\text{def}} - R(P + E_0)}{x}, \quad \eta = \frac{N_{\text{def}} - RP}{x} = y + R \frac{E_0}{x} \leq 0. \quad (395)$$

Direct differentiation of the exact pair objective (107), including the mutual pair and all future caps, gives the two nontrivial commutators

$$UM - MU = 1 + \left(1 + \eta - \frac{Rw}{x} \right) (y - c) a'_c(y), \quad (396)$$

$$UQ - QU = \left(\eta - \frac{cw}{x} \right) (y - c)a'_c(y). \quad (397)$$

Here $UX - XU$ means “new objective after moving U left” minus the old objective. The second quantity is nonnegative. For the first, if $B = 1 + \eta - Rw/x \leq 0$, its second term is nonnegative. If $B > 0$, then $B \leq 1$, and on $-1 < y \leq 0$

$$\frac{(c - y)B}{c - 2y} \leq 1.$$

Thus (396) is nonnegative as well. Below $y = -1$, the threshold is flat. Direct evaluation also gives

$$UZ - ZU = UE - EU = 1. \quad (398)$$

The suffix terms involving w are crucial: they make every adjacent exchange valid even while other caps remain to the right. Repeated swaps therefore put all caps into one prefix.

After that prefix there is no future cap. Let e, b, d be the preceding E -mass, substantive mass, and deferred substantive mass, and define

$$y = \frac{d - R(e + b)}{x}, \quad v = \frac{e}{x}, \quad h = d - Rb, \quad k = x + d - Rb.$$

The four ordinary commutators are

$$\begin{aligned} ME - EM &= a_c + a'_c(R - y)\frac{k}{x}, & QE - EQ &= 1 - a'_c(R - y)(-y - Rv), \\ MZ - ZM &= a_c - a'_c y \frac{k}{x}, & QZ - ZQ &= 1 - a'_c y \frac{h}{x}. \end{aligned} \quad (399)$$

They are nonnegative. Indeed, in the varying region $h/x = y + Rv \in [y, 0]$ and $k/x \in [0, 1]$. With $t = -y \in [0, 1]$,

$$a'_c(R - y)t = \frac{t(R + t)}{c + 2t} \leq 1, \quad a'_c t^2 = \frac{t^2}{c + 2t} \leq 1; \quad (400)$$

the first inequality is equivalent to $(t - 1)(t + c) \leq 0$. In the flat region $a'_c = 0$. Hence Z, E can be moved to the tail, and a worst word has the form

$$U^\mu \{M, Q\}^* \{Z, E\}^*. \quad (401)$$

On the middle block use the simpler potential

$$\widehat{W}(x, N, N_{\text{def}}) = xN_{\text{def}} + x^2 f_c \left(\frac{N_{\text{def}} - RN}{x} \right).$$

The identity

$$2f_c - yf'_c = \max\{0, -Rf'_c, -Rf'_c + a_c(1 + y), 1 - cf'_c + a_c y\} \quad (402)$$

is a direct rewriting of (120); it pays the entire ordinary continuation. If the cap prefix has mass μ , then after it $x = 1 - \mu$, $N = N_{\text{def}} = \mu$, and $y = -c\mu/(1 - \mu)$. Its internal cap-pair reward plus its worst continuation is therefore

$$F_{c,u}(\mu) = \frac{2 - c(u - 1)}{2} \mu^2 + \mu(1 - \mu) + (1 - \mu)^2 f_c \left(-\frac{c\mu}{1 - \mu} \right). \quad (403)$$

For $0 < \mu < 1/R$, differentiation gives

$$F'_{c,u}(\mu) = c\mu [Z(c) - u - \operatorname{artanh} \mu]. \quad (404)$$

On the mixed curve the unique maximizer in this range is therefore $\mu = m$, and

$$F_{c,u}(m) = f_c(0) + \frac{c}{2}(\operatorname{artanh} m - m) = \frac{c}{2}. \quad (405)$$

Here $m \leq 1/R$, since $c, m \leq 1/\varphi$. On the flat part $\mu \geq 1/R$,

$$F_{c,u}(\mu) = \mu - \frac{c(u-1)}{2}\mu^2. \quad (406)$$

Its derivative at $1/R$ is

$$\frac{c(1+m) - 1}{R} \leq 0,$$

and only decreases thereafter. Thus $F_{c,u}(\mu) \leq c/2$ on the whole unit interval. At $c = r, m = 0, u = u_5^{\det}$, the same calculation is maximized at $\mu = 0$, and increasing u only lowers (403).

To make this independent exchange proof finite, fix $\varepsilon > 0$ and first replace the last $\varepsilon n + O(1)$ outcomes by genuine zeros. Earlier states and actions are unchanged, and only $O(\varepsilon n^2 + n)$ pair terms are affected. All cap and ordinary exchanges in the retained prefix then occur with $x \geq \varepsilon$; their displayed fluid commutators have uniform Taylor remainders there. More precisely, after assigning mass $1/n$ to every job, a normalized adjacent exchange has remainder $O_{u,\varepsilon}(n^{-3})$; there are at most n^2 exchanges. At the join $y = -1$, split an exchange at the join and use the two one-sided formulae, whose values agree. The artificial zeros already form the terminal block, so no unubbled suffix cap invalidates (399). The normalized one-job remainder in the continuation-potential calculation is $O_{u,\varepsilon}(n^{-2})$. For fixed ε , first let $n \rightarrow \infty$, and then let $\varepsilon \downarrow 0$. Equations (402)–(405) give

$$\Gamma_n \leq c \binom{n}{2} + o(n^2),$$

which is an independent size-asymptotic certificate for the two upper bounds above.

D Machine-checked deterministic theorem and its correspondence

This appendix records the public theorem exported by the Lean 4 development [dMU21] associated with this manuscript and explains how its formal predicates correspond to the deterministic statements in Theorems 3.1 and 3.13. The development uses the mathlib library [The20] and is pinned to Lean and mathlib version 4.30.0. Its root module imports `SchedulingPaper.MainTheorems`, whose final theorem is reproduced below. Apart from the typographic rendering of Unicode symbols, this is a verbatim excerpt.

```
theorem exact_ratio_main
  : (∀ {u : ℝ}, 0 < u →
    FiniteExactRatioConclusion u (exactCurve u)) ∧
    ObligatoryExactRatioConclusion RStar := by
  constructor
  · intro u hu
    exact finite_exact_ratio hu
  · exact obligatory_exact_ratio
```

The two conclusion records occurring in the theorem are:

```

structure FiniteExactRatioConclusion (u ratio :  $\mathbb{R}$ ) : Prop where
  lower : LowerBound.FixedSizeLowerBound (.finite u) ratio
  upper :
     $\exists$  strategy :  $\forall$  n, Online.Strategy n,
      UpperBound.CompletesByAnalysisFuel (.finite u) strategy  $\wedge$ 
      SizeAsymptoticUpper
        (UpperBound.strategyCost (.finite u) strategy)
        (UpperBound.fixedOfflineCost (.finite u)) ratio

structure ObligatoryExactRatioConclusion (ratio :  $\mathbb{R}$ ) : Prop where
  lower : LowerBound.FixedSizeLowerBound .infinite ratio
  upper :
     $\exists$  strategy :  $\forall$  n, Online.Strategy n,
      UpperBound.CompletesByAnalysisFuel .infinite strategy  $\wedge$ 
      SizeAsymptoticUpper
        (UpperBound.strategyCost .infinite strategy)
        (UpperBound.fixedOfflineCost .infinite) ratio

```

D.1 Objects on the two sides

The following table gives the direct dictionary between the formal objects and the notation of the manuscript.

Lean object	Manuscript object	Meaning
<code>Cap.finite u</code>	finite common upper limit u	processing times lie in $[0, u]$; testing and raw execution are available
<code>Cap.infinite</code>	obligatory testing	processing times are nonnegative and raw execution is unavailable
<code>exactCurve u</code>	$R_{\text{RO}}^{\text{det}}(u)$	the six branches in Equation (75), including all five verified joins
<code>RStar</code>	$R_{\text{OT}}^{\text{det}}$	$1 + 2/(u_5^{\text{det}} - 1)$, where $u_5^{\text{det}} > 1$ is the unique solution of $u_5^{\text{det}} - 3 = \ln u_5^{\text{det}}$
<code>fixedOfflineCost</code>	OPT	the clairvoyant shortest-first cost of the model-dependent effective lengths in Equation (3)
<code>strategyCost</code>	ALG	the total completion cost of the operational online transcript

A formal strategy on n jobs is a deterministic function from the public transcript to its next action. A test publishes the revealed processing time; raw execution publishes no hidden value. Thus the strategy has exactly the information available to an online algorithm in the manuscript. The formal action type uses the standard no-idling normalization appropriate when all jobs are released at time zero. Processing times and comparisons are exact real numbers.

D.2 Expansion of the lower and upper fields

For a cap and a candidate value R , `FixedSizeLowerBound` expands to the following statement. For every size-indexed family of deterministic strategies $\mathcal{B}_n$, every $\varepsilon > 0$, and every N , there exist $n \geq N$, a genuine fixed admissible n -job input I , and a finite settled execution such that either the execution leaves a job incomplete or

$$(R - \varepsilon) \text{OPT}(I) \leq \text{ALG}_{\mathcal{B}_n}(I). \quad (407)$$

An incomplete settled run represents infinite cost. In particular, for algorithms that complete all inputs, (407) is the usual lower-bound sequence on arbitrarily large fixed instances. The lower constructions are first described by adaptive revelation oracles, but the formal replay theorem freezes every such interaction into the fixed input quantified here; this is the machine-checked counterpart of Lemma 2.10.

The predicate **SizeAsymptoticUpper** is the epsilon formulation

$$\forall \varepsilon > 0 \exists N \forall n \geq N \forall I, \quad \text{ALG}_{\mathcal{A}_n}(I) \leq (R + \varepsilon) \text{OPT}(I). \quad (408)$$

The accompanying completion field proves that every job is done after the common analysis fuel $2n + 1$, rather than merely assigning a cost to a partial trace. Consequently, Equations (407) and (408) are the two matching operational statements that characterize the optimal value in Equation (35). The Lean theorem packages these two directions; it does not introduce a separate real-valued infimum or `limsup` object.

Before projecting to **SizeAsymptoticUpper**, the checked upper proof constructs a stronger **FixedUpperCertificate**: for the chosen cap it contains a constant $C \geq 0$ such that, for every n and every fixed input,

$$\text{ALG}_{\mathcal{A}_n}(I) \leq R \text{OPT}(I) + Cn. \quad (409)$$

The quadratic lower bounds Equations (16) and (17) then imply (408). For $u \geq u_5^{\text{det}}$, the separately verified plateau certificate uses the same strategy and the same C , independent of u , including the obligatory limit. This uniformity is stronger than the quantifier order displayed by `exact_ratio_main`, which only exposes one strategy after a cap has been fixed.

D.3 Correspondence with the two deterministic theorems

The first conjunct of `exact_ratio_main` quantifies over every finite $u > 0$. Substituting $R = \text{exactCurve } u = R_{\text{RO}}^{\text{det}}(u)$, its lower field gives the fixed-instance lower direction of Theorem 3.1, while its upper field supplies a completing strategy and the matching upper direction. The definition of `exactCurve` is literally the six-way case split in Equation (75); the Lean development also proves its joins, continuity, plateau, and global maximum at u_2^{det} .

The second conjunct substitutes $R = \text{RStar} = R_{\text{OT}}^{\text{det}}$ in the obligatory model. Its lower field gives the lower half of Theorem 3.13; its completing upper strategy together with (409) gives the upper half. The common formal infrastructure also proves that the offline benchmark is the shortest-first schedule of effective lengths and that the transcript cost is exactly its suffix-weighted pair-accounting expression.

D.4 Proof boundary

The root Lean module imports the complete lower- and upper-bound assemblies. The public theorem has no operational interface hypotheses, and the development contains no `sorry`, `admit`, or project-defined `axiom`. It does use `mathlib`'s classical real analysis and noncomputable choices of exact real parameters. Thus it certifies deterministic mathematical strategy terms, not an extracted floating-point implementation.

The theorem recorded here is specifically the size-asymptotic result. It does not assert the pointwise inequality $\text{ALG} \leq R \text{OPT}$ without a remainder. Its checked remainder is the linear term in (409), and is therefore $o(n^2)$. The machine-checked mixed-regime upper bound follows the cap-reserve potential proof; the bubbling argument in Appendix C remains an independent manuscript cross-check.

References

- [ABC17] Peyman Afshani, Jérémy Barbay, and Timothy M. Chan. Instance-optimal geometric algorithms. *Journal of the ACM*, 64(1):3:1–3:38, 2017. doi:10.1145/3046673.
- [AE21] Susanne Albers and Alexander Eckl. Explorable uncertainty in scheduling with non-uniform testing times. In Christos Kaklamanis and Asaf Levin, editors, *Approximation and Online Algorithms: 18th International Workshop, WAOA 2020, Revised Selected Papers*, volume 12806 of *Lecture Notes in Computer Science*, pages 127–142. Springer, 2021. doi:10.1007/978-3-030-80879-2_9.
- [Azu67] Kazuoki Azuma. Weighted sums of certain dependent random variables. *Tohoku Mathematical Journal*, 19(3):357–367, 1967. doi:10.2748/tmj/1178243286.
- [BS26] Felix Buld and Andreas S. Schulz. Scheduling with testing: Competitive algorithms for minimizing the total weighted completion time in the adversarial model, 2026. arXiv:2606.25166.
- [Che52] Herman Chernoff. A measure of asymptotic efficiency for tests of a hypothesis based on the sum of observations. *The Annals of Mathematical Statistics*, 23(4):493–507, 1952. doi:10.1214/aoms/1177729330.
- [DDN⁺22] Fanny Dufossé, Christoph Dürr, Noël Nadal, Denis Trystram, and Óscar C. Vázquez. Scheduling with a processing time oracle. *Applied Mathematical Modelling*, 104:701–720, 2022. doi:10.1016/j.apm.2021.12.020.
- [DEL24] Konstantinos Dogeas, Thomas Erlebach, and Ya-Chun Liang. Scheduling with obligatory tests. In Timothy Chan, Johannes Fischer, John Iacono, and Grzegorz Herman, editors, *32nd Annual European Symposium on Algorithms (ESA 2024)*, volume 308 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 48:1–48:14, Dagstuhl, Germany, 2024. Schloss Dagstuhl – Leibniz-Zentrum für Informatik. doi:10.4230/LIPIcs.ESA.2024.48.
- [DEMM20] Christoph Dürr, Thomas Erlebach, Nicole Megow, and Julie Meißner. An adversarial model for scheduling with testing. *Algorithmica*, 82(12):3630–3675, 2020. doi:10.1007/s00453-020-00742-2.
- [DKL⁺23] Christoph Damerius, Peter Kling, Minming Li, Chenyang Xu, and Ruilong Zhang. Scheduling with a limited testing budget: Tight results for the offline and oblivious settings. In *31st Annual European Symposium on Algorithms (ESA 2023)*, volume 274 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 38:1–38:15. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2023. doi:10.4230/LIPIcs.ESA.2023.38.
- [dMU21] Leonardo de Moura and Sebastian Ullrich. The Lean 4 theorem prover and programming language. In *Automated Deduction – CADE 28*, volume 12699 of *Lecture Notes in Computer Science*, pages 625–635. Springer, 2021. doi:10.1007/978-3-030-79876-5_37.
- [Doo53] Joseph L. Doob. *Stochastic Processes*. Wiley, New York, 1953.

- [GCH24] Mingyang Gong, Zhi-Zhong Chen, and Kuniteru Hayashi. Approximation algorithms for multiprocessor scheduling with testing to minimize the total job completion time. *Algorithmica*, 86(5):1400–1427, 2024. doi:10.1007/s00453-023-01198-w.
- [GHK⁺26] Jan Grebík, Pavel Hubáček, Martin Koutecký, Matěj Kripner, Václav Rozhoň, Robert Šámal, and Adrián Zámečník. Bolzano: Case studies in LLM-assisted mathematical research, 2026. URL: <https://arxiv.org/abs/2604.16989>, arXiv:2604.16989.
- [Hoe63] Wassily Hoeffding. Probability inequalities for sums of bounded random variables. *Journal of the American Statistical Association*, 58(301):13–30, 1963. doi:10.1080/01621459.1963.10500830.
- [Kah91] Simon Kahan. A model for data in motion. In *Proceedings of the 23rd Annual ACM Symposium on Theory of Computing (STOC 1991)*, pages 265–277. Association for Computing Machinery, 1991. doi:10.1145/103418.103449.
- [KLL⁺26] Bob Krekelberg, Alison Hsiang-Hsuan Liu, Fu-Hong Liu, Prudence W. H. Wong, and Xiao-Ou Zhang. The power of amortization on minimizing total completion time with explorable uncertainty, 2026. Version 2, revised 8 May 2026. arXiv:2503.13357.
- [LL26] Kao-Chuan Liang and Ya-Chun Liang. The completion-threshold framework for obligatory-test scheduling on multiple machines, 2026. Version 2, revised 14 July 2026. arXiv:2606.02029.
- [LLWZ23] Alison Hsiang-Hsuan Liu, Fu-Hong Liu, Prudence W. H. Wong, and Xiao-Ou Zhang. The power of amortization on scheduling with explorable uncertainty. In Jaroslav Byrka and Andreas Wiese, editors, *Approximation and Online Algorithms: 21st International Workshop, WAOA 2023*, volume 14297 of *Lecture Notes in Computer Science*, pages 90–103. Springer, 2023. doi:10.1007/978-3-031-49815-2_7.
- [LMS19] Retsef Levi, Thomas L. Magnanti, and Yaron Shaposhnik. Scheduling with testing. *Management Science*, 65(2):776–793, 2019. doi:10.1287/mnsc.2017.2973.
- [LMS24] Retsef Levi, Thomas L. Magnanti, and Yaron Shaposhnik. Scheduling with testing of heterogeneous jobs. *Management Science*, 70(5):2934–2953, 2024. doi:10.1287/mnsc.2023.4833.
- [MPT94] Rajeev Motwani, Steven J. Phillips, and Eric Torng. Nonclairvoyant scheduling. *Theoretical Computer Science*, 130(1):17–47, 1994. doi:10.1016/0304-3975(94)90151-1.
- [NRTT24] Shyam Narayanan, Václav Rozhoň, Jakub Tětek, and Mikkel Thorup. Instance-optimality in I/O-efficient sampling and sequential estimation. In *65th IEEE Annual Symposium on Foundations of Computer Science (FOCS 2024)*, pages 658–688. IEEE, 2024. doi:10.1109/FOCS61266.2024.00048.
- [The20] The mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and*

Proofs (CPP 2020), pages 367–381. Association for Computing Machinery, 2020. doi:10.1145/3372885.3373824.

- [Yao77] Andrew Chi-Chih Yao. Probabilistic computations: Toward a unified measure of complexity. In *18th Annual Symposium on Foundations of Computer Science*, pages 222–227. IEEE, 1977. doi:10.1109/SFCS.1977.24.