

Scheduling with Uniform Tests: Information and Optimization

Václav Rozhoň

August 2026

Abstract

A machine faces many jobs whose lengths are hidden. Spending one unit of time to inspect a job may reveal a short job that should be finished now, or it may reveal nothing useful while every other job waits. When should the machine keep looking, and when should it start working?

We study this question for one machine and total completion time. We consider four natural rules: every job must be tested; a unit-time revealing optimization can replace a raw time u by a shorter revealed length; an untested job can be run blindly for its true hidden length; or a unit-time optimization can shorten it without revealing the new length. Together these models separate the two possible benefits of spending one unit before execution: learning which job to do next and making the job faster.

We give exact asymptotic answers for these models, both as competitive ratios and as instance-optimal guarantees. The resulting algorithms are often quite simple, which may make them useful in practice.

Contents

1	Introduction	3
1.1	Warmup: every job has to be tested	3
1.2	One algorithm for every bag	4
1.3	Generalizations of the model	5
1.4	Information without speedup: blind execution	6
1.5	Blind optimization	7
1.6	Revealing optimization	7
1.7	Techniques	9
1.8	Related work	10
2	Models and common preliminaries	10
2.1	Information and adversaries	11
2.2	Offline optima and pair accounting	11
2.3	Competitive-ratio conventions	13
2.4	Test-order adversaries	13
3	Deterministic obligatory testing: the exact ratio	13
3.1	Harmonic core and matching lower bound	14
3.2	Upper bound: a self-contained proof	19
4	Randomized obligatory testing: the sharp ratio is $4/3$	27
4.1	The normalized offline objective	27
4.2	Stationary prefixes	27
4.3	Learning the prefix from a sublinear sample	29

4.4	Good samples, fallback, and bad-event excess	30
4.5	The balanced binary lower bound	32
5	Obligatory-testing instance optimality	32
5.1	The maximum-density benchmark	34
5.2	The announced upper bound	35
5.3	No adaptive announced policy has a better leading cost	36
5.4	The same unannounced policy for every bag	37
6	Blind execution: instance-optimal two-phase policies	38
6.1	The distributional benchmark	39
6.2	The fixed-fraction completion envelope	41
6.3	Adaptive first touches are a uniform urn	42
6.4	The announced-multiset lower bound	43
6.5	A growing grid and arbitrary bounded multisets	44
6.6	Removing the announcement	44
6.7	Why the medium block is necessary	45
7	Blind optimization: speedup without revelation	46
7.1	Proof of instance optimality	46
7.2	The exact worst-case curve	48
8	Deterministic revealing optimization: the exact curve	50
8.1	The shape of the claimed curve	52
8.2	The binary stopping game and its lower envelope	52
8.3	Matching upper bounds through the binary regime	54
8.4	The unrestricted mixed lower bound	58
8.5	The large-upper-limit lower bound by cap-free transfer	60
8.6	Completion of the curve and consistency at the joins	61
8.7	Matching upper bound for revealing optimization	62
9	Revealing-optimization instance optimality	68
9.1	The fixed-fraction fluid benchmark	69
9.2	Adaptive policies and the announced lower bound	71
9.3	A universal policy without the multiset	73
10	Randomized revealing optimization: the exact four-piece curve	74
10.1	Offline and stationary fluid coefficients	75
10.2	Flattening the survival function	77
10.3	The two scalar maximizations	78
10.4	Learning the announced policy	79
10.5	Binary oblivious lower bounds	79
	Acknowledgements	82
A	Machine-checked verification	82
A.1	Deterministic exact ratios	82
A.2	Randomized obligatory testing	82
A.3	Blind-execution instance optimality	83
B	Finite rational checkpoint for the cap-free lower bound	83

C Independent certificate by bubbling caps to a prefix	84
D Machine-checked deterministic theorem and its correspondence	86
D.1 Objects on the two sides	87
D.2 Expansion of the lower and upper fields	87
D.3 Correspondence with the two deterministic theorems	88
D.4 Proof boundary	88

1 Introduction

Suppose that we have many jobs waiting for one machine. We want to minimize the sum of the times at which they finish. The length of each job is written on a folded piece of paper. If we knew all the lengths, the right schedule would be clear: we would simply run the jobs from shortest to longest. But the papers are folded, and opening one of them takes one unit of machine time.

Should we open another paper, or should we start working?

Opening a paper may reveal a tiny job that we can finish immediately. It may also reveal a huge job that is best left for the end. On the other hand, looking has a cost. If one hundred jobs are still unfinished, one unit spent opening a paper adds one hundred units to the sum of completion times. The machine must therefore balance two uses of its time: learning which jobs are short and completing jobs that are already understood.

Variants of this question have appeared in both stochastic and adversarial scheduling with testing [LMS19, DEMM20, DEL24]. We start with its cleanest version, in which every paper has to be opened. We then allow a job to be run without opening its paper. Depending on what this means, we obtain three further variants. The same simple ideas—random sampling, thresholding, and shortest-first execution—give exact answers in all of them.

1.1 Warmup: every job has to be tested

There are n jobs. Testing job i takes one unit of time and reveals a number $p_i \geq 0$. The job then needs another p_i units of processing; if $p_i = 0$, it completes with the test. For now every job has to be tested: testing is *obligatory*. The formal model is given in Section 2.

We compare an online scheduler with the shortest-first schedule that knows all values p_i in advance. The *competitive ratio* is the online cost divided by this offline optimum. We care about its asymptotic value, so lower-order additive terms of size $o(n^2)$ do not matter.

The obligatory-testing model for total completion time was introduced by Dogeas, Erlebach, and Liang [DEL24]. Their work and a later lower bound of Liang and Liang [LL26] left a gap between the best deterministic upper and lower bounds. They also left the randomized problem open. We close both gaps.

Theorem 1.1 (Obligatory testing, informal version of Theorems 3.1 and 4.1). *In the obligatory-testing model, the best deterministic asymptotic competitive ratio is the constant $R = 1.570574\dots$. Against an adversary that fixes the input before the algorithm draws its random bits, the best randomized ratio is $4/3$. The two optimal policies are sketched in algorithms 1 and 2.*

Here and below, *shortest first* means processing revealed jobs in nondecreasing order of p_i . The following pseudocode gives a little more detail about the two policies. The formulas are not meant to be memorable; the shape of the algorithms is.

Algorithm 1 Deterministic threshold

- 1: Put $\rho = 0.57 \dots$ and $R = 1 + \rho$.
- 2: Before test $i + 1$, where $0 \leq i < n$, let c_i be the number of positive jobs processed immediately and d_i the number deferred.
- 3: Put

$$y_i = \frac{d_i - R(c_i + d_i)}{n - i}.$$

- 4: Use threshold $T_i = 1$ if $y_i \leq -1$; otherwise use

$$T_i = 1 + \frac{1}{2} \log \frac{\rho + 2}{\rho - 2y_i}.$$
 - 5: Test the next job. Process it now if $p \leq T_i$; otherwise defer it.
 - 6: After all tests, finish deferred jobs shortest first.
-

Algorithm 2 Learned randomized threshold

- 1: Shuffle the jobs and test $o(n)$ of them to estimate the random variable X of job lengths distributions.
- 2: Treat this estimate as exact. For each threshold $0 \leq t \leq 100$ put

$$a(t) = \Pr(X \leq t), \quad m(t) = \mathbb{E}[X \mathbf{1}_{X \leq t}].$$

$$\hat{\rho} = \max_t \frac{a(t)}{1 + m(t)}.$$

- 3: If $\hat{\rho} < 1/100$, test every remaining job and then finish all revealed jobs shortest first.
 - 4: Otherwise set $T = 1/\hat{\rho}$. Process the sampled jobs with $p \leq T$. Keep testing jobs, processing them immediately when $p \leq T$ and deferring them otherwise.
 - 5: After all tests, finish deferred jobs shortest first.
-

Both algorithms follow the same principle: process short jobs as soon as possible and leave long jobs for the end. The difficult part is choosing the threshold. A deterministic algorithm has no representative view of the unseen jobs, so its threshold has to change with the history. The exact update is given in steps 3 and 4 of algorithm 1. The lower bound explains this moving threshold. It reveals positive jobs in decreasing levels while hiding many zero jobs until the end. Processing a positive level immediately delays the hidden zeros; postponing it instead creates a costly tail. The levels can be chosen so that these two losses balance at $1.57 \dots$; the formal construction is in Section 3.

Randomization gives the algorithm a global view of the input. It first tests a sublinear random sample and estimates the distribution of job lengths. This costs only $o(n^2)$ and therefore does not change the asymptotic ratio. Up to a small estimation error, the algorithm now knows what remains in the bag and can keep one fixed threshold for the rest of the schedule. The earlier obligatory-testing work studied deterministic algorithms and explicitly left randomization open [DEL24].

The matching randomized lower bound is correspondingly simple. Half of the jobs have length zero and half have length two, and their labels are shuffled. Until the shuffle is uncovered, one unit of work completes only about half a job. Counting the area under the number of unfinished jobs gives the lower bound $4/3$. The formal argument is in Section 4.

1.2 One algorithm for every bag

Let us return to algorithm 2. The ratio $4/3$ describes only the hardest bag. It says much less on an easy one. Suppose, for example, that all processing times are very large compared with the unit test. Then the simple strategy that first tests everything and subsequently runs the jobs shortest first is nearly optimal. A worst-case guarantee still tells us only that our cost is at most $1.33 \dots$ times the optimum.

A slight variant of algorithm 2 satisfies a much stronger bag-by-bag guarantee: it is *instance-optimal*. The variant maximizes over thresholds up to $O(n^{1/20})$, rather than up to a fixed constant. It is still worst-case optimal, with competitive ratio $4/3$; we presented

the fixed-cutoff algorithm first because its worst-case analysis is simpler.

Theorem 1.2 (Obligatory-testing instance optimality, informal version of Theorem 5.1). *Fix $L < \infty$. There is a variant $\mathcal{A}^*$ of algorithm 2 such that, for every bag $M \subseteq [0, L]$ and every online algorithm $\mathcal{A}'$,*

$$\text{cost}(\mathcal{A}^*) \leq \text{cost}(\mathcal{A}') + o_L(n^2),$$

where both costs are measured on M and include the algorithms' randomness. The competing algorithm $\mathcal{A}'$ may even be designed specifically for M and told the entire bag in advance.

Both assumptions in the theorem are necessary. Without a fixed bound L there is no randomized instance-optimal algorithm, and without randomization there is no deterministic one; see Theorem 5.2.

Two points about the theorem are worth emphasizing. First, the comparison is made separately on every bag. We must therefore compete not only with one fixed online algorithm, but also with algorithms optimized for this particular M . Allowing $\mathcal{A}'$ to know M makes this strongest possible online benchmark explicit.

Hence, the existence of some instance-optimal algorithm is perhaps not so surprising. One could imagine a meta-algorithm that samples the input to estimate M and then runs an optimal policy for that estimate. The substantive part of the theorem is that no complicated meta-algorithm is needed. The optimal leading-order policy for a known bag is itself a simple threshold policy, so $\mathcal{A}^*$ has essentially the same form as algorithm 2: sample the bag, choose the maximum-density threshold, and use it for the rest of the schedule.

1.3 Generalizations of the model

We next weaken the obligatory model in two directions. First, we may allow a job to run without being tested. Second, the unit preliminary action may have two different benefits. It may reveal the true running time p_i before the job is run. It may also be an optimization procedure that replaces a common raw running time u by a shorter time $p_i \leq u$. For example, a generic piece of code solves every task within time u , but an AI agent can spend one unit adapting the code to a particular input and make it faster.

These two benefits are independent. An optimization may reveal its new running time p_i , giving the scheduler both a faster job and information about where the job belongs. Or it may leave p_i hidden until the optimized job finishes.

These choices give the following four models. The table is a summary rather than four new sets of notation; in every row, the preliminary action in the third column costs one unit of machine time.

model	without a test	what the unit action does	where discussed
obligatory testing	cannot be run	reveals p_i and unlocks the job	Intro 1.1; Sections 3 to 5
blind execution	runs for its hidden length p_i	reveals p_i before execution	Intro 1.4; Section 6
blind optimization	runs for raw time u	replaces u by a still-hidden $p_i \leq u$	Intro 1.5; Section 7
revealing optimization	runs for raw time u	reveals p_i and replaces u by p_i	Intro 1.6; Sections 8 to 10

The last three rows separate the two benefits of the unit preliminary action. Blind execution studies information without speedup. Blind optimization studies speedup without

advance information. Revealing optimization contains both benefits. This model was introduced by Dürr et al. [DEMM20]. Blind execution is an adversarial-bag counterpart of stochastic scheduling with testing [LMS19]; its closest earlier adversarial relative is the binary processing-time oracle of Dufossé et al. [DDN⁺22]. Blind optimization is closest to the non-revealing optimization studied by Damerius et al. [DKL⁺23], although their optimization consumes a separate budget rather than machine time.

The three optional models have different optimal policies and different bag-specific benchmarks. We therefore state their results separately. The algorithms nevertheless share one simple design principle: privately shuffle the jobs, use a sublinear sample to estimate the bag, and optimize the rest of the schedule as if that estimate were exact.

1.4 Information without speedup: blind execution

Suppose an untested job can simply be run until it finishes. Its running time is its true hidden length p_i . Testing does not make the job shorter; it only tells us p_i before we decide when to run it. We call this *blind execution*.

Without an upper bound on the job lengths, no randomized algorithm has a finite competitive ratio, even if we ignore an additive $o(n^2)$ term; see Theorem 6.2. For every fixed bound L , however, sampling again gives one algorithm that is optimal on every bag.

Theorem 1.3 (Blind-execution instance optimality, informal version of Theorem 6.1). *Fix $L < \infty$. There is one randomized algorithm $\mathcal{A}^*$ such that, for every bag $M \subseteq [0, L]$ and every online algorithm $\mathcal{A}'$,*

$$\text{cost}(\mathcal{A}^*) \leq \text{cost}(\mathcal{A}') + o_L(n^2).$$

Both costs are measured on M and include the algorithms' randomness. The competing algorithm $\mathcal{A}'$ may even be designed specifically for M and told the entire bag in advance.

To describe the algorithm, let X be the job length of a random job in the bag and put $\mu = \mathbb{E}X$. Testing is initially worthwhile when the threshold τ determined by

$$\mathbb{E}(\tau - X)^+ = 1$$

is smaller than μ . Even then, it need not be optimal to test the entire bag. The leading cost of testing a fraction q is an explicit quadratic $F_X(q)$, given in Equation (148), and the algorithm chooses its minimum over $0 \leq q \leq 1$.

Algorithm 3 Instance-optimal blind execution

- 1: Shuffle the jobs, run $\lceil \sqrt{n} \rceil$ of them blindly, and use their observed lengths to estimate the random variable X .
 - 2: Treat the estimate as exact and put $\mu = \mathbb{E}X$.
 - 3: Find τ from $\mathbb{E}(\tau - X)^+ = 1$.
 - 4: If $\tau \geq \mu$, put $q = 0$. Otherwise choose $q \in \arg \min_{0 \leq z \leq 1} F_X(z)$.
 - 5: Test a random q -fraction of the remaining jobs. Whenever a test reveals $p < \tau$, process that job immediately; defer all other outcomes.
 - 6: After the tests stop, process the deferred jobs with $\tau \leq p < \mu$, shortest first.
 - 7: Run every untouched job blindly.
 - 8: Finish the remaining tested jobs shortest first.
-

The block between testing and blind execution is important. A job can be too long to interrupt the test phase and still be shorter than an untouched job on average. Such a job waits until testing stops and is then processed before the blind block.

1.5 Blind optimization

The two remaining models start from the same operation. Every job takes a known raw time u , and one unit of optimization reduces its running time to $p_i \leq u$. They differ only in whether optimization reveals p_i . We start with the case in which it does not. Before execution, all optimized jobs still look the same.

This is the simplest case. Its deterministic version has no useful learning mechanism: an adversarial labeling can make any fixed sample unrepresentative. In the randomized model, a private sample estimates the one number that matters, the mean μ of the optimized lengths. The scheduler then compares the two average times per job, u and $1 + \mu$, and uses the better action for all remaining jobs. We give the formal discussion in Section 7.

Theorem 1.4 (Blind-optimization instance optimality, informal version of Theorem 7.1). *Fix $u < \infty$. There is one randomized algorithm $\mathcal{A}^*$ such that, for every bag $M \subseteq [0, u]$ and every online algorithm $\mathcal{A}'$,*

$$\text{cost}(\mathcal{A}^*) \leq \text{cost}(\mathcal{A}') + o_u(n^2).$$

Both costs are measured on M and include the algorithms' randomness. The competing algorithm $\mathcal{A}'$ may even be designed specifically for M and told the entire bag in advance.

More precisely, if μ_M is the mean optimized length in M , the best possible leading value is

$$\frac{n^2}{2} \min\{u, 1 + \mu_M\}.$$

Algorithm 4 Instance-optimal blind optimization

- 1: Shuffle the jobs, optimize and run $\lceil n^{2/3} \rceil$ of them, and let $\hat{\mu}$ be the mean observed length.
 - 2: Treat $\hat{\mu}$ as the true mean of the bag.
 - 3: If $1 + \hat{\mu} < u$, optimize and then run every remaining job.
 - 4: Otherwise run every remaining job raw for time u .
-




Figure 1: Randomized ratios for blind and revealing optimization.

Knowing the instance-optimal algorithm does not yet give its competitive ratio: we must still find the worst bag. The resulting ratio is unbounded and grows on the order of $\sqrt{u}$.

Theorem 1.5 (Blind-optimization curve, informal version of Theorem 7.2). *The exact randomized asymptotic competitive ratio is 1 for $u \leq 1$ and is given by the explicit two-branch curve in Figure 1 for $u > 1$. It grows as $\sqrt{u}/2 + 1/2 + o(1)$ as $u \rightarrow \infty$.*

1.6 Revealing optimization

Now assume that optimization also reveals p_i . Thus the unit action buys both information and speed. We call this *revealing optimization*.

The two endpoints are easy to understand. If $u \leq 1$, the test alone costs at least as much as raw execution, so we should never test. If u is very large, raw execution is unattractive, so the problem approaches obligatory testing. The interesting question is what happens between these endpoints.

This model was introduced by Dürr et al. [DEMM20]. Their results already showed that intermediate values of u are the difficult ones. We determine the exact answer for every u , for both deterministic and randomized algorithms.

A deterministic algorithm uses three basic shapes.

Algorithm 5 Deterministic revealing-optimization policies, schematically

- 1: For small u , run every job raw.
 - 2: For intermediate u , test every job and use a fixed threshold.
 - 3: For large u , use the history-dependent obligatory-testing threshold.
 - 4: Whenever jobs are deferred, finish them shortest first.
-

For randomized algorithms we can say more. On each fixed bag, the sampling principle from blind execution finds the best threshold and the best tested fraction q . Unlike blind execution, every tested processing time is at most the raw time u , so all tested jobs belong before the untouched raw block.

Algorithm 6 Instance-optimal revealing-optimization policy

- 1: Shuffle the jobs, test $\lceil \sqrt{n} \rceil$ of them, and use the observed lengths to estimate the bag.
 - 2: Compute the best threshold and the best tested fraction q .
 - 3: Test a random q -fraction; finish outcomes below the threshold at once.
 - 4: Finish all other tested jobs, shortest first.
 - 5: Run every untouched job raw for time u .
-

Theorem 1.6 (Revealing-optimization instance optimality, informal version of Theorem 9.1). *Fix $u < \infty$. There is one randomized algorithm $\mathcal{A}^*$ such that, for every bag $M \subseteq [0, u]$ and every online algorithm $\mathcal{A}'$,*

$$\text{cost}(\mathcal{A}^*) \leq \text{cost}(\mathcal{A}') + o_u(n^2).$$

Both costs are measured on M and include the algorithms' randomness. The competing algorithm $\mathcal{A}'$ may even be designed specifically for M and told the entire bag in advance. The best leading cost is $n^2 \Phi_{\text{RO},u}(M)$, where $\Phi_{\text{RO},u}$ is obtained by minimizing an explicit quadratic over the tested fraction q .

For each bag, this policy attains its best online leading cost. We obtain the competitive ratio by asking which bag maximizes the ratio between that cost and the clairvoyant optimum. The worst bag can be reduced to two job lengths, leaving a one-variable optimization.

Theorem 1.7 (Revealing-optimization curves, informal version of Theorems 8.1 and 10.1). *For every $u > 0$, the exact deterministic and randomized asymptotic competitive ratios are the six-piece and four-piece curves in Figure 2. The deterministic curve reaches 1.87... and then falls to the obligatory value 1.57.... The randomized curve reaches 1.63... and eventually becomes 4/3.*




Figure 2: The exact asymptotic ratios as a function of the raw execution time u . Testing is pointless on the left. On the right, both curves reach their obligatory-testing values.

The curves are not monotone. This is not a contradiction: increasing u changes both players' options. For intermediate u , the clairvoyant scheduler makes especially good use of the choice between raw execution and testing. When u becomes large, both schedulers effectively test everything.

The hard bags change along the curves. In the first regimes, mixtures of zero jobs and jobs of length u are enough. Later, the deterministic lower bound becomes a capped form of the decreasing-level construction from obligatory testing. The exact formulas and matching constructions are in Sections 8 and 10.

1.7 Techniques

The proofs repeatedly use three pictures.

The area of unfinished jobs. During one unit of machine work, the objective increases by the number of jobs that are still unfinished. Thus the sum of completion times is the area under the curve “number of unfinished jobs versus time.” This picture is often easier to reason about than individual completion times.

Jobs completed per unit of work. A round of tests together with the short jobs completed during that round can be viewed as one large action. It spends some amount of time and completes some fraction of the jobs. Higher-density actions—those completing more jobs per unit of work—should come first. Sorting these actions produces the thresholds, the ordered blocks, and the one-variable quadratics in the instance-optimal results.

A private shuffle. The universal algorithms start by shuffling the labels and inspecting a small prefix. The same idea is needed in the lower bound, where the policy may adaptively decide which label to touch next. If equal job lengths are temporarily distinguished as separate tokens, the order in which any deterministic policy first reveals these tokens is itself just a permutation of them. The revealed sequence is therefore sampling without replacement. One concentration event can then control every possible stopping point of the policy.

The deterministic proofs use a complementary idea. Their lower bounds make the algorithm choose between completing the current jobs and waiting for shorter future jobs. Their upper bounds maintain a threshold and charge each wrongly ordered pair to a small potential. The exact constants appear when the two possible losses are equal.

1.8 Related work

The closest papers have already appeared next to the models they introduced. We give a broader comparison here.

Scheduling with tests belongs to optimization with *explorable uncertainty*, where uncertain data can be queried at a cost; the query viewpoint goes back at least to Kahan [Kah91]. Levi, Magnanti, and Shaposhnik study a stochastic scheduling model and derive prefix-testing policies followed by an order based on expected processing time per unit weight [LMS19, LMS24]. Our four-block blind-execution policy has a similar shape, but our input is an adversarial bag, need not converge to a distribution, and is not announced to the universal algorithm.

Dürr et al. [DEMM20] introduced the adversarial revealing-optimization model and obtained deterministic bounds, including the $1.87\dots$ guarantee for binary values. We close the deterministic gap for arbitrary values $p_i \in [0, u]$ and give the exact randomized curve against an oblivious adversary. Related work allows nonuniform test times [AE21, LLWZ23, KLL⁺26], multiple machines [GCH24], or weights [BS26].

Dogéas, Erlebach, and Liang [DEL24] introduced obligatory testing for total completion time. For uniform tests they proved the SIDLE upper bound $1.58\dots$ and the deterministic lower bound $\sqrt{2}$. Liang and Liang [LL26] later raised the lower bound to $3/2$. Our matching value is $1.57\dots$, and our randomized result gives the oblivious value $4/3$. More strongly, its stationary threshold is asymptotically optimal separately for every bounded bag.

The processing-time oracle of Dufossé et al. [DDN⁺22] is close to blind execution, but allows only two possible hidden durations. If no tests are available and processing times are revealed only at completion, one obtains the classical nonclairvoyant model of Motwani, Phillips, and Torng [MPT94]. These semantics differ from raw execution in revealing optimization, which always lasts the declared time u .

The rest of the paper gives precise versions and proofs. We first fix the models and offline benchmarks in Section 2. Each main section then starts with the theorem it proves and a short overview of the proof. The machine-checked part of the argument is summarized in Appendix A.

2 Models and common preliminaries

This section makes the four models from the introduction precise and records the common facts used throughout the paper. There are n jobs, all available at time zero, and one machine. Operations are nonpreemptive, and at most one operation can run at a time. Job j has a hidden value $p_j \geq 0$. The four models differ in what can be done before this value is known.

Obligatory testing OT.

The values p_j are arbitrary finite nonnegative numbers. Every job must first be tested for one unit. The test reveals p_j , after which its processing operation of length p_j may be performed immediately or delayed.

Blind execution BE(L).

Here $p_j \in [0, L]$. A job may be run blindly for its actual duration p_j ; the value becomes known only when that execution finishes. Alternatively, the job may be tested for one unit, revealing p_j , and its processing operation may then be delayed. Testing provides information but does not shorten the job.

Blind optimization $\text{BO}(u)$.

Here $p_j \in [0, u]$. A job may be run raw for the known time u . Alternatively, one unit of optimization changes its running time from u to p_j without revealing p_j ; the optimized job is subsequently run blindly, possibly after a delay. Its value becomes observable only when that execution finishes. Raw execution reveals no hidden value.

Revealing optimization $\text{RO}(u)$.

Again $p_j \in [0, u]$, and raw execution takes the known time u . The other option is a unit test that reveals p_j , followed, possibly after a delay, by processing for p_j . Thus the preliminary action provides both the speedup and its size.

The different letters L and u are intentional. In blind execution, L is only a known bound used by the uniform instance-optimal theorem. In the last two models, u is itself an available running time and hence part of every scheduling decision. The number n and the relevant parameter L or u are known to the algorithm.

The completion time C_j is the time at which job j finishes. If a test reveals $p_j = 0$, the job completes at the end of that test. A blind zero job in $\text{BE}(L)$ completes as soon as it is selected, whereas an optimized zero job in $\text{BO}(u)$ first pays its unit optimization cost. The objective is

$$\sum_{j=1}^n C_j.$$

Obligatory testing can be viewed as the limit of revealing optimization in which raw execution becomes unavailable. We use this only as intuition: the obligatory theorems and their algorithms are proved directly, without interchanging the limits $u \rightarrow \infty$ and $n \rightarrow \infty$.

We write $\text{cost}(\mathcal{A}, I)$ for the objective value of an online algorithm $\mathcal{A}$ on an instance I , and $\text{cost}(\text{OPT}, I)$ for the value of an offline optimum that knows all processing times. In calculations we use the shorter abbreviations

$$\text{ALG}_{\mathcal{A}}(I) = \text{cost}(\mathcal{A}, I), \quad \text{OPT}(I) = \text{cost}(\text{OPT}, I),$$

and omit algorithms or instances when they are clear. Thus OPT denotes an algorithm inside $\text{cost}(\text{OPT}, I)$ and its cost only in the abbreviated notation $\text{OPT}(I)$.

2.1 Information and adversaries

The deterministic revealing-optimization results allow the usual adversarial input and use adaptive revelation only as a lower-bound device that is frozen by Lemma 2.2. The randomized results use an *oblivious* adversary: the labeled processing-time vector is fixed before the algorithm draws its private random seed. Unless explicitly stated otherwise, the algorithm is not told the multiset of values. An *announced* policy is additionally given that multiset but not its assignment to labels. Announcing the multiset can only help and is therefore a valid strengthening in lower bounds.

All randomized expectations are over private algorithmic randomness. A uniform random relabeling is private randomness, not a distributional assumption on the adversarial multiset. For lower bounds we use a uniform distribution over fixed labelings as a finite Yao distribution and then select one labeling whose cost is at least the average.

2.2 Offline optima and pair accounting

The offline optimum knows all values p_j . It therefore chooses the shorter available way to complete each job and orders the resulting job blocks shortest first. For later use, define

$$q_u(p) := \min\{u, 1 + p\} \quad (0 < u < \infty), \quad (1)$$

and associate with job j the model-dependent *effective length*

$$q_j := \begin{cases} 1 + p_j, & \text{OT}, \\ p_j, & \text{BE}(L), \\ q_u(p_j), & \text{BO}(u) \text{ or } \text{RO}(u). \end{cases} \quad (2)$$

Lemma 2.1 (Unified offline order). *Fix any one of the four models and let $q_{(1)} \leq \dots \leq q_{(n)}$ be its effective lengths from Equation (2). A clairvoyant optimum completes the jobs in this order and has value*

$$\text{OPT} = \sum_{k=1}^n \sum_{i=1}^k q_{(i)}, \quad (3)$$

$$= \sum_i q_i + \sum_{i < j} \min\{q_i, q_j\}. \quad (4)$$

In OT it tests every job. In BE(L) it runs every job blindly. In BO(u) and RO(u) it uses raw mode when $u \leq 1 + p_j$ and the unit preliminary action followed by processing otherwise.

Proof. Let $C_{[1]} \leq \dots \leq C_{[n]}$ be the ordered completion times of any feasible schedule. Completing job j consumes at least q_j units of work belonging to that job. Hence, by time $C_{[k]}$, the machine has performed the required work of some k jobs, and

$$C_{[k]} \geq \sum_{i=1}^k q_{(i)}.$$

Choose the mode specified in the lemma, keep each resulting block contiguous, and execute the blocks in nondecreasing order of q_j . This schedule is feasible in the relevant model and attains equality for every k . Summing over k and then collecting the contribution of each unordered pair gives the two formulas. $\square$

The revealing-optimization and blind-optimization models have the same offline effective lengths. Their difference is entirely online: a test reveals p_j , whereas optimization does not. Blind execution is different even offline, because a clairvoyant scheduler can run a job for p_j without paying the unit test. Thus

$$\text{OPT}_{\text{BE}}(p) = \sum_i p_i + \sum_{i < j} \min\{p_i, p_j\}. \quad (5)$$

This is not the principal benchmark in Theorem 6.1. The clairvoyant value may be $o(n^2)$, whereas that theorem compares policies at the n^2 scale with the optimal announced-multiset online value $\Phi(D_M)$.

For obligatory testing, Equations (3) and (4) specialize to

$$\text{OPT}_{\text{OT}} = \sum_{k=1}^n \left(k + \sum_{i=1}^k p_{(i)} \right), \quad (6)$$

$$= \sum_i (1 + p_i) + \sum_{i < j} (1 + \min\{p_i, p_j\}). \quad (7)$$

In particular,

$$\text{OPT}_{\text{OT}} \geq \frac{n(n+1)}{2}. \quad (8)$$

For $\text{BO}(u)$ and $\text{RO}(u)$, $q_j \geq \min\{u, 1\}$ gives

$$\text{OPT}_u \geq \min\{u, 1\} \frac{n(n+1)}{2}. \quad (9)$$

There is deliberately no analogous positive lower bound for blind execution: a bag of zero jobs has clairvoyant cost zero.

2.3 Competitive-ratio conventions

Every asymptotic statement in this paper allows a uniform additive $o(n^2)$ term. Formally, a policy family $\mathcal{A}$ has asymptotic competitive ratio at most R if there is a deterministic sequence $\varepsilon(n) \rightarrow 0$ such that every n -job instance satisfies

$$\mathbb{E} \text{ALG}_{\mathcal{A}}(I) \leq R \text{OPT}(I) + \varepsilon(n)n^2. \quad (10)$$

For a deterministic policy the expectation is omitted. The optimal ratio in a model is the infimum of such R over all admissible policy families.

For obligatory testing and for either model with fixed raw time $u > 0$, Equations (8) and (9) show that $\text{OPT} = \Omega_u(n^2)$. In these models the definition above is equivalent to the usual worst-case ratio after taking $n \rightarrow \infty$. This equivalence fails for blind execution, where OPT can be $o(n^2)$. Accordingly, Theorem 6.1 makes the absolute statement $\mathbb{E} \text{ALG} = n^2\Phi(D_M) + o_L(n^2)$ and does not claim a bounded ratio to the clairvoyant optimum on every sparse bag.

2.4 Test-order adversaries

The lower bound will assign processing times according to the order in which tests finish. This is only a proof device.

Lemma 2.2 (Replay). *Fix a deterministic online algorithm. Suppose all untested jobs have the same known parameters, and an adversary assigns processing times deterministically from the transcript and test-completion order. The resulting interaction is reproduced by a fixed input instance.*

Proof. Run the interaction once and record the value revealed for every tested job label. Assign those values to the corresponding labels in advance. An induction over operation completions shows that the deterministic algorithm chooses the same tests, processing operations, and idle intervals on the fixed instance. If the algorithm fails to test or complete some job forever, its competitive ratio is already infinite; otherwise every relevant label is recorded. $\square$

3 Deterministic obligatory testing: the exact ratio

The goal of this section is to prove Theorem 3.1, the rigorous deterministic part of the informal obligatory-testing result Theorem 1.1 from the introduction. Let $z_\star > 1$ be the unique solution of

$$z_\star - 3 = \log z_\star, \quad (11)$$

and put

$$R_\star = 1 + \frac{2}{z_\star - 1} = 1.570574096649395 \dots \quad (12)$$

Theorem 3.1 (Deterministic obligatory testing). *For every deterministic online algorithm $\mathcal{B}$ and every $\varepsilon > 0$, there are arbitrarily large obligatory-testing instances I such that*

$$\text{cost}(\mathcal{B}, I) \geq (R_\star - \varepsilon) \text{cost}(\text{OPT}, I).$$

Conversely, there exist a deterministic algorithm $\mathcal{A}$ and an absolute constant C such that every n -job instance satisfies

$$\text{cost}(\mathcal{A}, I) \leq R_\star \text{cost}(\text{OPT}, I) + Cn.$$

Consequently the optimal deterministic size-asymptotic competitive ratio is exactly $R_\star$.

The proof has two independent halves. For the lower bound we reveal many equal-sized groups of positive jobs from longest to shortest and hide a large group of zero jobs until the end. Harmonic spacing makes the leading lower bound independent of the algorithm's choices. For the upper bound, ADAPTIVETHRESHOLD compares each revealed job with a threshold computed from the current history. A four-endpoint reduction and a nonnegative bank account certify its cost. These are also the two pieces reused in the finite- u analysis. The first subsection proves the lower half of the theorem; the second proves the upper half.

3.1 Harmonic core and matching lower bound

This subsection proves the lower bound in one self-contained calculation. Its notation is local to this subsection. The construction has many equally large groups of positive jobs, revealed from longest to shortest, followed by one large group of zero jobs. The only design choice is the gap between two consecutive positive processing times. Choosing these gaps harmonically will make all decisions of the online algorithm disappear from the leading term.

Theorem 3.2 (Lower bound). *For every deterministic online algorithm $\mathcal{B}$ and every $\varepsilon > 0$, there are arbitrarily large fixed instances I with unit tests and rational processing times such that*

$$\text{ALG}_{\mathcal{B}}(I) \geq (R_\star - \varepsilon) \text{OPT}(I).$$

The revelation experiment. Fix an integer K , a rational number $\xi > 0$, and a rational slack $\gamma \in (0, 1)$; one may keep $\gamma = 1/2$ throughout. For a large scaling integer H , the instance will contain H jobs of each of K positive types $0, 1, \dots, K-1$ and ξH zero jobs. We choose H to clear all denominators.

The adversary assigns processing times according to the order in which tests are started. The first H tested jobs receive processing time p_0 , the next H receive p_1 , and so on; after the K positive blocks, every remaining job receives processing time zero. Thus

$$p_0 > p_1 > \dots > p_{K-1} > 1,$$

so the algorithm sees the positive types from longest to shortest and learns about the zero jobs only at the end.

This adaptive description is only a convenient way to construct the input. After running the interaction against a fixed deterministic algorithm, assign to every job label the value revealed when that label was tested. Replaying the algorithm on this fixed assignment produces exactly the same transcript. If the algorithm never tests or completes some job, its cost is infinite and the lower bound is immediate; otherwise every job label is recorded. Hence it is enough to lower-bound the cost in the revelation experiment.

Why each phase can be relaxed to shortest processing time. Let T_ℓ be the start of the first test in positive block ℓ , and let T_K be the start of the first test in the zero block. Phase ℓ contains the completions in $(T_\ell, T_{\ell+1}]$; a completion exactly at T_ℓ belongs to the preceding phase. Each T_ℓ is the boundary between two nonpreemptive operations. Consequently, a previously tested job that is unfinished at T_ℓ has not started processing and still needs its entire processing time.

We repeatedly use the following elementary observation. Suppose jobs with remaining work $v_1, \dots, v_m$ complete after some time T . If their completion order is $q_1, \dots, q_m$, then the r th relative completion time is at least $\sum_{h \leq r} v_{q_h}$. Summing these prefix inequalities shows that their total relative completion time is minimized by arranging the v_i in nondecreasing order. This is merely a lower bound; the online algorithm need not actually use that order.

For normalized group masses $q_1, \dots, q_m$ and remaining lengths $v_1 \leq \dots \leq v_m$, this shortest-first bound has leading coefficient

$$\mathcal{S}((q_g, v_g)_{g=1}^m) = \sum_{g=1}^m \left(\frac{v_g q_g^2}{2} + q_g \sum_{h < g} v_h q_h \right). \quad (13)$$

More precisely, the corresponding contribution for groups of size $q_g H$ is $H^2 \mathcal{S} + O(H)$. The quadratic term is simply the familiar sum $v_g(1 + \dots + q_g H)$ inside one group; the cross term is the waiting caused by all shorter groups.

We ensure that

$$p_0 - p_{K-1} < 1. \quad (14)$$

In phase ℓ , an unfinished job of an older type $j < \ell$ therefore has remaining length p_j , whereas a type- ℓ job completed in its own phase needs both its test and processing, of total length $1 + p_\ell$. Their order in the shortest-first relaxation is

$$p_{\ell-1} < p_{\ell-2} < \dots < p_0 < 1 + p_\ell.$$

At T_K , an untested zero job has remaining length 1, so the relaxed tail starts with all zero jobs and then uses the positive types from shortest to longest. These are the only two places where Equation (14) is used.

The finite accounting identity. We now record the online algorithm's possible behavior. Normalize all job counts by H . Let

- $s_j \in [0, 1]$ be the mass of type- j jobs completed in their own phase;
- $d_{j,\ell} \geq 0$ be the mass of type- j jobs completed in the later phase $\ell > j$; and
-

$$X_{j,\ell} = s_j + \sum_{r=j+1}^{\ell-1} d_{j,r} \leq 1 \quad (\ell > j)$$

be the mass of type j completed before phase ℓ . Thus $X_{j,\ell+1}$ is the mass completed by the end of phase ℓ .

At time T_ℓ , the machine has already performed the tests of the ℓH jobs in earlier positive blocks and has processed every earlier job that has completed. Thus

$$\frac{T_\ell}{H} \geq \ell + \sum_{j < \ell} p_j X_{j,\ell}. \quad (15)$$

For each phase, charge its completions their common start time from Equation (15), and charge their relative completion times using Equation (13). Do the same at T_K for the remaining positive jobs and the ξH zero jobs. Expanding and collecting equal variables gives

$$\begin{aligned} \frac{\text{ALG}}{H^2} &\geq C_K + \sum_{j=0}^{K-1} \left(\frac{s_j^2}{2} - \theta_j s_j \right) \\ &\quad + \sum_{j < \ell} d_{j,\ell} \left[\ell - j - \theta_j - \sum_{j < k < \ell} (p_j - p_k) X_{k,\ell+1} \right] - O(1/H), \end{aligned} \quad (16)$$

where K, ξ, γ are fixed while $H \rightarrow \infty$, and

$$\theta_j = 1 - \xi(p_j - 1) - \sum_{k > j} (p_j - p_k - 1), \quad (17)$$

$$C_K = \frac{\xi^2}{2} + 2\xi K + K^2 + P_K, \quad P_K = \sum_{j=0}^{K-1} \left(j + \frac{1}{2} \right) p_j. \quad (18)$$

For completeness, here is how to audit Equation (16) without guessing its origin. In phase ℓ , the masses in the shortest-first relaxation are

$$d_{\ell-1,\ell}, d_{\ell-2,\ell}, \dots, d_{0,\ell}, s_\ell$$

with respective lengths

$$p_{\ell-1}, p_{\ell-2}, \dots, p_0, 1 + p_\ell.$$

Multiply their total mass by the right-hand side of Equation (15) and add the functional Equation (13). In the final phase use masses

$$\xi, 1 - X_{K-1,K}, \dots, 1 - X_{0,K}$$

and lengths $1, p_{K-1}, \dots, p_0$. The constant terms sum to C_K . The terms containing only s_j are $s_j^2/2 - \theta_j s_j$; the coefficient of each $d_{j,\ell}$ is precisely the bracket in Equation (16). Thus the displayed identity is just the phasewise shortest-first bound with the phase start times retained. No optimality assumption about the online schedule is hidden in it.

The meaning of the expression is useful. Completing an s_j fraction of a type in its own phase creates a quadratic completion cost but saves later waiting cost, producing $s_j^2/2 - \theta_j s_j$. A completion postponed to a later positive phase is represented by $d_{j,\ell}$ and its bracket. We next choose the processing times so that every such bracket is nonnegative and all θ_j are equal.

The harmonic choice and its telescoping cancellation. Set

$$p_{K-1} = 1 + \frac{\gamma}{\xi}, \quad p_i = p_{i+1} + \frac{1}{\xi + K - 1 - i} \quad (0 \leq i < K - 1). \quad (19)$$

In other words, measured upward from the shortest type, the consecutive gaps are

$$\frac{1}{\xi + 1}, \frac{1}{\xi + 2}, \dots, \frac{1}{\xi + K - 1}.$$

This is the key design choice. If $m = K - 1 - j$, then

$$p_j - 1 = \frac{\gamma}{\xi} + \sum_{r=1}^m \frac{1}{\xi + r},$$

$$\sum_{k>j} (p_j - p_k) = \sum_{r=1}^m \frac{r}{\xi + r}.$$

Adding the first line multiplied by ξ to the second line makes each denominator cancel:

$$\xi(p_j - 1) + \sum_{k>j} (p_j - p_k) = \gamma + \sum_{r=1}^m \frac{\xi + r}{\xi + r} = \gamma + m. \quad (20)$$

Substitution in Equation (17) gives the same value for every type,

$$\theta_j = 1 - \gamma. \quad (21)$$

This is why the increments in Equation (19) are harmonic: the factor ξ coming from the zero block and the multiplicity r coming from the r shorter positive types add up to the denominator $\xi + r$.

The delayed-completion terms are now harmless as well. Since $X_{k,\ell+1} \leq 1$, their brackets are at least

$$\begin{aligned} \ell - j - (1 - \gamma) - \sum_{j < k < \ell} (p_j - p_k) \\ = \gamma + \sum_{j < k < \ell} (1 - (p_j - p_k)) > 0, \end{aligned}$$

where the final inequality follows from Equation (14). We may therefore drop every term containing $d_{j,\ell}$. The remaining minimization is not a multivariable optimization problem: complete one square for each j ,

$$\frac{s_j^2}{2} - (1 - \gamma)s_j = \frac{1}{2}(s_j - (1 - \gamma))^2 - \frac{1}{2}(1 - \gamma)^2.$$

Because $1 - \gamma \in (0, 1)$, this minimum is feasible for $s_j \in [0, 1]$. Hence every online algorithm obeys

$$\liminf_{H \rightarrow \infty} \frac{\text{ALG}}{H^2} \geq Q_K := C_K - \frac{K}{2}(1 - \gamma)^2. \quad (22)$$

It remains to compute the offline benchmark. Knowing all processing times, the optimum runs the zero jobs first and the positive types from shortest to longest. Equivalently, the pairwise contribution of two jobs is $1 + \min\{p_i, p_j\}$. Its leading coefficient is

$$D_K = \frac{\xi^2}{2} + \xi K + \frac{K^2}{2} + P_K, \quad \lim_{H \rightarrow \infty} \frac{\text{OPT}}{H^2} = D_K. \quad (23)$$

The four terms respectively account for zero-zero pairs, zero-positive pairs, the unit-test part of positive-positive pairs, and the processing part of positive-positive pairs. Thus the finite- K lower-bound certificate is simply Q_K/D_K .

Passing to a smooth curve. Write

$$\alpha = \frac{K}{\xi}$$

for the ratio of positive mass to zero mass, and keep α fixed while $K \rightarrow \infty$. We restrict to $0 < \alpha < e - 1$. Then

$$p_0 - p_{K-1} = \sum_{r=1}^{K-1} \frac{1}{\xi + r} < \log\left(1 + \frac{K}{\xi}\right) = \log(1 + \alpha) < 1,$$

which verifies Equation (14).

The processing contribution P_K has the exact form

$$P_K = \frac{K^2}{2} \left(1 + \frac{\gamma}{\xi}\right) + \frac{1}{2} \sum_{h=1}^{K-1} \frac{(K-h)^2}{\xi+h}. \quad (24)$$

Indeed, the common baseline $1 + \gamma/\xi$ contributes the first term, and swapping the order of summation shows that the increment $1/(\xi+h)$ receives total weight $(K-h)^2/2$. After division by ξ^2 , the sum becomes a Riemann integral:

$$\begin{aligned} \frac{P_K}{\xi^2} &\longrightarrow I(\alpha) = \frac{\alpha^2}{2} + \frac{1}{2} \int_0^\alpha \frac{(\alpha-t)^2}{1+t} dt \\ &= \int_0^\alpha (\alpha-t)(1+\log(1+t)) dt \\ &= \frac{1}{2}(1+\alpha)^2 \log(1+\alpha) - \frac{\alpha}{2} - \frac{\alpha^2}{4}. \end{aligned} \quad (25)$$

The second integral form follows either by integration by parts or by summing the limiting processing-time curve $1+\log(1+t)$ against the remaining positive mass $\alpha-t$. The subtraction in Equation (22) is only $O(K)$, whereas $C_K, D_K = \Theta(K^2)$. Since $C_K - D_K = \xi K + K^2/2$, we obtain

$$R(\alpha) := \lim_{K \rightarrow \infty} \frac{Q_K}{D_K} = 1 + \frac{\alpha + \alpha^2/2}{\frac{1}{2} + \frac{\alpha}{2} + \frac{\alpha^2}{4} + \frac{1}{2}(1+\alpha)^2 \log(1+\alpha)}. \quad (26)$$

This last one-variable maximization is where the constant in the theorem comes from. Put $z = (1+\alpha)^2$. Then

$$R(z) = 1 + \frac{2(z-1)}{1+z+z \log z}. \quad (27)$$

The derivative of the fraction has the sign of

$$3 - z + \log z.$$

For $z > 1$ this expression is strictly decreasing, is positive at $z = 1$, and tends to $-\infty$. Hence it has one zero $z_\star$, characterized by

$$z_\star - 3 = \log z_\star.$$

The derivative is positive before $z_\star$ and negative afterward, so this is the global maximum. At the root,

$$1 + z_\star + z_\star \log z_\star = (z_\star - 1)^2,$$

and therefore

$$R(z_\star) = 1 + \frac{2}{z_\star - 1} = R_\star.$$

Moreover, $\sqrt{z_\star} - 1 = 1.12255\dots < e - 1$, so the maximizer lies in the range where Equation (14) holds.

Proof of Theorem 3.2. Choose a rational α sufficiently close to $\alpha_\star = \sqrt{z_\star} - 1$, and choose any rational $\gamma \in (0, 1)$. For a sufficiently large K , set $\xi = K/\alpha$ and use Equation (19); then $Q_K/D_K > R_\star - \varepsilon/2$. All parameters are rational. Let H tend to infinity through multiples of their common denominators. Equations (22) and (23) make the actual ratio exceed $R_\star - \varepsilon$ for all sufficiently large H . Finally, the replay argument at the beginning of the proof converts the revelation experiment into a fixed instance for the given deterministic algorithm. $\square$

Lemma 3.3 (Scaled harmonic core). *Fix a total normalized mass $s > 0$ and a parameter $t \in [1, e)$. Against every deterministic obligatory-testing algorithm, there are scales $M_\nu \rightarrow \infty$ and finite rational multilevel instances with $sM_\nu + o(M_\nu)$ jobs such that*

$$\frac{\text{OPT}}{M_\nu^2} \rightarrow O_0 = \frac{s^2}{4}(1 + t^{-2} + 2 \log t), \quad \liminf_{\nu \rightarrow \infty} \frac{\text{ALG}}{M_\nu^2} \geq A_0^{\text{lb}} := O_0 + \frac{s^2}{2}(1 - t^{-2}). \quad (28)$$

Their normalized total positive processing mass converges to

$$\frac{1}{M_\nu} \sum_{j: p_j > 0} p_j \rightarrow L = s \log t, \quad (29)$$

and their largest processing time converges to $1 + \log t < 2$. The adaptive revelation is converted by replay into a fixed instance.

Proof. The case $t = 1$ is obtained by continuity from instances with vanishing positive mass, so suppose $1 < t < e$. In the construction above set $\alpha = t - 1 = K/\xi$. Before rescaling, the total normalized mass is $K + \xi = \xi t$. Give every unit of normalized mass the common factor $s/(\xi t)$.

The phase accounting, harmonic cancellation, and completion of squares in Equations (16), (20) and (22) are homogeneous of degree two in the job masses. The Riemann-sum limit Equation (25) substituted into Equation (23) therefore gives

$$O_0 = \frac{s^2}{4}(1 + t^{-2} + 2 \log t).$$

Likewise $C_K - D_K = \xi K + K^2/2$, while the $O(K)$ square-completion loss in Equation (22) disappears after division by the quadratic scale. After the same rescaling this gives the guaranteed lower benchmark

$$A_0^{\text{lb}} - O_0 = \frac{s^2}{2}(1 - t^{-2}).$$

Finally, summing the limiting processing-time curve $1 + \log(1 + x)$ over the positive mass gives $L = s \log t$, and Equation (19) gives $p_0 \rightarrow 1 + \log t$. Rational approximation, clearing denominators, and the replay argument used in the theorem produce a diagonal sequence of arbitrarily large fixed rational instances with the stated limit and liminf bounds. $\square$

3.2 Upper bound: a self-contained proof

We prove the following statement from scratch in this subsection.

Theorem 3.4 (Upper bound). *Let $z > 1$ be the solution of $z - 3 = \log z$, and set*

$$R = 1 + \rho, \quad \rho = \frac{2}{z - 1} = 0.5705740966 \dots$$

There is a deterministic online algorithm and an absolute constant C such that every instance with n jobs satisfies

$$\text{ALG} \leq R \text{OPT} + Cn.$$

The proof has three ingredients. First, pairwise accounting gives an exact formula for the cost of a schedule. Second, a simple convexity argument says that it is enough to consider four boundary outcomes for every test. Third, we give a nonnegative *bank balance* W : whenever a boundary outcome creates positive excess cost, the balance drops by at least that amount, up to a constant rounding loss. Thus the proof below uses derivatives only as shorthand for the first-order change of this bank balance; no optimization background is needed.

The algorithm. For a parameter $c > 0$, the formal procedure $\text{ADAPTIVETHRESHOLD}(c)$ is given in algorithm 7. Immediately before a test, let

- x be the number of jobs not yet tested, including the job about to be tested;
- N be the number of earlier tests that revealed a strictly positive processing time; and
- D be the number of these N jobs that the algorithm deferred.

Define the normalized state

$$y = \frac{D - RN}{x}. \quad (30)$$

Because $D \leq N$ and $R > 1$, we always have $y \leq 0$. For later reuse define the parameterized threshold

$$a_c(y) = \begin{cases} 1, & y \leq -1, \\ 1 + \frac{1}{2} \log\left(\frac{c+2}{c-2y}\right), & -1 \leq y \leq 0. \end{cases} \quad (31)$$

The theorem uses $c = \rho$; throughout this subsection we abbreviate $A(y) = a_\rho(y)$ and $R = 1 + \rho$. The defining equation for $z = (\rho + 2)/\rho$ says

$$\log \frac{\rho + 2}{\rho} = \frac{2}{\rho} - 2. \quad (32)$$

Consequently, A is continuous at -1 , increases from $A(-1) = 1$ to $A(0) = 1/\rho = 1.7526207478\dots$, and

$$A'(y) = \frac{1}{\rho - 2y} \quad (-1 < y < 0). \quad (33)$$

Algorithm 7 $\text{ADAPTIVETHRESHOLD}(J, c)$

Input: jobs J and parameter $c > 0$.

- 1: Fix an arbitrary test order $j_1, \dots, j_n$.
 - 2: Set $N \leftarrow 0$, $D \leftarrow 0$, and $Q \leftarrow \emptyset$.
 - 3: **For** $i = 1, \dots, n$ **do**:
 - 4: Set $x \leftarrow n - i + 1$ and $y \leftarrow (D - (1 + c)N)/x$.
 - 5: Set $a \leftarrow a_c(y)$ according to Equation (31).
 - 6: Test j_i for one unit and observe p_{j_i} .
 - 7: **If** $p_{j_i} \leq a$, process j_i immediately.
 - 8: **Otherwise**, add j_i to Q and set $D \leftarrow D + 1$.
 - 9: **If** $p_{j_i} > 0$, set $N \leftarrow N + 1$.
 - 10: Process the jobs of Q in nondecreasing order of their revealed processing times.
-

For $c = \rho$, the state in the pseudocode is exactly (30). A genuine zero increments neither counter and completes at the end of its test. The procedure never uses raw execution; the same pseudocode will therefore apply to the large finite- u regimes.

The numerator $D - RN$ in (30) is best viewed as a balance, not as an estimate of a processing time. Every revealed positive job reduces the balance by R ; deferring it returns one unit, because every later unit test will also delay that deferred job. Dividing by x spreads this balance over the tests that remain. A state close to 0 can afford a threshold larger than one. Once the normalized balance falls below -1 , the proof can only support the conservative threshold one, so A becomes flat.

We will later derive (31), rather than asking the reader to guess why this particular logarithm appears.

Pairwise accounting. For two jobs $i < j$ in test order, charge to their unordered pair the duration of operations of either job performed while the other job is unfinished. If job i is processed immediately, this charge in the algorithm is $1 + p_i$. If i is deferred and j is immediate, it is $2 + p_j$. If both are deferred, it is $2 + \min\{p_i, p_j\}$. Hence

$$\text{ALG} = \sum_i (1 + p_i) + \sum_{i < j} g_{ij}, \quad g_{ij} = \begin{cases} 1 + p_i, & i \text{ immediate,} \\ 2 + p_j, & i \text{ deferred, } j \text{ immediate,} \\ 2 + \min\{p_i, p_j\}, & i, j \text{ deferred.} \end{cases} \quad (34)$$

The offline optimum tests and immediately processes jobs in nondecreasing order of p_i , and therefore

$$\text{OPT} = \sum_i (1 + p_i) + \sum_{i < j} (1 + \min\{p_i, p_j\}). \quad (35)$$

For completeness, the latter claim follows because the k th completed job cannot finish before the machine has performed k tests and the processing of the k shortest jobs. The shortest-first schedule attains all these lower bounds simultaneously.

We bound the excess

$$F = \text{ALG} - R \text{OPT}.$$

Why four boundary outcomes suffice. Fix only the symbolic decisions made along a run: for each job, specify whether its value is zero, positive and immediate, or positive and deferred. This fixes all later counters and thresholds. The allowed values of a coordinate are then, respectively,

$$\{0\}, \quad (0, A(y)], \quad (A(y), \infty).$$

If all processing times except p_i are fixed inside such a symbolic run, (34)–(35) express F as an affine function of p_i plus negative multiples of $\min\{p_i, h\}$. Since $p \mapsto \min\{p, h\}$ is concave, F is convex in p_i . A convex function on $(0, A(y)]$ has its supremum at one of the two ends. On $(A(y), \infty)$, the eventual slope is $1 - R = -\rho < 0$; convexity then implies that every earlier slope is also negative, so the supremum is at the lower end.

It is therefore enough to consider the following four formal outcomes.

type	limiting value	algorithmic decision	counter update
Z	0	immediate	none
E	0^+	immediate	$N \leftarrow N + 1$
I	$A(y)$	immediate	$N \leftarrow N + 1$
Q	$A(y)^+$	deferred	$N \leftarrow N + 1, D \leftarrow D + 1$

Here 0^+ is not a new processing time. It means that a positive immediate value tends to zero *after its symbolic decision has been fixed*. Although its processing contribution vanishes, it still increments N and therefore changes all future thresholds. This is why 0 and 0^+ cannot be identified. Similarly, $A(y)^+$ approaches the threshold from the deferred side. Applying the one-coordinate argument successively shows that a uniform bound on all formal four-type words implies the same bound on all genuine instances. Because F is a finite sum of affine and minimum terms on a fixed symbolic path, the simultaneous one-sided limit at every formal vertex exists and is independent of the rates at which its open coordinates approach their endpoints.

One more elementary property will simplify every minimum in the pairwise formulas. Put $u = D - RN \leq 0$. After an outcome of type Z, E or I, or Q, respectively, the pair

(u, x) changes as follows:

$$(u, x - 1), \quad (u - R, x - 1), \quad (u - R, x - 1), \quad (u - \rho, x - 1).$$

Whenever $x \geq 2$, the new quotient is at most u/x . Thus the states, and hence the thresholds, seen by successive tests never increase. When $x = 1$ there is no next test and therefore no next threshold to compare. The nonzero boundary values $A(y)$ consequently arrive in nonincreasing order. In particular, the minimum of the current nonzero boundary value and any earlier one is always the current value.

The proof state and the local excess. The algorithm needs only (x, N, D) . The analysis splits N into two kinds. Among earlier formal outcomes, let

- S be the number of *substantive* positive outcomes I, Q;
- e be the number of infinitesimal outcomes E; and
- d be the number of deferred outcomes Q.

Thus $N = S + e$ and $D = d$. Define

$$\eta = \frac{d - RS}{x}, \quad b = \frac{e}{x}, \quad y = \eta - Rb. \quad (36)$$

We have $d \leq S$, hence $\eta \leq 0$. The variable b records exactly the part of the algorithmic state that is invisible to limiting processing-cost minima: an infinitesimal value affects N , but its minimum with every other processing time tends to zero.

Consider the current boundary outcome at a pre-state (x, S, e, d) , and write $a = A(y)$. Pairwise accounting gives

$$F = -\rho \frac{n(n+1)}{2} + \sum_i \tilde{\ell}_i, \quad (37)$$

where the exact contribution of the current outcome is

$$\tilde{\ell}_Z = \tilde{\ell}_E = d, \quad (38)$$

$$\tilde{\ell}_I = d + a[x + d - RS - R], \quad (39)$$

$$\tilde{\ell}_Q = d + a[d - RS - \rho]. \quad (40)$$

Here is a quick derivation. The current unit test gives one extra unit of algorithmic pair cost for each earlier deferred job, producing d . Its own test and its pairs with all earlier jobs give a universal excess $-\rho(i+1)$ at step i ; summing this term produces the first term in (37). For an I outcome, the current processing operation contributes $a(x+d)$ to the algorithm and $Ra(S+1)$ to the scaled optimum. For a Q outcome, the corresponding quantities are $a(d+1)$ and $Ra(S+1)$. Monotonicity of the thresholds justifies using the current a in every earlier processing-time minimum.

For the potential calculation it is convenient to discard the two favorable diagonal terms $-Ra$ and $-\rho a$. Define the relaxed local rewards

$$\ell_Z = \ell_E = d, \quad (41)$$

$$\ell_I = d + a(x + d - RS) = d + ax(1 + \eta), \quad (42)$$

$$\ell_Q = d + a(d - RS) = d + ax\eta. \quad (43)$$

Then $\tilde{\ell}_c \leq \ell_c$ for all four types, and

$$F \leq -\rho \frac{n(n+1)}{2} + \sum_i \ell_i. \quad (44)$$

How the logarithmic threshold is found. Before giving the full bank balance, it is useful to see where the formula comes from. Ignore 0^+ outcomes for a moment, so $b = 0$, and look in the range $-1 \leq y \leq 0$. Suppose the nonlinear part of the balance has the form $x^2 f(y)$. Asking the immediate boundary I and the deferred boundary Q to be exactly paid by the first-order decrease of this balance gives

$$-2f + (y - R)f' + A(1 + y) = 0, \quad (45)$$

$$1 - 2f + (y - \rho)f' + Ay = 0. \quad (46)$$

Subtracting the equations gives $f' = A - 1$. Differentiating either equation then gives

$$1 + (2y - \rho)A'(y) = 0.$$

With the natural joining condition $A(-1) = 1$, this is exactly (31)–(33).

The universal negative term in (44) has leading part $-\rho n^2/2$. The initial balance will be $n^2 f(0)$, so exact cancellation requires $f(0) = \rho/2$. Since

$$f(y) = \int_{-1}^y (A(t) - 1) dt, \quad (47)$$

direct integration gives

$$f(0) = \frac{1}{2} - \frac{\rho}{4} \log\left(\frac{\rho + 2}{\rho}\right).$$

Thus $f(0) = \rho/2$ is equivalent to (32), or, after setting $z = (\rho + 2)/\rho$, to $z - 3 = \log z$. Thus both the threshold and the constant R are forced by two local balance equations and the initial accounting; they are not the output of an unexplained numerical optimization.

The bank balance. For $-1 \leq y \leq 0$, put

$$f(y) = \int_{-1}^y (A(t) - 1) dt, \quad h(y) = A(y)(1 + y). \quad (48)$$

The term xd in the balance below pays for the fact that each of the x remaining tests will delay each of the d already deferred jobs. The $x^2 f(y)$ term handles substantive outcomes. Finally, the terms involving b repair the discontinuity between a genuine zero and 0^+ .

Define

$$W(x, S, e, d) = xd + x^2 G(y, b), \quad (49)$$

using two formulas for G :

$$G(y, b) = \begin{cases} f(y) + bh(y) + \frac{R}{2}b^2, & y \geq -1, \\ g(\eta), & g(\eta) = \frac{(1 + \eta)_+^2}{2R}, \quad y \leq -1, \end{cases} \quad \eta = y + Rb. \quad (50)$$

The first formula is used while the threshold is changing. The second is used once the threshold has saturated at one. In that flat region the limiting processing-cost terms depend on the substantive balance η , not on the number of 0^+ outcomes separately, which explains why g contains no b .

This piecewise definition is smooth enough for a first-order argument. At $y = -1$, feasibility gives $0 \leq b \leq 1/R$ and $\eta = -1 + Rb$. Since

$$f(-1) = h(-1) = 0, \quad h'(-1) = 1,$$

the first formula in (50) has value $Rb^2/2$ and derivatives $G_y = b$, $G_b = Rb$. The second formula has exactly the same value and derivatives. This also checks the derivative in the raw x coordinate: at fixed (S, e, d) ,

$$\partial_x [x^2 G(y, b)] = x(2G - yG_y - bG_b),$$

which agrees on the two sides because G , G_y , and G_b agree. The remaining raw derivatives are linear combinations of G_y and G_b . The internal join of $(1 + \eta)_+^2$ at $\eta = -1$ also has a continuous first derivative. Hence W is continuously differentiable. Moreover, $W \geq 0$: in the first region $f \geq 0$, $h \geq 0$, and $b, d \geq 0$; in the second this is immediate from (50).

The choice $h = A(1 + y)$ also has a direct interpretation. At $b = 0$, it makes the first infinitesimal outcome 0^+ exactly neutral in the local balance. The quadratic correction $Rb^2/2$ then maintains the balance for all b and is precisely what makes the two regions meet smoothly.

Four local balance checks. We now verify that every formal outcome can be paid from the bank. In raw coordinate order (x, S, e, d) , the four unit-step directions are

$$v_Z = (-1, 0, 0, 0), \quad v_E = (-1, 0, 1, 0), \quad v_I = (-1, 1, 0, 0), \quad v_Q = (-1, 1, 0, 1). \quad (51)$$

For an endpoint type $q \in \{Z, E, I, Q\}$, define its first-order balance

$$J_q = \ell_q + \nabla W \cdot v_q. \quad (52)$$

The inequality $J_q \leq 0$ simply says: local reward plus first-order change in the bank balance is nonpositive.

We need only a few one-variable identities. On $-1 \leq y \leq 0$,

$$f' = A - 1, \quad f(-1) = 0, \quad f(0) = \frac{\rho}{2}, \quad (53)$$

$$-2f + (y - R)f' + A(1 + y) = 0, \quad (54)$$

$$1 - 2f + (y - \rho)f' + Ay = 0, \quad (55)$$

$$-2f + yf' = A(\rho - y) - R \leq 0, \quad (56)$$

$$h + (\rho - y)h' = AR + (\rho - y)(1 + y)A' \geq AR. \quad (57)$$

The first line follows from the definition and (32). For (54)–(55), differentiate the left-hand sides and use $A' = 1/(\rho - 2y)$; both derivatives vanish, and both expressions equal zero at $y = -1$. Equation (56) follows from (54). Its right-hand side is zero at -1 and decreases afterward, because

$$\frac{d}{dy} [A(y)(\rho - y)] = \frac{\rho - y}{\rho - 2y} - A(y) \leq 0.$$

Indeed, $y \leq 0$ gives $(\rho - y)/(\rho - 2y) \leq 1$, while $A(y) \geq 1$. Finally, (57) is obtained by differentiating $h = A(1 + y)$. Notice also that $h, h' \geq 0$.

In the changing-threshold region $y \geq -1$, direct differentiation of (49) first gives, for an arbitrary $G(y, b)$,

$$J_Z/x = -2G + yG_y + bG_b, \quad (58)$$

$$J_E/x = -2G + (y - R)G_y + (b + 1)G_b, \quad (59)$$

$$J_I/x = -2G + (y - R)G_y + bG_b + A(1 + y + Rb), \quad (60)$$

$$J_Q/x = 1 - 2G + (y - \rho)G_y + bG_b + A(y + Rb). \quad (61)$$

Substituting $G = f + bh + Rb^2/2$ and using (54)–(55) reduces the four expressions to

$$J_Z/x = (-2f + yf') + b(-h + yh'), \quad (62)$$

$$J_E/x = b[-h + (y - R)h' + R], \quad (63)$$

$$J_I/x = b[-h + (y - R)h' + AR], \quad (64)$$

$$J_Q/x = b[-h + (y - \rho)h' + AR]. \quad (65)$$

Now the signs are transparent. The Z line is nonpositive by (56), $h, h' \geq 0$, and $y \leq 0$. The Q line is nonpositive by (57). The I line is no larger because $y - R < y - \rho$ and $h' \geq 0$; the E line is no larger again because $A \geq 1$. Thus all four checks hold in the changing-threshold region.

In the flat region $y \leq -1$, we have $A = 1$ and $G = g(\eta)$. The four checks become

$$J_Z/x = J_E/x = -2g + \eta g', \quad (66)$$

$$J_I/x = -2g + (\eta - R)g' + 1 + \eta, \quad (67)$$

$$J_Q/x = 1 - 2g + (\eta - \rho)g' + \eta. \quad (68)$$

If $\eta \leq -1$, then $g = g' = 0$, and these expressions are 0, $1 + \eta$, and $1 + \eta$. If $-1 \leq \eta \leq 0$, then $g' = (1 + \eta)/R$, and they are, respectively,

$$-\frac{1 + \eta}{R}, \quad -\frac{1 + \eta}{R}, \quad 0.$$

They are therefore nonpositive in both cases. We have proved

$$J_Z, J_E, J_I, J_Q \leq 0 \quad (69)$$

at every feasible state.

From infinitesimal checks to unit steps. Equation (69) compares a local reward with the *derivative* of W , whereas one job changes the integer counters by one. This loses only a constant per job. More precisely, there is an absolute constant C_0 such that every admissible transition satisfies

$$W(s + v_q) - W(s) \leq \nabla W(s) \cdot v_q + C_0. \quad (70)$$

We include the bounded-curvature argument because it is what makes the $O(n)$ term uniform over all instances.

Suppose first that $x \geq 2$ and linearly interpolate the unit transition. The numerators of both y and η have the form $u - \kappa t$, where $u \leq 0$ and $\kappa \geq 0$, while the denominator is $x - t$. Their derivatives are therefore nonpositive, so the segment crosses each joining surface at most once. In the changing-threshold region, $y \geq -1$ implies

$$R(S + e) - d = \rho S + (S - d) + Re \leq x.$$

Consequently,

$$0 \leq \frac{S}{x} \leq \frac{1}{\rho}, \quad 0 \leq \frac{e}{x} \leq \frac{1}{R}, \quad 0 \leq \frac{d}{x} \leq \frac{1}{\rho}.$$

After normalizing by x , all reachable states in this region thus lie in a fixed compact set. The balance W is homogeneous of degree two, so its second derivatives in the raw

coordinates are homogeneous of degree zero and are uniformly bounded on this normalized set.

In the flat region the formula is even simpler:

$$W = xd + \frac{(x + d - RS)_+^2}{2R}. \quad (71)$$

Its gradient is globally Lipschitz. Because the two regional formulas have the same first derivative at their interface, an interpolated unit step has a uniformly bounded second derivative except at finitely many harmless joining points. More explicitly, let $\phi(t) = W(s + tv_q)$. The preceding bounds and the C^1 gluing make ϕ' Lipschitz with one uniform constant L along the whole segment, so

$$|\phi(1) - \phi(0) - \phi'(0)| \leq \int_0^1 |\phi'(t) - \phi'(0)| dt \leq \int_0^1 Lt dt = \frac{L}{2}.$$

This proves (70) for $x \geq 2$. When $x = 1$, the changing-threshold pre-states lie in the same compact normalized set, while (71) applies globally in the flat region. At $x = 0$ we define $W = 0$; this agrees with (71), since terminal feasibility gives $d \leq S$ and hence $d - RS \leq 0$.

Combining (52)–(70) gives the actual one-step bank inequality

$$\ell_i + W(s_{i+1}) - W(s_i) \leq C_0. \quad (72)$$

Telescoping the account. Initially $S = e = d = 0$, $x = n$, and $y = 0$. Therefore

$$W(s_0) = n^2 f(0) = \frac{\rho}{2} n^2. \quad (73)$$

The terminal balance is zero. Summing (72) over all n jobs yields

$$\sum_i \ell_i \leq \frac{\rho}{2} n^2 + C_0 n.$$

Substitution into (44) gives

$$\text{ALG} - R \text{OPT} = F \leq -\rho \frac{n(n+1)}{2} + \frac{\rho}{2} n^2 + C_0 n = \left(C_0 - \frac{\rho}{2}\right) n.$$

The constant is independent of the four-type word. Taking $C = \max\{0, C_0 - \rho/2\}$, the boundary reduction therefore extends the same estimate to every vector of nonnegative processing times, proving Theorem 3.4.

Finally, $\text{OPT} \geq n(n+1)/2$, so the additive $O(n)$ term is lower order and

$$\frac{\text{ALG}}{\text{OPT}} \leq R + O(1/n).$$

This proves the desired size-asymptotic upper bound. Notice exactly where the remaining finite-instance loss enters: it is only the uniform Taylor remainder in (70) (together with the deliberately discarded favorable diagonal terms), not any unproved limiting step in the endpoint reduction. In particular, this argument proves a linear additive term; it does not prove one fixed additive constant at the endpoint R .

4 Randomized obligatory testing: the sharp ratio is $4/3$

We keep the obligatory-testing model and now allow private randomization. The adversary fixes the labeled vector $p \in [0, \infty)^n$ before the algorithm draws its random bits. The multiset is not announced, and no upper bound on the processing times is assumed. The goal of this section is to prove Theorem 4.1, the rigorous randomized part of the informal obligatory-testing result Theorem 1.1 from the introduction.

Theorem 4.1 (Randomized obligatory testing). *There is a universal randomized nonanticipating algorithm $\mathcal{A}_n$, depending only on n , such that every fixed vector $p \in [0, \infty)^n$ satisfies*

$$\mathbb{E} \text{cost}(\mathcal{A}_n, p) \leq \frac{4}{3} \text{cost}(\text{OPT}, p) + 20378n^{7/4} \leq \left(\frac{4}{3} + 40756n^{-1/4} \right) \text{cost}(\text{OPT}, p). \quad (74)$$

Conversely, for every randomized algorithm $\mathcal{A}$ and every sufficiently large even n , some fixed labeling of $n/2$ zero jobs and $n/2$ jobs of processing time two satisfies

$$\mathbb{E} \text{cost}(\mathcal{A}, p) \geq n^2 - O(n^{3/2}), \quad \text{cost}(\text{OPT}, p) = \frac{3}{4}n^2 + n. \quad (75)$$

Hence the exact randomized size-asymptotic competitive ratio against an oblivious adversary is $4/3$.

For an announced multiset, the algorithmic core is the processing-time prefix with maximum completions per unit discovery work. Testing jobs in a uniformly random order gives an exact pair identity and a $4/3$ algebraic certificate. The universal algorithm learns an approximately separated prefix from a sublinear sample; a fixed cutoff and a fallback schedule protect it from unbounded values missed by the sample. For the lower bound, we randomly label half zero and half length-two jobs. A balanced-bridge discrepancy bound limits how fast any algorithm can complete them. The next subsection begins the formal proof.

4.1 The normalized offline objective

For a fixed vector p , put

$$K(p) = \frac{1}{n^2} \sum_{i,j} \min\{p_i, p_j\}, \quad O(p) = \frac{1 + K(p)}{2}, \quad (76)$$

$$D(p) = \frac{n + \sum_i p_i}{2}. \quad (77)$$

The pair identity Equation (7) is equivalently

$$\text{OPT}(p) = n^2 O(p) + D(p). \quad (78)$$

The linear correction $D(p)$ will be kept rather than hidden: this makes the analysis valid for unbounded processing times.

4.2 Stationary prefixes

A set E is a *processing-time prefix* if every value in E is at most every value outside it. We require that E contain every zero job; otherwise a supposedly deferred zero completes at its test and the following identity is false. Write $L = [n] \setminus E$ and

$$e = |E|, \quad a = \frac{e}{n}, \quad h = 1 - a, \quad m = \frac{1}{n} \sum_{i \in E} p_i, \quad K_X = \frac{1}{n^2} \sum_{i,j \in X} \min\{p_i, p_j\}. \quad (79)$$

Every early-late pair has minimum equal to its early value, so

$$K(p) = K_E + 2hm + K_L. \quad (80)$$

Define

$$P(p, E) = (1 + m) \left(1 - \frac{a}{2}\right) + \frac{K_L}{2}. \quad (81)$$

The stationary policy for E tests all jobs in a uniformly random order, processes a positive early job immediately after its test, and defers every late job. After the final test it processes the late jobs in SPT order.

Lemma 4.2 (Exact stationary cost). *For every processing-time prefix E containing all zero jobs,*

$$\mathbb{E} \text{ALG}_{\text{stat}}(p, E) = n^2 P(p, E) + \frac{e + \sum_i p_i}{2}. \quad (82)$$

Proof. Let

$$W = n + \sum_{i \in E} p_i, \quad L_E = e + \sum_{i \in E} p_i.$$

The discovery phase consists of n blocks: an early block has length $1 + p_i$ and completes its job, while a late block is its unit test and completes no job. For an early job, each other block precedes it with probability $1/2$. Its expected completion time is $W/2 + (1 + p_i)/2$, and the total early contribution is

$$\frac{eW + L_E}{2}. \quad (83)$$

Every late job sees the common offset W . Its tail contributes

$$(n - e)W + \sum_{i \in L} p_i + \sum_{\substack{i < j \\ i, j \in L}} \min\{p_i, p_j\}. \quad (84)$$

Adding Equations (83) and (84) and collecting diagonal terms gives Equation (82). $\square$

For a nonempty prefix define its completion density by

$$\rho(E) = \frac{a}{1 + m}. \quad (85)$$

Choose a maximum-density prefix and put

$$\theta = \frac{1 + m}{a} = \frac{1}{\rho(E)}. \quad (86)$$

All zeros belong to every maximizer: adding a zero increases the numerator without increasing the denominator.

Lemma 4.3 (Density threshold). *There is a maximum-density prefix for which every early value is at most θ and every late value is at least θ . Values equal to θ may be assigned to either side.*

Proof. Removing boundary mass δ at value x changes the density to $(a - \delta)/(1 + m - x\delta)$. Cross multiplication shows that this is larger exactly when $x > \theta$, so no selected boundary value can exceed θ . The symmetric addition argument excludes an unselected value below θ . Equality preserves the density. $\square$

The next identity is the algebraic source of the constant $4/3$.

Lemma 4.4 (Exact and robust certificates). *For every ordered prefix split with $a > 0$ and $\theta = (1 + m)/a$,*

$$2\theta(4O - 3P) = (\theta - 2)^2 + 4(\theta K_E - m^2) + \theta(K_L - \theta h^2). \quad (87)$$

Consequently:

- (i) *if early values are at most θ and late values are at least θ , then $P \leq 4O/3$;*
- (ii) *if early values are at most $\theta + s$ and late values are at least $\theta - s$, where $s \geq 0$, then*

$$P \leq \frac{4}{3}O + \frac{2}{3}s. \quad (88)$$

Proof. Expansion using $m = a\theta - 1$, $h = 1 - a$, and Equation (80) gives Equation (87). Under exact separation, $xy \leq \theta \min\{x, y\}$ for early x, y , hence $m^2 \leq \theta K_E$; for late x, y , one has $\min\{x, y\} \geq \theta$, hence $K_L \geq \theta h^2$. This proves (i).

Under the relaxed inequalities,

$$\theta K_E - m^2 \geq -sK_E, \quad K_L - \theta h^2 \geq -sh^2.$$

Moreover $K_E \leq am \leq a^2\theta$. Substitution into Equation (87) yields

$$P \leq \frac{4}{3}O + \frac{2}{3}sa^2 + \frac{1}{6}sh^2 \leq \frac{4}{3}O + \frac{2}{3}s,$$

because $(2/3)a^2 + (1/6)(1 - a)^2 \leq 2/3$ on $[0, 1]$. $\square$

Combining the exact certificate with Lemma 4.2 and Equation (78) already gives a finite statement useful below.

Corollary 4.5 (Announced multiset). *If the multiset is announced, the stationary policy for a maximum-density prefix satisfies*

$$\mathbb{E} \text{ALG}_{\text{stat}}(p, E) \leq \frac{4}{3} \text{OPT}(p).$$

Indeed, the correction in Equation (82) is at most $D(p)$. Equality in the quadratic coefficient of Equation (87) forces $\theta = 2$, zero mass $1/2$, and all positive mass at two. This predicts the lower-bound instance before any probabilistic argument is used.

We also use the following tail-safe estimate. Direct expansion gives

$$P - O = \frac{h + am - K_E}{2}. \quad (89)$$

If all early values are at most $R \geq 1$, then $m \leq aR$ and therefore

$$P(p, E) \leq O(p) + \frac{R}{2}. \quad (90)$$

4.3 Learning the prefix from a sublinear sample

Fix the universal cutoff

$$B = 32. \quad (91)$$

For an integer d , let $\eta = B/d$. Keep zero as a separate category, partition $(0, B]$ into the consecutive bins $((b-1)\eta, b\eta]$, and add one overflow category for values above B . A finite positive value is rounded upward to the endpoint $q_b = b\eta$.

Let S be a uniformly random subset of k labels. If μ_b and $\hat{\mu}_b$ are the population and sample fractions of category b , put

$$\Delta = \sum_b |\hat{\mu}_b - \mu_b|. \quad (92)$$

Lemma 4.6 (Histogram error). *For $1 \leq k < n$,*

$$\mathbb{E}\Delta \leq \sqrt{\frac{d+2}{k}}.$$

Proof. The count in category b is hypergeometric and its normalized variance is at most μ_b/k . Thus $\mathbb{E}|\hat{\mu}_b - \mu_b| \leq \sqrt{\mu_b/k}$. Summation and Cauchy–Schwarz give the claim. $\square$

For a finite category prefix J , define

$$\hat{a}(J) = \sum_{b \in J} \hat{\mu}_b, \quad \hat{m}(J) = \sum_{b \in J} q_b \hat{\mu}_b, \quad \hat{\rho}(J) = \frac{\hat{a}(J)}{1 + \hat{m}(J)}.$$

Let $\hat{\rho}$ be the maximum and $\hat{\theta} = 1/\hat{\rho}$. The learned early set is not an arbitrary maximizing category prefix. It is its *threshold closure*

$$\hat{J} = \{\text{finite categories } b : q_b \leq \hat{\theta}\}. \quad (93)$$

Lemma 4.7 (Threshold closure). *The prefix Equation (93) has the same maximum sample density. Every category in it has endpoint at most $\hat{\theta}$, and every finite category outside it has endpoint greater than $\hat{\theta}$, including categories with zero sample count.*

Proof. The removal/addition calculation in Lemma 4.3 applies to every positive-mass sample category. Empty categories do not change the sample statistics, and adding mass at equality preserves the identity $1 + \hat{m} = \hat{a}\hat{\theta}$. Closing the prefix therefore preserves optimality and makes the two pointwise assertions tautological. $\square$

For large n , set

$$k = \lfloor n^{3/4} \rfloor, \quad d = \lfloor n^{1/4} \rfloor, \quad \eta = B/d. \quad (94)$$

The algorithm draws a uniform permutation and first tests its first k jobs, leaving positive sample jobs pending. It computes the closed sample prefix. If

$$\hat{\rho} < \frac{2}{B}, \quad (95)$$

it enters *fallback mode*: test every remaining job and then process all positive jobs in SPT order. Otherwise it enters *learned mode*: process positive sampled early jobs in SPT order; test the remaining jobs in their random order, processing early outcomes immediately; and finish the deferred tail in SPT order. Overflow is never early.

4.4 Good samples, fallback, and bad-event excess

Put

$$\gamma = \frac{1}{B(B+1)}, \quad G = \{\Delta \leq \gamma\}. \quad (96)$$

Markov’s inequality and Lemma 4.6 give

$$\mathbb{P}(G^c) \leq B(B+1) \sqrt{\frac{d+2}{k}}. \quad (97)$$

Assume G and learned mode. Let E be the population set selected by the closed category prefix, and let $a, m, \theta = (1+m)/a$ be its true statistics. Since $\hat{\theta} \leq B/2$ and $\hat{a} = (1+\hat{m})/\hat{\theta} \geq 2/B$, we have $a \geq 1/B$. For this fixed category set,

$$|a - \hat{a}| \leq \Delta, \quad |m - \hat{m}| \leq B\Delta + \eta.$$

Consequently

$$|\theta - \hat{\theta}| \leq \frac{3}{2}B^2\Delta + B\eta. \quad (98)$$

Every early actual value is at most $\hat{\theta}$, while every late value is greater than $\hat{\theta} - \eta$. Thus the robust certificate applies with

$$s = \frac{3}{2}B^2\Delta + B\eta + \eta. \quad (99)$$

The sample-first order is not exactly the fully stationary comparator, but the difference is only lower order. Conditional on the unordered sample, write e_S, e_R for the early counts and

$$W_S = k + \sum_{i \in E \cap S} p_i, \quad W_R = n - k + \sum_{i \in E \setminus S} p_i.$$

Comparing the two independent random internal orders with one fully random order, the early-cost difference is

$$\frac{e_R W_S - e_S W_R}{2} \leq \frac{1}{2}nk(1 + B/2).$$

Postponing positive sampled early jobs until all sample tests finish adds at most $k^2(1 + B/2)$. The late tail is identical because every zero is early. Therefore, on a good learned sample,

$$\mathbb{E}[\text{ALG} \mid S] \leq \frac{4}{3}\text{OPT} + n^2 \left[B^2\Delta + \frac{2}{3}(B+1)\eta \right] + (1 + B/2)(nk/2 + k^2). \quad (100)$$

On a good fallback sample, comparison of sample and population densities shows that the true maximum-density reciprocal satisfies $\theta_* \geq B/4 = 8$. The exact threshold inequalities imply

$$K(p) \geq \frac{(\theta_* - 1)^2}{\theta_*}, \quad O(p) \geq \frac{\theta_* - 1 + 1/\theta_*}{2} \geq \frac{57}{16}. \quad (101)$$

For every test order, test-all-then-SPT satisfies

$$\text{ALG}_{\text{fb}} \leq \text{OPT} + \frac{n(n-1)}{2}. \quad (102)$$

Indeed, if there are z zeros, their total displacement from the first z SPT positions is at most $zn - z(z+1)/2$, while delaying the positive SPT tail adds $(n-z)(n-z-1)/2$; the sum is $n(n-1)/2$. Combining Equations (101) and (102) gives $\text{ALG}_{\text{fb}} < 4\text{OPT}/3$.

Because processing times are unbounded, a bad-event estimate must control excess over $4\text{OPT}/3$, not raw cost. In learned mode every early value is at most $B/2$, so Equation (90) bounds the stationary excess by $(B/4)n^2$; in fallback mode Equation (102) bounds it by $n^2/2$. Averaging Equation (100) and these bad-event bounds, and using Equation (97), yields the explicit uniform estimate

$$\mathbb{E} \text{ALG} \leq \frac{4}{3}\text{OPT} + n^2 \left[9472\sqrt{\frac{d+2}{k}} + \frac{704}{d} \right] + 17(nk/2 + k^2). \quad (103)$$

For $n \geq 12^4$, the floors in Equation (94) give $\sqrt{(d+2)/k} \leq 2n^{-1/4}$, $1/d \leq 2n^{-1/4}$, and $k \leq n^{3/4}$. The right-hand side of Equation (103) is at most

$$\frac{4}{3}\text{OPT} + 20378n^{7/4}. \quad (104)$$

For smaller n , test-all-then-SPT is covered by the same deliberately loose constant. Finally $\text{OPT} \geq n^2/2$ proves the multiplicative form in Equation (74).

4.5 The balanced binary lower bound

Let $n = 2m$ and assign m zeros and m twos uniformly to the labels. Fix a deterministic nonanticipating algorithm. In its adaptive order of new tests, let $X_r = 1$ for a zero and $X_r = -1$ for a two, and $S_k = \sum_{r \leq k} X_r$. Conditional on every history, the untested labels remain a uniform subset, so (S_k) is a uniform balanced sign bridge even though the test order is adaptive. Put

$$\Delta_0 = \frac{1}{2} \max_{0 \leq k \leq n} |S_k|. \quad (105)$$

Lemma 4.8 (Bridge discrepancy).

$$\mathbb{E} \Delta_0 \leq \frac{\sqrt{2}n}{\sqrt{n-1}}.$$

Proof. For $0 \leq k \leq m$, $M_k = S_k/(n-k)$ is a martingale. Doob's L^2 inequality and the hypergeometric variance $\mathbb{E} S_m^2 = n^2/[4(n-1)]$ give

$$\mathbb{E} \max_{k \leq m} |S_k|^2 \leq \frac{4n^2}{n-1}.$$

Time reversal gives the same bound for the second half, and Cauchy–Schwarz then yields $\mathbb{E} \max_k |S_k| \leq 2\sqrt{2}n/\sqrt{n-1}$. $\square$

At physical time t , suppose the algorithm has completed k tests and x processing operations of two-jobs. At most $k/2 + \Delta_0 + x$ jobs are complete, while $k + 2x \leq t$. Pathwise,

$$\# \text{completed by } t \leq t/2 + \Delta_0.$$

The area under the unfinished-job curve therefore gives

$$\text{ALG} \geq \int_0^{2(n-\Delta_0)} (n - \Delta_0 - t/2) dt = (n - \Delta_0)^2. \quad (106)$$

By Lemma 4.8, $\mathbb{E} \text{ALG} \geq n^2 - O(n^{3/2})$. Every input in the distribution has

$$\text{OPT} = 3m^2 + 2m = \frac{3}{4}n^2 + n.$$

Conditioning a randomized algorithm on its private seed and averaging over the finite placement distribution shows that one fixed labeling attains at least this average. That labeling is chosen before the private seed and is therefore an oblivious input. Together with Equation (104), this completes the proof of Theorem 4.1.

5 Obligatory-testing instance optimality

The goal of this section is to prove Theorem 5.1, the rigorous version of the “one algorithm for every bag” statement Theorem 1.2. The algorithm is not a new one: it is a slight variant of the policy from Section 4, obtained by replacing its fixed cutoff 32 by a slowly growing cutoff. The new point is that this variant attains, on every bounded bag, the best leading cost available even to a policy that is told the bag in advance.

For a multiset $M = \{p_1, \dots, p_n\}$, write D_M for its empirical probability measure. We define below an explicit number $\Phi_{\text{OT}}(D_M)$. It is obtained by scanning the processing times in increasing order, choosing the prefix with the largest number of completions per unit of discovery work, and evaluating the stationary policy for that prefix.

Theorem 5.1 (Obligatory-testing instance optimality). *There is a single randomized nonanticipating policy $\mathcal{A}^*$, depending only on n and not on the multiset, with the following property. For every fixed $L < \infty$ there is a deterministic sequence $\varepsilon_L(n) \rightarrow 0$ such that, for every size- n multiset $M \subseteq [0, L]$ and every fixed labeling σ ,*

$$\mathbb{E} \text{cost}(\mathcal{A}^*, \sigma(M)) \leq n^2 \Phi_{\text{OT}}(D_M) + \varepsilon_L(n) n^2. \quad (107)$$

Conversely, for every randomized policy $\mathcal{A}$, even one told M in advance, some fixed oblivious labeling σ satisfies

$$\mathbb{E} \text{cost}(\mathcal{A}, \sigma(M)) \geq n^2 \Phi_{\text{OT}}(D_M) - \varepsilon_L(n) n^2. \quad (108)$$

One may take $\varepsilon_L(n) = O_L(n^{-1/5})$. Thus $\mathcal{A}^$ is asymptotically instance-optimal, uniformly over all bounded bags.*

The upper estimate will have order $O_L(n^{-1/5})$. The announced lower bound is sharper, of order $O_L(n^{-1/2} \sqrt{\log(n+2)})$; we use the slower common rate only to keep the theorem statement symmetric.

Theorem 5.2 (When instance optimality is impossible). (i) *There is no deterministic instance-optimal policy, even when all job lengths lie in $[0, 2]$. More precisely, for every deterministic policy $\mathcal{A}$, even one told the bag, and every even n , there is a fixed labeling of the bag with $n/2$ zero jobs and $n/2$ length-two jobs for which*

$$\text{cost}(\mathcal{A}) \geq \frac{9}{8} n^2. \quad (109)$$

On the same bag, a randomized policy $\mathcal{A}'$ that is told the bag has

$$\text{cost}(\mathcal{A}') = n^2 + \frac{3}{4} n.$$

Thus $\text{cost}(\mathcal{A}) > \text{cost}(\mathcal{A}') + n^2/8 - O(n)$.

(ii) *If job lengths are not bounded by a fixed constant, there is no randomized instance-optimal policy with one uniform additive $o(n^2)$ term. More precisely, for every randomized policy $\mathcal{A}$ independent of the bag and every sufficiently large n , there are a bag with lengths in $[0, n^2]$, a fixed labeling, and a randomized policy $\mathcal{A}'$ told that bag such that*

$$\text{cost}(\mathcal{A}) \geq \text{cost}(\mathcal{A}') + \frac{1}{8} n^2. \quad (110)$$

Proof. As in the introduction, $\text{cost}(\mathcal{A})$ includes the expectation over the private randomness of $\mathcal{A}$. For part (i), write $n = 2m$. Against the fixed deterministic policy, reveal length two on the first m labels it tests and zero on the remaining m . As usual, replay freezes this interaction into one fixed oblivious labeling. If the policy does not finish, the claim is immediate. Otherwise, let r be the number of length-two jobs completed before the first zero is tested. By that time all m positive jobs have been tested, so at least $m + 2r$ work has elapsed. The first r completed positive jobs contribute at least $3r(r+1)/2$. At that point the remaining jobs consist of m unit tests of zero jobs and $m - r$ processing operations of length two. Even SPT order for this remaining work gives

$$\begin{aligned} \text{cost}(\mathcal{A}, M) \geq & \frac{3r(r+1)}{2} + (2m-r)(m+2r) + \frac{m(m+1)}{2} \\ & + m(m-r) + (m-r)(m-r+1) \end{aligned}$$

$$= \frac{9}{2}m^2 + \frac{1}{2}r^2 + \frac{3}{2}m + \frac{1}{2}r \geq \frac{9}{8}n^2.$$

The comparison policy $\mathcal{A}'$ tests in a uniformly random order, completes zeros at their tests, defers every length-two job, and finally processes the deferred jobs. This is the stationary policy with the zero jobs early. Its exact cost identity Equation (82) gives

$$\text{cost}(\mathcal{A}') = n^2 + \frac{3}{4}n.$$

This proves the first claim.

For part (ii), put $H = n^2$ and first run $\mathcal{A}$ on the bag M^0 in which every job has length H . For a fixed private seed, let X_k be the number of completed jobs immediately before the k th test. Every processing operation contributes the same $Hn(n+1)/2$ in total, regardless of its position. The k th test delays $n - X_k$ unfinished jobs. Hence the excess over the policy $\mathcal{A}'$ that tests and processes each job immediately satisfies

$$\text{cost}(\mathcal{A}) - \text{cost}(\mathcal{A}') = \mathbb{E} \sum_{k=1}^n (k - 1 - X_k). \quad (111)$$

If this sum has expectation at least $n^2/8$, we are done.

Otherwise, replace one uniformly random label by a zero and leave every other length equal to H ; call the resulting bag M^1 . Conditional on the seed, the rank K at which the zero label is tested is uniform on $\{1, \dots, n\}$. Until that test, the run agrees with the all- H run, and therefore X_K length- H jobs complete before the zero. From Equation (111),

$$\mathbb{E}X_K = \frac{1}{n} \mathbb{E} \sum_{k=1}^n X_k > \frac{n-1}{2} - \frac{n}{8} = \frac{3n}{8} - \frac{1}{2}. \quad (112)$$

Forcing the zero job behind X_K length- H jobs adds HX_K to the clairvoyant SPT lower bound. On the other hand, a policy told M^1 can test in a private random order, defer positive jobs until it finds the zero, and then finish the positive jobs. Its completion order is SPT, and moving its unit tests away from the corresponding processing operations costs at most n^2 . Thus

$$\text{cost}(\mathcal{A}) - \text{cost}(\mathcal{A}') \geq H \left(\frac{3n}{8} - \frac{1}{2} \right) - n^2.$$

For $H = n^2$ this exceeds $n^2/8$ for all sufficiently large n . Averaging over the possible zero labels shows that one fixed oblivious labeling has the same lower bound. This proves part (ii). $\square$

5.1 The maximum-density benchmark

We first define the benchmark for an arbitrary finitely supported probability distribution D on $[0, \infty)$. A discovery module is described by numbers $g(p) \in [0, 1]$ on the support, with $g(0) = 1$. It tests one unit mass of jobs and processes the selected positive outcomes. Its completed mass and work are

$$a(g) = \int g(p) dD(p), \quad w(g) = 1 + \int pg(p) dD(p). \quad (113)$$

Choose g_* maximizing $a(g)/w(g)$ and put

$$\rho_* = \frac{a_*}{w_*}, \quad \theta_* = \frac{1}{\rho_*}, \quad \nu = (1 - g_*)D. \quad (114)$$

The one-coordinate addition and removal calculation from Lemma 4.3 shows that $g_* = 1$ below θ_* and $g_* = 0$ above θ_* . Only an atom at equality may be split. For an empirical measure one may equivalently sort the occurrences and maximize

$$\frac{|E|/n}{1 + n^{-1} \sum_{i \in E} p_i} \quad (115)$$

over processing-time prefixes E containing all zeros. A tied boundary class may be placed on either side.

Define

$$\Phi_{\text{OT}}(D) = w_* \left(1 - \frac{a_*}{2}\right) + \frac{1}{2} \iint \min\{p, q\} d\nu(p) d\nu(q). \quad (116)$$

For $D = D_M$ and a whole-occurrence maximizing prefix E , this is precisely $P(p, E)$ from Equation (81). The next lemma proves both that the value is independent of choices at a tied atom and that it lower-bounds every fluid policy.

Lemma 5.3 (Pointwise completion envelope). *For scaled time $s \geq 0$, let*

$$F_D(s) = \max \left\{ D(\{0\})T + \sum_{p>0} c_p : \begin{array}{l} 0 \leq T \leq 1, \quad 0 \leq c_p \leq D(\{p\})T, \\ T + \sum_{p>0} p c_p \leq s \end{array} \right\}. \quad (117)$$

Then F_D is attained simultaneously at all times by the stationary maximum-density schedule: repeat infinitesimal full modules g_ until all jobs have been tested, then process the residual measure ν in SPT order. Moreover,*

$$\Phi_{\text{OT}}(D) = \int_0^\infty (1 - F_D(s)) ds. \quad (118)$$

Proof. Let F^* be the completion curve of the claimed schedule. Up to time w_* it is the line $F^*(s) = \rho_* s$. Afterwards its slopes are $1/p$ for the residual processing times in increasing order. Every residual value is at least $\theta_* = 1/\rho_*$, so the slopes do not increase at the join. Thus F^* is nondecreasing and concave, with $F^*(0) = 0$.

Take a feasible point of Equation (117). If $T = 0$ there is nothing to prove. Otherwise put $x_p = c_p/T$. At $T = 1$, a selection whose work is at most w_* completes at most ρ_* times its work, by maximality of g_* . Above w_* , fractional-knapsack optimality says to select processing times in increasing order; this is exactly the continuation traced by F^* . Hence the completed mass of the corresponding full-test point is at most $F^*(1 + \sum p x_p)$. Concavity and $F^*(0) = 0$ now give

$$F^*\left(T\left(1 + \sum p x_p\right)\right) \geq T F^*\left(1 + \sum p x_p\right) \geq T\left(D(\{0\}) + \sum x_p\right).$$

Monotonicity proves $F_D \leq F^*$. Conversely, fractional amounts of the full module, followed by fractional SPT processing of ν , are feasible in Equation (117), so equality holds pointwise.

During discovery, integrating $1 - \rho_* s$ from 0 to w_* gives $w_*(1 - a_*/2)$. The remaining SPT area is $\frac{1}{2} \iint \min\{p, q\} d\nu(p) d\nu(q)$. This proves Equation (118) and Equation (116). $\square$

5.2 The announced upper bound

Suppose the bag is announced but its labeling is not. Choose a maximizing whole-occurrence prefix E , test labels in a uniformly random order, process an E -job immediately, and finish the residual jobs in SPT order. The exact identity Equation (82) and Equation (116) give

$$\mathbb{E} \text{cost}(\text{stat}, M) = n^2 \Phi_{\text{OT}}(D_M) + \frac{1}{2} \left(|E| + \sum_i p_i \right). \quad (119)$$

In particular, if $M \subseteq [0, L]$, the remainder is at most $(1 + L)n/2$. The announced optimum is therefore at most the claimed fluid value plus $O_L(n)$.

5.3 No adaptive announced policy has a better leading cost

Assign the occurrences of M uniformly to the public labels and fix a deterministic nonanticipating policy. In the adaptive order in which it tests new labels, the revealed occurrences form a uniform random permutation $P_1, \dots, P_n$ of M . One way to see this without assuming distinct processing times is to give equal-valued occurrences temporary private tokens: the map from a placement to the order in which the policy first reveals these tokens is a bijection of the finite permutation space.

For every prefix length k and threshold z , put

$$N_k(z) = |\{j \leq k : P_j \leq z\}|, \quad N(z) = |\{i : p_i \leq z\}|, \quad (120)$$

and define

$$\Delta = \max_{0 \leq k \leq n} \max_z \left| N_k(z) - \frac{k}{n} N(z) \right|. \quad (121)$$

Only the at most $n+1$ thresholds immediately after values of M matter. The hypergeometric Hoeffding bound, followed by a union bound over k and z , gives universal constants $c, C > 0$ such that

$$\mathbb{P}(\Delta \geq u) \leq 2(n+1)^2 e^{-cu^2/n}, \quad \mathbb{E}\Delta \leq C\sqrt{n \log(n+2)}. \quad (122)$$

We now transfer a realized schedule prefix to the deterministic fluid envelope. For a finite subprobability measure λ , let

$$Q_\lambda(u) = \inf\{z : \lambda([0, z]) \geq u\}, \quad W_\lambda(c) = \int_0^c Q_\lambda(u) du. \quad (123)$$

Thus $W_\lambda(c)$ is the least work needed to process mass c from λ , with the endpoint fraction of an atom taken when necessary. After k tests define

$$R_k = \frac{1}{n} \sum_{j \leq k} \delta_{P_j}, \quad A_k = \frac{k}{n} D_M, \quad \delta = \frac{\Delta}{n}. \quad (124)$$

Both measures have mass k/n , and

$$\sup_z |R_k([0, z]) - A_k([0, z])| \leq \delta. \quad (125)$$

Consequently, for $u > \delta$, $Q_{A_k}(u - \delta) \leq Q_{R_k}(u)$. Indeed, whenever the R_k -CDF has reached u , Equation (125) says that the A_k -CDF has reached $u - \delta$; taking generalized inverses gives the inequality, including at atom endpoints. Thus, when $c \geq \delta$, integration yields

$$W_{A_k}((c - \delta)_+) \leq \int_\delta^c Q_{R_k}(u) du \leq W_{R_k}(c). \quad (126)$$

If $c < \delta$, the retained mass $(c - \delta)_+$ is zero and the corresponding claim is immediate.

At physical time t , suppose k tests have completed, the policy has completed mass c of jobs, and the positive processing work belonging to those completed jobs is nq . Replacing the completed submeasure by the shortest mass c of R_k can only reduce its work, so $W_{R_k}(c) \leq q$. Also $k + nq \leq t$. By Equation (126), the proportional supply A_k contains mass $(c - \delta)_+$ processable with work at most q . This is a feasible point of Equation (117) at scaled time t/n . Therefore, pathwise and simultaneously for every t ,

$$\#\{\text{jobs completed by time } t\} \leq nF_{D_M}(t/n) + \Delta. \quad (127)$$

Put $H = 1 + n^{-1} \sum_i p_i$. The fluid curve is one by time H . Integrating the unfinished-job area in Equation (127) gives the pathwise inequality

$$\begin{aligned} \text{cost}(\mathcal{A}, M) &\geq \int_0^{nH} [n(1 - F_{D_M}(t/n)) - \Delta] dt \\ &= n^2 \Phi_{\text{OT}}(D_M) - Hn\Delta. \end{aligned} \quad (128)$$

Taking expectations and using Equation (122), every deterministic announced policy has expected cost at least

$$n^2 \Phi_{\text{OT}}(D_M) - O\left(Hn^{3/2} \sqrt{\log(n+2)}\right). \quad (129)$$

Yao's principle gives the same lower bound for randomized policies: the uniform distribution over occurrence placements is an oblivious finite input distribution, so some fixed placement has at least the average cost. When $p_i \leq L$, one has $H \leq 1 + L$, uniformly over M .

5.4 The same unannounced policy for every bag

It remains to prove Equation (107). Start with the policy from Section 4, keep its sampling, threshold, and fallback rules, but replace its fixed cutoff 32 by $B_n = 32 + n^{1/20}$. The full parameter choice is

$$k = \lfloor n^{3/4} \rfloor, \quad d = \lfloor n^{1/4} \rfloor, \quad B_n = 32 + n^{1/20}, \quad \eta = B_n/d. \quad (130)$$

The policy privately permutes the labels, tests the first k , rounds positive values upward to the B_n -grid, and uses the threshold closure of a maximum-density sample prefix.

Fix L . For all sufficiently large n , $B_n > L$ and the grid has no overflow job. Moreover, the full sample prefix has density at least $1/(1+L+\eta) > 2/B_n$, so fallback never occurs. Let D^+ be the rounded population histogram, $\hat{D}$ the sample histogram, and $\Delta = \|D^+ - \hat{D}\|_1$. By Lemma 4.6,

$$\mathbb{E}\Delta \leq \sqrt{\frac{d+2}{k}}. \quad (131)$$

For a fixed grid prefix J , let $\Psi_D(J)$ be the right-hand side of Equation (116) when J is used as the early set, whether or not it is optimal for D . If D, E are grid histograms supported on $[0, L + \eta]$, then

$$|\Psi_D(J) - \Psi_E(J)| \leq 3(L+2)\|D - E\|_1. \quad (132)$$

Indeed, early mass changes by at most the ℓ_1 distance, early first moment by at most $(L + \eta)$ times that distance, and the residual pair kernel $\min\{p, q\}$ changes by at most twice this amount under the two product measures. The displayed loose constant covers the three terms.

By Lemma 5.3, a maximum-density prefix minimizes $\Psi_D(J)$ over grid prefixes. If $\hat{J}$ is the prefix learned from the sample, the standard empirical-optimization sandwich and Equation (132) give

$$\Psi_{D^+}(\hat{J}) \leq \Phi_{\text{OT}}(D^+) + 6(L+2)\Delta. \quad (133)$$

Rounding each processing time upward by at most η preserves zeros and changes the fluid unfinished-mass area by at most η under the coupled schedule. Hence

$$\Phi_{\text{OT}}(D^+) \leq \Phi_{\text{OT}}(D_M) + \eta. \quad (134)$$

Conditional on the unordered sample, compare the actual sample-first schedule with the fully stationary policy for $\hat{J}$. The calculation in Section 4 bounds the extra cost by $O_L(nk + k^2)$. The diagonal term in the exact stationary identity is $O_L(n)$, and using the rounded values can only increase the comparator's processing delays. Therefore

$$\mathbb{E}[\text{cost}(\mathcal{A}^*, M) \mid S] \leq n^2 \Psi_{D^+}(\hat{J}) + O_L(nk + k^2 + n). \quad (135)$$

Average Equation (135), then use Equations (131), (133) and (134). Since

$$\sqrt{d/k} = O(n^{-1/4}), \quad \eta = O(n^{-1/5}), \quad k/n = O(n^{-1/4}),$$

we obtain

$$\mathbb{E} \text{cost}(\mathcal{A}^*, \sigma(M)) \leq n^2 \Phi_{\text{OT}}(D_M) + O_L(n^{9/5}) \quad (136)$$

for every fixed labeling σ . Together with Equation (129), this proves Theorem 5.1.

Finally, the exact certificate Equation (87) says that $\Phi_{\text{OT}}(D)$ is at most $4/3$ times the leading clairvoyant SPT value for every D . Equality is attained by $D = \frac{1}{2}\delta_0 + \frac{1}{2}\delta_2$. Thus the sharp $4/3$ theorem is exactly the worst-bag consequence of the stronger instance-optimal description.

6 Blind execution: instance-optimal two-phase policies

We now turn to $\text{BE}(L)$. Running an untouched job blindly consumes its actual processing time and completes it; testing consumes one unit, reveals the value, and permits the processing operation to be delayed. The adversary fixes an arbitrary labeled vector in $[0, L]^n$ before the private seed is drawn. The goal of this section is to prove Theorem 6.1, the rigorous version of the informal blind-execution theorem Theorem 1.3.

Theorem 6.1 (Blind-execution instance optimality). *Fix $L < \infty$. There exist a randomized nonanticipating policy $\mathcal{A}^*$, which knows n, L but not the processing-time multiset, and a deterministic sequence $\varepsilon_L(n) \rightarrow 0$ with the following properties. For every multiset $M \subseteq [0, L]$ of size n and every fixed labeling σ ,*

$$\mathbb{E} \text{cost}(\mathcal{A}^*, \sigma(M)) \leq n^2 \Phi(D_M) + \varepsilon_L(n)n^2. \quad (137)$$

For every randomized policy $\mathcal{A}$, even one told M , some fixed oblivious labeling σ satisfies

$$\mathbb{E} \text{cost}(\mathcal{A}, \sigma(M)) \geq n^2 \Phi(D_M) - \varepsilon_L(n)n^2. \quad (138)$$

The benchmark Φ is defined explicitly in Section 6.1; it is the minimum of a one-variable quadratic. One may take $\varepsilon_L(n) = O_L(n^{-1/6} \sqrt{\log(n+2)})$.

The boundedness assumption in Theorem 6.1 cannot be removed. In fact, without it even the much weaker worst-case guarantee is impossible.

Theorem 6.2 (No finite ratio for unbounded blind execution). *There are no randomized policies $\mathcal{A}_n$, constant $R < \infty$, and sequence $\eta(n) \rightarrow 0$ such that every vector $p \in [0, \infty)^n$ satisfies*

$$\mathbb{E} \text{cost}(\mathcal{A}_n, p) \leq R \text{cost}(\text{OPT}, p) + \eta(n)n^2. \quad (139)$$

Proof. Suppose that such a guarantee exists, and run $\mathcal{A}_n$ on the all-zero instance. Its offline cost is zero, so if Z_n denotes the expected online cost, then $Z_n \leq \eta(n)n^2 = o(n^2)$. We may assume that the policy finishes every job, since otherwise its cost is infinite.

Fix its private seed and order the labels by their first touch on this zero instance. Every first touch completes its job: a blind zero takes no time, while a tested zero completes with its unit test. If the job at rank r is tested, that test is performed while $n - r + 1$ jobs are unfinished and hence contributes $n - r + 1$ to the objective. Consequently

$$Z_n \geq \mathbb{E} \sum_{r: \text{rank } r \text{ is tested}} (n - r + 1). \quad (140)$$

Now choose one label uniformly and change only its length from zero to $H = n^2$. Until this exceptional label is first touched, the policy sees exactly the zero-instance history. If its rank r is touched blindly, its nonpreemptive execution occupies the machine for H time units while $n - r + 1$ jobs are unfinished. Averaging over the exceptional label and the private seed therefore gives

$$\begin{aligned} \mathbb{E} \text{cost}(\mathcal{A}_n, p) &\geq \frac{H}{n} \mathbb{E} \sum_{r: \text{rank } r \text{ is blind}} (n - r + 1) \\ &\geq \frac{H}{n} \left(\frac{n(n+1)}{2} - Z_n \right) = H \left(\frac{n}{2} - o(n) \right). \end{aligned} \quad (141)$$

Some fixed choice of the exceptional label has at least this expected cost. For every such labeling, the clairvoyant schedule runs all zero jobs first and the long job last, so $\text{cost}(\text{OPT}, p) = H$. With $H = n^2$, Equation (141) contradicts Equation (139) for all sufficiently large n . $\square$

The proof first replaces a round of tests and its immediate short outcomes by one divisible completion module. For a fixed tested fraction q , a fractional-knapsack envelope then forces the four-block order from the introduction and gives its exact quadratic cost. To handle an arbitrary adaptive policy, we show that its first-touch order is a bijective relabeling of the occurrence permutation; predictable sampling controls all touch prefixes at once. A growing grid extends the bound to arbitrary bounded multisets, and a blind sample removes advance knowledge of the bag. The next subsection defines the benchmark and begins the formal proof.

6.1 The distributional benchmark

Let D be a probability distribution on $[0, L]$ and let $P \sim D$. Write

$$\mu = \mathbb{E}P. \quad (142)$$

Blind execution completes one unit of job mass using expected work μ . The best possible test module is described by

$$\rho(D) = \sup_{0 \leq g \leq 1} \frac{\mathbb{E}g(P)}{1 + \mathbb{E}[Pg(P)]}, \quad \tau(D) = \frac{1}{\rho(D)}. \quad (143)$$

The function g selects outcomes that will be processed immediately after a test. For every such selection rule,

$$\tau \mathbb{E}g(P) - \mathbb{E}[Pg(P)] = \mathbb{E}[(\tau - P)g(P)] \leq \mathbb{E}(\tau - P)^+. \quad (144)$$

The maximizing rule selects all values below τ , no values above it, and an arbitrary fraction of a boundary atom. At equality,

$$\mathbb{E}(\tau - P)^+ = 1. \quad (145)$$

This equation has a unique solution: its left side is continuous and, after becoming positive, strictly increasing. If

$$a = \Pr(P < \tau), \quad \ell = \mathbb{E}[P \mathbf{1}_{P < \tau}],$$

then, with the boundary convention just described,

$$a\tau - \ell = 1, \quad \tau = \frac{1 + \ell}{a}. \quad (146)$$

Thus a full unit of tests together with all selected low outcomes is a divisible module of completion mass a , work $1 + \ell$, and work per completion τ .

If $\tau \geq \mu$, blind work has at least the completion density of any test module, and pure blind execution is optimal at the fluid scale. Suppose now that $\tau < \mu$. After testing a fraction q , the canonical schedule is

$$\begin{aligned} \underbrace{\text{tests with immediate } p < \tau}_{\text{discovery}} &\longrightarrow \underbrace{\text{known } \tau < p < \mu \text{ in SPT}}_{\text{medium}} \\ &\longrightarrow \underbrace{\text{untouched jobs blind}}_{\text{blind block}} \longrightarrow \underbrace{\text{known } p > \mu \text{ in SPT}}_{\text{high tail}}. \end{aligned} \quad (147)$$

For the explicit cost, first assume no atom lies at τ or μ , and put

$$\begin{aligned} \mathcal{L} &= \{p < \tau\}, & \mathcal{M} &= \{\tau < p < \mu\}, & \mathcal{H} &= \{p > \mu\}, \\ a &= \Pr(P \in \mathcal{L}), & \ell &= \mathbb{E}[P \mathbf{1}_{\mathcal{L}}], & m &= \mathbb{E}[P \mathbf{1}_{\mathcal{M}}], \\ d &= \Pr(P \in \mathcal{H}), & K_M &= \mathbb{E}[\min(P, Q) \mathbf{1}_{P, Q \in \mathcal{M}}], & K_H &= \mathbb{E}[\min(P, Q) \mathbf{1}_{P, Q \in \mathcal{H}}], \end{aligned}$$

where P, Q are independent with law D . For $r = 1 - q$, remaining-mass accounting gives

$$\begin{aligned} F_D(q) &= (1 + \ell) \left(q - \frac{aq^2}{2} \right) + qm(r + qd) + \frac{q^2 K_M}{2} \\ &\quad + \mu \left(r q d + \frac{r^2}{2} \right) + \frac{q^2 K_H}{2}. \end{aligned} \quad (148)$$

Atoms at τ are split between the low module and residual stock; atoms at μ may be placed on either side of the blind block. The same formula then holds for the resulting virtual classes.

Each term in Equation (148) has a direct area interpretation. During the test module, work $(1 + \ell)dq$ reduces unfinished mass from 1 to $1 - aq$. The medium work qm delays the untouched mass r and high mass qd , while K_M is the internal SPT pair term. The blind block delays all high jobs and itself loses mass linearly. Finally K_H is the high-tail pair term. Expansion gives the quadratic

$$F_D(q) = \frac{\mu}{2} + Aq + Bq^2, \quad (149)$$

where

$$A = 1 + \ell + m + \mu(d - 1) = 1 - \mathbb{E}(\mu - P)^+, \quad (150)$$

$$B = -\frac{a(1 + \ell)}{2} + m(d - 1) + \frac{K_M}{2} + \mu \left(\frac{1}{2} - d \right) + \frac{K_H}{2}. \quad (151)$$

The envelope proof below implies $B \geq 0$. In the short-test regime $\tau < \mu$, monotonicity in Equation (145) gives $A < 0$, and the optimal tested fraction is therefore

$$q^* = \begin{cases} 1, & B = 0, \\ \min\{1, -A/(2B)\}, & B > 0. \end{cases} \quad (152)$$

We define

$$\Phi(D) = \min_{0 \leq q \leq 1} F_D(q), \quad (153)$$

with $\Phi(D) = \mu/2$ in the regime $\tau \geq \mu$.

6.2 The fixed-fraction completion envelope

We first prove that Equation (147) is globally optimal in a fluid relaxation. Let D have finite support $0 = p_0 < p_1 < \dots < p_s \leq L$, with masses D_i . Fix a final tested fraction q . At an arbitrary schedule prefix, let

$$\begin{aligned} t &= \text{tested mass}, & b &= \text{blindly completed mass}, \\ c_i &= \text{processed tested mass of class } i > 0. \end{aligned}$$

The completed mass and idealized work are

$$y = D_0 t + b + \sum_{i>0} c_i, \quad x = t + \mu b + \sum_{i>0} p_i c_i, \quad (154)$$

subject to

$$0 \leq t \leq q, \quad 0 \leq b \leq 1 - q, \quad 0 \leq c_i \leq D_i t. \quad (155)$$

Define $H_{D,q}(x)$ as the largest completed mass satisfying these constraints with used work at most x . Every fluid schedule that ultimately tests mass q obeys this envelope, hence

$$\text{cost} \geq \int_0^\infty (1 - H_{D,q}(x)) dx. \quad (156)$$

Lemma 6.3 (Fixed- q envelope). *For every $q \in [0, 1]$, the right side of Equation (156) equals $F_D(q)$ and is attained by the four-block fluid policy. If $\tau \geq \mu$, it is at least $\mu/2$ for every q , with equality at $q = 0$.*

Proof. Split a boundary atom at τ into selected and residual virtual classes. Let $\mathcal{L}$ be the selected positive classes and put

$$a = D_0 + \sum_{i \in \mathcal{L}} D_i, \quad w = 1 + \sum_{i \in \mathcal{L}} p_i D_i, \quad w/a = \tau.$$

For the low part of any feasible prefix, write

$$y_L = D_0 t + \sum_{i \in \mathcal{L}} c_i, \quad x_L = t + \sum_{i \in \mathcal{L}} p_i c_i.$$

The density optimality in Equation (144) gives $y_L/x_L \leq a/w$. With $\lambda = y_L/a$, the revelation constraints imply $\lambda \leq t \leq q$, while density gives $\lambda w \leq x_L$. We may therefore replace the entire low prefix by λ full test modules, preserving completions and weakly reducing work.

Relax each residual capacity from $c_i \leq D_i t$ to $c_i \leq D_i q$. The result is an ordinary divisible knapsack with items

item	completion capacity	work per completion
full test module	aq	τ
residual tested class i	qD_i^R	p_i
blind module	$1 - q$	μ

The relaxation is pointwise tight. While greedy is partway through the test item, run the corresponding fraction of tests and all selected low outcomes. After all q test modules have run, every residual tested capacity is truly available. Greedy therefore realizes the original envelope at every work budget.

When $\tau < \mu$, the increasing work-per-completion order is exactly Equation (147); integrating it gives Equation (148). When $\tau \geq \mu$, the blind item has no worse density than the best test module and every excluded class has $p_i \geq \tau \geq \mu$. One all-blind item of capacity one pointwise dominates every $q > 0$ solution. Finally, feasibility is affine in q , so $H_{D,q}(x)$ is concave in q for each x . Hence its area complement $F_D(q)$ is convex, proving $B \geq 0$. $\square$

6.3 Adaptive first touches are a uniform urn

The fluid proof fixed q , but an announced finite policy may choose whom to touch next and when to stop testing as arbitrary functions of its history. The next finite lemma is the deferred-decisions statement needed to control such a policy.

Distinguish occurrences even when their processing times coincide. Let J be the public labels, Ω the occurrence tokens, and $v : \Omega \rightarrow [0, L]$ their values. A placement is a bijection $\sigma : J \rightarrow \Omega$. Fix a deterministic proper policy, equivalently fix the private seed of a randomized policy. Its first-touch labels $\ell_1(\sigma), \dots, \ell_n(\sigma)$ form a permutation of J . Define

$$R_{\mathcal{A}}(\sigma)_k = \sigma(\ell_k(\sigma)). \quad (157)$$

Lemma 6.4 (Adaptive-reveal bijection). *The map $\sigma \mapsto R_{\mathcal{A}}(\sigma)$ is a bijection from placements to permutations of Ω . Moreover, whether first touch k is a test or a blind execution is determined by the preceding revealed values $v(R_1), \dots, v(R_{k-1})$.*

Proof. Suppose $R_{\mathcal{A}}(\sigma) = R_{\mathcal{A}}(\sigma')$. A lockstep induction on k shows that the two executions have the same observable history before touch k . They have seen the same preceding values, so the deterministic policy performs the same intervening known-job operations, chooses the same next new label, and chooses the same test/blind kind. This remains true after blind execution because its realized duration is recorded in the public history. Hence $\ell_k(\sigma) = \ell_k(\sigma')$ for all k . Now Equation (157) gives $\sigma(\ell_k) = \sigma'(\ell_k)$ for every k ; since the ℓ_k enumerate J , the placements are equal. The reveal map is injective between finite sets of the same cardinality $n!$, hence bijective. The same induction under only a common revealed prefix proves predictability of the touch kind. $\square$

Consequently a uniform placement is carried to a uniform random permutation of occurrence tokens, and the test indicator is predictable. No conditional suffix compiler is needed. Duplicate values cause no problem because the bijection acts on occurrences.

For finite support, standard prefix concentration and a predictable martingale give the following simultaneous event. Stop while at least δn labels remain untouched, and put

$$e_n = \sqrt{8n \log n}, \quad \Gamma_n = (1 + 1/\delta)e_n. \quad (158)$$

Except with probability $o(1)$, for every earlier touch prefix and every class i ,

$$|\#\text{tested class } i - D_{n,i}\#\text{tests}| \leq \Gamma_n, \quad (159)$$

$$|\text{blind work} - \mu_n \# \text{blind jobs}| \leq L\Gamma_n. \quad (160)$$

Indeed, Hoeffding's inequality for without-replacement prefixes controls the remaining urn proportions. Multiplying the centered next-draw indicator by the predictable test selector gives a martingale with bounded increments; maximal Azuma–Hoeffding controls all stopping prefixes at once. The same argument with the predictable blind selector and bounded values proves Equation (160). Importantly, we never condition on the eventual tested fraction.

6.4 The announced-multiset lower bound

Fix a policy seed. Stop after $\lfloor (1 - \delta)n \rfloor$ first touches and let $Q_{n,\delta}$ be the number of tests completed by then, divided by n . Edit the suffix: every still-untouched job that would later be tested is instead executed blindly at its first original operation, and its later processing record is deleted. At most $3\delta n + O(1)$ bounded operations are moved or deleted. Pairwise delay accounting changes total cost by at most

$$O((L + 1)\delta n^2). \quad (161)$$

The edited schedule performs exactly $Q_{n,\delta}n$ tests.

Work on the single event Equations (159) and (160) and only now set $q = Q_{n,\delta}$. Normalize mass and work by n , and let $\gamma_n = \Gamma_n/n$. At any operation boundary, let t, b, z, c_i denote tested mass, blindly completed mass, zero mass completed by tests, and processed known positive mass of class i . The event gives

$$z \leq D_{n,0}t + \gamma_n, \quad c_i \leq D_{n,i}t + \gamma_n.$$

Set $c'_i = (c_i - \gamma_n)^+$. Then

$$c'_i \leq D_{n,i}t, \quad t \leq q, \quad b \leq 1 - q.$$

If x is the actual normalized prefix work, then

$$t + \mu_n b + \sum_i p_i c'_i \leq x + L(\gamma_n + \delta), \quad (162)$$

$$z + b + \sum_i c_i \leq D_{n,0}t + b + \sum_i c'_i + (s + 1)\gamma_n. \quad (163)$$

By the definition of the empirical envelope,

$$\text{completed fraction by } x \leq H_{D_n,q}(x + L(\gamma_n + \delta)) + (s + 1)\gamma_n. \quad (164)$$

Horizontal shifting loses at most the shift itself because $0 \leq 1 - H \leq 1$; the vertical slack is multiplied by a horizon at most $L + 1$. After restoring Equation (161), the fixed- q envelope lemma therefore gives, pathwise on the good event,

$$\frac{\text{cost}(\mathcal{A}, M)}{n^2} \geq F_{D_n}(Q_{n,\delta}) - O_D(\gamma_n + \delta + 1/n) \geq \Phi(D_n) - O_D(\gamma_n + \delta + 1/n). \quad (165)$$

The long-test regime uses the $\mu_n/2$ dominance part of Lemma 6.3. Averaging first over the uniform placement and then over private seeds proves the same lower bound for every randomized announced policy.

The random-labeling and announced minimax values coincide. A uniform labeling is a valid finite Yao distribution, giving one inequality. For the other, privately relabel the physical jobs before running an optimal deterministic urn policy; against every fixed assignment, the policy sees a uniform assignment. Thus the lower bound above applies to the announced oblivious minimax problem, not merely to one randomized input model.

6.5 A growing grid and arbitrary bounded multisets

The number of distinct values may grow with n . Keep zero as a separate cell, partition $(0, L]$ into

$$K_n = \lfloor n^{1/6} \rfloor \quad (166)$$

bins, and round every positive value upward; let $h_n = L/K_n$ be the mesh. Zero preservation matters because a tested zero completes at its test.

Coupling each value with its rounded value changes total normalized processing work by at most h_n . Running corresponding operation words therefore changes normalized completion area by at most h_n . In particular,

$$|\Phi(D) - \Phi(D^{\text{grid}})| \leq h_n. \quad (167)$$

One way to see this without assuming a stable numerical threshold is to transport a policy through the coupling: attach a private auxiliary original mark to every rounded observation and make all decisions using that mark. The zero-preserving coupling makes the completion events coincide.

Apply the predictable urn event to all $K_n + 1$ cells and take $\delta_n = n^{-1/6}$. Repairing every class loses at most $(K_n + 1)\Gamma_n/n$ completion mass. The rounded analogue of Equation (164) is

$$\text{completed fraction by } x \leq H_q^{\text{grid}}(x + \zeta_n) + \zeta_n, \quad (168)$$

where

$$\zeta_n = O_L\left(\frac{(K_n + 1)\Gamma_n}{n} + h_n + \delta_n\right) = O_L(n^{-1/6}\sqrt{\log(n + 2)}). \quad (169)$$

Integrating exactly as above and using nonnegativity on the bad event yields the announced lower bound

$$\frac{\text{VAL}_{\text{ann}}(M)}{n^2} \geq \Phi(D_M) - O_L(n^{-1/6}\sqrt{\log(n + 2)}). \quad (170)$$

For the matching upper bound, compute an optimal canonical template for the rounded empirical multiset, choose $\lfloor q^*n \rfloor$ uniformly random labels to test, and execute the four blocks. Pairwise delay accounting expresses its expectation as bounded one-occurrence and two-distinct-occurrence kernels. The without-replacement pair coefficients differ from their product-law values by $O(1/n)$ uniformly in the tested count. Hence

$$\mathbb{E}C_n(\pi) = n^2\Psi_{D_M^{\text{grid}}}(\pi) + O_L(n) \quad (171)$$

uniformly over canonical templates π . Together with Equation (167), this proves

$$\left| \frac{\text{VAL}_{\text{ann}}(M)}{n^2} - \Phi(D_M) \right| \leq O_L(n^{-1/6}\sqrt{\log(n + 2)}). \quad (172)$$

6.6 Removing the announcement

The universal policy first draws a uniform random permutation and blindly executes

$$k_n = \lceil n^{1/2} \rceil \quad (173)$$

sample jobs. Their observed durations give a grid histogram $\hat{D}_n$. The policy minimizes the canonical quadratic over every grid threshold, blind-block insertion position, and $q \in [0, 1]$, then runs the selected four-block template on the remaining jobs in the unused random order.

Let D_n and D_n^R be the full and remaining grid histograms. Sampling without replacement gives

$$\mathbb{E}\|\hat{D}_n - D_n\|_1 \leq \sqrt{\frac{K_n + 1}{k_n}}, \quad \mathbb{E}\|\hat{D}_n - D_n^R\|_1 \leq \frac{n}{n - k_n} \sqrt{\frac{K_n + 1}{k_n}}. \quad (174)$$

For a fixed canonical template, formula Equation (148) consists only of bounded one- and two-draw expectations. Therefore, uniformly in the grid and template,

$$|\Psi_D(\pi) - \Psi_E(\pi)| \leq C_L \|D - E\|_1, \quad (175)$$

for example with $C_L = 20(L + 1)$. If $\hat{\pi}$ minimizes the sampled objective and π_R minimizes the remaining objective, the empirical optimization sandwich is

$$\begin{aligned} \Psi_{D_n^R}(\hat{\pi}) &\leq \Psi_{\hat{D}_n}(\hat{\pi}) + C_L \|\hat{D}_n - D_n^R\|_1 \\ &\leq \Psi_{\hat{D}_n}(\pi_R) + C_L \|\hat{D}_n - D_n^R\|_1 \\ &\leq \Phi(D_n^R) + 2C_L \|\hat{D}_n - D_n^R\|_1. \end{aligned} \quad (176)$$

Conditional on the sample, the remaining order is uniform and the learned template is fixed, so Equation (171) applies. The sample itself has work at most Lk_n and delays at most n jobs, adding at most $Lnk_n = O_L(n^{3/2})$. Combining Equations (167) and (174) with Equations (175) and (176) gives

$$\frac{\mathbb{E} \text{cost}(\mathcal{A}^*, p)}{n^2} \leq \Phi(D_M) + O_L \left(n^{-1/6} \sqrt{\log(n + 2)} \right) \quad (177)$$

uniformly over every fixed labeled vector $p \in [0, L]^n$.

For the converse, reveal M to an arbitrary policy for free and draw a uniform labeling. The announced lower bound Equation (170) applies seedwise. Some fixed labeling in the support has cost at least the average, which proves Equation (138). This completes the proof of Theorem 6.1.

6.7 Why the medium block is necessary

Let

$$\Pr(P = 0) = \frac{1}{5}, \quad \Pr(P = 9) = \frac{1}{5}, \quad \Pr(P = 16) = \frac{3}{5}.$$

Then $\mu = 57/5$ and $\tau = 5$. Thus tested zeros are immediate, known nine-jobs precede the blind block, and known sixteen-jobs follow it. Formula Equation (148) becomes

$$F_D(q) = \frac{57}{10} - \frac{44}{25}q + \frac{11}{10}q^2.$$

Its minimizer is $q^* = 4/5$ and its value is 4.996. If all tested nonzero jobs are instead deferred until after the blind block, the best value is 5.04. The gap is a positive constant times n^2 : the medium SPT block is not cosmetic, and a policy of the simpler form “threshold, then blind, then all tested residuals” is not instance optimal.

7 Blind optimization: speedup without revelation

We finally separate the two benefits that were bundled together in the revealing-optimization model. Raw execution still takes the known time u , and one unit of optimization still reduces the processing time to $p_i \in [0, u]$. The difference is that optimization does not reveal p_i . An optimized job must be started blindly and its duration becomes known only when it finishes. We call this the *blind-optimization* model $\text{BO}(u)$.

We study randomized policies against an oblivious adversary. This is the natural setting for learning the bag: a private random sample is representative under every fixed labeling, whereas an adversary can make any predetermined deterministic sample atypical. We do not pursue a separate deterministic worst-case curve for this model.

This small change removes the threshold decision entirely. It also shows that the bounded randomized ratio in the revealing model is genuinely an information effect.

The goal of this section is to prove two closely related statements. The first, Theorem 7.1, is the rigorous version of the informal instance-optimality theorem Theorem 1.4. The second, Theorem 7.2, is the rigorous version of Theorem 1.5 and maximizes that instance-specific value relative to the clairvoyant optimum. Keeping them together avoids repeating the same two-block reduction.

For a multiset M let

$$\mu_M = \frac{1}{n} \sum_{i=1}^n p_i$$

and define

$$\Phi_{\text{bo}}(M) = \frac{1}{2} \min\{u, 1 + \mu_M\}. \quad (178)$$

The two terms have an immediate meaning. Running every job raw uses u units per completion. Optimizing and then blindly processing every job uses $1 + \mu_M$ units per completion on average.

Theorem 7.1 (Blind-optimization instance optimality). *For every fixed $u < \infty$ there exist a randomized nonanticipating policy $\mathcal{A}^{\text{bo}}$, depending only on n, u , and a sequence $\varepsilon_u(n) \rightarrow 0$ such that every size- n multiset $M \subseteq [0, u]$ and every fixed labeling σ satisfy*

$$\mathbb{E} \text{cost}(\mathcal{A}^{\text{bo}}, \sigma(M)) \leq n^2 \Phi_{\text{bo}}(M) + \varepsilon_u(n) n^2. \quad (179)$$

Conversely, for every randomized nonanticipating policy $\mathcal{A}$, even one told M , some fixed oblivious labeling σ satisfies

$$\mathbb{E} \text{cost}(\mathcal{A}, \sigma(M)) \geq n^2 \Phi_{\text{bo}}(M) - \varepsilon_u(n) n^2. \quad (180)$$

Thus one policy is asymptotically optimal, at the n^2 scale, for every multiset simultaneously. Unlike the benchmarks in the preceding sections, Φ_{bo} depends only on the empirical mean.

The proof is simpler than in the revealing models. Because optimization reveals nothing, it can be moved next to the processing of its job. Every job is then either a raw block of known length u or an optimized blind block with average length $1 + \mu_M$. A predictable-sampling bound rules out an adaptive advantage, while a sample estimates the one number μ_M . The first subsection makes this argument formal; the second performs the outer worst-case optimization.

7.1 Proof of instance optimality

In the equivalent random-bag formulation, the labels are shuffled uniformly before the policy starts. Then the lower half simply says that no policy, even one given M , beats $n^2 \Phi_{\text{bo}}(M) - o_u(n^2)$ in expectation.

Normal form. An optimization that is separated from the processing of its job can be moved to immediately before that processing operation. The move reveals no information, leaves the completion time of that job and all later jobs unchanged, and makes every job completed in between one unit earlier. A policy can perform the same transformation online by remembering the optimized state and elapsed time in a virtual copy of its history while postponing the physical optimization. Hence we may regard every optimized job as one indivisible block of hidden length $1 + p_i$. A raw block has the known length u . Both blocks complete exactly one job.

The announced lower bound. Fix the private seed of a policy and place the announced occurrences uniformly behind the labels. The first-touch argument of Lemma 6.4 has a direct, slightly simpler analogue. Give equal-valued occurrences distinct hidden tokens and map a placement to the token order induced by the policy's first touches. If two placements induce the same token order, a lockstep induction gives the same observable histories and hence the same first-touch labels; the placements are equal. The map is therefore a bijection of the finite placement space. Notice that a raw block does not reveal its token's value to the policy. We expose that value only in the analysis, which gives a richer history; the policy's choice between raw and optimized mode is still predictable with respect to this richer prefix. Thus the order of analytically exposed occurrences is a uniform permutation and the selector of optimized blocks is predictable.

Fix $\delta > 0$ and ignore the last δn first touches for the moment. The same predictable without-replacement estimate as in Section 6.3 gives one event, of probability $1 - o(1)$, on which every earlier prefix containing tn completed optimized blocks has total hidden processing work

$$\sum_{i \text{ optimized in the prefix}} p_i = \mu_M tn + o_\delta(n). \quad (181)$$

The event is simultaneous over all prefixes, so a stopping fraction chosen adaptively by the policy may be substituted only after the event has been fixed. Editing the last δn jobs to either mode changes total completion cost by $O_u(\delta n^2)$.

This can be made uniform rather than diagonalized. Take $\delta_n = n^{-1/4}$. The bounded selected-work estimate used in Section 6.3 has deviation

$$O_u\left((1 + \delta_n^{-1})\sqrt{n \log(n+2)}\right) = O_u\left(n^{3/4}\sqrt{\log(n+2)}\right)$$

and failure probability $o(1)$, simultaneously over all relevant prefixes. After division by n , both this deviation and the edited suffix contribute $O_u(n^{-1/4}\sqrt{\log(n+2)})$ to the normalized completion-area bound.

Put $a = 1 + \mu_M$. If the final optimized fraction is q , the best possible fluid ordering puts the cheaper of the two block types first. When $a \leq u$ its area is

$$F_{a,u}(q) = a\left(q - \frac{q^2}{2}\right) + \frac{u(1-q)^2}{2}, \quad F'_{a,u}(q) = (a-u)(1-q). \quad (182)$$

It is minimized at $q = 1$, with value $a/2$. When $a \geq u$, raw blocks go first and the area is

$$F_{a,u}(q) = \frac{u(1-q^2)}{2} + \frac{aq^2}{2} = \frac{u}{2} + \frac{a-u}{2}q^2, \quad (183)$$

which is minimized at $q = 0$, with value $u/2$. An adjacent exchange proves that no interleaving has smaller area: swapping two consecutive blocks puts the shorter cost per

completion first. Equations (181)–(183), with the uniform choice $\delta = \delta_n$, give

$$\frac{1}{n^2} \mathbb{E} \text{cost}(\mathcal{A}, M) \geq \frac{1}{2} \min\{u, 1 + \mu_M\} - o_u(1).$$

The estimate is uniform in the deterministic policy. Averaging over its private seed and applying the same finite Yao/relabeling argument as in Section 6.4 proves (180). Together with the upper bound below, one may take $\varepsilon_u(n) = O_u(n^{-1/4} \sqrt{\log(n+2)})$ after enlarging its constant.

One policy for all multisets. Take $k = \lceil n^{2/3} \rceil$ uniformly random sample jobs. Optimize each sample job and process it immediately, thereby observing its duration, and let $\hat{\mu}$ be their sample mean. If $1 + \hat{\mu} < u$, optimize and immediately process every remaining job in fresh random order. Otherwise run every remaining job raw.

The sample contributes at most $O_u(nk) = o_u(n^2)$ beyond the main schedule. Sampling without replacement gives

$$\mathbb{E}|\hat{\mu} - \mu_M| = O_u(k^{-1/2} + k/n) = o_u(1).$$

Choosing the wrong mode can cost at most $O(n^2)|\hat{\mu} - \mu_M|$, while choosing the correct mode has leading cost $n^2\Phi_{\text{bo}}(M)$. This proves (179). In particular, the universal policy needs to learn only one number, not a histogram or a threshold.

7.2 The exact worst-case curve

The clairvoyant scheduler knows p_i and uses the shorter block

$$\ell_u(p) = \min\{u, 1 + p\}.$$

It orders these blocks shortest first. If P, Q are independent draws from a distribution D on $[0, u]$, its leading coefficient is

$$\Omega_u(D) = \frac{1}{2} \mathbb{E} \min\{\ell_u(P), \ell_u(Q)\}. \quad (184)$$

Consequently the instance-specific ratio from Theorem 7.1 is

$$\alpha_u^{\text{bo}}(D) = \frac{\min\{u, 1 + \mathbb{E}P\}}{\mathbb{E} \min\{\ell_u(P), \ell_u(Q)\}}. \quad (185)$$

Let $u_{\text{bo}} > 1$ denote the unique root greater than one of

$$u^3 - 2u^2 - u + 1 = 0, \quad u_{\text{bo}} = 2.2469796037 \dots \quad (186)$$

Theorem 7.2 (Exact blind-optimization curve). *The optimal randomized size-asymptotic competitive ratio in $\text{BO}(u)$ against an oblivious adversary is*

$$R_{\text{bo}}(u) = \begin{cases} 1, & 0 < u \leq 1, \\ \frac{u^3}{u^2 + (u-1)^3}, & 1 \leq u \leq u_{\text{bo}}, \\ \frac{1}{2} \left(1 + \sqrt{\frac{u^2 + u - 1}{u - 1}} \right), & u \geq u_{\text{bo}}. \end{cases} \quad (187)$$

The two branches agree at u_{bo} . Moreover,

$$R_{\text{bo}}(u) = \frac{\sqrt{u}}{2} + \frac{1}{2} + o(1) \quad (u \rightarrow \infty). \quad (188)$$

Proof. The claim is trivial for $u \leq 1$, because raw execution is no longer than the unit optimization operation. Assume $u > 1$ and let

$$H(t) = \Pr(P > t), \quad 0 \leq t \leq u.$$

The layer-cake identity applied to (184) gives

$$2\Omega_u(D) = 1 + \int_0^{u-1} H(t)^2 dt. \quad (189)$$

Put $S = \int_0^{u-1} H(t) dt$. Since H is nonincreasing,

$$\int_{u-1}^u H(t) dt \leq H(u-1) \leq \frac{S}{u-1}.$$

Therefore, writing $\mu = \mathbb{E}P = \int_0^u H(t) dt$,

$$S \geq \frac{u-1}{u} \mu. \quad (190)$$

Cauchy–Schwarz now yields

$$\int_0^{u-1} H(t)^2 dt \geq \frac{S^2}{u-1} \geq (u-1) \left(\frac{\mu}{u} \right)^2. \quad (191)$$

Set $b = \mu/u$. Equations (185)–(191) imply

$$\alpha_u^{\text{bo}}(D) \leq f_u(b) := \frac{\min\{u, 1 + ub\}}{1 + (u-1)b^2}. \quad (192)$$

Equality is attained by the binary distribution

$$\Pr(P = u) = b, \quad \Pr(P = 0) = 1 - b. \quad (193)$$

Indeed, its survival function is constant b on $[0, u)$, so both inequalities in (191) are equalities.

It remains to maximize the scalar function f_u . Put

$$b_0 = \frac{u-1}{u}.$$

For $b \geq b_0$ the numerator is u and f_u decreases, so this region is maximized at b_0 . For $b \leq b_0$, differentiation gives the unique unconstrained maximizer

$$b_*(u) = \frac{u}{u-1 + \sqrt{(u-1)(u^2 + u - 1)}}. \quad (194)$$

The equality $b_*(u) = b_0$ is equivalent to $u^3 - 2u^2 - u + 1 = 0$. Hence b_0 is optimal up to u_{bo} and $b_*(u)$ is optimal afterwards. Substitution gives

$$f_u(b_0) = \frac{u^3}{u^2 + (u-1)^3}, \quad f_u(b_*) = \frac{1}{2} \left(1 + \sqrt{\frac{u^2 + u - 1}{u-1}} \right).$$

This proves (187). The binary distributions (193), uniformly relabeled, are oblivious Yao distributions. Together with the universal upper policy of Theorem 7.1, they prove both sides of the competitive-ratio claim. If the maximizing mass b is irrational, finite multisets use $\lfloor bn \rfloor$ copies of u ; continuity gives the same asymptotic ratio. Finally, $(u^2 + u - 1)/(u - 1) = u + 2 + 1/(u - 1)$ gives (188). $\square$




Figure 3: Revealing p_i keeps the randomized revealing-optimization ratio bounded and eventually gives the $4/3$ plateau. If optimization changes the processing time but does not reveal it before execution, the exact ratio instead grows as $\sqrt{u}/2$.

The hard distribution explains the separation. Most jobs have optimized length zero, while a carefully chosen small fraction have length u . Blind optimization is attractive on average, so the online policy optimizes the jobs. It cannot recognize a long outcome before committing to it, and an early long job delays many zeros. The clairvoyant optimum optimizes the zeros, runs the long jobs raw, and places them last. In the revealing model, by contrast, the same long outcomes can be deferred after their tests. The gap between the two curves therefore measures the scheduling value of the revealed information itself.

8 Deterministic revealing optimization: the exact curve

The goal of this section is to prove Theorem 8.1, the rigorous deterministic part of the informal revealing-optimization curve theorem Theorem 1.7. We now allow raw execution for a finite common time u . All jobs have this same known upper limit, and a test reveals $p_j \in [0, u]$. Thus the offline effective length of job j is

$$q_j = \min\{u, 1 + p_j\}, \quad (195)$$

and the offline optimum schedules the q_j in nondecreasing order. We write $V(u)$ for the optimal deterministic size-asymptotic ratio at this fixed common upper limit.

The answer has six branches. We first define their transition points. Put

$$\varphi = \frac{1 + \sqrt{5}}{2}, \quad u_\diamond = \frac{1 + \sqrt{3 + 2\sqrt{5}}}{2} = 1.866760399173862\dots \quad (196)$$

Let $s_0 > 1$ be the root of

$$s_0^3 = (s_0 + 1)^2, \quad u_0 = 1 + s_0 = 3.147899035704787\dots \quad (197)$$

For $s = u - 1$, put

$$T_s(z) = sz^2 + (s^3 + s^2 + 1)z - (s^2 + s + 1)(s + 2), \quad (198)$$

and let $\rho_I(u)$ be the positive root of

$$s\rho^2 + (s^3 + s^2 + 1)\rho - (s^2 + s + 1)(s + 2) = 0. \quad (199)$$

Denote the left-hand side by $T_s(\rho)$. Equivalently,

$$\rho_I(u) = \frac{-1 - u + 2u^2 - u^3 + \sqrt{u^6 - 4u^5 + 10u^4 - 6u^3 - 3u^2 + 6u - 3}}{2(u - 1)}. \quad (200)$$

At the other end, let $z_* > 1$ solve $z_* - 3 = \log z_*$ and define

$$r = \frac{2}{z_* - 1}, \quad R_{\text{obl}} = 1 + r = 1.570574096649395\dots, \quad z_* = 4.505241495792883\dots \quad (201)$$

For $\varphi + 2 \leq u \leq z_*$, define $C_{\text{mix}}(u) = 1 + c$ by the unique solution (c, m) of

$$\text{artanh } m - m = 1 - \frac{1}{c} + \frac{1}{2} \log\left(1 + \frac{2}{c}\right), \quad u = 1 + \frac{2}{c} - m, \quad (202)$$

with $r \leq c \leq 1/\varphi$ and $0 \leq m \leq 1/\varphi$.

Theorem 8.1 (Exact revealing-optimization curve). *For every fixed $u > 0$, the optimal deterministic size-asymptotic ratio in the revealing-optimization model is*

$$V(u) = \begin{cases} 1, & 0 < u \leq 1, \\ u, & 1 \leq u \leq u_\diamond, \\ \rho_I(u), & u_\diamond \leq u \leq u_0, \\ 1 + \frac{1}{\sqrt{u-1}}, & u_0 \leq u \leq \varphi + 2, \\ C_{\text{mix}}(u), & \varphi + 2 \leq u \leq z_*, \\ R_{\text{obl}}, & u \geq z_*. \end{cases} \quad (203)$$

For every branch there is an explicit deterministic algorithm satisfying

$$\text{cost}(\mathcal{A}, I) \leq V(u) \text{cost}(\text{OPT}, I) + O_u(n).$$

On finite $u \geq z_*$ the remainder is one absolute $O(n)$ term, and Theorem 3.1 supplies the same algorithm at $u = \infty$. Conversely, for every deterministic algorithm $\mathcal{A}$ and every $\epsilon > 0$, there are arbitrarily large fixed instances I such that

$$\text{cost}(\mathcal{A}, I) \geq (V(u) - \epsilon) \text{cost}(\text{OPT}, I).$$

The proof combines two kinds of lower bounds. Binary hidden-stopping instances give the first four branches, while a continuous capped version of the harmonic construction gives the mixed branch and the plateau. Three explicit policies match them: raw execution, a forced-prefix uniform threshold, and the history-dependent threshold from obligatory testing. The upper-bound witness can be read from the following policy map. The algorithms themselves are specified formally in algorithms 7 to 9.

range	policy	parameter choice
$0 < u \leq u_\diamond$	RAW	none
$u_\diamond < u \leq u_0$	FORCEDPREFIXUTE	$b = b(u)$ from (220)
$u_0 < u \leq \varphi + 2$	FORCEDPREFIXUTE	$b = 0$
$\varphi + 2 < u < z_*$	ADAPTIVETHRESHOLD	$c = c(u)$ from (202)
$z_* \leq u \leq \infty$	ADAPTIVETHRESHOLD	$c = r$

8.1 The shape of the claimed curve

The global-maximum assertion follows directly from the displayed formulas. Only the algebraic branch needs a check. Implicit differentiation of $T_s(\rho_I(u)) = 0$, with $s = u - 1$, gives

$$\rho'_I(u) = -\frac{\rho^2 + (3s^2 + 2s)\rho - 3(s+1)^2}{2s\rho + s^3 + s^2 + 1}. \quad (204)$$

Using $T_s(\rho) = 0$, its numerator factors as

$$\frac{2s^3 + s^2 - 1}{s} \left(\rho - \frac{2s^3 + 3s^2 - 2}{2s^3 + s^2 - 1} \right).$$

On the relevant interval $s \geq u_\diamond - 1 > 2/3$, the prefactor is positive, while the fraction in parentheses is at most $5/3$ because $4s^3 - 4s^2 + 1 > 0$. The sign certificate (224) gives $\rho > 5/3$. Hence $\rho'_I(u) < 0$. The reciprocal branch and the mixed parametrization are also decreasing, so V attains its global maximum only at $u = u_\diamond$.

The lower bounds, including the fact that the middle construction is safe against raw execution, are proved next. We then give self-contained upper bounds through $u = \varphi + 2$. The parameterized upper bound for the last two branches is proved separately in Section 8.7.

8.2 The binary stopping game and its lower envelope

The first four branches already occur on binary instances $p_j \in \{0, u\}$. The stopping point must be hidden from the algorithm; announcing it in advance gives a strictly weaker lower bound.

Lemma 8.2 (Raw-safe hidden stopping). *Fix $u > 1$, a target $\rho \leq u$, and a stopping fraction $\alpha \in (0, 1)$. Against a deterministic algorithm, there is a legal binary adversary whose leading excess at its first crossing is*

$$(u - \rho)(1 - \sigma^2) + \sigma^2 f_{u,\rho}(y, b), \quad (205)$$

where $0 \leq b \leq y \leq 1$ and

$$0 \leq y - \alpha < \frac{1}{n - v}. \quad (206)$$

Moreover,

$$f_{u,\rho}(y, b) = 1 - \rho + 2y + (u - 1)(1 - \rho)y^2 + b^2 + 2b(u - 1 - uy). \quad (207)$$

Here $\sigma \in [0, 1]$ is the fraction of jobs not already executed raw. Consequently, if

$$\min_{0 \leq b \leq \alpha} f_{u,\rho}(\alpha, b) \geq 0,$$

then the adversary gives ratio at least $\rho - O_u(1/n)$.

Proof. Start in a long phase. A fresh job that is tested receives $p = u$, while a fresh job run raw is assigned the hidden value zero. If v jobs have been run raw and L tested-long jobs have been seen, stop at the first touch for which

$$L \geq \alpha(n - v). \quad (208)$$

All untouched jobs are then assigned value zero. Only for the purpose of lower-bounding the remaining cost, we grant the algorithm this future information and allow an optimal

continuation; in the actual interaction, values are still revealed only by tests. If the line is never crossed, every job was run raw: if even one job had been tested, then after the last first touch the tested fraction would be one. On the all-zero hidden instance the ratio is $u \geq \rho$.

At a crossing, let e of the L long jobs already be complete, let $d = L - e$ be pending, and let $x = n - v - e - d$ jobs remain untouched. The following favorable canonical schedule lower-bounds the algorithm: run the v raw jobs; test and immediately process the e completed long jobs; perform the d noncompleting long tests; test the x zero jobs; and finally process the d pending long jobs. Indeed, if a noncompleting unit test is immediately before an available length- u completion while h jobs remain, the two orders contribute $h + hu$ and $hu + (h - 1)$, respectively; moving the completion left saves one. Raw completions are available from time zero, and identical long jobs may be relabeled so that the e completed ones are tested first. After the switch, a zero test-completion has length one and therefore precedes a pending operation of length u . These exchanges respect precedence.

The exact costs of this relaxed schedule and of OPT are

$$A_{\text{ex}} = u \left(vn - \frac{v(v-1)}{2} \right) + (1+u) \left(e(n-v) - \frac{e(e-1)}{2} \right) + d(n-v-e) + x(d+x) - \frac{x(x-1)}{2} + u \frac{d(d+1)}{2}, \quad (209)$$

$$O_{\text{ex}} = \frac{n(n+1)}{2} + (u-1) \frac{(e+d)(e+d+1)}{2}. \quad (210)$$

Put $\nu = v/n$, $\delta = (v+e+d)/n$, and $\lambda = e/n$. Twice the quadratic coefficients in Equations (209) and (210) are

$$\begin{aligned} \mathcal{A} &= 1 + 2\delta(1 - u\nu) + (u-1)\delta^2 + 2\nu(\nu + u - 2) \\ &\quad + \lambda^2 + 2\lambda(\nu + u - 1 - u\delta), \\ \mathcal{O} &= 1 + (u-1)(\delta - \nu)^2. \end{aligned}$$

Now put $\sigma = 1 - \nu$, $y = (e+d)/(n-v)$, and $b = e/(n-v)$. Direct expansion gives exactly Equation (205) after normalizing by $n^2/2$. The first-crossing overshoot in Equation (208) is less than one: a test increases $e+d - \alpha(n-v)$ by one and a raw first touch increases it by α . This proves Equation (206), and in particular $\sigma(y - \alpha) < 1/n$. There is one small domain issue: the actual $b \leq y$ can exceed α . Put $\bar{b} = \min\{b, \alpha\}$. Then $0 \leq b - \bar{b} \leq y - \alpha$. Since $f_{u,\rho}$ is Lipschitz on the unit square,

$$\sigma^2 f_{u,\rho}(y, b) \geq \sigma^2 f_{u,\rho}(\alpha, \bar{b}) - O_u(1/n). \quad (211)$$

Thus the stated minimum over $0 \leq \bar{b} \leq \alpha$ suffices uniformly even when $n-v$ is small. The discarded diagonal terms are $O_u(n)$ in the unnormalized costs. Finally, the adaptive transcript defines a fixed instance by replay: every tested answer was fixed when revealed, and every raw job's zero value stayed hidden. This proves the lemma. $\square$

Minimizing Equation (207) over the algorithm's completion choice gives

$$b_*(y) = \max\{0, uy - (u-1)\}. \quad (212)$$

The upper constraint $b \leq y$ never binds, because $uy - (u-1) \leq y$ for $y \leq 1$. There are therefore two elementary quadratic branches. On the positive branch, put

$$A_u = u^2 - u + 1, \quad D_{u,\rho} = A_u + (u-1)\rho. \quad (213)$$

Then

$$\min_b f_{u,\rho}(y, b) = 1 - \rho - (u - 1)^2 + 2A_u y - D_{u,\rho} y^2. \quad (214)$$

Its maximizer is $\alpha = A_u/D_{u,\rho}$, and tangency to zero is

$$(1 - \rho - (u - 1)^2)D_{u,\rho} + A_u^2 = 0. \quad (215)$$

Expanding Equation (215) gives Equation (199). On the zero- b branch,

$$\min_b f_{u,\rho}(y, b) = 1 - \rho + 2y + (u - 1)(1 - \rho)y^2, \quad (216)$$

and tangency gives

$$\rho = 1 + \frac{1}{\sqrt{u - 1}}, \quad \alpha = \frac{1}{\sqrt{u - 1}}. \quad (217)$$

The latter point belongs to the zero- b branch iff $(u - 1)^3 \geq u^2$, with equality at $u = u_0$.

For $\rho = u$, the maximum in Equation (214) is nonnegative exactly while

$$[u(u - 1)]^2 - u(u - 1) - 1 \leq 0, \quad (218)$$

whose positive equality is $u = u_\diamond$. We have proved the following binary lower envelope.

Proposition 8.3 (Binary lower curve). *Every deterministic algorithm has size-asymptotic ratio at least*

$$B(u) = \begin{cases} 1, & 0 < u \leq 1, \\ u, & 1 \leq u \leq u_\diamond, \\ \rho_I(u), & u_\diamond \leq u \leq u_0, \\ 1 + \frac{1}{\sqrt{u - 1}}, & u \geq u_0. \end{cases} \quad (219)$$

The lower bound uses only $p_j \in \{0, u\}$.

Proof. For $1 < u \leq u_\diamond$, use $\rho = u$ and the maximizing point of Equation (214). For $u_\diamond \leq u \leq u_0$, use the positive root $\rho_I(u)$ and $\alpha = A_u/D_{u,\rho_I(u)}$; the tangency identity makes the maximum zero. The minimizing completion mass is positive in the interior and becomes zero at $u = u_0$. For $u \geq u_0$, use the zero- b tangency point from Equation (217). The branch conditions are precisely the two inequalities checked above. Finally, $V(u) \geq 1$ is trivial for $u \leq 1$. In each nontrivial case Lemma 8.2 supplies fixed binary instances by replay. $\square$

8.3 Matching upper bounds through the binary regime

The first policy is formalized in algorithm 8.

Algorithm 8 RAW(J, u)

Input: jobs J and a finite common upper limit u .

1: Fix an arbitrary order $j_1, \dots, j_n$.

2: **For** $i = 1, \dots, n$, execute j_i untested for u time.

For $u \leq u_\diamond$, use RAW. It is optimal when $u \leq 1$; when $u > 1$, every effective length in Equation (195) is at least one, so the ratio is at most u .

For the next interval, let $\rho = \rho_I(u)$, $s = u - 1$, and define

$$K = s^2 + s + 1, \quad D = K + s\rho, \quad b = \frac{K - s^2\rho}{D}, \quad \chi = \frac{1 - b}{\rho}, \quad \alpha = \frac{K}{D}. \quad (220)$$

The defining equation for ρ gives

$$0 \leq b < \chi < \alpha < 1, \quad \chi = \frac{s(s+1)}{D}, \quad b = (s+1)\alpha - s; \quad (221)$$

For completeness, these inequalities follow without numerical root isolation. Since T_s is increasing on positive arguments,

$$T_s\left(\frac{K}{s^2}\right) = \frac{K}{s^3}(-s^3 + s^2 + 2s + 1) \geq 0 \quad (s \leq s_0), \quad (222)$$

and hence $b = (K - s^2\rho)/D \geq 0$, with equality only at $s = s_0$. Moreover,

$$\alpha - \chi = \frac{1}{D}, \quad \chi - b = \frac{s^2\rho - 1}{D} > 0. \quad (223)$$

For the last sign,

$$9T_s(5/3) = f(s) := 6s^3 - 12s^2 - 2s - 3 < 0. \quad (224)$$

throughout the present interval. Indeed, on $[4/5, 1]$ the polynomial is decreasing and $f(4/5) = -1151/125$; on $[1, s_0]$ it has one interior minimum, while $f(1) = -11$ and $f(s_0) < f(13/6) = -95/36$. Here $s_0 < 13/6$ follows by substituting $13/6$ in $s^3 - (s+1)^2$. Thus $\rho > 5/3$, while $s > 4/5$, and $s^2\rho > 16/15$. Finally $K < D$ gives $\alpha < 1$. In particular, $b = 0$ occurs only at $u = u_0$. The remaining two elementary branches use the single procedure in algorithm 9.

Algorithm 9 FORCEDPREFIXUTE(J, u, b)

Input: jobs J , finite $u > 1$, and $b \in [0, 1]$.

- 1: Fix a test order $j_1, \dots, j_n$; set $k \leftarrow \lfloor bn \rfloor$, $\tau \leftarrow \min\{1, u - 1\}$, and $Q \leftarrow \emptyset$.
 - 2: **For** $i = 1, \dots, n$ **do**:
 - 3: Test j_i for one unit and observe p_{j_i} .
 - 4: **If** $i \leq k$ or $p_{j_i} \leq \tau$, process j_i immediately.
 - 5: **Otherwise**, add j_i to Q .
 - 6: Process the jobs of Q in nondecreasing revealed processing time.
-

On $u_\diamond \leq u \leq u_0$, use FORCEDPREFIXUTE with b from Equation (220). Rounding bn changes the cost by $O_u(n)$.

Here is the complete binary calculation. Adjacent exchanges place as many long jobs as possible in the forced prefix and put all suffix-long jobs before suffix zeros. If x is the total long mass, then the prefix-long mass is $a = \min\{b, x\}$ and the deferred-long mass is $d = (x - b)_+$. After division by n^2 , the leading excess $G(x) = A - \rho O$ is

$$G(x) = \begin{cases} \frac{1 - \rho}{2} + u \left(x - \frac{x^2}{2} \right) - \frac{\rho s}{2} x^2, & 0 \leq x \leq b, \\ \frac{1 - \rho}{2} + u \left(b - \frac{b^2}{2} \right) + (1 - b)(x - b) + \frac{s}{2}(x - b)^2 - \frac{\rho s}{2} x^2, & b \leq x \leq 1. \end{cases} \quad (225)$$

The second quadratic is strictly concave, has its stationary point at $x = \alpha$, and satisfies $G(\alpha) = 0$; after substituting Equation (220), the latter identity is exactly Equation (199).

Since $b < \alpha$, its right derivative at b is positive. The left derivative satisfies $G'_-(b) = G'_+(b) + s(1 - b) > 0$, and the first quadratic is concave, so it is increasing on $[0, b]$. Thus $G(x) \leq 0$ for every $x \in [0, 1]$, including the limiting case $b = 0$. This proves the binary upper bound without a hidden local-versus-global step.

For $u_\diamond \leq u \leq 2$, arbitrary values reduce to binary values by independent Bernoulli extremalization, in the following order. First raise *every* suffix-deferred value to u . This cannot lower ALG, while its offline effective length was already u . Only after this deterministic step, replace a prefix value p by u with probability p/u and by zero otherwise, and replace a suffix-immediate value $p \leq u - 1$ by u with probability p and by zero otherwise. Notice that the u realization of a suffix-immediate job becomes deferred, so status is not preserved. Nevertheless the online check is pairwise. For jobs i before j , the leading pair contribution is

$$g_{ij} = \begin{cases} 1 + p_i, & i \text{ is immediate,} \\ 2 + p_j, & i \text{ is deferred and } j \text{ is immediate,} \\ 2 + \min\{p_i, p_j\}, & i, j \text{ are deferred.} \end{cases} \quad (226)$$

A rounded prefix job stays immediate and preserves its expected value. If i is suffix-immediate, its rounded contribution is one when it becomes zero and at least two when it becomes u , hence its expectation is at least $1 + p_i$. If i is deferred and j is suffix-immediate, then i has been raised to u and the expected pair contribution is $2 + up_j \geq 2 + p_j$; a deferred-deferred pair can only increase. Thus expected ALG does not decrease, up to its $O_u(n)$ diagonal terms.

The expected effective lengths are, respectively,

$$1 + \frac{u-1}{u}p \leq q(p), \quad 1 + (u-1)p \leq 1 + p, \quad (227)$$

and u for a deferred value. Since the minimum is jointly concave, $\mathbb{E} \min(q'_i, q'_j) \leq \min(q_i, q_j)$. Thus some binary realization has competitive excess at least that of the original instance.

It remains to justify arbitrary values for $2 \leq u \leq u_0$. The following quadratic lemma records the entire endpoint calculation.

Lemma 8.4 (UTE endpoint game). *For $2 \leq u \leq u_0$, UTE with the parameters Equation (220) satisfies*

$$\text{ALG} \leq \rho_I(u) \text{OPT} + O_u(n). \quad (228)$$

Proof. Fix the symbolic prefix/immediate/deferred decisions. Convexity in one coordinate reduces a prefix job to $p \in \{0, u\}$, a suffix-immediate job to $p \in \{0, 1\}$, and a suffix-deferred job to $p \in \{1^+, u\}$. For the last claim, on $1 < p < u - 1$ the deferred-deferred term has coefficient $1 - \rho < 0$ multiplying a concave minimum; on $[u - 1, u]$, OPT is flat and ALG is nondecreasing. Adjacent exchanges give the canonical endpoint order.

Normalize counts by n . Let a be prefix- u mass, d suffix- u mass, t suffix- 1^+ mass, and m suffix-immediate-one mass. Their constraints are

$$0 \leq a \leq b, \quad d, t, m \geq 0, \quad d + t + m \leq 1 - b. \quad (229)$$

Pairwise accounting using Equation (226) gives

$$A = \frac{1}{2} + (s+1) \left(a - \frac{a^2}{2} \right) + (1-b)(d+t+m) - \frac{m^2}{2} + \frac{s}{2}d^2, \quad (230)$$

$$O = \frac{1}{2} + \frac{(a+d+t+m)^2}{2} + \frac{s-1}{2}(a+d)^2. \quad (231)$$

Moving m to t does not change O and increases A , so $m = 0$. Put $x = a + d$ and write $F(x) = A - \rho O$ after optimizing the other variables. The derivative of the gap in t is $1 - b - \rho(x + t)$, so its maximizing value is $(\chi - x)_+$. At fixed x, t , moving mass from d to a leaves O unchanged and changes A at rate $s(1 - x) + b - a \geq 0$. Hence

$$t = (\chi - x)_+, \quad a = \min\{b, x\}, \quad d = (x - b)_+. \quad (232)$$

For $x \geq \chi$, the gap $A - \rho O$ is the binary concave quadratic already discussed; it is tangent to zero at $x = \alpha$. For $x < \chi$, put

$$C_0 = -\frac{\rho - 1}{2} + \frac{(1 - b)^2}{2\rho}, \quad \gamma = s - \rho(s - 1). \quad (233)$$

Substitution gives

$$A - \rho O = C_0 + a(s + b - sx) - \frac{a^2}{2} + \frac{\gamma x^2}{2}. \quad (234)$$

One has $\gamma > 0$ on $1 \leq s \leq s_0$. Indeed, $T_s(2) = s(s^2 - s + 1) > 0$, so for $s \leq 2$ the equation $T_s(\rho) = 0$ gives $\rho < 2$, and hence $\gamma > 2 - s \geq 0$. For $2 \leq s \leq s_0 < 13/6$,

$$16T_s(7/4) = 12s^3 - 20s^2 + s - 4 > 0; \quad (235)$$

the polynomial is 14 at $s = 2$ and has positive derivative thereafter. Thus $\rho < 7/4$ and $\gamma > 7/4 - 3s/4 > 1/8$.

On $0 \leq x \leq b$, one has $a = x$, and

$$A - \rho O = C_0 + (s + b)x - \frac{s + 1 + \rho(s - 1)}{2}x^2. \quad (236)$$

Its derivative is decreasing, and the derivative at b multiplied by D is

$$\rho(1 + s^2 + \rho s^2(s - 1)) > 0, \quad (237)$$

so the maximum is at b . On $b \leq x \leq \chi$, Equation (234) is convex and its maximum is at b or χ . The endpoint χ lies on the binary branch. Finally,

$$F(b) - F(\chi) = \frac{\chi - b}{2} [2sb - \gamma(b + \chi)], \quad (238)$$

and direct use of Equation (220) gives

$$D[\gamma(b + \chi) - 2sb] = -s + (1 + s - s^3)\rho + s^2(s - 1)\rho^2 > 0. \quad (239)$$

For a rational sign certificate, T_s is increasing on positive arguments and

$$9T_s(5/3) = 6s^3 - 12s^2 - 2s - 3 < 0, \quad (240)$$

so $\rho > 5/3$. If the right-hand side of Equation (239) is denoted by $q_s(\rho)$, then

$$3 \partial_\rho q_s(5/3) = 7s^3 - 10s^2 + 3s + 3 > 0. \quad (241)$$

The last polynomial is increasing for $s \geq 1$, so q_s is increasing for $\rho \geq 5/3$. Finally,

$$9q_s(5/3) = 10s^3 - 25s^2 + 6s + 15 > 0. \quad (242)$$

For the last sign, write $s = 3/2 + z$; the expression becomes $3/2 - 3z/2 + 20z^2 + 10z^3$. It is at least $3/2$ for $-1/2 \leq z \leq 0$, and for $z \geq 0$ it is larger than $3/2 - 9/320$. These inequalities prove that every endpoint gap is nonpositive. Diagonals and prefix rounding contribute $O_u(n)$. $\square$

At $u = u_0$, $b = 0$. The same zero-prefix algorithm remains exact through $u = \varphi + 2$.

Lemma 8.5 (Zero-prefix extension). *For $u_0 \leq u \leq \varphi + 2$, FORCEDPREFIXUTE($J, u, 0$) from algorithm 9 is $(1 + 1/\sqrt{u-1})$ -competitive up to $O_u(n)$.*

Proof. The same coordinatewise reduction leaves capped-deferred mass d , boundary-deferred mass t , immediate-one mass m , and zeros. With $s = u - 1$, these masses satisfy $d, t, m \geq 0$ and $d + t + m \leq 1$, and the leading costs are

$$A = \frac{1}{2} + d + t + m - \frac{m^2}{2} + \frac{sd^2}{2}, \quad O = \frac{1}{2} + \frac{(d + t + m)^2}{2} + \frac{(s-1)d^2}{2}. \quad (243)$$

Move m to t and put $y = d + t$. For fixed y , the ratio is fractional-linear in d^2 , so its maximum is on one of two faces:

$$\frac{1+2y}{1+y^2} \leq \varphi \quad (d=0), \quad 1 + \frac{2y}{1+sy^2} \leq 1 + \frac{1}{\sqrt{s}} \quad (d=y). \quad (244)$$

The first one-variable function is maximized at $y = 1/\varphi$ and the second at $y = 1/\sqrt{s}$, which gives the two displayed constants. The second value dominates the first exactly when $u \leq \varphi + 2$. Binary instances attain it. $\square$

8.4 The unrestricted mixed lower bound

For $\varphi + 2 < u < z_*$, neither the binary family nor the pure cap-free family is tight. The correct adversary puts a capped block before a scaled harmonic core. Two lemmas ensure that optional raw execution and early processing of caps cannot escape it.

The cap-free construction needed below is exactly the scaled harmonic core from Lemma 3.3; no second phase-accounting proof is needed here. We now prepend capped jobs to that reusable core.

Lemma 8.6 (Cap deferral). *Prepend capped mass m with value $p = u$ to a later cap-free block of mass $s = 1 - m$ and total processing mass L . Suppose every later value satisfies $1 + p \leq u$ and*

$$s(u-1) - L - m \geq 0. \quad (245)$$

Then processing any capped job before all later jobs are complete cannot decrease the leading online cost.

Proof. A cap completed after its test block but before a later job of value p has mutual cost $1 + u$; leaving it pending until after that job gives $2 + p$. The exchange saves $u - 1 - p \geq 0$. If cap mass q is instead completed inside the cap-test block, its pairs with all later jobs add $q[s(u-1) - L]$. Its largest possible saving against not-yet-tested caps is $qm - q^2/2$. Relative to leaving every cap pending, the change is therefore at least

$$q[s(u-1) - L - m] + \frac{q^2}{2} \geq 0. \quad (246)$$

The finite diagonal discrepancy is lower order. $\square$

After cap deferral, pairwise accounting gives the offline coefficient O and the guaranteed online lower benchmark A^{lb} :

$$A^{\text{lb}} = A_0^{\text{lb}} + m(1 - m) + m(1 + L) + \frac{u}{2}m^2, \quad (247)$$

$$O = O_0 + m(1 - m + L) + \frac{u}{2}m^2. \quad (248)$$

Indeed, a cap-later pair contributes $2 + p$ online and $1 + p$ offline, whereas a cap-cap pair contributes $2 + u$ online and u offline; collecting these pairs yields the two cross terms and the cap squares above. In particular, $A^{\text{lb}} - O = (A_0^{\text{lb}} - O_0) + m$.

Lemma 8.7 (Raw-safe quota and replay). *Fix $m \in (0, 1)$ and a later cap-free block satisfying the hypotheses of Lemma 8.6. Against every deterministic optional algorithm there are fixed instances whose asymptotic excess is at least the cap-free hybrid benchmark $A^{\text{lb}} - CO$ from Equations (247) and (248) for every target $C \leq u$.*

Proof. In the initial phase, a tested fresh job receives $p = u$, while a job run raw is assigned hidden value zero. If v raw jobs and ℓ tested caps have been touched, stop at the first crossing

$$\frac{\ell}{n - v} \geq m. \quad (249)$$

With $S = n - v$, the overshoot is less than one, so $\ell = mS + O(1)$. If no crossing occurs, no test can have occurred: after all first touches, any positive ℓ would make the ratio in Equation (249) equal one. Thus every job was run raw and the all-zero hidden instance has ratio $u \geq C$.

At a crossing, scale the prescribed later instance to the remaining $S - \ell$ first touches, absorbing the one-job cap overshoot and integer rounding into its zero group. A later test reveals the next value in this fixed type stream; a later raw action consumes the next value but keeps it hidden from the optional algorithm.

Move every initial raw-zero completion to the start. It has the same length u as cap processing, and crossing a unit test only lowers completion cost. With

$$Z_v = vn - \frac{v(v-1)}{2}, \quad (250)$$

this prefix contributes uZ_v online and Z_v offline. Delete it. The remaining S labels consist, up to one-job rounding, of capped fraction m and the prescribed later cap-free block.

If the optional algorithm later runs a fresh assigned value p raw, a physical cap-free simulation tests and immediately processes that job in time $1 + p \leq u$, suppresses the answer from the virtual optional algorithm, and idles until its virtual u -block ends. Inductively every physical completion is no later than its virtual optional completion. Since the finite-cap and cap-free offline lengths are both $1 + p$ on the later block,

$$\text{ALG}_{\text{opt}} - C \text{OPT}_{\text{opt}} \geq (u - C)Z_v + (\text{ALG}_{\text{hyb}} - C \text{OPT}_{\text{hyb}}) - O_u(n). \quad (251)$$

The first term is nonnegative. Finally, record every assigned value on its job label. Replaying this fixed vector produces the same transcript because the values learned only by the physical simulation were hidden from the virtual algorithm. This proves legality. $\square$

Apply Lemma 3.3 with $s = 1 - m$. Substitution into Equations (247) and (248) gives the guaranteed two-parameter lower ratio

$$\mathcal{R}_u(m, t) = 1 + \frac{m + \frac{(1-m)^2}{2}(1 - t^{-2})}{\frac{(1-m)^2}{4}(1 + t^{-2} + 2 \log t) + m(1 - m)(1 + \log t) + \frac{u}{2}m^2}. \quad (252)$$

At an interior stationary point with value $1 + c$, differentiation and the zero-gap equation give

$$\log t = u - 2 - \frac{1}{c}, \quad t^2 = \frac{1 - m}{1 + m} \left(1 + \frac{2}{c}\right). \quad (253)$$

Eliminating t gives precisely Equation (202). Conversely, those equations make $\mathcal{R}_u(m, t) = 1 + c$.

The branch runs from

$$(u, m, t) = \left(\varphi + 2, \frac{1}{\varphi}, 1\right) \quad \text{to} \quad (u, m, t) = (z_*, 0, \sqrt{z_*}). \quad (254)$$

It satisfies $u \geq \varphi + 2$, $\log t < 1$, and $m \leq 1/\varphi$. Therefore

$$(1 - m)(u - 1 - \log t) > (1 - m)\varphi \geq \frac{1}{\varphi} \geq m, \quad (255)$$

so Equation (245) holds; also $1 + p < 3 < u$ for every later positive value in a sufficiently fine finite construction. The functions

$$g(c) = 1 - \frac{1}{c} + \frac{1}{2} \log\left(1 + \frac{2}{c}\right), \quad h(m) = \operatorname{artanh} m - m \quad (256)$$

have $g'(c) = 2/[c^2(c + 2)] > 0$ and $h'(m) = m^2/(1 - m^2) > 0$. Hence $m(c) = h^{-1}(g(c))$ is strictly increasing, while

$$u(c) = 1 + \frac{2}{c} - m(c) \quad (257)$$

is strictly decreasing. The identities $g(r) = h(0) = 0$ and $g(1/\varphi) = h(1/\varphi)$ give the two endpoints in Equation (254); consequently every $u \in [\varphi + 2, z_*]$ has exactly one parameter pair.

To pass to fixed finite instances, first approximate (m, t) rationally while retaining the strict cap inequalities, then choose a sufficiently fine harmonic family, and finally multiply all rational block sizes by a common integer H . For fixed harmonic depth, quota overshoot, diagonals, and rounding are $O_u(H)$ whereas the displayed costs are $\Theta(H^2)$. Send first H , then the harmonic depth, and finally the rational approximation to their limits. Together with Lemma 8.7, this proves

$$V(u) \geq C_{\text{mix}}(u) \quad (\varphi + 2 \leq u \leq z_*). \quad (258)$$

The construction above applies directly in the open interval. At $u = \varphi + 2$ and $u = z_*$ the same statement follows by approaching from the interior, so the endpoints are included.

8.5 The large-upper-limit lower bound by cap-free transfer

Lemma 8.8 (Bounded cap-free transfer). *Let $u \geq 1$, and let $V_\infty(P)$ denote the deterministic size-asymptotic value in the cap-free limit, restricted to $0 \leq p_j \leq P$. Then*

$$V(u) \geq V_\infty(u - 1). \quad (259)$$

In particular,

$$V(u) \geq R_{\text{obl}} \quad \left(u \geq 1 + \frac{1}{r} = 2.752620747896442 \dots\right). \quad (260)$$

Proof. Given a finite-cap algorithm, construct a cap-free algorithm on instances with $p_j \leq u - 1$ by running a virtual copy of the finite-cap algorithm. Copy ordinary tests and processing actions. When the virtual algorithm runs an untouched job raw for u time, physically test and immediately process it in $1 + p_j \leq u$ time, hide the answer from the virtual copy, and wait for the virtual clock. Every physical completion is no later

than its virtual completion. Moreover, $\min\{u, 1 + p_j\} = 1 + p_j$, so finite-cap and cap-free optima coincide. Thus we obtain the following game-value comparison. For every finite-cap algorithm $\mathcal{A}$, the simulation produces a cap-free algorithm $\mathcal{B}$ and, pointwise on every instance with $p_j \leq u - 1$,

$$\text{ALG}_{\mathcal{A}} \geq \text{ALG}_{\mathcal{B}}, \quad \text{OPT}_u = \text{OPT}_{\infty}.$$

Consequently

$$\text{CR}_{\infty}(\mathcal{A}) \geq \text{CR}_{\infty}(\mathcal{B}; p \leq u - 1) \geq V_{\infty}(u - 1).$$

Taking the infimum over $\mathcal{A}$ proves Equation (259).

The harmonic cap-free family attaining R_{obl} has limiting largest processing time

$$1 + \log(1 + y_*) = 1 + \frac{1}{2} \log z_* = \frac{z_* - 1}{2} = \frac{1}{r}, \quad y_* = \sqrt{z_*} - 1. \quad (261)$$

Approximation from below also gives the endpoint, proving Equation (260). $\square$

8.6 Completion of the curve and consistency at the joins

The lower-bound side of Theorem 8.1 is now complete: Proposition 8.3 supplies the first four branches, Equation (258) the fifth, and Equation (260) the last. The raw algorithm, Lemmas 8.4 and 8.5, supplies the matching upper bounds through $\varphi + 2$. The remaining upper bounds are proved in Section 8.7.

For clarity, all five joins are exact:

$$\begin{aligned} \rho_I(u_{\diamond}) &= u_{\diamond}, \\ \rho_I(u_0) &= 1 + \frac{1}{\sqrt{s_0}} = 1.682327803828019\dots, \\ 1 + \frac{1}{\sqrt{(\varphi + 2) - 1}} &= \varphi = C_{\text{mix}}(\varphi + 2), \\ C_{\text{mix}}(z_*) &= 1 + r = R_{\text{obl}}. \end{aligned}$$

At $u = 1$, both initial branches equal one. Representative values of the new mixed branch are

u	$V(u)$	m	t
3.800000	1.598851588	0.539725635	1.138984621
4.000000	1.583283633	0.428863572	1.330517832
4.071679486	1.579204806	0.381330539	1.412236296
4.300000	1.571692026	0.198387083	1.734651379

Remark 8.9 (Asymptotic scope). All statements above concern the fixed- u size-asymptotic ratio. Below the plateau, the upper bounds have additive error $O_u(n)$. On $u \in [z_*, \infty]$ the same algorithm has one uniform $O(n)$ error. We do not claim a finite-instance endpoint inequality with one additive constant.

The next subsection, Section 8.7, proves the matching middle- and high- u upper bounds and thereby completes the proof of Theorem 8.1.

8.7 Matching upper bound for revealing optimization

We now prove the upper bounds on the two ranges on which capped outcomes matter. Besides completing the upper-bound side of the revealing-optimization curve, the proof records explicitly why an outcome on the optional cap cannot be treated as an ordinary deferred outcome. Throughout this subsection,

$$C = 1 + c, \quad \varphi = \frac{1 + \sqrt{5}}{2},$$

and $z_* > 1$ is the solution of $z_* - 3 = \log z_*$. We also set

$$r = \frac{2}{z_* - 1}.$$

For $c \in [r, 1/\varphi]$, let $m = m(c) \in [0, 1/\varphi]$ be determined by

$$\operatorname{artanh} m - m = 1 - \frac{1}{c} + \frac{1}{2} \log \left(1 + \frac{2}{c} \right), \quad u = 1 + \frac{2}{c} - m. \quad (262)$$

The right-hand side of the first equation is strictly increasing in c , as is the left-hand side in m . Consequently $m(c)$ is increasing and $u(c)$ is decreasing. The endpoints are

$$(c, m, u) = \left(\frac{1}{\varphi}, \frac{1}{\varphi}, \varphi + 2 \right) \quad \text{and} \quad (c, m, u) = (r, 0, z_*).$$

Theorem 8.10. *For every $u \in [\varphi + 2, \infty]$, the formally specified algorithm ADAPTIVETHRESHOLD from algorithm 7 satisfies*

$$\text{ALG} \leq \begin{cases} (1 + c) \text{OPT} + O_u(n), & \varphi + 2 \leq u \leq z_*, \quad c \text{ given by (262)}, \\ (1 + r) \text{OPT} + O(n), & u \geq z_*. \end{cases}$$

The constant in the last $O(n)$ is absolute and uniform over the extended interval $u \in [z_*, \infty]$. In particular these are upper bounds on the deterministic size-asymptotic competitive ratio.

Proof.

At $u = \infty$ this is Theorem 3.4, with the same algorithm and the same four-endpoint bank. For the finite-cap calculations below, fix $u < \infty$.

The algorithm. Use ADAPTIVETHRESHOLD(J, c) from algorithm 7; on the high interval use $c = r$. For the analysis, immediately before a test let x be the number of jobs not yet tested, N the number of earlier strictly positive outcomes, and D the number of those outcomes deferred to the final tail. Thus

$$y = \frac{D - CN}{x} \leq 0 \quad (263)$$

and the threshold is

$$a_c(y) = \begin{cases} 1, & y \leq -1, \\ 1 + \frac{1}{2} \log \left(\frac{c + 2}{c - 2y} \right), & -1 \leq y \leq 0. \end{cases} \quad (264)$$

This is the same threshold as Equation (31). On the mixed interval $c \geq r$, and hence

$$a_c(y) \leq a_c(0) \leq a_r(0) = \frac{1}{r} < u - 1.$$

The same inequality holds on the high interval, where we use $c = r$. Thus every outcome in the capped region $[u - 1, u]$ is deferred.

The state y never increases. Indeed, the numerator in (263) changes by 0, $-C$, or $-c$ after a zero, a positive immediate outcome, or a positive deferred outcome, respectively, while x decreases. This monotonicity will allow us to allocate every pair cost when its relevant endpoint becomes known.

Exact pair objective and the five endpoints. Write

$$\bar{p}_i = \min\{p_i, u - 1\}.$$

The clairvoyant optional optimum is SPT on jobs of effective lengths $1 + \bar{p}_i = \min\{1 + p_i, u\}$. For two jobs tested in the order $i < j$, let

$$e_{ij} = \begin{cases} (p_i - p_j)_+, & i \text{ is processed immediately,} \\ 1 + (p_j - p_i)_+, & i \text{ is deferred and } j \text{ is immediate,} \\ 1, & i \text{ and } j \text{ are deferred.} \end{cases} \quad (265)$$

The first line applies whenever i is immediate, regardless of the action taken on j . Direct completion-time accounting gives

$$\text{ALG} = \sum_i (1 + p_i) + \sum_{i < j} (1 + \min\{p_i, p_j\} + e_{ij}), \quad (266)$$

$$\text{OPT} = \sum_i (1 + \bar{p}_i) + \sum_{i < j} (1 + \min\{\bar{p}_i, \bar{p}_j\}). \quad (267)$$

It is therefore natural to introduce

$$\Gamma_n = \sum_{i < j} (e_{ij} + \min\{p_i, p_j\} - C \min\{\bar{p}_i, \bar{p}_j\}). \quad (268)$$

Equations (266)–(268) yield the exact identity

$$\text{ALG} - C \text{OPT} = \Gamma_n - c \binom{n}{2} + \sum_i [(1 + p_i) - C(1 + \bar{p}_i)]. \quad (269)$$

The final sum is nonpositive. Below the cap its i -th summand is $-c(1 + p_i)$; on the cap it is at most $1 - cu \leq 0$. Thus it is enough to control Γ_n .

Fix a symbolic immediate/deferred path. Coordinatewise convexity in (268) reduces a worst outcome vector to the five formal endpoint types

type	outcome	action
Z	0	immediate
E	0^+	immediate
M	$a_c(y)$	immediate
Q	$a_c(y)^+$	deferred
U	u	deferred and capped.

(270)

For an immediate coordinate the two positive endpoints are 0^+ and the live threshold; the genuine zero is kept separate because it does not enter N . A deferred coordinate below $u - 1$ moves to the live threshold or to $u - 1$. On $[u - 1, u]$, the offline contribution is flat and the algorithmic contribution is nondecreasing, so the latter endpoint moves to u . The symbolic categories, and therefore all later threshold states, remain unchanged during each coordinate move. This is the only endpoint reduction used below.

A local pair allocation. Separate the earlier E -outcomes from the substantive outcomes M, Q, U . Let

$$P = \#\{\text{earlier } M, Q, U\}, \quad E = \#\{\text{earlier } E\}, \quad D = \#\{\text{earlier } Q, U\},$$

and let K be the number of earlier U -outcomes. Among the ordinary substantive outcomes, let I, d be the numbers of M, Q -outcomes and let A_I, A_D be the sums of their processing values. Thus $P = I + d + K$ and $D = d + K$.

At a current ordinary endpoint, with $a = a_c(y)$, allocate the relaxed local charges

$$\ell_Z = \ell_E = D, \quad \ell_M = D + a(x + D - CP), \quad \ell_Q = D + a(D - CP). \quad (271)$$

For a capped endpoint the exact charge is instead

$$\begin{aligned} \ell_U &= D + A_D + uK - C(A_I + A_D + (u - 1)K) \\ &= d - CA_I - cA_D + Kh_U, \quad h_U = 1 + u - C(u - 1). \end{aligned} \quad (272)$$

These formulae follow by checking the finitely many ordered endpoint pairs. In particular, an earlier immediate outcome already allocated its online processing delay when it appeared, so a current cap contributes only $-Cp$ against that outcome. Against an earlier ordinary deferred outcome it contributes $p - Cp = -cp$, and against an earlier cap it contributes h_U . The same check shows

$$\Gamma_n \leq \sum_i \ell_i; \quad (273)$$

the difference is exactly the sum of the nonnegative own-processing terms allocated by the M -endpoints. Formula (272), rather than a substitution $p = u$ into ℓ_Q , is essential before caps have been put into a prefix.

Threshold monotonicity gives $A_I \geq aI$ and $A_D \geq ad$. Comparing (272) with the hypothetical Q -charge at the same state gives the exact identity

$$\begin{aligned} \ell_U - \ell_Q &= C(aI - A_I) + c(ad - A_D) + K[1 - c(u - 1 - a)] \\ &\leq K[1 - c(u - 1 - a)]. \end{aligned} \quad (274)$$

The reusable four-endpoint bank. The next potential is the parameterized form of the cap-free bank from Section 3.2. The states Z, E, M, Q are the same four ordinary endpoints; only their target coefficient changes from r to c . At $c = r$ the formulas below coincide term for term with the obligatory certificate. Keeping the parameter visible lets us add the finite-cap reserve without proving a second ordinary endpoint theorem. Define

$$\eta = \frac{D - CP}{x}, \quad b = \frac{E}{x}, \quad y = \eta - Cb. \quad (275)$$

For $-1 \leq y \leq 0$, put

$$f_c(y) = \frac{c+2}{4} + \frac{c-2y}{4} \left(\log \frac{c-2y}{c+2} - 1 \right), \quad (276)$$

$$h_c(y) = a_c(y)(1 + y),$$

$$G_c(y, b) = f_c(y) + bh_c(y) + \frac{C}{2}b^2. \quad (277)$$

For $y \leq -1$, set instead

$$G_c(y, b) = g_c(\eta), \quad g_c(\eta) = \frac{(1 + \eta)_+^2}{2C}. \quad (278)$$

The two definitions are C^1 across $y = -1$. Indeed, feasibility implies $0 \leq b \leq 1/C$ there, and both sides have value $Cb^2/2$ and derivatives

$$(G_y, G_b) = (b, Cb).$$

They are also nonnegative: $f_c(-1) = 0$, $f'_c = a_c - 1 \geq 0$, and $h_c, b, g_c \geq 0$.

Use the base potential

$$W_{\text{base}}(x, P, E, D) = xD + x^2 G_c(y, b). \quad (279)$$

In the raw state (x, P, E, D, K) , the endpoint directions are

	x	P	E	D	K
v_Z	-1	0	0	0	0
v_E	-1	0	1	0	0
v_M	-1	1	0	0	0
v_Q	-1	1	0	1	0
v_U	-1	1	0	1	1.

(280)

Let $J_T = \ell_T + \nabla W_{\text{base}} \cdot v_T$, where for the moment $T \in \{Z, E, M, Q\}$. In Region I, direct differentiation and the identities

$$\begin{aligned} -2f_c + (y - C)f'_c + a_c(1 + y) &= 0, \\ 1 - 2f_c + (y - c)f'_c + a_c y &= 0 \end{aligned} \quad (281)$$

give

$$\begin{aligned} \frac{J_Z}{x} &= (-2f_c + yf'_c) + b(-h_c + yh'_c), \\ \frac{J_E}{x} &= b[-h_c + (y - C)h'_c + C], \\ \frac{J_M}{x} &= b[-h_c + (y - C)h'_c + a_c C], \\ \frac{J_Q}{x} &= b[-h_c + (y - c)h'_c + a_c C]. \end{aligned} \quad (282)$$

All four quantities are nonpositive. For the first one,

$$-2f_c + yf'_c = a_c(c - y) - C \leq 0; \quad (283)$$

the right-hand side is zero at $y = -1$ and is nonincreasing because $a'_c(c - y) - a_c \leq 1 - a_c \leq 0$. Also $h_c, h'_c \geq 0$. For the remaining three drifts, observe that

$$h_c + (c - y)h'_c = a_c C + (c - y)(1 + y)a'_c \geq a_c C. \quad (284)$$

The Q -bracket in (282) is therefore nonpositive; the M -bracket is the Q -bracket minus h'_c , and the E -bracket is the M -bracket minus $C(a_c - 1)$.

In Region II, the same calculation gives

$$\begin{aligned} J_Z/x &= J_E/x = -2g_c + \eta g'_c, \\ J_M/x &= -2g_c + (\eta - C)g'_c + 1 + \eta, \\ J_Q/x &= 1 - 2g_c + (\eta - c)g'_c + \eta. \end{aligned} \quad (285)$$

For $-1 \leq \eta \leq 0$, substitution of $g'_c = (1 + \eta)/C$ makes these values $-(1 + \eta)/C$, $-(1 + \eta)/C$, and 0, respectively. For $\eta \leq -1$ their signs are immediate from $g_c = g'_c = 0$. Hence the base bank pays every ordinary endpoint.

The cap reserve. Define

$$Z(c) = 2 + \frac{1}{c} + \frac{1}{2} \log\left(1 + \frac{2}{c}\right), \quad \delta = Z(c) - u. \quad (286)$$

On the mixed curve, (262) says precisely that

$$\delta = \operatorname{artanh} m. \quad (287)$$

For $0 \leq t \leq m$, set

$$\beta(t) = ct [\delta - \operatorname{artanh} t], \quad \mathcal{V}(t) = \int_t^m \beta(s) ds, \quad (288)$$

and extend $\mathcal{V}$ by zero for $t \geq m$. Since $\beta(m) = 0$, this extension is C^1 . With

$$q = \frac{K}{x}, \quad \mu = \frac{q}{1+q}, \quad \mathcal{H}(q) = (1+q)^2 \mathcal{V}(\mu), \quad (289)$$

define the full potential

$$W = xD + x^2(G_c(y, b) + \mathcal{H}(q)). \quad (290)$$

It is nonnegative.

For an ordinary endpoint, K is fixed while x decreases, so the reserve contributes, after division by x ,

$$L_0(q) = -2\mathcal{H}(q) + q\mathcal{H}'(q). \quad (291)$$

A cap also increments K , and its contribution is $L_0 + \mathcal{H}'$. On the active range $\mu \leq m$, differentiation of (289) gives

$$\mathcal{H}' = 2(1+q)\mathcal{V} - \beta(\mu), \quad L_0 = -2(1+q)\mathcal{V} - q\beta(\mu) \leq 0. \quad (292)$$

Thus the reserve can only improve every ordinary drift.

For a cap, feasibility $D \leq P$ and $P \geq K$ gives

$$y = \frac{D - C(P + E)}{x} \leq -\frac{cK}{x} = -cq. \quad (293)$$

While $\mu \leq m$, we have $q \leq m/(1-m) \leq \varphi$ and hence $cq \leq 1$. Monotonicity of a_c , together with (274), yields

$$\frac{\ell_U - \ell_Q}{x} \leq cq \left[\delta - \frac{1}{2} \log(1 + 2q) \right] =: B(q). \quad (294)$$

The change of variables in (289) satisfies

$$\operatorname{artanh} \mu = \frac{1}{2} \log(1 + 2q), \quad (1+q)\beta(\mu) = B(q). \quad (295)$$

Consequently the cap residual is cancelled exactly:

$$L_0 + \mathcal{H}' + B(q) = 0 \quad (\mu \leq m). \quad (296)$$

When $\mu \geq m$, the reserve vanishes. At $q_* = m/(1-m)$, the bracket in (274), evaluated at the largest possible threshold $a_c(-cq_*)$, is zero by (287) and (295). For $q \geq q_*$, (293) only decreases the threshold, so that bracket is nonpositive. The fifth Hamiltonian inequality therefore holds for every state and for arbitrary interleavings of capped and ordinary outcomes.

Initial calibration. Initially $y = b = q = 0$, and

$$f_c(0) = \frac{1}{2} - \frac{c}{4} \log\left(1 + \frac{2}{c}\right). \quad (297)$$

Elementary integration in (288) gives

$$\mathcal{V}(0) = \frac{c}{2}(\delta - m). \quad (298)$$

Indeed,

$$\int_0^m t \operatorname{artanh} t \, dt = \frac{(m^2 - 1) \operatorname{artanh} m + m}{2}.$$

Using (262) and (287), we obtain the exact calibration

$$f_c(0) + \mathcal{H}(0) = f_c(0) + \mathcal{V}(0) = \frac{c}{2}. \quad (299)$$

Thus $W_{\text{initial}} = cn^2/2$.

Finite jobs, and uniformity on the plateau. We spell out the regularity needed to pass from the fluid Hamiltonian to unit job transitions. It is important here to restrict attention to feasible states. In Region I, $y \geq -1$ implies

$$-xy = C(P + E) - D = cP + (P - D) + CE \leq x. \quad (300)$$

All terms in the middle expression are nonnegative. Hence $P/x, D/x, q \leq 1/c$ and $b \leq 1/C$, so every normalized derivative of the Region-I perspective in (279) is bounded on the feasible cone. In Region II the raw base bank is simply

$$xD + \frac{(x + D - CP)_+^2}{2C}, \quad (301)$$

which has bounded one-sided Hessians. The two pieces are C^1 at $y = -1$, as verified above, and the positive-part square is C^1 at $\eta = -1$.

For the reserve $R(x, K) = x^2 \mathcal{H}(K/x)$, on its active cone $0 \leq q \leq q_*$ one has bounded $\mathcal{H}''$, and

$$R_{xx} = 2\mathcal{H} - 2q\mathcal{H}' + q^2\mathcal{H}'', \quad R_{xK} = \mathcal{H}' - q\mathcal{H}'', \quad R_{KK} = \mathcal{H}''. \quad (302)$$

For $q \geq q_*$, $R = 0$; moreover $\mathcal{H}(q_*) = \mathcal{H}'(q_*) = 0$. Thus the join is C^1 with bounded one-sided Hessians. At $x = 0$, set $R = 0$: for $K > 0$ it is already locally zero, while at $(x, K) = (0, 0)$ two-homogeneity gives zero gradient. Altogether W is $C^{1,1}$ on the feasible state cone. On the mixed interval its regularity constant may depend on the fixed value of u . When $c = r$ and $m = 0$, however, the reserve is identically zero and the base bank depends only on the absolute constant r ; its regularity constant is therefore uniform over all $u \geq z_*$. This includes the last transition from $x = 1$ to $x = 0$.

It follows that a unit transition has the Taylor remainder

$$\ell_i + W_{i+1} - W_i \leq O_u(1). \quad (303)$$

The endpoint $E = 0^+$ is interpreted one-sidedly. For a fixed symbolic word, replace all E -coordinates by positive values ϵ . Its state trajectory is unchanged, and the exact finite pair objective, being a finite sum of affine and minimum terms, converges as $\epsilon \downarrow 0$.

Equivalently, retain an $O(\epsilon n^2)$ error and take this limit for each fixed n . Therefore the formal E -Hamiltonian transfers to genuine outcome vectors without an asymptotic loss.

Summing (303), using $W \geq 0$, and applying (299) gives

$$\sum_i \ell_i \leq \frac{c}{2} n^2 + O_u(n). \quad (304)$$

Together with (269) and (273), this yields

$$\text{ALG} - C \text{OPT} \leq -c \binom{n}{2} + \frac{c}{2} n^2 + O_u(n) = O_u(n). \quad (305)$$

At $u = z_*$, one has $c = r$, $m = 0$, and $Z(r) = z_*$. The cap reserve is identically zero, while (299) remains valid. If u is increased above z_* , the only adverse fifth-endpoint bracket $1 - r(u - 1 - a_r(y))$ decreases, and the capped diagonal term in (269) also decreases.

Here is the promised uniform finite calculation. The base bank and its feasible cone contain no u , so there is one constant κ , depending only on r , for which every ordinary or capped transition with $u \geq z_*$ satisfies

$$\ell_i + W_{i+1} - W_i \leq \kappa.$$

Because $W_0 = rn^2/2$ and $W_n = 0$, telescoping gives

$$\sum_i \ell_i \leq \frac{r}{2} n^2 + \kappa n. \quad (306)$$

In the diagonal term of Equation (269), a value below the cap contributes $-r(1 + p_i)$. A capped value contributes at most $1 - ru \leq 0$. With the convention $\bar{p}_i = p_i$ at $u = \infty$, the first formula applies there and the capped endpoint is absent. Hence the diagonal sum is nonpositive uniformly on $[z_*, \infty]$. Combining Equations (269), (273) and (306) now yields

$$\text{ALG} - (1 + r) \text{OPT} \leq -r \binom{n}{2} + \frac{r}{2} n^2 + \kappa n = \left(\kappa + \frac{r}{2} \right) n. \quad (307)$$

Thus, with $B = \kappa + r/2$,

$$\text{ALG} \leq (1 + r) \text{OPT} + Bn \quad \text{for every } u \in [z_*, \infty], \quad (308)$$

and B is independent of u . The estimate is genuinely uniform rather than the result of sending a parameter-dependent remainder to infinity. This proves Theorem 8.10. Finally, because $u > 1$ on the present range, every effective offline length is at least one and $\text{OPT} \geq n(n + 1)/2$. The additive term in (305) vanishes in the size-asymptotic ratio. $\square$

9 Revealing-optimization instance optimality

Fix $u < \infty$. An untouched job may be run raw for the public time u , or tested for one unit; a test reveals $p_i \in [0, u]$, after which the processing operation of length p_i may be delayed. The goal of this section is to prove Theorem 9.1, the rigorous version of the informal revealing-optimization theorem Theorem 1.6.

For a distribution D on $[0, u]$, we define an explicit value $\Phi_u(D)$ in Equation (320). It is obtained by choosing one threshold and minimizing a quadratic over the tested fraction q . For a multiset M , write D_M for its empirical distribution.

Theorem 9.1 (Revealing-optimization instance optimality). *For every fixed $u > 0$ there exist randomized nonanticipating policies $\mathcal{A}_{n,u}^*$, depending only on n, u , and a deterministic sequence $\varepsilon_u(n) \rightarrow 0$ such that every size- n multiset $M \subseteq [0, u]$ has the following properties. For every fixed labeling σ ,*

$$\mathbb{E} \text{cost}(\mathcal{A}_{n,u}^*, \sigma(M)) \leq n^2 \Phi_u(D_M) + \varepsilon_u(n) n^2. \quad (309)$$

Conversely, for every randomized nonanticipating policy $\mathcal{A}$, even one told M , some fixed oblivious labeling σ satisfies

$$\mathbb{E} \text{cost}(\mathcal{A}, \sigma(M)) \geq n^2 \Phi_u(D_M) - \varepsilon_u(n) n^2. \quad (310)$$

One may take

$$\varepsilon_u(n) = O_u\left(n^{-1/6} \sqrt{\log(n+2)}\right). \quad (311)$$

The proof has three layers. First, for a fixed final tested fraction q , a completion envelope reduces every fluid schedule to an ordinary fractional knapsack. Second, an adaptive first-touch trace turns every deterministic policy under a uniform placement into predictable sampling without replacement. One event then controls every touch prefix, so the realized stopping fraction can be substituted pathwise. Finally, a growing grid and a sublinear sample make the result uniform over arbitrary bounded multisets without announcing the bag. The first subsection begins the formal proof.

9.1 The fixed-fraction fluid benchmark

Let D initially have finite support, let $P \sim D$, and put

$$D_0 = \Pr(P = 0), \quad \mu = \mathbb{E}P.$$

A test automatically completes a zero outcome. For a rule $0 \leq g \leq 1$ selecting positive outcomes to process immediately, define

$$A(g) = D_0 + \mathbb{E}[g(P) \mathbf{1}_{P>0}], \quad W(g) = 1 + \mathbb{E}[Pg(P) \mathbf{1}_{P>0}]. \quad (312)$$

Choose g^* maximizing $A(g)/W(g)$, put

$$a = A(g^*), \quad \ell = \mathbb{E}[Pg^*(P) \mathbf{1}_{P>0}], \quad \tau = \frac{1 + \ell}{a}, \quad (313)$$

and interpret the ratio as zero when its numerator vanishes.

Lemma 9.2 (Maximum-density threshold). *There is a maximizer which selects every $p < \tau$, rejects every $p > \tau$, and may split an atom at equality. It satisfies*

$$\mathbb{E}(\tau - P)^+ = 1, \quad \tau(D_0 + \mathbb{E}[g(P) \mathbf{1}_{P>0}]) \leq 1 + \mathbb{E}[Pg(P) \mathbf{1}_{P>0}] \quad (314)$$

for every $0 \leq g \leq 1$.

Proof. If selected mass d at a positive value p is added to a candidate with density ρ , cross multiplication says that the density increases exactly when $1/p > \rho$. Removing mass gives the reverse implication. A maximizer may therefore be chosen as a threshold with boundary $1/\rho = \tau$. The identity $1 + \ell = a\tau$ expands to

$$1 = \tau D_0 + \mathbb{E}[(\tau - P)g^*(P) \mathbf{1}_{P>0}] = \mathbb{E}(\tau - P)^+.$$

Maximality gives $A(g)/W(g) \leq 1/\tau$ for every g ; clearing the positive denominators proves the second inequality in Equation (314). $\square$

Split a boundary atom according to g^* . Let R_i be the residual mass at positive support value p_i . Thus

$$\sum_i R_i = 1 - a, \quad p_i \geq \tau \text{ whenever } R_i > 0. \quad (315)$$

For a fixed final tested fraction $q \in [0, 1]$, consider any prefix of a fluid schedule and write t for tested mass, b for raw-completed mass, and c_i for processed known mass of class i . Its completed mass and work are

$$y = D_0 t + b + \sum_i c_i, \quad x = t + ub + \sum_i p_i c_i, \quad (316)$$

and every feasible prefix satisfies

$$0 \leq t \leq q, \quad 0 \leq b \leq 1 - q, \quad 0 \leq c_i \leq D_i t. \quad (317)$$

Let $H_{D,q}^u(x)$ be the maximum completed mass subject to Equations (316) and (317) and work at most x .

Lemma 9.3 (Fixed- q completion envelope). *The envelope $H_{D,q}^u$ is the divisible-knapsack curve with items*

<i>item</i>	<i>completion capacity</i>	<i>work per completion</i>
<i>maximum-density test module</i>	aq	τ
<i>residual tested class i</i>	qR_i	p_i
<i>raw module</i>	$1 - q$	u

If $\tau < u$, the greedy order is the test module, the residual classes in nondecreasing p_i , and the raw module. If $\tau \geq u$, the all-raw curve pointwise dominates every choice with $q > 0$.

Proof. For a feasible prefix, let y_L, x_L be the completion and work contributed by the tests, tested zeros, and processed selected positive outcomes. Apply Equation (314) to the processed fractions within the first t units of tests. It gives

$$\tau y_L \leq x_L, \quad y_L \leq at.$$

With $\lambda = y_L/a$, replacing this part by λ full test modules preserves completed mass, weakly decreases work, and uses $0 \leq \lambda \leq t \leq q$. Residual class i has capacity at most qR_i , and the raw mass has capacity at most $1 - q$. Thus every feasible prefix maps to the displayed knapsack.

Conversely, while greedy is partway through the test-module item, run the corresponding fraction of tests and process their selected outcomes. Once all q tests have run, every residual capacity is available; the other items can then be processed in greedy order. A partial final item is realized fractionally. The knapsack curve is therefore the closure of the original fluid trajectories. Fractional knapsack orders items by increasing work per completion. In the short-test regime, $\tau \leq p_i \leq u$, which gives the claimed order. In the other regime all test and residual items cost at least u , while one raw item of capacity one costs exactly u . $\square$

Put

$$m_R = \sum_i p_i R_i = \mu - \ell, \quad K_R = \sum_{i,j} \min\{p_i, p_j\} R_i R_j. \quad (318)$$

Integrating the envelope of Lemma 9.3 gives

$$F_{u,D}(q) = (1 + \ell) \left(q - \frac{aq^2}{2} \right) + qm_R(1 - q) + \frac{q^2 K_R}{2} + \frac{u(1 - q)^2}{2}. \quad (319)$$

Indeed, the four terms are respectively the test-module area, residual work delaying the raw block, internal residual SPT pairs, and the homogeneous raw block. Define

$$\Phi_u(D) = \begin{cases} \min_{0 \leq q \leq 1} F_{u,D}(q), & \tau < u, \\ u/2, & \tau \geq u. \end{cases} \quad (320)$$

This is an explicit one-dimensional optimization. In fact

$$F_{u,D}(q) = \frac{u}{2} + Aq + Bq^2, \quad A = 1 + \mu - u, \quad B = -\frac{a(1 + \ell)}{2} - m_R + \frac{K_R}{2} + \frac{u}{2}. \quad (321)$$

For each work budget, convexly combining feasible points for q_1, q_2 gives a feasible point for the corresponding convex combination of the two fractions. Hence $H_{D,q}^u(x)$ is concave in q , its remaining-mass area is convex, and $B \geq 0$. Moreover,

$$\tau < u \iff \mathbb{E}(u - P) > 1 \iff A < 0.$$

Thus the minimizer is

$$q^* = 1 \quad (B = 0), \quad q^* = \min\{1, -A/(2B)\} \quad (B > 0). \quad (322)$$

We have proved pathwise at the fluid level that a policy with realized final tested fraction q has area at least $F_{u,D}(q) \geq \Phi_u(D)$.

9.2 Adaptive policies and the announced lower bound

We now transfer the fluid envelope to a finite policy which may choose labels and its stopping time adaptively. Equal-valued jobs are first distinguished as occurrence tokens. Let J be the public labels, Ω the tokens, $v : \Omega \rightarrow [0, u]$ their values, and let a placement $\sigma : J \rightarrow \Omega$ be a bijection. Fix the private seed of a completing deterministic policy. In its run on σ , list the labels in order of first touch,

$$\ell_1(\sigma), \dots, \ell_n(\sigma),$$

where a first touch is a test or a raw execution, and define

$$R_{\mathcal{A}}(\sigma)_k = \sigma(\ell_k(\sigma)). \quad (323)$$

Lemma 9.4 (Adaptive first-touch bijection). *The map $\sigma \mapsto R_{\mathcal{A}}(\sigma)$ is a bijection from placements to permutations of Ω . Whether touch k is a test or a raw execution is determined by the preceding hidden values $v(R_{\mathcal{A}}(\sigma)_1), \dots, v(R_{\mathcal{A}}(\sigma)_{k-1})$.*

Proof. Suppose two placements have the same occurrence order. A lockstep induction shows that the two runs have the same observable history before every first touch. Between touches, equal histories prescribe the same known-job operations. At the next touch they prescribe the same fresh label and the same test/raw choice. A test then reveals the same next token value; a raw operation lasts the same public time u and reveals no value. Thus the first-touch labels agree at every position. Equality of the occurrence orders now gives equality of the placements on a permutation of all labels. The map is injective between finite sets of the same cardinality $n!$, hence bijective. The same induction stopped after a common value prefix proves predictability. The hidden value of a raw occurrence is extra analytical information and therefore cannot invalidate that conclusion. $\square$

Under a uniform placement, the occurrence order is consequently a uniform permutation. Partition $[0, u]$ into $K + 1$ cells, keeping zero separate, and let D_j be the exact mass of cell j . If X_k is the value at touch k and A_k is the predictable indicator that this touch is a test, the following event is uniform over the policy.

Lemma 9.5 (Predictable urn discrepancy). *Fix $\delta \in (0, 1)$ and put*

$$e_n = \sqrt{8n \log(n(K + 2))}, \quad \Gamma_n = (1 + \delta^{-1})e_n. \quad (324)$$

With probability $1 - o(1)$, simultaneously for every cell j and every touch prefix $m \leq (1 - \delta)n$,

$$\left| \sum_{k \leq m} A_k \mathbf{1}_{X_k \in j} - D_j \sum_{k \leq m} A_k \right| \leq \Gamma_n. \quad (325)$$

If K is polynomial in n , the failure probability is inverse polynomial.

Proof. Hoeffding's inequality for sampling without replacement, followed by a union bound over all prefixes and cells, gives $|N_j(k) - kD_j| \leq e_n$ simultaneously, except with inverse-polynomial probability. While $k \leq (1 - \delta)n$, the remaining proportion $R_{k,j}$ therefore obeys $|R_{k,j} - D_j| \leq e_n/(\delta n)$. Since A_k is predictable,

$$\sum_{k \leq m} A_k (\mathbf{1}_{X_k \in j} - R_{k-1,j})$$

is a martingale with increments bounded by one. Maximal Azuma–Hoeffding, again union-bounded over the cells, places its running maximum below e_n . Adding the predictable drift gives Equation (325). $\square$

Stop the first-touch sequence when only $\delta n + O(1)$ labels remain. In the suffix, replace every future test of an untouched job by raw execution at its first original operation and delete its later known processing. Only $O(\delta n)$ bounded operations change, so

$$C_{\text{original}} \geq C_{\text{edited}} - O_u(\delta n^2 + n). \quad (326)$$

The edited schedule has a fixed final tested fraction q , but we do not condition on its value.

On the single event of Lemma 9.5, consider any operation prefix. Normalize counts and work by n ; let t, b, z, c_j denote tested mass, raw mass, tested-zero completions, and processed known mass from positive cell j . With $\gamma_n = \Gamma_n/n$,

$$z \leq D_0 t + \gamma_n, \quad c_j \leq D_j t + \gamma_n.$$

Set $c'_j = (c_j - \gamma_n)^+$. Then

$$c'_j \leq D_j t, \quad t \leq q, \quad b \leq 1 - q. \quad (327)$$

Raw work is exactly ub , and reducing known mass reduces work. Thus the repaired point is feasible for the empirical envelope at the actual work budget, while completed mass decreases by at most $(K + 1)\gamma_n$. Hence

$$\text{completed fraction at work } x \leq H_{D,q}^u(x) + (K + 1)\gamma_n. \quad (328)$$

Integrating the remaining-mass curve, substituting the realized q only after Equation (328) is established, and restoring Equation (326) gives the fixed-grid lower bound

$$\frac{C_{\text{original}}}{n^2} \geq \Phi_u(D) - O_{u,K}(\gamma_n + \delta + 1/n). \quad (329)$$

For an arbitrary multiset, take

$$K_n = \lfloor n^{1/6} \rfloor, \quad h_n = u/K_n, \quad \delta_n = n^{-1/6}, \quad (330)$$

keep zero separate, and round every positive value upward. If two distributions admit a zero-preserving coupling with pointwise displacement at most h , transporting the same fluid policy changes total known processing work by at most h and hence changes its remaining-mass area by at most h . Therefore

$$|\Phi_u(D) - \Phi_u(D')| \leq h. \quad (331)$$

For the finite repair, raising a processed value to its bin endpoint shifts the envelope horizontally by at most h_n ; deleting γ_n from every cell creates vertical error $(K_n + 1)\gamma_n$. With the parameters above,

$$(K_n + 1) \frac{\Gamma_n}{n} = O\left(n^{-1/6} \sqrt{\log(n+2)}\right). \quad (332)$$

On the bad event we use nonnegativity and $0 \leq \Phi_u \leq u/2$. Averaging Equation (329), using Equations (331) and (332), and then averaging over the private seed proves that every announced policy has uniform-placement cost at least

$$n^2 \Phi_u(D_M) - O_u\left(n^{11/6} \sqrt{\log(n+2)}\right). \quad (333)$$

At least one placement has seed-expected cost at least this average. It is chosen before the seed and is therefore a fixed oblivious labeling. This proves Equation (310).

9.3 A universal policy without the multiset

It remains to attain the benchmark without being told M . We first record the finite cost of one canonical template: select r labels uniformly, test them in random order while immediately processing outcomes in a fixed low prefix, then process the tested residual stock in SPT order and finally run the other $n - r$ jobs raw. If $q = r/n$, direct pair accounting gives, uniformly over the empirical grid histogram and the chosen prefix,

$$\mathbb{E}C_n(\pi) = n^2 \Psi_D(\pi) + O_u(n), \quad (334)$$

where $\Psi_D(\pi)$ is the right-hand side of Equation (319). To verify this, expand the testing phase into one-job terms and relative-order kernels for pairs of tested jobs; expand the residual SPT block into its work against raw jobs and its minimum kernel; and add the exact raw-block cost $u(n - r)(n - r + 1)/2$. For distinct occurrences the finite coefficients

$$\frac{r(r-1)}{n(n-1)}, \quad \frac{r(n-r)}{n(n-1)}, \quad \frac{(n-r)(n-r-1)}{n(n-1)}$$

differ from $q^2, q(1-q), (1-q)^2$ by $O(1/n)$. Every kernel is bounded in terms of u , so fewer than n^2 pair terms create only $O_u(n)$ error.

Use the grid from Equation (330) and take a private random permutation. Test its first

$$k_n = \lceil n^{1/2} \rceil \quad (335)$$

labels, keep positive sample jobs pending, and form their rounded histogram $\hat{D}_n$. Among all grid prefixes containing zero and all $q \in [0, 1]$, minimize the quadratic Equation (319).

Apply the selected template to the remaining labels in their unused random order, then finish the positive sample jobs.

Let D_n and D_n^R be the full and remaining rounded histograms. Hypergeometric coordinate variances and Cauchy–Schwarz give

$$\mathbb{E}\|\hat{D}_n - D_n\|_1 \leq \sqrt{\frac{K_n + 1}{k_n}}, \quad \|D_n^R - D_n\|_1 \leq \frac{2k_n}{n - k_n}. \quad (336)$$

For a fixed template, the quantities a, ℓ, m_R in Equation (319) are bounded one-draw expectations and K_R is a bounded two-draw expectation. Inserting an intermediate product measure shows

$$|\Psi_D(\pi) - \Psi_E(\pi)| \leq 4(u + 1)\|D - E\|_1, \quad (337)$$

uniformly over the grid and template. The same bound holds for the minima. The threshold lemma and completion envelope show that the minimum over this common template family is exactly $\Phi_u(D)$.

Condition on the sample. The learned template is fixed and the unused labels are uniformly ordered, so Equation (334) applies. The usual empirical-minimization sandwich, twice using Equation (337), bounds the main cost by

$$(n - k_n)^2 \left(\Phi_u(D_n^R) + O_u(\|\hat{D}_n - D_n^R\|_1) \right) + O_u(n).$$

The sample and its final tail contribute $O_u(nk_n)$. Now apply Equation (336), replace the remaining histogram by the full one, and use Equation (331). Since

$$h_n = O_u(n^{-1/6}), \quad \sqrt{(K_n + 1)/k_n} = O(n^{-1/6}), \quad k_n/n = O(n^{-1/2}),$$

the resulting bound is

$$\mathbb{E} \text{cost}(\mathcal{A}_{n,u}^*, \sigma(M)) \leq n^2 \Phi_u(D_M) + O_u(n^{11/6}) \quad (338)$$

for every fixed labeling. Together with the announced lower bound Equation (333), this proves Theorem 9.1. The larger logarithmic lower-bound error gives the common sequence in Equation (311).

10 Randomized revealing optimization: the exact four-piece curve

The previous section identified the optimal leading online cost for every fixed bag in the revealing-optimization model. We now compare that cost with the clairvoyant optimum and maximize over all bags. The cap u is fixed, but the algorithm is not told the processing-time multiset. This section proves Theorem 10.1, the rigorous randomized half of the informal curve theorem Theorem 1.7. Let $u_r > 1$ be the unique root of

$$u_r^3 - 6u_r^2 + 5u_r - 1 = 0, \quad u_r = 5.048917339522306 \dots \quad (339)$$

Theorem 10.1 (Randomized revealing-optimization curve). *For every fixed $u > 0$, the optimal randomized size-asymptotic competitive ratio against an oblivious adversary is*

$$R_{\text{rand}}(u) = \begin{cases} 1, & 0 < u \leq 1, \\ \frac{u^3}{u^2 + (u-1)^3}, & 1 \leq u \leq u_r, \\ 1 + \frac{1}{2(\sqrt{u}-1)}, & u_r \leq u \leq 25/4, \\ \frac{4}{3}, & u \geq 25/4. \end{cases} \quad (340)$$

More precisely, there are randomized nonanticipating algorithms $\mathcal{A}_{n,u}$ and numbers $\varepsilon_u(n) \rightarrow 0$ such that every fixed labeled vector $p \in [0, u]^n$ satisfies

$$\mathbb{E} \text{cost}(\mathcal{A}_{n,u}, p) \leq R_{\text{rand}}(u) \text{cost}(\text{OPT}_u, p) + \varepsilon_u(n)n^2. \quad (341)$$

Conversely, for every randomized online algorithm $\mathcal{A}$ and every $\varepsilon > 0$, there are arbitrarily large n and a fixed vector $p \in [0, u]^n$ for which

$$\mathbb{E} \text{cost}(\mathcal{A}, p) \geq (R_{\text{rand}}(u) - \varepsilon) \text{cost}(\text{OPT}_u, p). \quad (342)$$

The curve is continuous and has global maximum

$$\max_{u>0} R_{\text{rand}}(u) = \frac{27 + 6\sqrt{3}}{23} = 1.625752384583185\dots, \quad (343)$$

attained at $u = (3 + \sqrt{3})/2$.

The preceding instance-optimal theorem identifies the best policy on each bag. Here we compare that value with the clairvoyant minimum kernel and maximize over all empirical distributions. A survival-function flattening argument reduces the outer optimization to binary distributions. Two scalar maximizations give the four branches above. A sublinear sample removes advance knowledge of the bag, and binary random labelings give matching Yao lower bounds. The next subsection starts the formal calculation.

10.1 Offline and stationary fluid coefficients

Assume initially that $u > 1$ and write

$$s = u - 1. \quad (344)$$

Let D be a finite distribution on $[0, u]$, let P, P' be independent with law D , and use the survival function

$$H(t) = \Pr(P \geq t), \quad H(t) = 0 \quad (t > u). \quad (345)$$

The convention at atoms does not affect any integral below. Since a clairvoyant job has effective length $1 + \min\{s, P\}$, the layer-cake identity and Lemma 2.1 give the leading offline coefficient

$$\Omega_u(D) := \frac{1}{2} \mathbb{E} \min\{q_u(P), q_u(P')\} = \frac{1}{2} \left(1 + \int_0^s H(t)^2 dt \right), \quad (346)$$

$$\text{OPT}_u(M) = n^2 \Omega_u(D_M) + O_u(n) \quad (347)$$

for every empirical distribution D_M .

We next recall the maximum-density stationary policy in a form suited to the cap. For a processing-time prefix of mass a and first moment m , its completion density is $a/(1+m)$. Choose a maximum-density prefix, include all zeros, and put

$$\tau = \frac{1+m}{a}. \quad (348)$$

The one-point addition/removal test puts every value below τ in the prefix and every value above τ outside it; an atom at equality may be split. Equivalently,

$$\mathbb{E}(\tau - P)^+ = 1, \quad \int_0^\tau H(t) dt = \tau - 1. \quad (349)$$

The equation has a unique solution $\tau \geq 1$.

The stationary policy tests all jobs in a uniform random order, processes a positive prefix outcome immediately after its test, and after the last test processes the residual jobs in SPT order. The following formula is the finite-cap analogue of Lemma 4.2.

Lemma 10.2 (Stationary survival formula). *The doubled leading coefficient of the stationary policy is*

$$\mathcal{T}_u(D) = \begin{cases} \tau + \int_\tau^u H(t)^2 dt, & \tau \leq u, \\ \tau, & \tau > u. \end{cases} \quad (350)$$

Consequently the better of the raw and stationary policies has leading ratio

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s H(t)^2 dt}. \quad (351)$$

Proof. Let $h = 1 - a$ be the residual mass and let K_L be the minimum kernel of two residual draws, with their subprobability masses left unnormalized. The random-order pair calculation gives the stationary coefficient

$$(1+m) \left(1 - \frac{a}{2}\right) + \frac{K_L}{2}. \quad (352)$$

If $\tau \leq u$, the threshold property gives

$$K_L = \tau h^2 + \int_\tau^u H(t)^2 dt, \quad (1+m) \left(1 - \frac{a}{2}\right) = \frac{\tau}{2}(1 - h^2).$$

Their sum is one half of the first line of (350). If $\tau > u$, the whole distribution is in the prefix and (352) equals $\tau/2$. Raw execution has leading coefficient $u/2$, and division by (346) proves (351). $\square$

This is where the instance-optimal theorem enters the curve calculation. In Equation (319), $q = 0$ is the raw policy and $q = 1$ is precisely the stationary maximum-density policy. Hence

$$2\Phi_u(D) \leq \min\{u, \mathcal{T}_u(D)\}, \quad (353)$$

and Equation (351) is an upper bound on the instance-specific ratio $\Phi_u(D)/\Omega_u(D)$. Interior values of q can strictly improve individual bags; the point of the next two subsections is that the two endpoint policies already give the sharp value after maximizing over all D . The binary lower bounds at the end apply to every policy, not only to these endpoints.

10.2 Flattening the survival function

We now maximize (351) over arbitrary D . Suppose first that $1 \leq \tau \leq u$ and put

$$y = \frac{\tau - 1}{\tau}. \quad (354)$$

By (349), y is the average of H on $[0, \tau]$. Since H is nonincreasing, its average on every initial subinterval is at least y . Thus Cauchy–Schwarz gives

$$\int_0^a H(t)^2 dt \geq ay^2 \quad (0 \leq a \leq \tau), \quad (355)$$

and $H(t) \leq y$ almost everywhere for $t > \tau$.

Replace H on $(0, \tau)$ by the constant y and leave its tail unchanged. This remains a nonincreasing survival function, preserves (349) and $\mathcal{T}_u(D)$, and by (355) can only decrease the offline coefficient. Hence it can only increase the ratio. We may therefore assume

$$H(t) = y \quad (0 < t < \tau). \quad (356)$$

If $u - 1 \leq \tau \leq u$, the offline integral lies entirely in the flat part, and $H(t)^2 \leq y^2$ on the remaining tail. It follows that

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s H^2} \leq B_u(\tau) := \frac{\tau + (u - \tau)y^2}{1 + (u - 1)y^2}. \quad (357)$$

The numerator displayed on the right is at most u , so the raw minimum is inactive.

Now suppose $\tau < u - 1$. Put

$$L = u - 1 - \tau, \quad E = \int_{\tau}^{u-1} H(t)^2 dt, \quad G = \int_{u-1}^u H(t)^2 dt. \quad (358)$$

Monotonicity gives

$$0 \leq E \leq Ly^2, \quad G \leq E/L. \quad (359)$$

Consequently

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s H^2} \leq \frac{\tau + \frac{u - \tau}{u - 1 - \tau} E}{1 + \tau y^2 + E}. \quad (360)$$

For fixed τ this is linear-fractional in E , so its derivative has a constant sign. Its maximum on $[0, Ly^2]$ occurs at an endpoint. The two endpoint values are

$$A(\tau) := \frac{\tau}{1 + \tau y^2} = \frac{\tau^2}{\tau^2 - \tau + 1}, \quad (361)$$

$$B_u(\tau) = \frac{\tau + (u - \tau)y^2}{1 + (u - 1)y^2}. \quad (362)$$

They are attained by rectangular survival functions: respectively a binary distribution on $\{0, \tau\}$ and one on $\{0, u\}$.

It remains to cover $\tau > u$. Equation (349) and the zero extension of H give $\int_0^u H = \tau - 1 \geq u - 1$. The average of a nonincreasing function on $[0, u - 1]$ is no smaller than its average on $[0, u]$, and hence

$$\int_0^{u-1} H(t)^2 dt \geq (u - 1) \left(\frac{\tau - 1}{u} \right)^2 \geq \frac{(u - 1)^3}{u^2}. \quad (363)$$

Here the raw policy is chosen, and its ratio is at most $B_u(u)$. We have proved the reduction

$$\sup_D \frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s H(t)^2 dt} \leq \max \left\{ \sup_{1 \leq \tau \leq u-1} A(\tau), \sup_{1 \leq \tau \leq u} B_u(\tau) \right\}, \quad (364)$$

where the first supremum is omitted for $u < 2$. In fact equality holds in (364), because the rectangular survival functions described above attain both families.

10.3 The two scalar maximizations

The first family satisfies

$$A'(\tau) = \frac{\tau(2 - \tau)}{(\tau^2 - \tau + 1)^2}. \quad (365)$$

Thus its maximum is attained at $\tau = \min\{2, u - 1\}$ and equals $4/3$ as soon as $u \geq 3$.

For the second family, substitution of (354) gives

$$B_u(\tau) = \frac{\tau^2 u + 2\tau^2 - 2\tau u - \tau + u}{\tau^2 u - 2\tau u + 2\tau + u - 1}. \quad (366)$$

The sign of its derivative is the sign of

$$g_u(\tau) = (2\tau - 1)^2 - u(\tau - 1)^2. \quad (367)$$

For $u \leq 4$ this is positive on $[1, \infty)$. For $u > 4$, its unique root above one is

$$\tau_*(u) = \frac{u - 2 + \sqrt{u}}{u - 4} = \frac{\sqrt{u} - 1}{\sqrt{u} - 2}, \quad (368)$$

and B_u increases before this point and decreases afterwards. The identity $\tau_*(u) = u$ is equivalent to

$$u^3 - 6u^2 + 5u - 1 = 0. \quad (369)$$

The cubic is strictly increasing beyond its last critical point and changes sign there exactly once; its unique root above one is u_r from (339). Hence

$$\max_{1 \leq \tau \leq u} B_u(\tau) = B_u(u) = \frac{u^3}{u^3 - 2u^2 + 3u - 1} = \frac{u^3}{u^2 + (u - 1)^3}, \quad u \leq u_r, \quad (370)$$

$$= B_u(\tau_*(u)) = 1 + \frac{1}{2(\sqrt{u} - 1)}, \quad u \geq u_r. \quad (371)$$

There is no missing branch from A . On $2 \leq u \leq 3$, clearing the positive denominators in $B_u(u) - A(u - 1)$ and putting $v = u - 2$ leaves

$$v^4 + 3v^3 + 3v^2 + 3v + 3 > 0.$$

On $3 \leq u \leq u_r$, one has $B_u(u) > 4/3$. On $u_r \leq u \leq 25/4$, (371) is at least $4/3$, with equality exactly at $u = 25/4$; afterwards $A(2) = 4/3$ is larger. Combining these observations with (364) proves the announced upper bound

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s H(t)^2 dt} \leq R_{\text{rand}}(u) \quad (372)$$

for every finite distribution D .

The junction statements now follow directly. Equations (369) to (371) give continuity at u_r , and the square-root branch equals $4/3$ at $u = 25/4$. Differentiating (370) shows that its only interior maximum above one occurs at $u = (3 + \sqrt{3})/2$; substitution gives (343).

10.4 Learning the announced policy

The policy used for the upper bound must not know D . The next lemma removes this announcement at lower-order cost.

Lemma 10.3 (Sublinear-sample transfer). *For every fixed $u > 1$, there are randomized nonanticipating policies $\mathcal{A}_{n,u}$ such that every fixed labeled input with empirical distribution D satisfies*

$$\mathbb{E} \text{ALG}_{\mathcal{A}_{n,u}} \leq \frac{n^2}{2} \min\{u, \mathcal{T}_u(D)\} + O_u(n^{11/6}). \quad (373)$$

Proof. Let $k = \lfloor n^{2/3} \rfloor$, $d = \lfloor n^{1/3} \rfloor$, and $\eta = u/d$. Keep zero as a separate category and round each positive value upward to one of the d grid endpoints. The algorithm privately permutes the labels, tests the first k , and forms the sample histogram. On that histogram it compares the raw template with every stationary grid-prefix template. It applies a minimizing template to the remaining labels; in the stationary case, prefix outcomes are processed immediately and the residual tail in SPT order. Sampled positive jobs are inserted in the corresponding known part of the schedule.

Let $\hat{D}$ and D^g be the sample and population grid histograms. Hypergeometric variance and Cauchy–Schwarz over the $d + 1$ categories give

$$\mathbb{E} \|\hat{D} - D^g\|_1 \leq \sqrt{\frac{d+1}{k}} = O(n^{-1/6}). \quad (374)$$

For a fixed prefix, the stationary coefficient is a quadratic polynomial in the category masses, with coefficients bounded in terms of u . It is therefore $O_u(1)$ -Lipschitz in total variation, uniformly over all grid prefixes. The empirical-minimization sandwich loses at most twice the right-hand side of (374). Zero-preserving upward rounding changes every normalized work and pair kernel by $O_u(1/d)$.

Finally, placing the sample before the main random-order template affects at most $O(nk + k^2)$ bounded operation–job pairs. The already tested sample has the same bound when raw is selected. After normalization by n^2 , the total error is

$$O_u \left(\sqrt{d/k} + 1/d + k/n \right) = O_u(n^{-1/6}).$$

This proves (373). The initial private permutation makes the law identical for every fixed public labeling. $\square$

Together with (347) and (372), Lemma 10.3 yields

$$\mathbb{E} \text{ALG}_{\mathcal{A}_{n,u}} \leq R_{\text{rand}}(u) \text{OPT}_u + O_u(n^{11/6}), \quad (375)$$

which is (341) with $\varepsilon_u(n) = O_u(n^{-1/6})$. For $0 < u \leq 1$, raw execution is exactly clairvoyantly optimal, so no learning is required.

10.5 Binary oblivious lower bounds

It remains to show that the two endpoint policies used above cannot be improved in the worst case by a more adaptive algorithm. We first record a binary random-labeling envelope.

Lemma 10.4 (Binary cap envelope). *Fix $u > 1$, $x \in [0, 1]$, and put $z = 1 - x$. Let M_n have $xn + O(1)$ jobs of value u and $zn + O(1)$ zero jobs. For every randomized nonanticipating algorithm, even one told M_n , some fixed labeling satisfies*

$$\frac{\mathbb{E} \text{ALG}}{n^2} \geq \begin{cases} \frac{u}{2} - o_u(1), & x \geq (u-1)/u, \\ \frac{1+x+ux^2}{2} - o_u(1), & x \leq (u-1)/u. \end{cases} \quad (376)$$

Every labeling has

$$\frac{\text{OPT}_u}{n^2} = \frac{1+(u-1)x^2}{2} + O_u(1/n). \quad (377)$$

Proof. First fix a deterministic policy and place the occurrences uniformly on the public labels. List labels in the order in which the policy first touches them, by either a test or a raw execution, and expose analytically also the hidden value of a raw-touched occurrence. The map from placements to this occurrence word is a bijection: equal word prefixes give equal observable histories, hence the same next label and touch kind, and induction recovers the whole placement. The word is therefore a uniform permutation, and the test/raw choice before its next entry is predictable from the preceding word.

Take $\delta_n = n^{-1/4}$. Uniform-prefix hypergeometric concentration, followed by maximal Azuma–Hoeffding for the predictably selected tests, gives one event of probability $1 - o(1)$ on which every first-touch prefix ending with at least $\delta_n n$ untouched labels and containing t tests satisfies

$$|N_0(t) - zt| \leq \Delta_n, \quad \Delta_n = O(\delta_n^{-1} \sqrt{n \log n}) = o(n). \quad (378)$$

Here $N_0(t)$ is the number of tested zeros. The event is simultaneous, so we do not condition on the policy’s adaptive stopping time.

Edit the suffix by running raw every still-untouched job that the original schedule would later test, deleting its later known-processing operation. Only $\delta_n n + O(1)$ jobs change, so the edit changes total completion cost by $o_u(n^2)$. Let q be the final tested fraction of the edited schedule. On (378), every prefix of its completion curve is dominated, up to $o(1)$ vertical error, by the divisible items

item	completion capacity	work per completion
zero-test module	zq	$1/z$
tested cap processing	xq	u
raw jobs	$1 - q$	u

When $z = 0$, the zero-test module is simply omitted. Indeed, t tests complete at most $zt + \Delta_n$ zeros and expose at most $xt + \Delta_n$ cap jobs; raw and known cap processing both use u work per completion. The ordinary divisible-knapsack exchange therefore upper-bounds completed mass at every work value. Integrating the remaining mass gives a lower bound on total completion time.

If $z \leq 1/u$, every displayed item costs at least u per completion, so the normalized area is at least $u/2 - o_u(1)$. If $z \geq 1/u$, the zero-test module comes first. For fixed q its remaining-mass area is

$$F_z(q) = q - \frac{zq^2}{2} + \frac{u}{2}(1 - zq)^2, \quad F'_z(q) = (1 - zq)(1 - uz). \quad (379)$$

It is minimized at $q = 1$, giving $F_z(1) = (1 + x + ux^2)/2$. The suffix edit, vertical repair, and bad event all contribute $o_u(n^2)$.

The argument holds for each deterministic policy. For a randomized policy, fix its seed, average the result over seeds and the finite uniform placement space, and select a placement whose seed-expected cost is at least the joint average. This placement is fixed before the seed and is therefore an oblivious input. Finally, the effective lengths of the two job types are 1 and u , so (377) follows from Lemma 2.1. $\square$

We now select the binary masses. If $1 < u \leq u_r$, take

$$x = \frac{u-1}{u}. \quad (380)$$

This is the boundary $z = 1/u$ in Lemma 10.4, and its online and offline coefficients have ratio

$$\frac{u}{1 + (u-1)((u-1)/u)^2} = \frac{u^3}{u^2 + (u-1)^3}. \quad (381)$$

If $u_r \leq u \leq 25/4$, take

$$x = \frac{1}{\sqrt{u}-1}. \quad (382)$$

The definition of u_r is precisely the point from which $x \leq (u-1)/u$. Since $(u-1)x^2 = 1+2x$, the second line of (376) divided by (377) becomes

$$1 + \frac{x}{2} = 1 + \frac{1}{2(\sqrt{u}-1)}. \quad (383)$$

Irrational masses are approximated by empirical rational frequencies; all rounding and diagonal terms are $O_u(n)$.

For the last branch, the cap distribution is replaced by a fixed bounded one.

Lemma 10.5 (Balanced zero–two lower bound). *For every $u \geq 25/4$ and every randomized algorithm, arbitrarily large even n admit a fixed labeling with $n/2$ jobs of value zero and $n/2$ jobs of value two such that*

$$\mathbb{E} \text{ALG} \geq n^2 - o_u(n^2), \quad \text{OPT}_u = \frac{3}{4}n^2 + O(n). \quad (384)$$

Proof. Use the same trace bijection, predictable-urn event, and suffix edit as in Lemma 10.4. For a realized tested fraction q , the three fluid items now have capacities and costs

item	completion capacity	work per completion
zero-test module	$q/2$	2
tested two processing	$q/2$	2
raw jobs	$1 - q$	u

Every item costs at least two per completion. Thus the completion envelope lies below $\min\{1, t/2\}$ independently of q , whose remaining-mass area is one. This proves the online estimate after the $o_u(n^2)$ repairs and the finite Yao selection. Since $u > 3$, the two effective lengths are one and three; their equally weighted minimum kernel gives offline coefficient $3/4$. $\square$

Equations (381) and (383), together with Lemma 10.5, match (375) in every range. For $0 < u \leq 1$, raw execution is both an online and a clairvoyant optimum. This completes the proof of the exact curve in Theorem 10.1.

Acknowledgements

An initial improvement over the then-current deterministic lower bound for obligatory testing was obtained with the Bolzano open-source multi-agent research system [GHK⁺26]. That preliminary result motivated the present project. Most of the subsequent conjecture generation, proof development, counterexample search, exposition, and Lean formalization was carried out with substantial assistance from OpenAI’s GPT-5.6 through interactive Codex sessions. In particular, the randomized revealing-optimization curve was developed in a separate model run and then integrated with the instance-optimal framework. The author selected the research directions, reviewed and revised the arguments, and takes responsibility for the claims in this paper.

A Machine-checked verification

The accompanying Lean 4 development [dMU21] uses mathlib [The20] and is pinned to version 4.30.0. Its root module imports the complete deterministic, randomized-obligatory, and blind-execution developments. The source contains no `sorry`, `admit`, or project-defined axiom. This section records the precise proof boundary; Appendix D gives the detailed dictionary for the deterministic exact curve. The revealing-optimization instance theorem of Section 9 and its randomized curve in Section 10 are currently analytic only and are not included in the machine-checked claims below. The blind-optimization instance theorem and exact curve of Section 7 are also currently analytic only.

A.1 Deterministic exact ratios

The theorem `SchedulingPaper.exact_ratio_main` packages both directions of the six-piece finite-cap curve and the obligatory limit. Its upper fields contain literal transcript-only strategies, termination proofs, and $O(n)$ certificates; its lower fields quantify over arbitrary deterministic strategy families and return genuine fixed inputs after the formal replay theorem. The formal curve includes all five joins, continuity, the plateau, and the global maximum.

A.2 Randomized obligatory testing

The module `SchedulingPaper.RandomizedFourThirds` exports the operational upper theorem

```
theorem unknownMultiset_operational_expectedCost_le_20378 ... :  
  uniformAverage (physicalSampledRunCost ...) <=  
    4 / 3 * finiteObligatoryOPT p + 20378 * R ^ 7
```

At the concrete parameter choice $n \leq R^4$, this is Equation (104). The proof below this interface includes the maximum-density optimizer, exact and robust $4/3$ certificates, threshold closure for sample-empty bins, finite hypergeometric concentration, the cutoff-32 fallback, sample-first accounting, bad-event excess, and termination of the sampled transcript strategy.

The same module exports `oblivious_iid_binary_lower_online_actualOPT`. It accepts an arbitrary finite family of public-transcript strategies, assumes only that the supplied fuel completes every fair binary input, and returns a fixed binary input whose average cost has coefficient $4n/(3n+5)$. The formal development identifies its denominator

with the literal clairvoyant obligatory optimum. Since $4n/(3n+5) \rightarrow 4/3$, the upper and lower exports give the machine-checked sharp asymptotic value.

The instance-optimal variant in Section 5 replaces the checked cutoff 32 by the slowly growing cutoff from Equation (130). Its full assembly is currently analytic. Lean verifies the shared finite stationary-cost identity, maximum-density algebra, and fluid 4/3 certificate in `RandomizedAnnouncedFluid` and the randomized operational modules. It does not currently export the whole-measure quantile transfer Equation (126) or an end-to-end theorem for Theorem 5.1; those are not included in the machine-checked claim.

A.3 Blind-execution instance optimality

This formalization uses an operation semantics in which completion of a blind job records its realized duration. The trace-bijection and observed-trace modules prove the occurrence-token reveal bijection of Lemma 6.4, its lockstep causality theorem, the predictable test selector, and the exact selected-count and blind-work identities.

The deterministic analytic layer checks the full-module replacement, fractional-knapsack envelope, long-test dominance, revelation repair, horizontal integral shift, formula Equation (148), its quadratic normal form and vertex minimizer, and the three-point witness. The finite layer checks all-grid repair, bounded one- and two-job kernels, histogram stability, the canonical physical operation word, a blind-sample compiler, and the announced adaptive lower bound.

The end-to-end theorem is exported as `boundedUniform_blindPilot_concrete_rate` in `RandomizedOptionalUnknownRates`. In mathematical notation, for $m = \lfloor n^{1/16} \rfloor \geq 2$, every $p \in [0, L]^n$, and every complete announced policy π , it proves

$$\frac{\mathbb{E} \text{cost}(\text{learned}, p)}{n^2} \leq \frac{\mathbb{E} \text{cost}(\pi, p)}{n^2} + \frac{7830(L+1)^2}{m}. \quad (385)$$

The learned algorithm depends only on n, L and its private permutations, not on the multiset. Lean separately proves $\lfloor n^{1/16} \rfloor \rightarrow \infty$ and hence that the displayed error tends to zero. It also discharges the zero-mean case for the same algorithm. This is a slower rate than the exponential-tail analysis in Equation (177), but it proves the same uniform $o_L(n^2)$ theorem using finite second-moment and checkpoint bounds.

The formal statements use finite uniform averages rather than measure-valued random seeds. This is sufficient for the advertised algorithms and Yao distributions, all of which use finite permutations, finite sample choices, and finite tie-breaking coins. Classical real analysis and noncomputable exact minimizers are used in certificates; no floating-point program is extracted.

B Finite rational checkpoint for the cap-free lower bound

The lower bound at $u = \infty$ takes two limits: first the scaling $H \rightarrow \infty$, and then the number K of positive levels tends to infinity. This appendix is not needed for that proof. It gives one exact finite instance showing that the leading-term calculation already predicts the true ratio accurately.

Take

$$K = 1000, \quad \alpha = \frac{9}{8}, \quad \gamma = \frac{679}{1000}, \quad \xi = \frac{K}{\alpha} = \frac{8000}{9}, \quad H = 9000.$$

There are H jobs at each positive level and ξH zero jobs. All multiplicities are integral, and all processing times defined by Equation (19) are rational.

Using exact rational arithmetic in Equations (18) and (22) to (24) gives the leading-coefficient comparison

$$\frac{Q_K}{D_K} = 1.5705287442294085 \dots > 1.5705.$$

The displayed decimal is only for orientation; all comparisons below are between integers or rational numbers.

For the offline schedule, retaining the terms that are linear in H gives

$$\text{OPT}_H = H^2 D_K + H L_{\text{off}}, \quad (386)$$

where

$$\begin{aligned} L_{\text{off}} &= \frac{1}{2} \left(\xi + \sum_{i=0}^{K-1} (1 + p_i) \right) \\ &= \frac{1}{2} \left(\xi + 2K + \frac{K\gamma}{\xi} + \sum_{s=1}^{K-1} \frac{K-s}{\xi+s} \right) = 1656.440800628232 \dots \end{aligned} \quad (387)$$

Again, the middle expression is the exact rational value used in the comparison.

The discrete shortest-prefix calculation behind Equation (16) also retains favorable diagonal terms that vanish after division by H^2 . Before dropping them it gives

$$\text{ALG}_H \geq H^2 Q_K + \frac{H}{2} \left(\xi + \sum_{i=0}^{K-1} p_i + \sum_{i=0}^{K-1} s_i \right) \geq H^2 Q_K.$$

Here the s_i 's are the nonnegative completion fractions from the lower proof. Exact rational arithmetic verifies

$$2000H^2 Q_K > 3141(H^2 D_K + H L_{\text{off}}). \quad (388)$$

Consequently, the fixed instance obtained by the replay argument in Section 3.1 satisfies

$$\frac{\text{ALG}_H}{\text{OPT}_H} \geq \frac{H^2 Q_K}{H^2 D_K + H L_{\text{off}}} = 1.5705286254871234 \dots > \frac{3141}{2000} = 1.5705.$$

Thus the checkpoint includes the finite- H correction; it is not merely a comparison of the two leading coefficients.

C Independent certificate by bubbling caps to a prefix

For completeness, we give a second proof of the quadratic estimate. This argument is also a useful audit of the fifth-endpoint calculation: it never inserts $p = u$ into an ordinary deferred formula before the caps have actually been moved.

Work first in normalized fluid mass. Immediately before an adjacent pair XU , let E_0 be the preceding E -mass, P the preceding substantive mass, D its deferred submass, x the remaining mass, and w the mass of U -endpoints strictly after the pair. Put

$$y = \frac{D - C(P + E_0)}{x}, \quad \eta = \frac{D - CP}{x} = y + C \frac{E_0}{x} \leq 0. \quad (389)$$

Direct differentiation of the exact pair objective (268), including the mutual pair and all future caps, gives the two nontrivial commutators

$$UM - MU = 1 + \left(1 + \eta - \frac{Cw}{x} \right) (y - c) a'_c(y), \quad (390)$$

$$UQ - QU = \left(\eta - \frac{cw}{x}\right)(y - c)a'_c(y). \quad (391)$$

Here $UX - XU$ means “new objective after moving U left” minus the old objective. The second quantity is nonnegative. For the first, if $B = 1 + \eta - Cw/x \leq 0$, its second term is nonnegative. If $B > 0$, then $B \leq 1$, and on $-1 < y \leq 0$

$$\frac{(c - y)B}{c - 2y} \leq 1.$$

Thus (390) is nonnegative as well. Below $y = -1$, the threshold is flat. Direct evaluation also gives

$$UZ - ZU = UE - EU = 1. \quad (392)$$

The suffix terms involving w are crucial: they make every adjacent exchange valid even while other caps remain to the right. Repeated swaps therefore put all caps into one prefix.

After that prefix there is no future cap. Let e, b, d be the preceding E -mass, substantive mass, and deferred substantive mass, and define

$$y = \frac{d - C(e + b)}{x}, \quad v = \frac{e}{x}, \quad h = d - Cb, \quad k = x + d - Cb.$$

The four ordinary commutators are

$$\begin{aligned} ME - EM &= a_c + a'_c(C - y)\frac{k}{x}, & QE - EQ &= 1 - a'_c(C - y)(-y - Cv), \\ MZ - ZM &= a_c - a'_c y \frac{k}{x}, & QZ - ZQ &= 1 - a'_c y \frac{h}{x}. \end{aligned} \quad (393)$$

They are nonnegative. Indeed, in the varying region $h/x = y + Cv \in [y, 0]$ and $k/x \in [0, 1]$. With $t = -y \in [0, 1]$,

$$a'_c(C - y)t = \frac{t(C + t)}{c + 2t} \leq 1, \quad a'_c t^2 = \frac{t^2}{c + 2t} \leq 1; \quad (394)$$

the first inequality is equivalent to $(t - 1)(t + c) \leq 0$. In the flat region $a'_c = 0$. Hence Z, E can be moved to the tail, and a worst word has the form

$$U^\mu \{M, Q\}^* \{Z, E\}^*. \quad (395)$$

On the middle block use the simpler potential

$$\widehat{W}(x, N, D) = xD + x^2 f_c \left(\frac{D - CN}{x} \right).$$

The identity

$$2f_c - yf'_c = \max\{0, -Cf'_c, -Cf'_c + a_c(1 + y), 1 - cf'_c + a_c y\} \quad (396)$$

is a direct rewriting of (281); it pays the entire ordinary continuation. If the cap prefix has mass μ , then after it $x = 1 - \mu$, $N = D = \mu$, and $y = -c\mu/(1 - \mu)$. Its internal cap-pair reward plus its worst continuation is therefore

$$F_{c,u}(\mu) = \frac{2 - c(u - 1)}{2} \mu^2 + \mu(1 - \mu) + (1 - \mu)^2 f_c \left(-\frac{c\mu}{1 - \mu} \right). \quad (397)$$

For $0 < \mu < 1/C$, differentiation gives

$$F'_{c,u}(\mu) = c\mu [Z(c) - u - \operatorname{artanh} \mu]. \quad (398)$$

On the mixed curve the unique maximizer in this range is therefore $\mu = m$, and

$$F_{c,u}(m) = f_c(0) + \frac{c}{2}(\operatorname{artanh} m - m) = \frac{c}{2}. \quad (399)$$

Here $m \leq 1/C$, since $c, m \leq 1/\varphi$. On the flat part $\mu \geq 1/C$,

$$F_{c,u}(\mu) = \mu - \frac{c(u-1)}{2}\mu^2. \quad (400)$$

Its derivative at $1/C$ is

$$\frac{c(1+m) - 1}{C} \leq 0,$$

and only decreases thereafter. Thus $F_{c,u}(\mu) \leq c/2$ on the whole unit interval. At $c = r, m = 0, u = z_*$, the same calculation is maximized at $\mu = 0$, and increasing u only lowers (397).

To make this independent exchange proof finite, fix $\varepsilon > 0$ and first replace the last $\varepsilon n + O(1)$ outcomes by genuine zeros. Earlier states and actions are unchanged, and only $O(\varepsilon n^2 + n)$ pair terms are affected. All cap and ordinary exchanges in the retained prefix then occur with $x \geq \varepsilon$; their displayed fluid commutators have uniform Taylor remainders there. More precisely, after assigning mass $1/n$ to every job, a normalized adjacent exchange has remainder $O_{u,\varepsilon}(n^{-3})$; there are at most n^2 exchanges. At the join $y = -1$, split an exchange at the join and use the two one-sided formulae, whose values agree. The artificial zeros already form the terminal block, so no unubbled suffix cap invalidates (393). The normalized one-job remainder in the continuation-potential calculation is $O_{u,\varepsilon}(n^{-2})$. For fixed ε , first let $n \rightarrow \infty$, and then let $\varepsilon \downarrow 0$. Equations (396)–(399) give

$$\Gamma_n \leq c \binom{n}{2} + o(n^2),$$

which is an independent size-asymptotic certificate for the two upper bounds above.

D Machine-checked deterministic theorem and its correspondence

This appendix records the public theorem exported by the Lean 4 development [dMU21] associated with this manuscript and explains how its formal predicates correspond to the deterministic statements in Theorems 3.1 and 8.1. The development uses the mathlib library [The20] and is pinned to Lean and mathlib version 4.30.0. Its root module imports `SchedulingPaper.MainTheorems`, whose final theorem is reproduced below. Apart from the typographic rendering of Unicode symbols, this is a verbatim excerpt.

```
theorem exact_ratio_main
  : (∀ {u : ℝ}, 0 < u →
    FiniteExactRatioConclusion u (exactCurve u)) ∧
    ObligatoryExactRatioConclusion RStar := by
  constructor
  · intro u hu
    exact finite_exact_ratio hu
  · exact obligatory_exact_ratio
```

The two conclusion records occurring in the theorem are:

```

structure FiniteExactRatioConclusion (u ratio :  $\mathbb{R}$ ) : Prop where
  lower : LowerBound.FixedSizeLowerBound (.finite u) ratio
  upper :
     $\exists$  strategy :  $\forall$  n, Online.Strategy n,
      UpperBound.CompletesByAnalysisFuel (.finite u) strategy  $\wedge$ 
      SizeAsymptoticUpper
        (UpperBound.strategyCost (.finite u) strategy)
        (UpperBound.fixedOfflineCost (.finite u)) ratio

structure ObligatoryExactRatioConclusion (ratio :  $\mathbb{R}$ ) : Prop where
  lower : LowerBound.FixedSizeLowerBound .infinite ratio
  upper :
     $\exists$  strategy :  $\forall$  n, Online.Strategy n,
      UpperBound.CompletesByAnalysisFuel .infinite strategy  $\wedge$ 
      SizeAsymptoticUpper
        (UpperBound.strategyCost .infinite strategy)
        (UpperBound.fixedOfflineCost .infinite) ratio

```

D.1 Objects on the two sides

The following table gives the direct dictionary between the formal objects and the notation of the manuscript.

Lean object	Manuscript object	Meaning
<code>Cap.finite u</code>	finite common upper limit u	processing times lie in $[0, u]$; testing and raw execution are available
<code>Cap.infinite</code>	obligatory testing	processing times are nonnegative and raw execution is unavailable
<code>exactCurve u</code>	$V(u)$	the six branches in Equation (203), including all five verified joins
<code>RStar</code>	R_*	$1 + 2/(z_* - 1)$, where $z_* > 1$ is the unique solution of $z_* - 3 = \log z_*$
<code>fixedOfflineCost</code>	OPT	the clairvoyant shortest-first cost of the model-dependent effective lengths in Equation (2)
<code>strategyCost</code>	ALG	the total completion cost of the operational online transcript

A formal strategy on n jobs is a deterministic function from the public transcript to its next action. A test publishes the revealed processing time; raw execution publishes no hidden value. Thus the strategy has exactly the information available to an online algorithm in the manuscript. The formal action type uses the standard no-idling normalization appropriate when all jobs are released at time zero. Processing times and comparisons are exact real numbers.

D.2 Expansion of the lower and upper fields

For a cap and a candidate value R , `FixedSizeLowerBound` expands to the following statement. For every size-indexed family of deterministic strategies $\mathcal{B}_n$, every $\varepsilon > 0$, and every N , there exist $n \geq N$, a genuine fixed admissible n -job input I , and a finite settled execution such that either the execution leaves a job incomplete or

$$(R - \varepsilon) \text{OPT}(I) \leq \text{ALG}_{\mathcal{B}_n}(I). \quad (401)$$

An incomplete settled run represents infinite cost. In particular, for algorithms that complete all inputs, (401) is the usual lower-bound sequence on arbitrarily large fixed instances. The lower constructions are first described by adaptive revelation oracles, but the formal replay theorem freezes every such interaction into the fixed input quantified here; this is the machine-checked counterpart of Lemma 2.2.

The predicate **SizeAsymptoticUpper** is the epsilon formulation

$$\forall \varepsilon > 0 \ \exists N \ \forall n \geq N \ \forall I, \quad \text{ALG}_{\mathcal{A}_n}(I) \leq (R + \varepsilon) \text{OPT}(I). \quad (402)$$

The accompanying completion field proves that every job is done after the common analysis fuel $2n + 1$, rather than merely assigning a cost to a partial trace. Consequently, Equations (401) and (402) are the two matching operational statements that characterize the optimal value in Equation (10). The Lean theorem packages these two directions; it does not introduce a separate real-valued infimum or `limsup` object.

Before projecting to **SizeAsymptoticUpper**, the checked upper proof constructs a stronger **FixedUpperCertificate**: for the chosen cap it contains a constant $C \geq 0$ such that, for every n and every fixed input,

$$\text{ALG}_{\mathcal{A}_n}(I) \leq R \text{OPT}(I) + Cn. \quad (403)$$

The quadratic lower bounds Equations (8) and (9) then imply (402). For $u \geq z_*$, the separately verified plateau certificate uses the same strategy and the same C , independent of u , including the obligatory limit. This uniformity is stronger than the quantifier order displayed by `exact_ratio_main`, which only exposes one strategy after a cap has been fixed.

D.3 Correspondence with the two deterministic theorems

The first conjunct of `exact_ratio_main` quantifies over every finite $u > 0$. Substituting $R = \text{exactCurve } u = V(u)$, its lower field gives the fixed-instance lower direction of Theorem 8.1, while its upper field supplies a completing strategy and the matching upper direction. The definition of `exactCurve` is literally the six-way case split in Equation (203); the Lean development also proves its joins, continuity, plateau, and global maximum at u_∞ .

The second conjunct substitutes $R = \text{RStar} = R_*$ in the obligatory model. Its lower field gives the lower half of Theorem 3.1; its completing upper strategy together with (403) gives the upper half. The common formal infrastructure also proves that the offline benchmark is the shortest-first schedule of effective lengths and that the transcript cost is exactly its suffix-weighted pair-accounting expression.

D.4 Proof boundary

The root Lean module imports the complete lower- and upper-bound assemblies. The public theorem has no operational interface hypotheses, and the development contains no `sorry`, `admit`, or project-defined `axiom`. It does use `mathlib`'s classical real analysis and noncomputable choices of exact real parameters. Thus it certifies deterministic mathematical strategy terms, not an extracted floating-point implementation.

The theorem recorded here is specifically the size-asymptotic result. It does not assert the pointwise inequality $\text{ALG} \leq R \text{OPT}$ without a remainder. Its checked remainder is the linear term in (403), and is therefore $o(n^2)$. The machine-checked mixed-regime upper bound follows the cap-reserve potential proof; the bubbling argument in Appendix C remains an independent manuscript cross-check.

References

- [AE21] Susanne Albers and Alexander Eckl. Explorable uncertainty in scheduling with non-uniform testing times. In Christos Kaklamanis and Asaf Levin, editors, *Approximation and Online Algorithms: 18th International Workshop, WAOA 2020, Revised Selected Papers*, volume 12806 of *Lecture Notes in Computer Science*, pages 127–142. Springer, 2021. doi:10.1007/978-3-030-80879-2_9.
- [BS26] Felix Buld and Andreas S. Schulz. Scheduling with testing: Competitive algorithms for minimizing the total weighted completion time in the adversarial model, 2026. arXiv:2606.25166.
- [DDN⁺22] Fanny Dufossé, Christoph Dürr, Noël Nadal, Denis Trystram, and Óscar C. Vázquez. Scheduling with a processing time oracle. *Applied Mathematical Modelling*, 104:701–720, 2022. doi:10.1016/j.apm.2021.12.020.
- [DEL24] Konstantinos Dogeas, Thomas Erlebach, and Ya-Chun Liang. Scheduling with obligatory tests. In Timothy Chan, Johannes Fischer, John Iacono, and Grzegorz Herman, editors, *32nd Annual European Symposium on Algorithms (ESA 2024)*, volume 308 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 48:1–48:14, Dagstuhl, Germany, 2024. Schloss Dagstuhl – Leibniz-Zentrum für Informatik. doi:10.4230/LIPIcs.ESA.2024.48.
- [DEMM20] Christoph Dürr, Thomas Erlebach, Nicole Megow, and Julie Meißner. An adversarial model for scheduling with testing. *Algorithmica*, 82(12):3630–3675, 2020. doi:10.1007/s00453-020-00742-2.
- [DKL⁺23] Christoph Damerius, Peter Kling, Minming Li, Chenyang Xu, and Ruilong Zhang. Scheduling with a limited testing budget: Tight results for the offline and oblivious settings. In *31st Annual European Symposium on Algorithms (ESA 2023)*, volume 274 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 38:1–38:15. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2023. doi:10.4230/LIPIcs.ESA.2023.38.
- [dMU21] Leonardo de Moura and Sebastian Ullrich. The Lean 4 theorem prover and programming language. In *Automated Deduction – CADE 28*, volume 12699 of *Lecture Notes in Computer Science*, pages 625–635. Springer, 2021. doi:10.1007/978-3-030-79876-5_37.
- [GCH24] Mingyang Gong, Zhi-Zhong Chen, and Kuniteru Hayashi. Approximation algorithms for multiprocessor scheduling with testing to minimize the total job completion time. *Algorithmica*, 86(5):1400–1427, 2024. doi:10.1007/s00453-023-01198-w.
- [GHK⁺26] Jan Grebík, Pavel Hubáček, Martin Koutecký, Matěj Kripner, Václav Rozhoň, Robert Šámal, and Adrián Zámečník. Bolzano: Case studies in LLM-assisted mathematical research, 2026. URL: <https://arxiv.org/abs/2604.16989>, arXiv:2604.16989.
- [Kah91] Simon Kahan. A model for data in motion. In *Proceedings of the 23rd Annual ACM Symposium on Theory of Computing (STOC 1991)*, pages 265–277. Association for Computing Machinery, 1991. doi:10.1145/103418.103449.

- [KLL⁺26] Bob Krekelberg, Alison Hsiang-Hsuan Liu, Fu-Hong Liu, Prudence W. H. Wong, and Xiao-Ou Zhang. The power of amortization on minimizing total completion time with explorable uncertainty, 2026. Version 2, revised 8 May 2026. [arXiv:2503.13357](#).
- [LL26] Kao-Chuan Liang and Ya-Chun Liang. The completion-threshold framework for obligatory-test scheduling on multiple machines, 2026. Version 2, revised 14 July 2026. [arXiv:2606.02029](#).
- [LLWZ23] Alison Hsiang-Hsuan Liu, Fu-Hong Liu, Prudence W. H. Wong, and Xiao-Ou Zhang. The power of amortization on scheduling with explorable uncertainty. In Jaroslaw Byrka and Andreas Wiese, editors, *Approximation and Online Algorithms: 21st International Workshop, WAOA 2023*, volume 14297 of *Lecture Notes in Computer Science*, pages 90–103. Springer, 2023. doi:10.1007/978-3-031-49815-2_7.
- [LMS19] Retsef Levi, Thomas L. Magnanti, and Yaron Shaposhnik. Scheduling with testing. *Management Science*, 65(2):776–793, 2019. doi:10.1287/mnsc.2017.2973.
- [LMS24] Retsef Levi, Thomas L. Magnanti, and Yaron Shaposhnik. Scheduling with testing of heterogeneous jobs. *Management Science*, 70(5):2934–2953, 2024. doi:10.1287/mnsc.2023.4833.
- [MPT94] Rajeev Motwani, Steven J. Phillips, and Eric Torng. Nonclairvoyant scheduling. *Theoretical Computer Science*, 130(1):17–47, 1994. doi:10.1016/0304-3975(94)90151-1.
- [The20] The mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020)*, pages 367–381. Association for Computing Machinery, 2020. doi:10.1145/3372885.3373824.