

Scheduling with Uniform Tests: Information and Optimization

Václav Rozhoň

August 2026

Abstract

A machine faces many jobs whose lengths are hidden. Spending one unit of time to inspect a job may reveal a short job that should be finished now, or it may reveal nothing useful while every other job waits. When should the machine keep looking, and when should it start working?

We study this question for one machine and total completion time. We consider four natural rules: every job must be tested; a unit-time revealing optimization can replace a raw time u by a shorter revealed length; an untested job can be run blindly for its true hidden length; or a unit-time optimization can shorten it without revealing the new length. Together these models separate the two possible benefits of spending one unit before execution: learning which job to do next and making the job faster.

We give exact asymptotic answers for these models, both as competitive ratios and as instance-optimal guarantees. The resulting algorithms are often quite simple, which may make them useful in practice.

Contents

1	Introduction	3
1.1	Warmup: every job has to be tested	3
1.2	One algorithm for every input	5
1.3	Generalizations of the model	6
1.4	Information without speedup: blind execution	6
1.5	Blind optimization	7
1.6	Revealing optimization	9
1.7	Techniques	11
1.8	Related work	12
1.9	Roadmap	12
2	Models and common preliminaries	13
2.1	Models	13
2.2	Input, information and adversaries	14
2.3	Offline optima and pair accounting	15
2.4	Competitive-ratio conventions	16
2.5	Probabilistic tools	16
3	Fluid benchmarks and instance optimality	18
3.1	Fluid blocks and schedules	19
3.2	The best testing block	20
3.3	Completion curves of finite block collections	21
3.4	The optimal schedule at a fixed tested fraction	22

3.5	From the fluid theorem to finite inputs	24
3.6	A common finite-to-fluid transfer	24
3.7	From an adaptive run to the fluid envelope	25
3.8	Learning and implementing the fluid optimum	27
3.9	Revealing-optimization instance optimality	28
3.9.1	The explicit revealing benchmark	28
3.10	Obligatory-testing instance optimality	29
3.10.1	The explicit obligatory benchmark	30
3.10.2	A learner that does not know the input bound	31
3.10.3	Limits of the theorem	32
3.11	Blind-execution instance optimality	33
3.11.1	The explicit four-block benchmark	34
3.11.2	Why all four blocks can be necessary	35
3.11.3	Why bounded processing times are necessary	36
4	Deterministic revealing optimization and its obligatory endpoint	36
4.1	Upper bounds	38
4.1.1	Raw execution	38
4.1.2	Forced-prefix uniform thresholds	39
4.1.3	Adaptive thresholds for the final two parameter intervals	45
4.2	Lower bounds	53
4.2.1	The binary stopping game and its piecewise lower bound	54
4.2.2	The harmonic construction for short jobs	57
4.2.3	Combining jobs with $p = u$ and a harmonic block	63
4.2.4	Transferring the lower bound when raw execution is no shorter	67
4.3	Assembly of the curve	67
4.3.1	Shape of the candidate curve	67
4.3.2	Exact joins and completion of the proof	68
4.4	The obligatory endpoint	69
5	Randomized revealing optimization: the exact four-piece curve	69
5.1	Offline and stationary fluid coefficients	70
5.2	Flattening the survival function	72
5.3	The two scalar maximizations	73
5.4	Worst-case upper bound from instance optimality	74
5.5	Binary oblivious lower bounds	74
5.6	The randomized obligatory endpoint	76
5.6.1	Normalization of the offline pair identity	77
5.6.2	Stationary prefixes	78
5.6.3	Learning the prefix from a sublinear sample	80
5.6.4	Analysis on good and bad samples	81
5.6.5	Transfer of the zero-two lower bound	82
6	Blind optimization: speedup without revelation	82
6.1	The deterministic worst-case curve	84
6.2	Proof of randomized instance optimality	88
6.3	The exact worst-case curve	89
	Acknowledgements	91

A Auditing the main statements in Lean	91
A.1 The common asymptotic meaning	91
A.2 Obligatory testing: exact competitive ratios	91
A.3 Obligatory-testing instance optimality	92
A.4 Why the obligatory instance assumptions are necessary	93
A.5 Blind execution	94
A.6 Blind optimization: the two worst-case curves	95
A.7 Blind-optimization instance optimality	96
A.8 Revealing optimization: exact competitive-ratio curves	97
A.9 Revealing-optimization instance optimality	98
A.10 Certified numerical summaries	99
A.11 Proof boundary and reproduction	99

1 Introduction

Suppose that we have many jobs waiting for one machine. We want to minimize the sum of the times at which they finish. The length of each job is written on a folded piece of paper. If we knew all the lengths, the right schedule would be clear: we would simply run the jobs from shortest to longest. But the papers are folded, and opening one of them takes one unit of machine time. Should we open another paper, or should we start working?

Opening a paper may reveal a tiny job that we can finish immediately. It may also reveal a huge job that is best left for the end. On the other hand, looking has a cost. If one hundred jobs are still unfinished, one unit spent opening a paper adds one hundred units to the sum of completion times. The machine must therefore balance two uses of its time: learning which jobs are short and completing jobs that are already understood.

Variants of this question have appeared in both stochastic and adversarial scheduling with testing [LMS19, DEMM20, DEL24]. This work improves earlier results and studies new models. We start with the cleanest version, in which every paper has to be opened. We then allow a job to be run without opening its paper. Depending on what this means, we obtain three further variants. The same algorithmic ideas—random sampling, thresholding, and shortest-first execution—give exact asymptotic answers in all four models.

1.1 Warmup: every job has to be tested

Consider the following problem. There are n jobs. Testing job i takes one unit of time and reveals a number $p_i \geq 0$. The job then needs another p_i units of processing; if $p_i = 0$, it completes with the test. For now every job has to be tested: testing is *obligatory*. The formal model is given in Section 2.

We compare an online scheduler with the shortest-first schedule that knows all values p_i in advance. The *competitive ratio* is the online cost divided by this offline optimum. We care about its asymptotic value. Since every job must be tested, even the optimum has cost $\Omega(n^2)$, so we ignore additive terms of size $o(n^2)$.

The obligatory-testing model for total completion time was introduced by Dogeas, Erlebach, and Liang [DEL24]. Their work and a later lower bound of Liang and Liang [LL26] left a gap between the best deterministic upper and lower bounds. They also left the randomized case open. We close both gaps.

Theorem 1.1 (Obligatory testing, informal version of Theorems 4.2 and 5.5). *In the obligatory-testing model, the best deterministic asymptotic competitive ratio is $R_{\text{OT}}^{\text{det}} = 1.570574\dots$. Against an adversary that fixes the input before the algorithm draws its*

random bits, the best randomized ratio is $R_{\text{OT}}^{\text{rand}} = 4/3$. The two optimal algorithms are sketched in algorithms 1 and 2.

Here and below, *shortest first* means processing revealed jobs in nondecreasing order of p_i . The following pseudocode gives a little more detail about the two algorithms. The formulas are not meant to be memorable.

Algorithm 1 Deterministic threshold

- 1: Put $c = 0.570574\dots$ and $R = 1 + c = R_{\text{OT}}^{\text{det}}$.
- 2: Before test $i + 1$, where $0 \leq i < n$, let c_i be the number of positive jobs processed immediately and d_i the number deferred.

- 3: Put

$$y_i = \frac{d_i - R(c_i + d_i)}{n - i}.$$

- 4: Use threshold $T_i = 1$ if $y_i \leq -1$; otherwise use

$$T_i = 1 + \frac{1}{2} \ln \frac{c + 2}{c - 2y_i}.$$

- 5: Test the next job. Process it now if $p \leq T_i$; otherwise defer it.
 - 6: After all tests, finish deferred jobs shortest first.
-

Algorithm 2 Lean-audited learned randomized thresholds

- 1: For the $4/3$ guarantee, set $B = 32$, $d = \lfloor n^{1/4} \rfloor$, $\eta = B/d$, and $k = \lfloor n^{3/4} \rfloor$. For bounded-input instance optimality, set $m = \lfloor n^{1/32} \rfloor$, $B = m - 1$, $d = m^2 - 1$, $\eta = 1/m$, and $k = \lfloor n/m^4 \rfloor$. Test k random jobs and form the rounded empirical distribution $\widehat{D}$ on the resulting d positive bins (plus zero and overflow).
- 2: For each candidate threshold $0 \leq t \leq B$, put

$$a(t) = \widehat{D}([0, t]), \quad m(t) = \int_{[0, t]} p d\widehat{D}(p),$$

and define

$$\widehat{\tau} = \min_{\substack{0 \leq t \leq B \\ a(t) > 0}} \frac{1 + m(t)}{a(t)}.$$

- 3: If $\widehat{\tau} > B$, test every remaining job and then finish all revealed jobs shortest first.
 - 4: Otherwise process the sampled jobs with $p \leq \widehat{\tau}$. Keep testing jobs, processing them immediately when $p \leq \widehat{\tau}$ and deferring them otherwise.
 - 5: After all tests, finish deferred jobs shortest first.
-

Both algorithms follow the same principle: process short jobs as soon as possible and leave long jobs for the end. The difficult part is choosing the threshold. A deterministic algorithm has no representative view of the unseen jobs. Its decisions are therefore more complicated: in fact, its threshold must change with the history. The exact update is given in steps 3 and 4 of algorithm 1. The matching lower bound in Section 4.2.2 uses an adversary that reveals jobs with decreasing positive values of p_i while hiding many zero jobs until the end.

Randomization gives the algorithm a global view of the input. It first tests a sublinear random sample and estimates the distribution of job lengths. Because the labels are privately shuffled, the remaining empirical distribution stays close to the sampled one

throughout the run. The algorithm therefore faces essentially the same distribution at every step and can use one fixed threshold.

The choice of that threshold has a simple interpretation. For a candidate t , a full round of tests together with the immediate processing of outcomes at most t uses work $1 + m(t)$ per original job and completes a fraction $a(t)$ of the jobs. We call this repeated test-and-process action the *testing module* for t . Thus

$$\frac{1 + m(t)}{a(t)}$$

is the work per immediate completion of this testing module; the unit term includes the tests of *all* jobs, including those deferred to the tail. The algorithm minimizes this quantity and uses its optimum $\hat{t}$ as the actual threshold. Equivalently, adding an outcome of length p improves the module exactly when $p < \hat{t}$.

The matching randomized lower bound is correspondingly simple: half of the jobs have length zero and half have length two, and their labels are shuffled. Until the shuffle is uncovered, one unit of work completes only about half a job. Counting the area under the number of unfinished jobs gives the lower bound $4/3$. The formal argument is in Section 5.6.

1.2 One algorithm for every input

Let us return to algorithm 2. The ratio $4/3$ describes only the hardest input. It says much less on an easy one. Suppose, for example, that all processing times are very large compared with the unit test—a plausible regime in many applications. Then the straightforward strategy that first tests everything and subsequently runs the jobs shortest first is nearly optimal. The worst-case guarantee of algorithm 2 still tells us only that its cost is at most $1.33 \dots$ times the optimum.

The growing-cutoff companion in the same pseudocode satisfies a much stronger per-input guarantee: it is *instance-optimal* on every class with a fixed upper bound on job lengths. It has the same sample-learn-fallback structure as the fixed-cutoff $4/3$ algorithm, but the two Lean exports use the distinct parameter sequences displayed above.

Theorem 1.2 (Obligatory-testing instance optimality, informal version of Theorem 3.15). *Fix $L < \infty$. The growing-cutoff algorithm in algorithm 2 has the following property: for every vector $p \in [0, L]^n$ and every online algorithm $\mathcal{A}'$,*

$$\mathbb{E} \text{ALG}_{\mathcal{A}_{\text{grow}}} \leq \mathbb{E} \text{ALG}_{\mathcal{A}'} + o_L(n^2),$$

where both costs are measured on p and include the private shuffle and the algorithms' remaining randomness. The competing algorithm $\mathcal{A}'$ may even be designed specifically for p and told all its entries in advance, though not their shuffled assignment to the public labels.

Here and in the instance-optimal statements below, randomized cost uses the input convention formalized in the model section: the displayed vector is privately shuffled before the algorithm starts.

Both assumptions in the theorem are necessary. Without a fixed bound L there is no randomized instance-optimal algorithm, and without randomization there is no deterministic one; see Theorem 3.18.

Two points about the theorem are worth emphasizing. First, the comparison is made separately on every input vector. We must therefore compete not only with one fixed

online algorithm, but also with algorithms optimized for this particular p . Allowing $\mathcal{A}'$ to know the entries of p makes this strongest possible online benchmark explicit.

Second, the existence of some instance-optimal algorithm is perhaps not surprising. One could imagine a meta-algorithm that samples the input to estimate the empirical distribution of the entries of p and then runs an optimal algorithm for that estimate. The substantive part of the theorem is that no complicated meta-algorithm is needed. The optimal leading-order algorithm for a known input is itself a simple threshold algorithm, so $\mathcal{A}^{\text{grow}}$ has essentially the same form as algorithm 2: sample the input distribution, choose the threshold that completes the most job mass per unit of work, and use it for the rest of the schedule.

1.3 Generalizations of the model

We next weaken the obligatory model in two directions. First, we may allow a job to run without being tested. Second, the unit preliminary action may have two different benefits. It may reveal the true running time p_i before the job is run. It may also be an optimization procedure that replaces a common raw running time u by a shorter time $p_i \leq u$. For example, a generic piece of code solves every task within time u , but an AI agent can spend one unit of time adapting the code to a particular input and make it faster.

These two benefits are independent. An optimization may reveal its new running time p_i , giving the scheduler both a faster job and information about where the job belongs. Or it may leave p_i hidden until the optimized job finishes.

These choices give the following four models.

model	without the unit action	with the unit action	where discussed
obligatory testing	cannot be run	reveals p_i and unlocks the job	Intro 1.1; Sections 3.10, 4.4 and 5.6
blind execution	runs for its hidden length p_i	reveals p_i before execution	Intro 1.4; Section 3.11
blind optimization	runs for raw time u	replaces u by a still-hidden $p_i \leq u$	Intro 1.5; Section 6
revealing optimization	runs for raw time u	reveals p_i and replaces u by p_i	Intro 1.6; Sections 3.9, 4 and 5

The last three rows separate the two benefits of the unit preliminary action. Blind execution studies information without speedup and is related to the stochastic model of Levi, Magnanti, and Shaposhnik [LMS19]. Blind optimization studies speedup without advance information and is related to the model of Damerius et al. [DKL⁺23]. Revealing optimization, introduced by Dürr et al. [DEMM20], contains both benefits.

The three optional models have somewhat different optimal algorithms and input-specific benchmarks, so we state their results separately. Their randomized algorithms nevertheless share one design principle: privately shuffle the jobs, use a sublinear sample to estimate the input distribution, and optimize the rest of the schedule using that information.

1.4 Information without speedup: blind execution

Suppose an untested job can simply be run until it finishes. Its running time is its true hidden length p_i . Testing does not make the job shorter; it only tells us p_i before we decide when to run it. We call this *blind execution*.

Without an upper bound on the job lengths, no randomized algorithm has a finite competitive ratio, even if we ignore an additive $o(n^2)$ term; see Theorem 3.20 for a simple counterexample. For every fixed bound L , however, sampling again gives one algorithm that is optimal on every bounded input vector.

Theorem 1.3 (Blind-execution instance optimality, informal version of Theorem 3.19). *Fix $L < \infty$. There is one randomized algorithm $\mathcal{A}^*$ such that, for every vector $p \in [0, L]^n$ and every online algorithm $\mathcal{A}'$,*

$$\mathbb{E} \text{ALG}_{\mathcal{A}^*} \leq \mathbb{E} \text{ALG}_{\mathcal{A}'} + o_L(n^2).$$

To describe the algorithm, let $\hat{D}$ denote the estimated job-length distribution and put $\hat{\mu} = \int p d\hat{D}(p)$. As in algorithm 2, define

$$a(t) = \hat{D}([0, t]), \quad m(t) = \int_{[0, t]} p d\hat{D}(p), \quad \hat{\tau} = \min_{\substack{t \geq 0 \\ a(t) > 0}} \frac{1 + m(t)}{a(t)}.$$

The optimal algorithm is more subtle than algorithm 2, because running a job blindly can be better than testing it. Sometimes, after the initial sample, the best choice is to run every remaining job blindly. Even when further testing is useful, it need not be useful all the way to the end of the input.

For a simple illustration, suppose that jobs have only two possible lengths. The obligatory-testing algorithm would process short outcomes immediately and defer long ones. Now consider the last untouched job. If it is short, we want to run it next. If it is long, it can still be run next, as the first job in the arbitrarily ordered long tail. Thus in either case the useful action is to run the job; testing it first only adds one unit of work. This is why the value of testing can disappear before every job has been tested.

In general, the algorithm tests only a fraction q of the jobs and uses the threshold $\hat{\tau}$ from algorithm 2 to decide which tested jobs to process immediately. After testing stops, deferred jobs with $\hat{\tau} \leq p < \hat{\mu}$ are processed before the untouched jobs: their known lengths are smaller than the expected duration $\hat{\mu}$ of a blind execution. The untouched jobs are then run blindly, and tested jobs with $p \geq \hat{\mu}$ form the final tail. The optimal fraction q minimizes the explicit quadratic $F_{\text{BE}, \hat{D}}$ from Equation (80).

Algorithm 3 Instance-optimal blind execution

- 1: Shuffle the jobs and run $o(n)$ of them blindly; use their observed lengths to estimate $\hat{D}$ and $\hat{\mu}$.
 - 2: Compute $\hat{\tau} = \min_{t \geq 0, a(t) > 0} (1 + m(t))/a(t)$.
 - 3: If $\hat{\tau} \geq \hat{\mu}$, put $q = 0$. Otherwise choose $q \in \arg \min_{0 \leq z \leq 1} F_{\text{BE}, \hat{D}}(z)$.
 - 4: Test a random q -fraction of the remaining jobs. Whenever a test reveals $p < \hat{\tau}$, process that job immediately; defer all other outcomes.
 - 5: After the tests stop, process the deferred jobs with $\hat{\tau} \leq p < \hat{\mu}$, shortest first.
 - 6: Run every untouched job blindly.
 - 7: Finish the remaining tested jobs shortest first.
-

1.5 Blind optimization

The two remaining models start from the same operation. Every job takes a known raw time u , and one unit of optimization reduces its running time to $p_i \leq u$. They differ only in whether optimization reveals p_i . We call u the *raw-execution cap*, or simply the *cap*.

A clairvoyant scheduler completes a job by the shorter of raw execution in time u and optimization followed by processing in time $1 + p_i$; thus u is also the *offline cap* on the resulting job length. We start with the case in which optimization does not reveal p_i . Before execution, all optimized jobs still look the same.

This is the simpler case. Let us first describe the hardest inputs for both deterministic and randomized algorithms. Roughly a $1/\sqrt{u}$ fraction of the jobs cannot be shortened at all: they have $p_i = u$. The remaining jobs are optimized to length zero. Even if an online algorithm knows these proportions, it cannot distinguish the two types before committing to a job. An optimized job therefore has expected length of order $(1/\sqrt{u})u = \sqrt{u}$, giving total completion cost $\Theta(n^2\sqrt{u})$. The clairvoyant optimum first optimizes the zero-length jobs and then runs the hard jobs raw, paying only $\Theta(n^2)$.

Randomization improves the asymptotic ratio by a factor of two, because a private shuffle prevents the hard jobs from being concentrated at the beginning of the schedule. The exact ratios are as follows.

Theorem 1.4 (Blind-optimization worst-case curves, informal version of Theorems 6.3 and 6.6). *The exact deterministic and randomized asymptotic competitive ratios satisfy*

$$R_{\text{BO}}^{\text{det}}(u) = \sqrt{u} + o(1), \quad R_{\text{BO}}^{\text{rand}}(u) = \frac{\sqrt{u}}{2} + \frac{1}{2} + o(1).$$

Their complete formulas are given in Theorems 6.3 and 6.6.

This poor worst-case behavior is driven by a very specific family of inputs, which again raises the question of whether the worst-case ratio is the right way to evaluate this model. Instance optimality gives a more informative answer. For every fixed u , the simple algorithm in algorithm 4 is instance-optimal. The structure of the optimal algorithm is particularly simple: at the leading n^2 scale, it always either optimizes every job or optimizes none. The algorithm therefore samples the input and chooses between these two options.

Theorem 1.5 (Blind-optimization instance optimality, informal version of Theorem 6.5). *Consider the blind-optimization model with some $u < \infty$. There is one randomized algorithm $\mathcal{A}^*$ such that, for every vector $p \in [0, u]^n$ and every online algorithm $\mathcal{A}'$,*

$$\mathbb{E} \text{ALG}_{\mathcal{A}^*} \leq \mathbb{E} \text{ALG}_{\mathcal{A}'} + o_u(n^2).$$

Algorithm 4 Instance-optimal blind optimization

- 1: Shuffle the jobs, optimize and run $o(n)$ of them, and let $\hat{\mu}$ be the mean observed length.
 - 2: If $1 + \hat{\mu} < u$, optimize and then run every remaining job.
 - 3: Otherwise run every remaining job raw for time u .
-

More precisely, if μ_p is the mean optimized length in p , the best possible leading value is

$$\frac{n^2}{2} \min\{u, 1 + \mu_p\}.$$

The two terms correspond respectively to running every job raw and to optimizing every job. Thus the mean μ_p is the only statistic that the algorithm needs to estimate.

1.6 Revealing optimization

Now assume that optimization also reveals p_i . Thus the unit action buys both information and speed. We call this *revealing optimization*.

The two extreme regimes are easy to understand. If $u \leq 1$, the test alone costs at least as much as raw execution, so we should never test. If u gets very large, raw execution is unattractive, so the problem approaches obligatory testing. The interesting question is what happens between these regimes.

This model was introduced by Dürr et al. [DEMM20]. Their results showed that intermediate values of u are the difficult ones. We determine the exact competitive ratio for every u , for both deterministic and randomized algorithms.

Theorem 1.6 (Revealing-optimization curves, informal version of Theorems 4.1 and 5.1). *For every constant $u > 0$, the exact deterministic and randomized asymptotic competitive ratios $R_{\text{RO}}^{\text{det}}(u)$ and $R_{\text{RO}}^{\text{rand}}(u)$ are the six-piece and four-piece curves derived in Sections 4 and 5, respectively. The deterministic curve reaches $1.87\dots$ and then falls to the obligatory value $1.57\dots$. The randomized curve reaches $1.63\dots$ and eventually becomes $4/3$.*

To describe the deterministic curve, define the following five transition points:

$$u_1^{\text{det}} = 1, \quad u_2^{\text{det}} = 1.86\dots, \quad u_3^{\text{det}} = 3.14\dots, \quad u_4^{\text{det}} = 3.61\dots, \quad u_5^{\text{det}} = 4.50\dots$$

They delimit the six parameter intervals in the piecewise definition of $R_{\text{RO}}^{\text{det}}(u)$.




Figure 1: The three deterministic algorithms, each shown only on the range where it gives the best guarantee.

The deterministic curve is realized by three explicit algorithm families, used on consecutive ranges of u . We describe them here and defer the exact parameters to Section 4.

The first algorithm, $\text{RAW}(u)$, simply fixes an arbitrary job order and executes every job untested for the public raw time u . On its relevant range, its ratio is 1 for $u \leq u_1^{\text{det}}$ and u for $u_1^{\text{det}} \leq u \leq u_2^{\text{det}}$; see Figure 1.

The second algorithm is FORCEDPREFIXUTE , where UTE stands for *uniform-threshold execution*. As specified in algorithm 5, it tests every job and normally processes only outcomes below a fixed threshold. It makes one deliberate exception: every outcome in an

initial $b(u)$ -fraction of the test order is processed immediately. Intuitively, a deterministic adversary may reveal long jobs first and shorter jobs only later. Deferring all early long jobs would create a tail that waits through most of the later tests. The forced prefix limits this risk by testing and immediately processing a controlled initial fraction, at the cost of sometimes processing a long job too early. After the forced prefix, the same threshold is applied to every tested job.

The exact parameter $b(u)$ is defined in Equation (98); it is nonnegative on $[u_2^{\det}, u_3^{\det}]$ and vanishes at $u_3^{\det}$. For $u_3^{\det} \leq u \leq u_4^{\det}$, the zero-prefix specialization gives the formula $1 + 1/\sqrt{u-1}$.

Algorithm 5 FORCEDPREFIXUTE(J, u)

Range: $u_2^{\det} \leq u \leq u_4^{\det}$.

- 1: Fix the test order $1, \dots, n$ and set $Q \leftarrow \emptyset$.
 - 2: Set $\theta_u \leftarrow \min\{1, u-1\}$. If $u_2^{\det} \leq u \leq u_3^{\det}$, set $b \leftarrow b(u)$; for $u_3^{\det} \leq u \leq u_4^{\det}$, set $b \leftarrow 0$.
 - 3: For $i = 1, \dots, n$, test job i and observe p_i .
 - 4: If $i \leq \lfloor bn \rfloor$ or $p_i \leq \theta_u$, process job i immediately; otherwise add it to Q .
 - 5: Process the jobs of Q in nondecreasing revealed processing time.
-

Finally, ADAPTIVETHRESHOLD is the same history-dependent algorithm as algorithm 1. For finite u , only its parameter c changes: on $u_4^{\det} \leq u \leq u_5^{\det}$ it is chosen by the equations in Equation (95), after which it equals the obligatory-testing value $0.570574\dots$. The three algorithms cover the deterministic curve in Figure 1 in that order: RAW first, FORCEDPREFIXUTE in the middle, and ADAPTIVETHRESHOLD for large u . The exact formulas and matching constructions are in Section 4.

For randomized algorithms, private sampling again gives instance optimality with an algorithm similar in shape to algorithm 3. Unlike blind execution, every tested processing time is at most the raw time u , so all tested jobs belong before the untouched raw block.

Theorem 1.7 (Revealing-optimization instance optimality, informal version of Theorem 3.14). *Fix $u < \infty$. There is one randomized algorithm $\mathcal{A}^*$ such that, for every vector $p \in [0, u]^n$ and every online algorithm $\mathcal{A}'$,*

$$\mathbb{E} \text{ALG}_{\mathcal{A}^*} \leq \mathbb{E} \text{ALG}_{\mathcal{A}'} + o_u(n^2).$$

For the randomized curve, write

$$u_1^{\text{rand}} = 1, \quad u_2^{\text{rand}} = 5.048917\dots, \quad u_3^{\text{rand}} = \frac{25}{4} = 6.25.$$

These three points delimit the four parameter intervals in the piecewise definition of $R_{\text{RO}}^{\text{rand}}(u)$. The exact formulas and matching constructions are in Section 5.

Algorithm 6 Instance-optimal revealing-optimization algorithm

- 1: Put $m = \lfloor n^{1/16} \rfloor$. Shuffle the jobs, test $k = \lfloor n/m^2 \rfloor = \Theta(n^{7/8})$ of them, and use the observed lengths to estimate the input distribution.
 - 2: Choose a threshold t maximizing $\hat{a}/(1 + \hat{\ell})$, where $\hat{a} = \hat{D}([0, t])$ and $\hat{\ell} = \int_{[0, t]} p d\hat{D}(p)$.
 - 3: Put $\hat{\tau} = (1 + \hat{\ell})/\hat{a}$. If $\hat{\tau} \geq u$, set $q = 0$; otherwise choose q minimizing $F_{\text{RO}, u, \hat{D}}(q)$ from Equation (57).
 - 4: Test a random q -fraction; finish outcomes below $\hat{\tau}$ at once.
 - 5: Finish all other tested jobs, shortest first.
 - 6: Run every untouched job raw for time u .
-

Notice that the two exact revealing-optimization curves are not monotone. This is not a contradiction. For very small u , testing is useless, while as $u \rightarrow \infty$ the model converges to the obligatory-testing setting of Section 4.4. Both extreme cases are comparatively easy: the online cost can be charged directly to the optimum. Intermediate values are harder because raw execution gives the clairvoyant scheduler additional useful options.

1.7 Techniques

The proofs repeatedly use similar ideas. We summarize the main ones here.

Modules and completion density. At the leading n^2 scale, lower-order rounding effects are negligible, so we can replace numbers of jobs by fractions of the input. The private shuffle keeps the normalized distribution of untouched jobs close to the input distribution throughout all but a negligible final part of the schedule. In the resulting fluid model, testing any amount of job mass therefore reveals the corresponding multiple of one fixed, finitely supported distribution. A schedule becomes a finite sequence of divisible blocks, or *modules*. Each module spends some work and completes some mass; its *completion density* is the completed mass per unit of work. The finite fluid problem and its exact shortest-first optimum are developed in Section 3. The same section proves a common finite-to-fluid transfer, then derives the revealing-optimization, obligatory-testing, and blind-execution instance-optimality theorems. The actual algorithms learn the block parameters from a sublinear pilot and approximate the fluid blocks with finite batches.

Remaining-job and pair accounting. Every unit of machine work delays every job that is still unfinished. Total completion time can therefore be computed by tracking the number of unfinished jobs throughout the schedule. Equivalently, each operation can be charged to the jobs that wait for it, giving the one-job and pairwise cost identities used throughout the paper. For example, if an algorithm tests a fraction q of the jobs, a pair may contain two tested jobs, one tested and one untested job, or two untested jobs. The numbers of such pairs are proportional to q^2 , $q(1 - q)$, and $(1 - q)^2$, respectively. The leading cost is therefore often a quadratic function of q .

From announcement to sampling. Our randomized instance-optimality proofs first study an announced problem: the multiset of processing times is known, but its assignment to the shuffled labels remains hidden. We identify an optimal leading-order template that depends on only a few parameters, such as a threshold and a tested fraction. To remove the announcement, the actual algorithm uses a sublinear prefix of a private random shuffle as a pilot sample and applies the best template for its empirical distribution to the remaining jobs. Concentration and stability show that this changes the total cost by only $o(n^2)$. The same shuffle is useful in lower bounds. Even if an algorithm adaptively chooses the next untouched label, the value attached to it is uniform among the remaining values, so one concentration event can control all relevant prefixes and stopping times.

Extremal reductions. After identifying an instance-specific benchmark, we still have to determine which input maximizes its ratio to the clairvoyant optimum. After a suitable reformulation, pairwise formulas frequently make this extremal problem convex in each processing time, so its maximum is attained at boundary values. In our distributional arguments, flattening and Cauchy–Schwarz play the same role. These reductions often leave a binary input and only a few scalar parameters, such as the fraction of long jobs.

Different constraints and boundary configurations become active for different values of u , however, so several cases must still be analyzed. Their switching points produce the piecewise exact-ratio curves in this paper.

1.8 Related work

The closest papers have already appeared next to the models they introduced. We give a broader comparison here.

Scheduling with tests belongs to optimization with *explorable uncertainty*, where uncertain data can be queried at a cost; the query viewpoint goes back at least to Kahan [Kah91]. Levi, Magnanti, and Shaposhnik study a stochastic scheduling model and derive prefix-testing algorithms followed by an order based on expected processing time per unit weight [LMS19, LMS24]. Our four-block blind-execution algorithm has a similar shape, but our inputs are adversarial, need not converge to a distribution, and are not announced to the universal algorithm.

Dürr et al. [DEMM20] introduced the adversarial revealing-optimization model and obtained deterministic bounds, including the $1.87\dots$ guarantee for binary values. We close the deterministic gap for arbitrary values $p_i \in [0, u]$ and give the exact randomized curve against an oblivious adversary. Related work allows nonuniform test times [AE21, LLWZ23, KLL⁺26], multiple machines [GCH24], or weights [BS26].

Dogeas, Erlebach, and Liang [DEL24] introduced obligatory testing for total completion time. For uniform tests they proved a deterministic upper bound of $1.58\dots$ and the lower bound $\sqrt{2}$. Liang and Liang [LL26] later raised the lower bound to $3/2$. Our matching value is $1.57\dots$, and our randomized result gives the oblivious value $4/3$. More strongly, the resulting fixed-threshold algorithm is asymptotically optimal separately for every bounded input.

The processing-time oracle of Dufossé et al. [DDN⁺22] is close to blind execution, but allows only two possible hidden durations. If no tests are available and processing times are revealed only at completion, one obtains the classical nonclairvoyant model of Motwani, Phillips, and Torng [MPT94], in which the scheduler does not know a job’s length before it completes. These semantics differ from raw execution in revealing optimization, which always lasts the declared time u .

1.9 Roadmap

Section 2 formally defines the four models, their information structures and adversaries, the corresponding offline optima, and our asymptotic competitive-ratio convention.

Section 3 develops all three bounded instance-optimality results in one place. It first defines fluid schedules as finite sequences of divisible blocks and proves that one explicit maximum-density, shortest-first schedule maximizes completed mass after every amount of work. A common transfer theorem then compares this fluid optimum with arbitrary adaptive algorithms on shuffled finite inputs and implements it by learning a finite template. Its specializations give the explicit revealing-optimization, obligatory-testing, and blind-execution benchmarks in Sections 3.9 to 3.11.

Section 4 then proves the exact deterministic revealing-optimization curve. Its multilevel lower-bound instance, in which every job satisfies $1 + p_i < u$ and raw execution is therefore never shorter, is reused both for $u_4^{\text{det}} \leq u \leq u_5^{\text{det}}$ and for $u \geq u_5^{\text{det}}$, while a bookkeeping potential handles jobs with $p_i = u$ in the matching upper bound. Only after the finite curve is complete does Section 4.4 derive the deterministic obligatory ratio using bounds whose constants do not depend on the raw time u .

Section 5 maximizes the revealing benchmark against the clairvoyant optimum and proves the exact four-piece randomized curve. Its final subsection, Section 5.6, transfers the lower bound to the unbounded obligatory endpoint; for the upper bound it truncates very long observations and switches to a safe all-testing schedule when the sample is unreliable. Section 6 studies speedup without revelation, proves the deterministic lower bound using an adversary that hides when its first phase will stop, and obtains the matching two-algorithm curve. It then proves the randomized instance-optimal theorem and maximizes its mean-based benchmark to obtain the exact randomized worst-case curve.

The appendix provides a reader-facing audit of the Lean 4 development. For each main claim it first gives a self-contained English statement, then shows the decisive public Lean signature and explains how its quantified inputs, algorithms, completion premise, cost comparison, and error term map to the statement in the paper. It also records the exact proof boundary where the current formalization covers only part of a paper theorem.

2 Models and common preliminaries

The four models differ in small but consequential ways. This section collects their common building blocks so that the later arguments need not repeat them. We first define the models and adversaries, then develop the offline pair identities and asymptotic conventions, and finally record the probabilistic tools used throughout the paper. The common fluid problem is developed separately in Section 3.

2.1 Models

There are n jobs, all available at time zero, and one machine. Operations are nonpreemptive: once an operation starts, it runs to completion. At most one operation can run at a time. Job j has a hidden value $p_j \geq 0$. The four models differ in what can be done before this value is known.

Obligatory testing OT.

The values p_j are arbitrary finite nonnegative numbers. Every job must first be tested for one unit. The test reveals p_j , after which its processing operation of length p_j may be performed immediately or delayed.

Blind execution BE.

The values p_j are arbitrary finite nonnegative numbers. A job may be run blindly for its actual duration p_j ; the value becomes known only when that execution finishes. Alternatively, the job may be tested for one unit, revealing p_j , and its processing operation may then be delayed. Testing provides information but does not shorten the job.

Blind optimization BO(u).

Here $p_j \in [0, u]$. A job may be run raw for the known time u . Alternatively, one unit of optimization changes its running time from u to p_j without revealing p_j ; the optimized job is subsequently run blindly, possibly after a delay. Its value becomes observable only when that execution finishes. Raw execution reveals no hidden value.

Revealing optimization RO(u).

Again $p_j \in [0, u]$, and raw execution takes the known time u . The other option is a unit test that reveals p_j , followed, possibly after a delay, by processing for p_j . Thus the preliminary action provides both the speedup and information about its size.

The notation BE has no model parameter: blind execution permits arbitrary finite nonnegative processing times. For its bounded instance-optimal theorem we restrict the inputs to $[0, L]$ and make L public. In the last two models, by contrast, u is itself an available running time and hence part of every scheduling decision. We call it the *raw-execution cap*, or simply the *cap*. Since the offline effective length in these models is $\min\{u, 1 + p_j\}$, we call it the *offline cap* u when discussing the clairvoyant benchmark. The number n and any theorem-specific bound L or model parameter u are known to the algorithm.

The completion time C_j is the time at which job j finishes. If a test reveals $p_j = 0$, the job completes at the end of that test. A blind zero job in BE completes as soon as it is selected, whereas an optimized zero job in BO(u) first pays its unit optimization cost. The objective is

$$\sum_{j=1}^n C_j.$$

Obligatory testing can be viewed as the limit of revealing optimization in which raw execution becomes unavailable. We use this only as intuition. The obligatory theorems and their algorithms are proved directly; deriving them from revealing optimization would require arguing about both u and n going to infinity, which we avoid.

We write $\text{ALG}_{\mathcal{A}}(I)$ for the objective value of an online algorithm $\mathcal{A}$ on an instance I , and $\text{OPT}(I)$ for the value of an offline optimum that knows all processing times. We omit algorithm subscripts or instance arguments when they are clear. When a sequence of costs has the form $cn^2 + o(n^2)$, we call c its *leading coefficient*; terms contributed by individual jobs rather than pairs are typically only $O(n)$.

2.2 Input, information and adversaries

For a deterministic algorithm, the adversary may choose the processing-time vector and its assignment to labels with full knowledge of the algorithm. The randomized results use an *oblivious* adversary: the processing-time vector is fixed before the algorithm draws its private random seed.

An input is a vector $p = (p_1, \dots, p_n)$ of job lengths chosen by the adversary. For a randomized algorithm, this vector is then relabeled by a uniform permutation. We abbreviate the expectation over this shuffle and all further private randomness by $\mathbb{E} \text{ALG}_{\mathcal{A}}(p)$. Thus randomized performance depends only on the entries of p , not on their displayed order. We distinguish three levels of information.

- A *clairvoyant offline algorithm* knows the entire labeled vector, including the assignment of processing times to labels. Its minimum cost is $\text{OPT}(p)$.
- An *announced online algorithm* knows the multiset $\{p_1, \dots, p_n\}$, but not its assignment to labels. Thus it knows the input distribution exactly, but not which value belongs to the next job it chooses.
- An *unannounced*, or *fully online*, algorithm initially knows only the public parameters, such as n , L , and u , and receives no input-specific information. Our instance-optimal algorithms belong to this class.

Both online classes are *nonanticipating*: each action may depend on the public parameters, private randomness, and everything revealed so far, but not on a hidden value that the model has not yet exposed. The *transcript* of a run is the resulting ordered sequence of actions and observations.

For some lower bounds it is convenient to describe a finite distribution over inputs before selecting one fixed input. If every deterministic algorithm has average payoff at least L under this distribution, then the same is true after conditioning on the private seed of a randomized algorithm. Averaging over that seed shows that at least one fixed input in the support gives the randomized algorithm expected payoff at least L . This is the finite one-sided form of Yao's principle [Yao77].

We write

$$D[p] := \frac{1}{n} \sum_{i=1}^n \delta_{p_i} \quad (1)$$

for the empirical distribution of the input, where δ_x denotes one unit of probability concentrated at x . It is unchanged by the shuffle.

Our instance-optimality results compare a single unannounced algorithm against all announced online algorithms on this shuffled input. This is ordinary instance optimality when the input is viewed as a multiset, and order-oblivious instance optimality when it is viewed as an ordered vector; see [ABC17, NRTT24].

2.3 Offline optima and pair accounting

The offline optimum knows all values p_j . It therefore chooses the shorter available way to complete each job and orders the resulting job blocks shortest first. Associate with job j the model-dependent *effective length*

$$\lambda_j := \begin{cases} 1 + p_j, & \text{OT,} \\ p_j, & \text{BE,} \\ \min\{u, 1 + p_j\}, & \text{BO}(u) \text{ or RO}(u). \end{cases} \quad (2)$$

Lemma 2.1 (Effective lengths and the offline pair identity). *Fix any one of the four models and let $\lambda_{(1)} \leq \dots \leq \lambda_{(n)}$ be its effective lengths from Equation (2). A clairvoyant (fully informed) optimum completes the jobs in this order and has value*

$$\text{OPT} = \sum_{k=1}^n \sum_{i=1}^k \lambda_{(i)}, \quad (3)$$

$$= \sum_i \lambda_i + \sum_{i < j} \min\{\lambda_i, \lambda_j\}, \quad (4)$$

$$= \frac{1}{2} \left(\sum_i \lambda_i + \sum_{i,j} \min\{\lambda_i, \lambda_j\} \right). \quad (5)$$

In OT it tests every job. In BE it runs every job blindly. In BO(u) and RO(u) it uses raw mode when $u \leq 1 + p_j$ and the unit preliminary action followed by processing otherwise.

Proof. Let $C_{[1]} \leq \dots \leq C_{[n]}$ be the ordered completion times of any feasible schedule. Completing job j consumes at least λ_j units of work belonging to that job. Hence, by time $C_{[k]}$, the machine has performed the required work of some k jobs, and

$$C_{[k]} \geq \sum_{i=1}^k \lambda_{(i)}.$$

Choose the mode specified in the lemma, keep each resulting block contiguous, and execute the blocks in nondecreasing order of λ_j . This schedule is feasible in the relevant model and

attains equality for every k . Summing over k and then collecting the contribution of each unordered pair gives the first two formulas. Separating the terms with $i = j$ from the two orientations of every pair with $i \neq j$ gives the symmetric form. $\square$

For later pair calculations, if $X \subseteq [n]$, we write

$$\text{SPT}(X) := \sum_{i \in X} p_i + \sum_{\substack{i, j \in X \\ i < j}} \min\{p_i, p_j\} \quad (6)$$

be the exact cost of processing that block in shortest-processing-time (SPT) order. Thus $\text{SPT}(X)$ is the processing-time part of the offline pair formula when the jobs in X are executed without preliminary actions.

2.4 Competitive-ratio conventions

Every asymptotic statement in this paper allows a uniform additive $o(n^2)$ term. Formally, an algorithm family $\mathcal{A}$ has asymptotic competitive ratio at most R if there is a deterministic sequence $\varepsilon(n) \rightarrow 0$ such that every n -job instance satisfies

$$\mathbb{E} \text{ALG}_{\mathcal{A}}(I) \leq R \cdot \text{OPT}(I) + \varepsilon(n)n^2. \quad (7)$$

For a deterministic algorithm the expectation is omitted. The optimal ratio in a model is the infimum of such R over all admissible algorithm families. We also call this the *size-asymptotic* ratio to emphasize that the model parameters are fixed while the number of jobs n tends to infinity.

In obligatory testing every effective length is at least one, and in either optimization model it is at least $\min\{u, 1\}$. Hence Equation (3) gives $\text{OPT} = \Omega(n^2)$ in these models, and the definition above is equivalent to the usual worst-case ratio after taking $n \rightarrow \infty$. This equivalence fails for blind execution, where OPT can be $o(n^2)$ in the special case where most of the jobs have length zero. Accordingly, Theorem 3.19 makes the absolute statement $\mathbb{E} \text{ALG} = n^2 \Phi_{\text{BE}}(D[p]) + o_L(n^2)$, where the benchmark Φ_{BE} is defined in Section 3.11.1. It does not claim a bounded ratio to the clairvoyant optimum when most jobs have length zero.

2.5 Probabilistic tools

We collect here the concentration estimates used in the randomized sections. All of them are finite-space statements. We use without further mention the union bound, Markov's inequality $\mathbb{P}(Y \geq t) \leq \mathbb{E}Y/t$ for $Y \geq 0$, and the tail identity $\mathbb{E}Y = \int_0^\infty \mathbb{P}(Y \geq t) dt$.

We begin with Chernoff bounds for sums of independent Bernoulli variables.

Lemma 2.2 (Chernoff bounds). *Let $X = \sum_{i=1}^m X_i$, where the X_i are independent Bernoulli random variables, and write $\mu = \mathbb{E}X$. For every $\delta > 0$,*

$$\mathbb{P}(X \geq (1 + \delta)\mu) \leq \left(\frac{e^\delta}{(1 + \delta)^{1+\delta}} \right)^\mu \leq \exp\left(-\frac{\delta^2 \mu}{2 + \delta} \right). \quad (8)$$

For $0 \leq \delta \leq 1$,

$$\mathbb{P}(X \leq (1 - \delta)\mu) \leq \exp\left(-\frac{\delta^2 \mu}{2} \right). \quad (9)$$

These are the standard exponential-moment bounds of Chernoff [Che52].

Most of our samples are drawn from a fixed input without replacement. The next lemma gives both a tail bound and the finite-population variance used for such samples.

Lemma 2.3 (Sampling without replacement). *Let $x_1, \dots, x_N \in [0, 1]$, let $X_1, \dots, X_N$ be a uniformly random permutation of these numbers, and put $\bar{x} = N^{-1} \sum_i x_i$. For $1 \leq k \leq N$ and $t \geq 0$,*

$$\mathbb{P} \left(\left| \sum_{i=1}^k X_i - k\bar{x} \right| \geq t \right) \leq 2 \exp \left(-\frac{2t^2}{k} \right). \quad (10)$$

Moreover, if $\hat{x}_k = k^{-1} \sum_{i=1}^k X_i$ and $\sigma^2 = N^{-1} \sum_i (x_i - \bar{x})^2$, then, for $N > 1$,

$$\text{Var}(\hat{x}_k) = \frac{N-k}{k(N-1)} \sigma^2 \leq \frac{1}{4k}. \quad (11)$$

Proof. The estimate Equation (10) is Hoeffding's sampling-without-replacement inequality [Hoe63, Section 6]. The variance identity is the elementary finite-population correction. $\square$

Applying the scalar variance bound to class indicators gives a concise estimate for an empirical histogram.

Lemma 2.4 (Histogram error without replacement). *Partition a population of N occurrences into d classes. Let p be the vector of class frequencies in the population and let $\hat{p}$ be the corresponding frequency vector among the first k occurrences in a uniformly random permutation. Then*

$$\mathbb{E} \|\hat{p} - p\|_1 \leq \sqrt{\frac{d}{k}}. \quad (12)$$

Proof. For a class of mass p_j , Equation (11) and Cauchy–Schwarz give $\mathbb{E} |\hat{p}_j - p_j| \leq \sqrt{p_j/k}$. Summing over the classes and using $\sum_j \sqrt{p_j} \leq \sqrt{d}$ proves the claim. $\square$

An adaptive algorithm may stop at a data-dependent time. We therefore also need one event that controls every class at every possible prefix.

Lemma 2.5 (Uniform prefix histogram bounds). *In the same population, let $H_{k,j}$ count class j among the first k entries of a uniformly random permutation. Then*

$$\mathbb{P} \left(\max_{0 \leq k \leq N} \max_{1 \leq j \leq d} |H_{k,j} - kp_j| \geq t \right) \leq 2d(N+1) \exp \left(-\frac{2t^2}{N} \right), \quad (13)$$

$$\mathbb{E} \max_{0 \leq k \leq N} \max_{1 \leq j \leq d} |H_{k,j} - kp_j| \leq C \sqrt{N \ln(2d(N+1))} \quad (14)$$

for a universal constant C .

Proof. Apply Equation (10) to every binary class indicator and union-bound over k and j to obtain Equation (13). Integrating that tail bound proves Equation (14). $\square$

Predictable sampling introduces dependent increments rather than a fixed prefix. The following maximal inequalities provide the needed uniform control.

Lemma 2.6 (Martingale maximal inequalities). *Let $(M_k, \mathcal{F}_k)_{k=0}^N$ be a real martingale with $M_0 = 0$; that is, $\mathcal{F}_k$ records the information available by time k and $\mathbb{E}[M_k | \mathcal{F}_{k-1}] = M_{k-1}$. If $\mathbb{E}|M_N|^2 < \infty$, then Doob's inequality gives*

$$\mathbb{E} \max_{0 \leq k \leq N} |M_k|^2 \leq 4\mathbb{E}|M_N|^2. \quad (15)$$

If instead $|M_k - M_{k-1}| \leq b_k$ almost surely for deterministic b_k , then maximal Azuma–Hoeffding gives, for every $t \geq 0$,

$$\mathbb{P} \left(\max_{0 \leq k \leq N} |M_k| \geq t \right) \leq 2 \exp \left(- \frac{t^2}{2 \sum_{k=1}^N b_k^2} \right). \quad (16)$$

We use the classical forms due to Doob [Doo53] and Azuma [Azu67]; the maximal version of the latter follows by stopping the usual exponential bound when the displayed threshold is first crossed.

The final estimate allows predictable weights and is uniform over prefixes.

Lemma 2.7 (Predictable sampling from a random permutation). *Let $X_1, \dots, X_N$ be a uniform random permutation of fixed vectors $x_1, \dots, x_N \in [0, B]^d$, and let $\bar{x} = N^{-1} \sum_i x_i$. Suppose $A_k \in [0, 1]$ is chosen using only the previous draws, before X_k is exposed. Then for $\delta, \eta \in (0, 1)$, with probability at least $1 - \eta$, simultaneously for every coordinate ℓ and every $m \leq (1 - \delta)N$,*

$$\left| \sum_{k \leq m} A_k (X_{k,\ell} - \bar{x}_\ell) \right| \leq (1 + \delta^{-1}) E_{N,d,B,\eta}, \quad E_{N,d,B,\eta} := B \sqrt{2N \ln \left(\frac{4d(N+1)}{\eta} \right)}. \quad (17)$$

The statement remains valid after conditioning on independent private randomness used to choose the predictable sequence.

Proof. For each coordinate, apply Equation (10) after scaling by B , and union-bound over coordinates and prefix lengths. Except with probability at most $\eta/2$, every centered prefix sum has absolute value at most $E = E_{N,d,B,\eta}$. Before draw $k \leq (1 - \delta)N$, the mean $\tilde{x}_{k-1,\ell}$ of the remaining population therefore satisfies

$$|\tilde{x}_{k-1,\ell} - \bar{x}_\ell| \leq \frac{E}{\delta N}.$$

For each ℓ ,

$$M_{m,\ell} = \sum_{k \leq m} A_k (X_{k,\ell} - \tilde{x}_{k-1,\ell})$$

is a martingale with increments bounded by B . The maximal Azuma–Hoeffding inequality Equation (16), followed by a union bound over ℓ , puts all $|M_{m,\ell}|$ below E except with probability at most $\eta/2$. The accumulated difference between the remaining-population mean and the original mean is at most E/δ , which proves Equation (17). Conditioning and then averaging proves the final assertion. $\square$

3 Fluid benchmarks and instance optimality

The randomized instance-optimality results all have the same leading-order explanation. Fix an input vector and privately shuffle its values among the public job labels. Uniformly over predictable prefixes, the cumulative number of encountered jobs of each type differs

by only $o(n)$ from the prefix length times its full-input proportion. Thus, if a fraction t of the jobs has been tested, the revealed mass of type p_i is approximately $tD[p](\{p_i\})$. As long as a nonnegligible fraction remains, the normalized composition of the untouched jobs is consequently still close to $D[p]$.

The fluid model makes this approximation exact. It replaces n jobs by one unit of divisible job mass and divides machine work by n . Testing mass 0.01, for example, reveals exactly a 0.01 copy of the input distribution. This removes the random fluctuations and the rounding of job counts, leaving a deterministic scheduling problem for the leading n^2 term of total completion time.

Despite the name, a fluid schedule will not be a policy that makes a new decision at every real time. The distribution below has finitely many job types, and a schedule is a finite sequence of divisible blocks. Within one block, work and completions accrue proportionally. This linear traversal is the limit of dividing the block into many finite batches; the later subsections justify the corresponding approximation for actual jobs. The fluid optimization developed first contains no sampling or limiting argument: it solves the finite-dimensional problem exactly when the distribution is known. Only afterward do we transfer that solution to shuffled finite inputs.

The same problem covers three models. We prescribe a total tested fraction q and let v be the work per job in a completion mode that does not reveal the processing time in advance. Obligatory testing has $q = 1$, revealing optimization has $v = u$, and blind execution has $v = \int p dD(p)$. Blind optimization is different because its preliminary operation does not reveal the processing time, so it is not represented by the testing blocks below.

3.1 Fluid blocks and schedules

We first define the fluid problem operationally. This makes clear both what an arbitrary fluid schedule may do and why its state remains finite. Let

$$D = \sum_{i=0}^m d_i \delta_{p_i}, \quad 0 = p_0 < p_1 < \dots < p_m, \quad \sum_{i=0}^m d_i = 1, \quad (18)$$

where d_0 may be zero. Fix $q \in [0, 1]$ and $v \geq 0$.

A fluid schedule is a finite sequence of the following operations.

Testing block of mass z with selector g .

The vector $g = (g_1, \dots, g_m) \in [0, 1]^m$ specifies which fraction of each positive type is processed directly after being revealed. The block tests mass z , completes all revealed zero jobs and the selected positive jobs, and places the remaining revealed positive jobs in deferred queues. It uses work

$$z \left(1 + \sum_{i=1}^m p_i d_i g_i \right)$$

and completes mass

$$z \left(d_0 + \sum_{i=1}^m d_i g_i \right).$$

It adds $z d_i (1 - g_i)$ to the deferred queue of type i . The total mass of all testing blocks may not exceed q .

Deferred-processing block of type i .

Processing mass z from the currently available deferred queue of type i uses work $p_i z$ and completes mass z .

Alternative-completion block.

Completing mass z in the other mode uses work vz and completes mass z . The total mass of these blocks may not exceed $1 - q$.

The schedule is complete if it tests total mass q , uses the other mode on mass $1 - q$, and eventually empties every deferred queue. A positive-work block is traversed linearly; a zero-work block completes its mass at once. Consequently every finite fluid schedule has a piecewise-linear completion curve $C(x)$, the mass completed after x units of work.

The area under the unfinished mass is its total completion cost:

$$\text{Area}(C) := \int_0^\infty (1 - C(x)) dx. \quad (19)$$

Indeed, if a mass element z completes after work $T(z)$, the elementary layer-cake identity gives

$$\int T(z) dz = \int_0^\infty \text{mass}\{z : T(z) > x\} dx = \text{Area}(C).$$

Thus continuous work is only a convenient coordinate for drawing the completion curve; the schedule itself makes finitely many block decisions.

3.2 The best testing block

We next determine which revealed positive types should be completed inside a testing block. For the finite fluid model, all integrals below are finite sums under Equation (18). The density lemma is no more difficult for an arbitrary finite-mean distribution, so we state it in that form for later use.

For a selector g , let

$$a_D(g) := D(\{0\}) + \int_{(0,\infty)} g(p) dD(p), \quad w_D(g) := 1 + \int_{(0,\infty)} pg(p) dD(p). \quad (20)$$

These are the completed mass and work when one unit of mass is tested. Their ratio $a_D(g)/w_D(g)$ is the block's *completion density*.

For example, suppose half the mass has length zero and half has length one. Tests alone complete mass $1/2$ using work 1 . Processing the length-one outcomes as well completes the full mass using work $3/2$, improving the completion density from $1/2$ to $2/3$. For longer positive outcomes the same decision can reduce the density. At the optimum, a positive type is included precisely when its length is below the work per completion of the resulting block.

Lemma 3.1 (Maximum-density testing block). *Let D be a probability distribution on $[0, \infty)$ with finite mean. There is a unique $\tau_D \geq 1$ satisfying*

$$\int_{[0,\infty)} (\tau_D - p)^+ dD(p) = 1. \quad (21)$$

Choose g_ to select every positive $p < \tau_D$, reject every $p > \tau_D$, and select any fraction of the mass at $p = \tau_D$. Write $a_* = a_D(g_*)$ and $w_* = w_D(g_*)$. Then*

$$\frac{a_*}{w_*} = \max_g \frac{a_D(g)}{w_D(g)} = \frac{1}{\tau_D}, \quad \tau_D a_D(g) \leq w_D(g) \quad \text{for every } g. \quad (22)$$

Proof. The function $h(t) = \int (t - p)^+ dD(p)$ is continuous and nondecreasing, satisfies $h(0) = 0$, and tends to infinity because $h(t) \geq t - \int p dD(p)$. Once positive it is strictly increasing, so $h(t) = 1$ has a unique solution. The inequality $h(t) \leq t$ gives $\tau_D \geq 1$.

For an arbitrary selector g ,

$$\begin{aligned} \tau_D a_D(g) - \int_{(0,\infty)} p g(p) dD(p) &= \tau_D D(\{0\}) + \int_{(0,\infty)} (\tau_D - p) g(p) dD(p) \\ &\leq \int_{[0,\infty)} (\tau_D - p)^+ dD(p) = 1. \end{aligned}$$

This is the second inequality in Equation (22). The threshold selector g_* attains equality because it keeps exactly the positive terms; selecting mass at $p = \tau_D$ changes both sides equally. Hence $w_* = \tau_D a_*$ and g_* has maximum density $1/\tau_D$. $\square$

For the rest of this section, return to the finite distribution in Equation (18). The positive mass not completed in the maximum-density testing block forms the finite residual measure

$$d\nu(p) = (1 - g_*(p)) \mathbf{1}_{p>0} dD(p), \quad \nu(\{0\}) = 0. \quad (23)$$

It has total mass $1 - a_*$ and support in $[\tau_D, \infty)$. Thus every deferred type has at least as much work per completion as the testing block.

3.3 Completion curves of finite block collections

Once testing activity has been compressed into one block, only an ordering problem remains. The next elementary lemma is the divisible version of the shortest-processing-time rule.

Represent a finite block of completion mass c_k and work v_k per completion by $c_k \delta_{v_k}$. For a collection $\mathcal{M} = \sum_k c_k \delta_{v_k}$ of total mass one, define

$$K_{\mathcal{M}}(x) := \max \left\{ \sum_k z_k : 0 \leq z_k \leq c_k, \sum_k v_k z_k \leq x \right\}. \quad (24)$$

This finite fractional-knapsack problem asks for the largest mass that the blocks can complete using work x . Its value as a function of x is their completion curve. We also write

$$\text{SPT}(\xi) := \frac{1}{2} \iint \min\{v, w\} d\xi(v) d\xi(w). \quad (25)$$

Lemma 3.2 (Shortest-first order for divisible blocks). *The curve $K_{\mathcal{M}}$ is realized by processing the blocks in nondecreasing order of v_k , and*

$$\text{Area}(K_{\mathcal{M}}) = \text{SPT}(\mathcal{M}). \quad (26)$$

Proof. Consider two consecutive blocks with masses c, c' and work per completion $v \leq v'$. Putting the v -block first changes the area, relative to the opposite order, by $cc'(v - v') \leq 0$. Repeated neighboring exchanges sort all blocks by nondecreasing work per completion.

In a block of mass c_k and work per completion v_k , unfinished mass decreases linearly. Once the blocks are sorted, its area contribution is

$$v_k c_k \left(1 - \sum_{h < k} c_h - \frac{c_k}{2} \right).$$

Summing these terms and collecting each unordered pair gives Equation (26). $\square$

3.4 The optimal schedule at a fixed tested fraction

We can now state the common optimality theorem. The proof compares complete curves, not merely their final areas: after every possible amount of work, one explicit schedule has completed at least as much mass as any competitor.

First describe the accounting state of an arbitrary schedule after it has used work x . Let t be its tested mass, b its mass completed in the other mode, and

$$c = (c_1, \dots, c_m)$$

the vector of tested positive mass already processed in each class. The postponed vector is not missing from this state: its i th coordinate is $d_i t - c_i$. An algorithm may retain a much richer internal history, but its aggregate state necessarily satisfies

$$0 \leq t \leq q, \quad 0 \leq b \leq 1 - q, \quad 0 \leq c_i \leq d_i t.$$

It has completed mass $d_0 t + b + \sum_i c_i$ and used work $t + vb + \sum_i p_i c_i$. Therefore define the finite-dimensional envelope

$$\mathcal{C}_{D,q}^v(x) := \max \left\{ d_0 t + b + \sum_{i=1}^m c_i : \begin{array}{l} 0 \leq t \leq q, \quad 0 \leq b \leq 1 - q, \\ 0 \leq c_i \leq d_i t \quad (1 \leq i \leq m), \\ t + vb + \sum_{i=1}^m p_i c_i \leq x \end{array} \right\}. \quad (27)$$

The candidate schedule is completely explicit.

Algorithm 7 Shortest-first fluid schedule FLUIDSPT(D, q, v)

- 1: Compute the maximum-density selector g_* , its completed mass a_* , and its work per completion τ_D .
 - 2: Create a testing block of completion mass qa_* and work per completion τ_D ; running it tests mass q and uses selector g_* .
 - 3: For every $p_i > 0$, create a deferred block of mass $qd_i(1 - g_*(p_i))$ and work per completion p_i .
 - 4: Create an alternative-completion block of mass $1 - q$ and work per completion v .
 - 5: Order all nonempty blocks by nondecreasing work per completion, placing the testing block before a deferred block in case of a tie, and execute them in that order.
-

Equivalently, the collection of blocks created by the algorithm is

$$\mathcal{M}_{D,q,v} := qa_*\delta_{\tau_D} + q\nu + (1 - q)\delta_v. \quad (28)$$

Theorem 3.3 (Pointwise optimality of the finite fluid schedule). *For every complete fluid schedule S that tests mass q ,*

$$C_S(x) \leq \mathcal{C}_{D,q}^v(x) = K_{\mathcal{M}_{D,q,v}}(x) \quad \text{for every } x \geq 0. \quad (29)$$

The schedule FLUIDSPT(D, q, v) attains equality for every x . Consequently it minimizes fluid total completion time and

$$\text{Area}(\mathcal{C}_{D,q}^v) = \text{SPT}(\mathcal{M}_{D,q,v}). \quad (30)$$

The proof has three short parts. First, the envelope contains the aggregate state of every schedule. Second, the density inequality compresses every feasible envelope state into the blocks of Equation (28). Finally, the support property of ν makes the shortest-first block order operationally feasible.

Claim 3.4 (Every schedule obeys the envelope). *For every complete fluid schedule S and every $x \geq 0$, $C_S(x) \leq \mathcal{C}_{D,q}^v(x)$.*

Proof. At work x , let $t, b, c_1, \dots, c_m$ be the aggregate state described above. Every tested mass has type distribution D , so at most $d_i t$ mass of positive type i has been revealed and hence $c_i \leq d_i t$. The test, alternative, and positive-processing work is respectively t, vb , and $\sum_i p_i c_i$. Thus the aggregate state is feasible in Equation (27), whose objective is exactly its completed mass. $\square$

Claim 3.5 (Testing activity compresses into the canonical blocks). *For every $x \geq 0$, $\mathcal{C}_{D,q}^v(x) \leq K_{\mathcal{M}_{D,q,v}}(x)$.*

Proof. Fix a feasible point $(t, b, c_1, \dots, c_m)$ in Equation (27). For each positive type, put

$$s_i = \min\{c_i, d_i t g_*(p_i)\}, \quad r_i = c_i - s_i.$$

The tests, tested zeros, and masses s_i complete

$$y_{\text{test}} = d_0 t + \sum_i s_i$$

using work

$$x_{\text{test}} = t + \sum_i p_i s_i.$$

If $t > 0$, apply Equation (22) to the selector $s_i/(d_i t)$, taking it to be zero when $d_i = 0$. This gives

$$\tau_D y_{\text{test}} \leq x_{\text{test}}, \quad y_{\text{test}} \leq a_* t \leq a_* q.$$

The same inequalities are trivial for $t = 0$. Hence the completed mass y_{test} fits inside the canonical testing block and requires no more work there.

Moreover,

$$0 \leq r_i \leq d_i t (1 - g_*(p_i)) \leq q d_i (1 - g_*(p_i)),$$

so the remaining processed mass fits inside the deferred block of type i . The mass b fits inside the alternative block. We have represented the same total completed mass by items from $\mathcal{M}_{D,q,v}$ using no more than the original work x . The definition of $K_{\mathcal{M}_{D,q,v}}$ proves the claim. $\square$

Claim 3.6 (The shortest-first curve is feasible). *The schedule $\text{FLUIDSPT}(D, q, v)$ is a complete fluid schedule and has completion curve $K_{\mathcal{M}_{D,q,v}}$.*

Proof. The alternative block can be executed wherever its value v places it in the shortest-first order. Running any fraction of the testing block means testing the corresponding fraction of its mass and using g_* on the revealed types. By Equation (23), every nonempty deferred block has processing time at least τ_D . The prescribed tie rule therefore places the testing block before all deferred work, so each deferred queue has been exposed before the schedule uses it. All block capacities are respected, and their total completion mass is

$$q a_* + q(1 - a_*) + (1 - q) = 1.$$

Thus the schedule is complete. By lemma 3.2, its shortest-first execution realizes $K_{\mathcal{M}_{D,q,v}}$. $\square$

Proof of Theorem 3.3. The first inequality in Equation (29) is claim 3.4. Claims 3.5 and 3.6 give the equality and show that one schedule realizes it simultaneously for every work budget. Integrating the pointwise inequality and applying lemma 3.2 proves Equation (30) and optimality. $\square$

3.5 From the fluid theorem to finite inputs

The theorem above deliberately fixes D , q , and v . We next prove one transfer theorem showing that its value is also the correct leading-order cost for bounded shuffled inputs. The remaining subsections then specialize that result and perform the model-specific algebra.

In obligatory testing every job must be tested, so $q = 1$ and the alternative block is absent. In revealing optimization with raw time u , set $v = u$ and minimize the area in Equation (30) over $q \in [0, 1]$. In blind execution, predictably chosen untouched jobs retain distribution D , so blind execution of unit mass uses mean work

$$\mu_D = \int p dD(p);$$

there we set $v = \mu_D$ and again minimize over q . The resulting model-specific values are denoted by $\Phi_{\text{OT}}(D)$, $\Phi_{\text{RO},u}(D)$, and $\Phi_{\text{BE}}(D)$ in their respective sections.

For an actual vector p , its empirical distribution $D[p]$ is already finitely supported. Gridding merely reduces the number of coordinates so that one concentration event controls them simultaneously. The next three subsections show that every finite competitor induces a fluid state obeying Equation (27), up to $o(1)$ normalized error, and that a sampled finite algorithm implements FLUIDSPT with the same accuracy. Thus the probabilistic transfer is a theorem following the exact fluid optimization, not an assumption used inside it.

3.6 A common finite-to-fluid transfer

The fluid theorem deliberately fixed a known distribution and made proportional revelation exact. We now show that learning and private shuffling reproduce both features up to $o(n^2)$ cost on bounded finite inputs. Doing this once also exposes the precise common structure of obligatory testing, revealing optimization, and blind execution.

Fix $L < \infty$ and a known function

$$r : [0, L] \longrightarrow [0, L]$$

with $|r(p) - r(p')| \leq |p - p'|$. A job first touched in the *direct mode* runs nonpreemptively for $r(p)$ and completes. The choice of this mode must be made before p is known; revealing p after completion may be granted to the algorithm for free. Alternatively, the job may be tested for one unit, after which its processing operation of length p can be scheduled.

Let Q be either $[0, 1]$, when both first-touch modes are available, or $\{1\}$, when every job must first be tested. For a distribution D , put

$$v_r(D) := \int r(p) dD(p). \quad (31)$$

Predictably chosen direct jobs have asymptotic mean work $v_r(D)$ per job. We therefore define the common fluid value

$$\Phi_{r,Q}(D) := \min_{q \in Q} \text{Area}(\mathcal{C}_{D,q}^{v_r(D)}). \quad (32)$$

When $Q = \{1\}$, the alternative block has zero mass and r is immaterial.

The three specializations used later are

model	$r(p)$	Q
OT	0	$\{1\}$
RO(u)	u	$[0, 1]$
BE	p	$[0, 1]$.

(33)

Thus the distinction between the last two models is entirely in the direct work: it is fixed in revealing optimization and sampled from the input in blind execution.

Theorem 3.7 (Common bounded instance-optimality transfer). *Fix L , r , and Q as above. There are randomized nonanticipating algorithms $\mathcal{L}_n$, depending only on n, L, r, Q , and a deterministic sequence*

$$\varepsilon_L(n) = O_L\left(n^{-1/6}\sqrt{\ln(n+2)}\right) \rightarrow 0 \quad (34)$$

such that every vector $p \in [0, L]^n$ satisfies

$$\mathbb{E} \text{ALG}_{\mathcal{L}_n}(p) \leq n^2 \Phi_{r,Q}(D[p]) + \varepsilon_L(n)n^2. \quad (35)$$

Conversely, every randomized nonanticipating algorithm $\mathcal{A}$ whose allowed first-touch modes are described by Q , even if it is told the multiset of entries of p , satisfies

$$\mathbb{E} \text{ALG}_{\mathcal{A}}(p) \geq n^2 \Phi_{r,Q}(D[p]) - \varepsilon_L(n)n^2. \quad (36)$$

Both expectations include the private shuffle and the algorithms' remaining randomness.

The proof is split into a lower transfer and a learning upper transfer. The first maps every finite adaptive run to the envelope from Equation (27). The second estimates the empirical distribution and implements the finite block schedule FLUIDSPT. The theorem then follows immediately by using the same fluid value between them.

3.7 From an adaptive run to the fluid envelope

The central fact is that adaptivity cannot bias the hidden value at a first touch. After fixing an algorithm's private seed, list jobs in the order in which their labels are first touched. Their values form a uniform permutation $X_1, \dots, X_n$ of the input. The indicator A_k that the k th job is tested is determined before X_k is exposed. Hence A_k and $1 - A_k$ are predictable selectors in lemma 2.7.

Keep zero as a separate class, partition $(0, L]$ into

$$K_n = \lfloor n^{1/6} \rfloor \quad (37)$$

cells, and round positive values upward to their cell endpoints. Write D^+ for the resulting empirical distribution and $h_n = L/K_n$ for the mesh. Take $\delta_n = n^{-1/6}$ and stop the first-touch order while $\delta_n n + O(1)$ labels remain untouched.

Apply the predictable-permutation estimate simultaneously to the vector of cell indicators selected by A_k and to the scalar $r(X_k)$ selected by $1 - A_k$. With probability at least $1 - n^{-2}$, every earlier prefix obeys

$$|\#\{\text{tested jobs in cell } j\} - d_j \#\{\text{tests}\}| \leq \Gamma_n, \quad (38)$$

$$|\text{direct work} - v_r(D[p])\#\{\text{direct jobs}\}| \leq \Gamma_n, \quad (39)$$

for all cells and all such prefixes, where

$$\frac{(K_n + 1)\Gamma_n}{n} = O\left(n^{-1/6}\sqrt{\ln(n+2)}\right). \quad (40)$$

For $Q = \{1\}$ the second estimate is vacuous.

Edit the last $\delta_n n + O(1)$ first touches to one fixed legal mode: test them when $Q = \{1\}$ and complete them directly when $Q = [0, 1]$. Delete any operations superseded by this edit. At most $\delta_n n + O(1)$ jobs and a constant number of operations per job change. Since every legal job uses at most $1 + L$ work, operation-by-operation remaining-job accounting gives

$$C_{\text{original}} \geq C_{\text{edited}} - O_L(\delta_n n^2 + n). \quad (41)$$

Let q be the final tested fraction of the edited run; then $q \in Q$.

Claim 3.8 (Repairing a finite prefix). *On the event Equations (38) and (39), there is*

$$\zeta_n = O_L\left(\frac{(K_n + 1)\Gamma_n}{n} + h_n + \delta_n\right) \quad (42)$$

such that, simultaneously throughout the edited run,

$$\text{completed fraction after normalized work } x \leq \mathcal{C}_{D^+, q}^{v_r(D^+)}(x + \zeta_n) + \zeta_n. \quad (43)$$

Proof. At an operation boundary, let t be tested mass, b directly completed mass, and c_j the processed tested mass in positive cell j . The cell estimate gives

$$c_j \leq d_j t + \Gamma_n/n,$$

and gives the analogous bound for tested zeros. Replace each c_j by $(c_j - \Gamma_n/n)^+$. The resulting class capacities are exact and the removed completed mass is at most $(K_n + 1)\Gamma_n/n$.

The work estimate replaces the actual direct work by $v_r(D[p])b$ with error at most Γ_n/n . Upward rounding changes tested processing work by at most h_n and, because r is one-Lipschitz, changes the direct mean by at most h_n . The edited suffix has mass $O(\delta_n)$. After these horizontal repairs, the variables t, b, c_j satisfy every constraint in Equation (27) with $v = v_r(D^+)$. This proves the displayed inequality with Equation (42). $\square$

We shall also use the following elementary stability observation. It avoids any claim that the optimizing threshold itself changes continuously.

Lemma 3.9 (Zero-preserving rounding stability). *Suppose D, D' admit a coupling that preserves zero and satisfies $|P - P'| \leq h$ almost surely. Then*

$$|\Phi_{r, Q}(D) - \Phi_{r, Q}(D')| \leq 2h. \quad (44)$$

Proof. Use the same tested fraction, selectors, and block order for coupled mass elements. At every prefix, at most one unit of tested mass receives processing, so its work changes by at most h . Direct work per unit mass also changes by at most h because r is one-Lipschitz. Thus either completion curve lies within a horizontal $2h$ shift of the other. Their areas differ by at most $2h$. Exchange D, D' and minimize over schedules. $\square$

Claim 3.10 (Common announced lower bound). *Every announced randomized algorithm in the common finite model satisfies*

$$\frac{\mathbb{E} \text{ALG}_{\mathcal{A}}(p)}{n^2} \geq \Phi_{r, Q}(D[p]) - O_L\left(n^{-1/6} \sqrt{\ln(n+2)}\right). \quad (45)$$

Proof. The envelope reaches one after at most $1 + L$ normalized work. Integrating Equation (43) therefore loses only $O_L(\zeta_n)$ from its area. By Theorem 3.3 and Equation (32), this area is at least $\Phi_{r, Q}(D^+)$. The sampling event fails with probability at most n^{-2} , while every fluid value is at most $(1 + L)/2$. Finally apply the suffix edit and lemma 3.9. Equations (40) and (42) give the stated rate. The argument is uniform after fixing the algorithm seed, so averaging proves the randomized claim. $\square$

3.8 Learning and implementing the fluid optimum

For the upper transfer, a *grid template* records a tested fraction, a selector on the grid classes, and a feasible order of the resulting testing, deferred, and direct blocks. For a grid distribution D and template π , let $F_{r,D}(\pi)$ be its fluid completion cost. Theorem 3.3 says that

$$\min_{\pi} F_{r,D}(\pi) = \Phi_{r,Q}(D). \quad (46)$$

Two finite facts make this template learnable. We state them separately so that the model-specific corollaries can reuse them.

Claim 3.11 (Finite implementation of a fixed template). *For every fixed grid template π , selecting the prescribed number of test labels uniformly and executing its blocks gives*

$$\mathbb{E}C_n(\pi) = n^2 F_{r,D[p]}(\pi) + O_L(n), \quad (47)$$

uniformly over the grid, its frequencies, and π .

Proof. Expand total completion time into one-job and unordered-pair contributions. For two distinct jobs, the probabilities that zero, one, or both are tested differ from the corresponding products of the tested fraction by $O(1/n)$. Every operation and pair contribution is bounded in terms of L . Summing the diagonal terms and these finite-population corrections gives $O_L(n)$; the remaining terms are exactly the fluid block area. $\square$

Claim 3.12 (Uniform stability of fixed templates). *There is $C_L < \infty$ such that any two distributions D, E on the same grid and every common template π satisfy*

$$|F_{r,D}(\pi) - F_{r,E}(\pi)| \leq C_L \|D - E\|_1. \quad (48)$$

The same inequality holds after minimizing over templates.

Proof. For a fixed template, the block-area expansion is a finite sum of single and double integrals whose kernels are bounded by a constant depending only on L . In a double integral, replace the first distribution and then the second. This gives the displayed total-variation bound uniformly over the number of grid classes and over the template. Applying the bound first to a minimizer for D and then to one for E proves the assertion for minima. $\square$

The learner uses the grid from Equation (37), privately permutes the labels, and tests the first

$$k_n = \lceil n^{1/2} \rceil \quad (49)$$

jobs. It keeps positive sample jobs pending and forms their rounded histogram $\widehat{D}_n$. It chooses a template minimizing $F_{r,\widehat{D}_n}$ and applies it to the unused labels, then finishes the positive sample jobs.

Let D_n and D_n^{rem} be the full and remaining rounded histograms. The sampling-without-replacement estimate Equation (12) and deletion of the pilot give

$$\mathbb{E}\|\widehat{D}_n - D_n\|_1 \leq \sqrt{\frac{K_n + 1}{k_n}}, \quad \|D_n^{\text{rem}} - D_n\|_1 \leq \frac{2k_n}{n - k_n}. \quad (50)$$

Claim 3.13 (Common learning upper bound). *The learner above satisfies*

$$\frac{\mathbb{E} \text{ALG}_{\mathcal{L}_n}(p)}{n^2} \leq \Phi_{r,Q}(D[p]) + O_L(n^{-1/6}) \quad (51)$$

uniformly over $p \in [0, L]^n$.

Proof. Condition on the pilot. The unused labels remain uniformly permuted, so claim 3.11 applies to the selected template. The pilot and its final positive tail use $O_L(k_n)$ work and therefore add $O_L(nk_n)$ to total completion time.

The fixed-template kernel is applied to the upward-rounded values. Coupling the physical and rounded executions with the same decisions changes tested processing work by at most h_n per unit mass and direct work by at most h_n per unit mass, because r is one-Lipschitz. Remaining-job accounting therefore changes the normalized template cost by at most $O(h_n)$.

Apply claim 3.12 first to the learned template, then to an optimal template for D_n^{rem} , and finally to an optimal template for D_n . This bounds the conditional fluid cost by

$$\Phi_{r,Q}(D_n) + O_L(\|\hat{D}_n - D_n\|_1 + \|D_n^{\text{rem}} - D_n\|_1).$$

Average and use Equation (50). The three remaining errors are

$$h_n = O_L(n^{-1/6}), \quad \sqrt{(K_n + 1)/k_n} = O(n^{-1/6}), \quad k_n/n = O(n^{-1/2}).$$

The rounding term is controlled by lemma 3.9, proving the claim. $\square$

Proof of Theorem 3.7. Claim 3.13 gives Equation (35). Claim 3.10 gives Equation (36), since announcing the multiset only strengthens the competing algorithm. Taking the larger of their vanishing errors gives Equation (34). $\square$

3.9 Revealing-optimization instance optimality

We first specialize the common transfer to revealing optimization. Here a direct first touch is raw execution: it always occupies the machine for the public time u , independently of the hidden value. Thus $r(p) = u$, and the only model-specific work is to expand the fluid block area into an explicit quadratic in the tested fraction.

Theorem 3.14 (Revealing-optimization instance optimality). *For every fixed $u > 0$ there exist randomized nonanticipating algorithms $\mathcal{A}_{n,u}^*$, depending only on n, u , and a deterministic sequence $\varepsilon_u(n) \rightarrow 0$ such that every vector $p \in [0, u]^n$ satisfies*

$$\mathbb{E} \text{ALG}_{\mathcal{A}_{n,u}^*}(p) \leq n^2 \Phi_{\text{RO},u}(D[p]) + \varepsilon_u(n)n^2. \quad (52)$$

Conversely, every randomized nonanticipating algorithm $\mathcal{A}$, even an announced one, satisfies

$$\mathbb{E} \text{ALG}_{\mathcal{A}}(p) \geq n^2 \Phi_{\text{RO},u}(D[p]) - \varepsilon_u(n)n^2. \quad (53)$$

One may take

$$\varepsilon_u(n) = O_u\left(n^{-1/6} \sqrt{\ln(n+2)}\right). \quad (54)$$

3.9.1 The explicit revealing benchmark

Let D be a probability distribution on $[0, u]$. Apply lemma 3.1, choose its threshold selector g_* , and write

$$a = a_D(g_*), \quad \ell = \int_{(0,u]} pg_*(p) dD(p), \quad \tau_D = \frac{1+\ell}{a}. \quad (55)$$

Let ν be the residual measure from Equation (23). It has mass $1 - a$, first moment

$$\ell_\nu = \int p d\nu(p) = \int p dD(p) - \ell, \quad \text{SPT}(\nu) = \frac{1}{2} \iint \min\{p, p'\} d\nu(p) d\nu(p'), \quad (56)$$

and support in $[\tau_D, u]$.

For a fixed tested fraction q , revealing optimization is the common fluid model with $v = u$. Hence Theorem 3.3 gives the blocks

$$qa\delta_{\tau_D} + q\nu + (1 - q)\delta_u.$$

When $\tau_D < u$, shortest-first order runs the testing block, the residual types in SPT order, and finally the raw block. Expanding Equation (30) gives

$$F_{\text{RO},u,D}(q) = (1 + \ell) \left(q - \frac{aq^2}{2} \right) + q\ell_\nu(1 - q) + q^2 \text{SPT}(\nu) + \frac{u(1 - q)^2}{2}. \quad (57)$$

The four terms are, respectively, the testing block, the residual work delaying raw jobs, the internal residual SPT cost, and the raw block.

If $\tau_D \geq u$, every testing or residual block has work per completion at least u , so pure raw execution pointwise dominates every positive tested fraction. We therefore define

$$\Phi_{\text{RO},u}(D) = \begin{cases} \min_{0 \leq q \leq 1} F_{\text{RO},u,D}(q), & \tau_D < u, \\ u/2, & \tau_D \geq u. \end{cases} \quad (58)$$

The minimization is explicit. Expanding gives

$$F_{\text{RO},u,D}(q) = \frac{u}{2} + Aq + Bq^2, \quad A = 1 + \int p dD(p) - u, \quad B = -\frac{a(1 + \ell)}{2} - \ell_\nu + \text{SPT}(\nu) + \frac{u}{2}. \quad (59)$$

For each work budget, convexly combining feasible states for two tested fractions gives a feasible state for their convex combination. Thus the completion envelope is concave in q , its unfinished-mass area is convex, and $B \geq 0$. Moreover,

$$\tau_D < u \iff \int (u - p) dD(p) > 1 \iff A < 0.$$

Consequently a minimizing tested fraction is

$$q^* = 1 \quad (B = 0), \quad q^* = \min\{1, -A/(2B)\} \quad (B > 0). \quad (60)$$

Proof of Theorem 3.14. Take $r(p) = u$ and $Q = [0, 1]$ in Theorem 3.7. Then $v_r(D) = u$, and Equations (57) and (58) show that the common value $\Phi_{r,Q}(D)$ is exactly $\Phi_{\text{RO},u}(D)$. The upper and lower conclusions are Equations (52) and (53), and Equation (34) gives Equation (54). $\square$

3.10 Obligatory-testing instance optimality

In obligatory testing the tested fraction is fixed at $q = 1$, so the common fluid optimum contains only the maximum-density testing block and its deferred SPT tail. This specialization admits a stronger learner than the generic bounded transfer: a slowly growing cutoff yields one algorithm that does not know the bound L and is nevertheless optimal on every fixed bounded class.

Theorem 3.15 (Obligatory-testing instance optimality). *There is a single randomized nonanticipating algorithm $\mathcal{A}^*$, depending only on n , with the following property. For every fixed $L < \infty$ there is a deterministic sequence $\varepsilon_L(n) \rightarrow 0$ such that every vector $p \in [0, L]^n$ satisfies*

$$\mathbb{E} \text{ALG}_{\mathcal{A}^*}(p) \leq n^2 \Phi_{\text{OT}}(D[p]) + \varepsilon_L(n) n^2. \quad (61)$$

Conversely, every randomized nonanticipating algorithm $\mathcal{A}$, even if it is told the multiset of entries of p , satisfies

$$\mathbb{E} \text{ALG}_{\mathcal{A}}(p) \geq n^2 \Phi_{\text{OT}}(D[p]) - \varepsilon_L(n) n^2. \quad (62)$$

One may take $\varepsilon_L(n) = O_L(n^{-1/5})$. Thus the same algorithm is asymptotically instance optimal, uniformly on every bounded input class.

3.10.1 The explicit obligatory benchmark

Let D be a finitely supported distribution on $[0, \infty)$. Apply lemma 3.1, choose its threshold selector g_* , and put

$$a_* = a_D(g_*), \quad w_* = w_D(g_*) = a_* \tau_D. \quad (63)$$

The residual measure ν from Equation (23) has mass $1 - a_*$ and support in $[\tau_D, \infty)$. Since $q = 1$, the block collection in Equation (28) is

$$a_* \delta_{\tau_D} + \nu.$$

The pointwise fluid optimality theorem therefore gives

$$\boxed{\Phi_{\text{OT}}(D) = w_* \left(1 - \frac{a_*}{2}\right) + \text{SPT}(\nu).} \quad (64)$$

Indeed, the first term is the area contributed by the testing block, both over itself and over the residual mass, and the second is the internal SPT area of the residual block. This is exactly $\text{Area}(\mathcal{C}_{D,1}^0)$, so Theorem 3.3 proves that no other fluid schedule has smaller cost.

For an empirical distribution, the selector has a direct finite description. Sort the job occurrences by processing time and, among prefixes E containing all zero jobs, maximize

$$\frac{|E|/n}{1 + n^{-1} \sum_{i \in E} p_i}. \quad (65)$$

The addition and removal inequalities include every occurrence below τ_D , exclude every occurrence above τ_D , and allow either choice at equality. Test jobs in a uniformly random order, immediately process the positive jobs in E , and finally process the remaining jobs in SPT order. Direct remaining-job accounting gives the exact identity

$$\mathbb{E} \text{ALG}_{\text{stat}}(p) = n^2 \Phi_{\text{OT}}(D[p]) + \frac{1}{2} \left(|E| + \sum_i p_i \right). \quad (66)$$

Thus the fluid value is already attainable up to $O_L(n)$ when the multiset is announced.

The common lower transfer also becomes sharper because every job is first touched by a test. There is no predictable choice between two first-touch modes and no unobserved suffix that has to be edited.

Claim 3.16 (Sharper obligatory lower transfer). *For every $p \in [0, L]^n$ and every announced randomized obligatory-testing algorithm $\mathcal{A}$,*

$$\frac{\mathbb{E} \text{ALG}_{\mathcal{A}}(p)}{n^2} \geq \Phi_{\text{OT}}(D[p]) - O_L(n^{-1/5}). \quad (67)$$

Proof. Follow the proof of claim 3.10, but partition $(0, L]$ into $K_n = \lfloor n^{1/5} \rfloor$ cells and use the uniform-prefix estimate Equation (13). With probability at least $1 - n^{-2}$, every prefix of the test order has discrepancy $O(\sqrt{n \ln(n+2)})$ in every cell. In the repair argument from claim 3.8, the removed completion mass is therefore

$$O\left(K_n \sqrt{\frac{\ln(n+2)}{n}}\right) = o(n^{-1/5}),$$

whereas the zero-preserving rounding mesh is $O_L(n^{-1/5})$. There is no direct-work discrepancy and no suffix edit. Integrating the repaired completion curve and applying lemma 3.9 proves the displayed bound. The argument is uniform after conditioning on the algorithm's private seed, so it also holds for randomized algorithms. $\square$

3.10.2 A learner that does not know the input bound

We now specify the single algorithm promised in Theorem 3.15. It uses a cutoff that tends to infinity and a grid whose mesh tends to zero.

Algorithm 8 Growing-cutoff obligatory learner $\mathcal{A}^*$

- 1: Set $k = \lfloor n^{3/4} \rfloor$, $d = \lfloor n^{1/4} \rfloor$, $B_n = 32 + n^{1/20}$, and $\eta = B_n/d$.
 - 2: Privately permute the labels and test labels $1, \dots, k$, leaving their positive processing operations pending.
 - 3: Keep zero as its own category, round values in $(0, B_n]$ upward to the B_n -grid of mesh η , and put values above B_n in an overflow category.
 - 4: Among finite category prefixes, choose one with maximum sample completion density as in Equation (65), and close it by including every finite category up to its reciprocal density.
 - 5: If the maximum sample density is less than $1/B_n$, test all remaining jobs and process every positive job in SPT order.
 - 6: Otherwise, process the pending sample jobs in the chosen categories; then test labels $k+1, \dots, n$ in order and immediately process each positive outcome in a chosen category.
 - 7: Finally process all remaining positive jobs in SPT order.
-

The overflow category is never selected. The closure step also includes finite categories absent from the sample, preventing an unseen short value from being incorrectly placed in the final queue.

Claim 3.17 (Growing-cutoff learning upper bound). *For every fixed $L < \infty$ and every $p \in [0, L]^n$,*

$$\mathbb{E} \text{ALG}_{\mathcal{A}^*}(p) \leq n^2 \Phi_{\text{OT}}(D[p]) + O_L(n^{9/5}). \quad (68)$$

Proof. Fix L . For all sufficiently large n , $B_n > L$, so the overflow category is empty. The prefix containing the whole sample has density at least $1/(1 + L + \eta) > 1/B_n$, and hence fallback does not occur.

Let D^+ be the rounded full histogram, D^{rem} the rounded histogram of the $n - k$ unused jobs, and $\hat{D}$ the sample histogram. The sampling-without-replacement estimate Equation (12) and deletion of the pilot give

$$\mathbb{E} \|\hat{D} - D^+\|_1 \leq \sqrt{\frac{d+2}{k}}, \quad \|D^{\text{rem}} - D^+\|_1 \leq \frac{2k}{n-k}. \quad (69)$$

The selected category prefix is the optimal grid template for $\widehat{D}$ by lemma 3.1 and Theorem 3.3. Conditional on the sample, the remaining labels are uniformly permuted. The fixed-template implementation claim 3.11 therefore evaluates their main schedule, while testing the pilot first and finishing its pending tail adds only $O_L(nk + k^2)$ to total completion time. Uniform template stability claim 3.12, applied to the learned template and then to an optimizer, yields

$$\frac{\mathbb{E}[\text{ALG}_{\mathcal{A}^*}(p) \mid \widehat{D}]}{n^2} \leq \Phi_{\text{OT}}(D^+) + O_L\left(\|\widehat{D} - D^+\|_1 + \|D^{\text{rem}} - D^+\|_1 + \frac{k}{n}\right).$$

Using actual rather than upward-rounded processing times can only decrease the cost. Moreover, zero-preserving rounding and lemma 3.9 give $\Phi_{\text{OT}}(D^+) \leq \Phi_{\text{OT}}(D[p]) + 2\eta$. Average, use Equation (69), and observe that

$$\sqrt{d/k} = O(n^{-1/4}), \quad k/n = O(n^{-1/4}), \quad \eta = O(n^{-1/5}).$$

Multiplication by n^2 proves the claim. Enlarging the constant covers the finitely many smaller values of n . $\square$

Proof of Theorem 3.15. Claim 3.17 proves Equation (61). Claim 3.16 proves Equation (62), since announcing the multiset can only help the competing algorithm. Both use an $O_L(n^{-1/5})$ normalized error, and the algorithm in algorithm 8 depends only on n . $\square$

3.10.3 Limits of the theorem

Randomization and bounded inputs are both essential to the preceding statement.

Theorem 3.18 (When instance optimality is impossible). (i) *There is no deterministic instance-optimal algorithm, even when all job lengths lie in $[0, 2]$. More precisely, for every deterministic algorithm $\mathcal{A}$, even an announced one, and every even n , there is a fixed labeling with $n/2$ zero jobs and $n/2$ length-two jobs for which*

$$\text{ALG}_{\mathcal{A}} \geq \frac{9}{8}n^2. \quad (70)$$

On the same input, an announced randomized algorithm $\mathcal{A}'$ has

$$\text{ALG}_{\mathcal{A}'} = n^2 + \frac{3}{4}n.$$

(ii) *If processing times are not bounded by a fixed constant, there is no randomized instance-optimal algorithm with one uniform additive $o(n^2)$ term. More precisely, for every randomized algorithm $\mathcal{A}$ independent of the input and every sufficiently large n , there is an input with values in $[0, n^2]$ and an announced randomized algorithm $\mathcal{A}'$ such that*

$$\text{ALG}_{\mathcal{A}} \geq \text{ALG}_{\mathcal{A}'} + \frac{1}{8}n^2. \quad (71)$$

Proof. For part (i), write $n = 2m$. Against the fixed deterministic algorithm, reveal length two on the first m labels it tests and zero on the remaining m . Record each answer on the label where it was given. Rerunning the algorithm on this fixed labeling reproduces exactly the same transcript. If it does not finish, the claim is immediate. Otherwise, let h be the number of length-two jobs completed before the first zero is tested. By then all m

positive jobs have been tested, so at least $m + 2h$ work has elapsed. Even if the remaining work is put in SPT order,

$$\begin{aligned} \text{ALG}_{\mathcal{A}}(p) &\geq \frac{3h(h+1)}{2} + (2m-h)(m+2h) + \frac{m(m+1)}{2} \\ &\quad + m(m-h) + (m-h)(m-h+1) \\ &= \frac{9}{2}m^2 + \frac{1}{2}h^2 + \frac{3}{2}m + \frac{1}{2}h \geq \frac{9}{8}n^2. \end{aligned}$$

The comparison algorithm tests in a uniformly random order, completes zeros at their tests, defers all length-two jobs, and finally processes them. Remaining-job accounting gives cost $n^2 + 3n/4$.

For part (ii), put $H = n^2$ and first run $\mathcal{A}$ on the input in which every job has length H . For a fixed private seed, let X_k be the number of completed jobs immediately before the k th test. Every processing operation contributes the same $Hn(n+1)/2$ in total, regardless of its position. Relative to the algorithm $\mathcal{A}'$ that tests and processes each job immediately, the expected excess is

$$\text{ALG}_{\mathcal{A}} - \text{ALG}_{\mathcal{A}'} = \mathbb{E} \sum_{k=1}^n (k-1 - X_k). \quad (72)$$

If this is at least $n^2/8$, we are done.

Otherwise replace one uniformly random label by a zero. Conditional on the seed, its test rank K is uniform on $\{1, \dots, n\}$. Until that test the run agrees with the all- H run, and Equation (72) gives

$$\mathbb{E}X_K > \frac{3n}{8} - \frac{1}{2}. \quad (73)$$

Forcing the zero behind X_K long jobs adds HX_K to the clairvoyant SPT lower bound. An announced algorithm can instead test in a private random order, defer positive jobs until it finds the zero, and then finish them; moving its unit tests away from the corresponding processing operations costs at most n^2 . Hence

$$\text{ALG}_{\mathcal{A}} - \text{ALG}_{\mathcal{A}'} \geq H \left(\frac{3n}{8} - \frac{1}{2} \right) - n^2,$$

which exceeds $n^2/8$ for all sufficiently large n . Averaging over the possible zero labels is exactly the private-shuffle expectation. $\square$

3.11 Blind-execution instance optimality

We next specialize the common transfer to blind execution. A direct first touch now runs the job for its actual processing time, so $r(p) = p$ and the direct work per unit mass is the mean processing time. The fluid optimum has four blocks: a maximum-density testing block, tested medium jobs, jobs run blindly, and tested long jobs. We first state instance optimality and then expand this four-block schedule into a one-variable quadratic.

Theorem 3.19 (Blind-execution instance optimality). *Fix $L < \infty$. There exist randomized nonanticipating algorithms $\mathcal{A}_{n,L}^*$, depending only on n, L , and a deterministic sequence $\varepsilon_L(n) \rightarrow 0$ such that every vector $p \in [0, L]^n$ satisfies*

$$\mathbb{E} \text{ALG}_{\mathcal{A}_{n,L}^*}(p) \leq n^2 \Phi_{\text{BE}}(D[p]) + \varepsilon_L(n)n^2. \quad (74)$$

Conversely, every randomized nonanticipating algorithm $\mathcal{A}$, even if it is told the multiset of entries of p , satisfies

$$\mathbb{E} \text{ALG}_{\mathcal{A}}(p) \geq n^2 \Phi_{\text{BE}}(D[p]) - \varepsilon_L(n) n^2. \quad (75)$$

The benchmark Φ_{BE} is defined explicitly below, and one may take

$$\varepsilon_L(n) = O_L\left(n^{-1/6} \sqrt{\ln(n+2)}\right).$$

3.11.1 The explicit four-block benchmark

Let D be a probability distribution on $[0, L]$ and put

$$\mu = \int p dD(p). \quad (76)$$

Thus blindly executing one unit of well-mixed job mass uses work μ . Apply lemma 3.1, choose its threshold selector g_* , and let ν be the residual measure from Equation (23). Write

$$a = a_D(g_*), \quad \ell = \int_{(0, L]} p g_*(p) dD(p).$$

The maximum-density identity is

$$a\tau_D - \ell = 1, \quad \tau_D = \frac{1 + \ell}{a}. \quad (77)$$

Hence a full unit of tests together with the selected short outcomes is a divisible block of completion mass a , work $1 + \ell$, and work per completion τ_D .

For a fixed tested fraction q , blind execution is the common fluid model with alternative work $v = \mu$. By Theorem 3.3, its optimal block collection is

$$qa\delta_{\tau_D} + q\nu + (1 - q)\delta_{\mu}. \quad (78)$$

If $\tau_D \geq \mu$, the direct block is no slower per completion than any testing block, so the optimum is pure blind execution and has cost $\mu/2$. Suppose now that $\tau_D < \mu$. Sorting the blocks in Equation (78) by work per completion gives, in order,

$$\begin{aligned} \underbrace{\text{tests and selected } p < \tau_D}_{\text{testing block}} &\longrightarrow \underbrace{\text{tested } \tau_D < p < \mu}_{\text{SPT order}} \\ &\longrightarrow \underbrace{\text{blindly executed jobs}}_{\text{work } \mu} \longrightarrow \underbrace{\text{tested } p > \mu}_{\text{SPT order}}. \end{aligned} \quad (79)$$

We next write the area of this completion curve explicitly. To avoid inessential boundary notation, first suppose that D has no mass at τ_D or μ , and set

$$\begin{aligned} \mathcal{L} &= \{p < \tau_D\}, & \mathcal{M} &= \{\tau_D < p < \mu\}, & \mathcal{H} &= \{p > \mu\}, \\ a &= D(\mathcal{L}), & \ell &= \int_{\mathcal{L}} p dD(p), & \ell_M &= \int_{\mathcal{M}} p dD(p), \\ d &= D(\mathcal{H}), & D_M &= D|_{\mathcal{M}}, & D_H &= D|_{\mathcal{H}}. \end{aligned}$$

For $\bar{q} = 1 - q$, remaining-mass accounting gives

$$F_{\text{BE}, D}(q) = (1 + \ell) \left(q - \frac{aq^2}{2} \right) + q\ell_M(\bar{q} + qd) + q^2 \text{SPT}(D_M)$$

$$+ \mu \left(\bar{q}qd + \frac{\bar{q}^2}{2} \right) + q^2 \text{SPT}(D_H). \quad (80)$$

The five terms describe, respectively, the testing block, the external and internal costs of the tested medium jobs, the blind block, and the internal cost of the tested long jobs. If D has mass at either boundary, split that mass between the adjacent blocks; the value is unchanged.

Expansion yields

$$F_{\text{BE},D}(q) = \frac{\mu}{2} + Aq + Bq^2, \quad (81)$$

where

$$A = 1 + \ell + \ell_M + \mu(d - 1) = 1 - \int (\mu - p)^+ dD(p), \quad (82)$$

$$B = -\frac{a(1 + \ell)}{2} + \ell_M(d - 1) + \text{SPT}(D_M) + \mu \left(\frac{1}{2} - d \right) + \text{SPT}(D_H). \quad (83)$$

The feasible set in Equation (27) is affine in q . Consequently its completion envelope is concave in q , its unfinished-mass area is convex, and $B \geq 0$. Moreover, when $\tau_D < \mu$, Equation (21) gives $A < 0$. A minimizing tested fraction is therefore

$$q^* = 1 \quad (B = 0), \quad q^* = \min\{1, -A/(2B)\} \quad (B > 0). \quad (84)$$

We define

$$\Phi_{\text{BE}}(D) = \begin{cases} \min_{0 \leq q \leq 1} F_{\text{BE},D}(q), & \tau_D < \mu, \\ \mu/2, & \tau_D \geq \mu. \end{cases} \quad (85)$$

Proof of Theorem 3.19. Take $r(p) = p$ and $Q = [0, 1]$ in Theorem 3.7. Then r is one-Lipschitz, $v_r(D) = \mu$, and Equations (78) and (85) show that the common value $\Phi_{r,Q}(D)$ equals $\Phi_{\text{BE}}(D)$. The two conclusions and the stated rate now follow from Equations (34) to (36). $\square$

3.11.2 Why all four blocks can be necessary

Let

$$D(\{0\}) = \frac{1}{5}, \quad D(\{9\}) = \frac{1}{5}, \quad D(\{16\}) = \frac{3}{5}.$$

Then $\mu = 57/5$ and $\tau_D = 5$. Thus tested zeros complete during their tests, tested jobs of length nine precede the jobs run blindly, and tested jobs of length sixteen follow them. Formula Equation (80) becomes

$$F_{\text{BE},D}(q) = \frac{57}{10} - \frac{44}{25}q + \frac{11}{10}q^2.$$

Its minimizer is $q^* = 4/5$ and its value is 4.996. If all tested nonzero jobs are instead deferred until after the blind block, the best value is 5.04. The gap is a positive constant at the fluid scale and therefore a positive constant times n^2 for large finite inputs. In particular, the simpler order “testing block, blind block, all tested residuals” is not instance optimal.

3.11.3 Why bounded processing times are necessary

The common transfer assumes a fixed upper bound L . For blind execution this is essential even for obtaining any finite worst-case ratio.

Theorem 3.20 (No finite ratio for unbounded blind execution). *There are no randomized algorithms $\mathcal{A}_n$, constant $R < \infty$, and sequence $\eta(n) \rightarrow 0$ such that every vector $p \in [0, \infty)^n$ satisfies*

$$\mathbb{E} \text{ALG}_{\mathcal{A}_n}(p) \leq R \cdot \text{OPT}(p) + \eta(n)n^2. \quad (86)$$

Proof. Suppose that such a guarantee exists, and run $\mathcal{A}_n$ on the all-zero input. Its offline cost is zero, so if Z_n denotes the expected online cost, then $Z_n \leq \eta(n)n^2 = o(n^2)$. We may assume that the algorithm finishes every job.

Fix its private seed and order the labels by their first touch. Every first touch completes its zero job. If the job at rank h is tested, that test is performed while $n - h + 1$ jobs are unfinished and contributes $n - h + 1$ to the objective. Consequently

$$Z_n \geq \mathbb{E} \sum_{h: \text{rank } h \text{ is tested}} (n - h + 1). \quad (87)$$

Now choose one label uniformly and change only its length from zero to $H = n^2$. Until this exceptional label is first touched, the algorithm sees the zero-input history. If its rank h is touched blindly, its execution occupies the machine for H time units while $n - h + 1$ jobs are unfinished. Averaging over the exceptional label and the private seed gives

$$\begin{aligned} \mathbb{E} \text{ALG}_{\mathcal{A}_n}(p) &\geq \frac{H}{n} \mathbb{E} \sum_{h: \text{rank } h \text{ is blind}} (n - h + 1) \\ &\geq \frac{H}{n} \left(\frac{n(n+1)}{2} - Z_n \right) = H \left(\frac{n}{2} - o(n) \right). \end{aligned} \quad (88)$$

Some fixed choice of the exceptional label has at least this expected cost. Its offline optimum runs all zero jobs first and the long job last, and hence $\text{OPT}(p) = H$. With $H = n^2$, Equation (88) contradicts Equation (86) for all sufficiently large n . $\square$

4 Deterministic revealing optimization and its obligatory endpoint

This section proves two exact deterministic results: the revealing-optimization curve in Theorem 4.1 and the obligatory-testing ratio in Theorem 4.2, obtained as its uniform unbounded endpoint. These are the rigorous deterministic parts of the informal statements in Theorems 1.1 and 1.6. We begin with revealing optimization at a finite common raw-execution time u . All jobs have this same known upper limit, and a test reveals $p_j \in [0, u]$. By Equation (2) and lemma 2.1, the offline effective length is $\min\{u, 1 + p_j\}$, and the offline optimum schedules these effective lengths in nondecreasing order. We write $R_{\text{RO}}^{\text{det}}(u)$ for the optimal deterministic size-asymptotic ratio at this fixed common upper limit.

The exact ratio is given by six formulas on six intervals separated by five transition points. Put

$$\varphi = \frac{1 + \sqrt{5}}{2}, \quad u_1^{\text{det}} = 1, \quad u_2^{\text{det}} = \frac{1 + \sqrt{3 + 2\sqrt{5}}}{2} = 1.866760399173862 \dots \quad (89)$$

Let $s_0 > 1$ be the root of

$$s_0^3 = (s_0 + 1)^2, \quad u_3^{\det} = 1 + s_0 = 3.147899035704787 \dots \quad (90)$$

The fourth transition point is

$$u_4^{\det} = \varphi + 2 = 3.618033988749895 \dots \quad (91)$$

For $s = u - 1$, define

$$T_s(z) = sz^2 + (s^3 + s^2 + 1)z - (s^2 + s + 1)(s + 2),$$

and let $R_{\text{quad}}(u)$ be its positive root, so

$$T_s(R_{\text{quad}}(u)) = 0. \quad (92)$$

Equivalently,

$$R_{\text{quad}}(u) = \frac{-1 - u + 2u^2 - u^3 + \sqrt{u^6 - 4u^5 + 10u^4 - 6u^3 - 3u^2 + 6u - 3}}{2(u - 1)}. \quad (93)$$

At the other end, let $u_5^{\det} > 1$ solve $u_5^{\det} - 3 = \ln u_5^{\det}$ and define

$$\begin{aligned} r &= \frac{2}{u_5^{\det} - 1}, \quad R_{\text{OT}}^{\det} = 1 + r = 1.570574096649395 \dots, \\ u_5^{\det} &= 4.505241495792883 \dots \end{aligned} \quad (94)$$

For each $u_4^{\det} \leq u \leq u_5^{\det}$, let $c = c(u)$ and $m = m(u)$ be the unique solution of

$$\operatorname{artanh} m - m = 1 - \frac{1}{c} + \frac{1}{2} \ln \left(1 + \frac{2}{c} \right), \quad u = 1 + \frac{2}{c} - m, \quad (95)$$

with $r \leq c \leq 1/\varphi$ and $0 \leq m \leq 1/\varphi$.

Theorem 4.1 (Exact revealing-optimization curve). *For every fixed $0 < u < \infty$, the optimal deterministic size-asymptotic ratio in the revealing-optimization model is*

$$R_{\text{RO}}^{\det}(u) = \begin{cases} 1, & 0 < u \leq u_1^{\det}, \\ u, & u_1^{\det} \leq u \leq u_2^{\det}, \\ R_{\text{quad}}(u), & u_2^{\det} \leq u \leq u_3^{\det}, \\ 1 + \frac{1}{\sqrt{u-1}}, & u_3^{\det} \leq u \leq u_4^{\det}, \\ 1 + c(u), & u_4^{\det} \leq u \leq u_5^{\det}, \\ R_{\text{OT}}^{\det}, & u \geq u_5^{\det}. \end{cases} \quad (96)$$

On every interval in Equation (96), there is an explicit deterministic algorithm satisfying

$$\text{ALG}_{\mathcal{A}}(I) \leq R_{\text{RO}}^{\det}(u) \cdot \text{OPT}(I) + O_u(n).$$

On $u \geq u_5^{\det}$ the remainder is one absolute $O(n)$ term, uniform over all finite u . Conversely, for every deterministic algorithm $\mathcal{A}$ and every $\varepsilon > 0$, there are arbitrarily large fixed instances I such that

$$\text{ALG}_{\mathcal{A}}(I) \geq (R_{\text{RO}}^{\det}(u) - \varepsilon) \cdot \text{OPT}(I).$$

For $u \geq u_5^{\det}$, these lower-bound instances may be chosen with $p_j < u - 1$ for every job, so the offline cap u is inactive. Thus both witnesses on the plateau are uniform in the sense needed to pass separately to the obligatory endpoint.

Theorem 4.2 (Deterministic obligatory testing). *Let $u_5^{\text{det}} > 1$ be the unique solution of $u_5^{\text{det}} - 3 = \ln u_5^{\text{det}}$, and set*

$$R_{\text{OT}}^{\text{det}} = 1 + \frac{2}{u_5^{\text{det}} - 1} = 1.570574096649395 \dots$$

For every deterministic online algorithm $\mathcal{B}$ and every $\varepsilon > 0$, there are arbitrarily large obligatory-testing instances I such that

$$\text{ALG}_{\mathcal{B}}(I) \geq (R_{\text{OT}}^{\text{det}} - \varepsilon) \cdot \text{OPT}(I).$$

Conversely, there are a deterministic algorithm $\mathcal{A}$ and an absolute constant B such that every n -job obligatory instance satisfies

$$\text{ALG}_{\mathcal{A}}(I) \leq R_{\text{OT}}^{\text{det}} \cdot \text{OPT}(I) + Bn.$$

Consequently the optimal deterministic size-asymptotic competitive ratio is exactly $R_{\text{OT}}^{\text{det}}$.

The proof of Theorem 4.1 combines two kinds of lower bounds. Binary hidden-stopping instances give the formulas on $0 < u \leq u_4^{\text{det}}$, while a continuous version of the harmonic construction with an initial block of jobs satisfying $p_j = u$ gives the lower bound on $u_4^{\text{det}} \leq u \leq u_5^{\text{det}}$ and the constant value for $u \geq u_5^{\text{det}}$. Three explicit algorithms match them: raw execution, a forced-prefix uniform threshold, and a history-dependent adaptive threshold. They are introduced informally in Section 1.6 and specified formally in algorithms 9 to 11. The matching lower bounds are developed in Section 4.2. The following table maps each interval to its upper-bound witness.

range	algorithm	parameter choice
$0 < u \leq u_2^{\text{det}}$	RAW	none
$u_2^{\text{det}} < u \leq u_3^{\text{det}}$	FORCEDPREFIXUTE	$b = b(u)$ from (98)
$u_3^{\text{det}} < u \leq u_4^{\text{det}}$	FORCEDPREFIXUTE	$b = 0$
$u_4^{\text{det}} < u < u_5^{\text{det}}$	ADAPTIVETHRESHOLD	$c = c(u)$ from (95)
$u_5^{\text{det}} \leq u < \infty$	ADAPTIVETHRESHOLD	$c = r$

4.1 Upper bounds

We prove the six interval guarantees using three algorithms. Raw execution covers $0 < u \leq u_2^{\text{det}}$, a forced-prefix uniform threshold covers $u_2^{\text{det}} \leq u \leq u_4^{\text{det}}$, and an adaptive threshold covers $u \geq u_4^{\text{det}}$. Each of the following subsections treats one algorithm and ends with its complete guarantee on the relevant range of u .

4.1.1 Raw execution

We first handle the range in which testing is unnecessary. Running every job raw is optimal for $u \leq 1$ and gives the matching factor u until the first nontrivial transition point u_2^{det} .

Algorithm 9 $\text{RAW}(J, u)$

Input: jobs J and a finite common upper limit u .

1: Fix the job order $1, \dots, n$.

2: **For** $i = 1, \dots, n$, execute job i untested for u time.

Lemma 4.3 (Raw-execution guarantee). *For every $0 < u \leq u_2^{\det}$, RAW satisfies*

$$\text{ALG}_{\text{RAW}}(I) \leq \max\{1, u\} \cdot \text{OPT}(I). \quad (97)$$

In particular, it is optimal for $0 < u \leq u_1^{\det}$ and is u -competitive for $u_1^{\det} \leq u \leq u_2^{\det}$.

Proof. The raw blocks all have length u , so $\text{ALG}_{\text{RAW}} = un(n+1)/2$. If $u \leq 1$, every offline effective length is $\min\{u, 1 + p_j\} = u$, and equality holds. If $u > 1$, every offline effective length is at least one, so $\text{OPT} \geq n(n+1)/2$ and the ratio is at most u . $\square$

4.1.2 Forced-prefix uniform thresholds

The next algorithm tests every job. It processes a forced prefix regardless of the observed values and thereafter processes only short outcomes immediately. We prove its complete guarantee on $[u_2^{\det}, u_3^{\det}]$ and then analyze the zero-prefix specialization, namely $b = 0$, which is optimal up to $u_4^{\det}$.

Here UTE stands for *uniform-threshold execution*: outside the forced prefix, the algorithm uses the same threshold for every tested job.

Algorithm 10 FORCEDPREFIXUTE(J, u, b)

- Input:** jobs J , finite $u > 1$, and $b \in [0, 1]$.
- 1: Fix the test order $1, \dots, n$; set $k \leftarrow \lfloor bn \rfloor$, $\theta_u \leftarrow \min\{1, u - 1\}$, and $Q \leftarrow \emptyset$.
 - 2: **For** $i = 1, \dots, n$ **do**:
 - 3: Test job i for one unit and observe p_i .
 - 4: **If** $i \leq k$ or $p_i \leq \theta_u$, process job i immediately.
 - 5: **Otherwise**, add job i to Q .
 - 6: Process the jobs of Q in nondecreasing revealed processing time.
-

In this subsection, the first $k = \lfloor bn \rfloor$ positions in the fixed test order are the *prefix*, and the remaining positions are the *suffix*. We call a tested job *immediate* if it is processed at once, and *deferred* if it is placed in Q for the final queue.

Recall from the beginning of Section 4 that $R_{\text{quad}}(u)$ denotes the positive root specified by Equation (92).

Lemma 4.4 (Forced-prefix UTE guarantee). *For every $u_2^{\det} \leq u \leq u_3^{\det}$, define the algorithm parameter*

$$b = b(u) := \frac{u^2 - u + 1 - (u - 1)^2 R_{\text{quad}}(u)}{u^2 - u + 1 + (u - 1) R_{\text{quad}}(u)}. \quad (98)$$

Then FORCEDPREFIXUTE(J, u, b) satisfies

$$\text{ALG} \leq R_{\text{quad}}(u) \cdot \text{OPT} + O_u(n). \quad (99)$$

We prove the lemma in three steps. The first claim verifies that the root R is well-defined and that the prefix parameter is admissible. The next two claims prove the guarantee below and above $u = 2$ by reducing the possible processing times to boundary values in two different ways. Throughout these claims, put $R = R_{\text{quad}}(u)$ and $s = u - 1$. In this notation, the parameter from Equation (98) is

$$b = \frac{s^2 + s + 1 - s^2 R}{s^2 + s + 1 + s R}.$$

Claim 4.5 (Parameter admissibility). *The positive root R is well-defined. Moreover, $0 \leq b < 1$, with $b = 0$ only at $u = u_3^{\text{det}}$, and $R > 5/3$ and $s^2 R > 16/15$ throughout the interval.*

Proof. Since $s > 0$, we have $T_s(0) < 0$ and $T_s(z) \rightarrow \infty$ as $z \rightarrow \infty$. Moreover, T_s is strictly increasing for $z > 0$, because

$$T'_s(z) = 2sz + s^3 + s^2 + 1 > 0.$$

It therefore has a unique positive root R .

Since T_s is increasing on positive arguments,

$$T_s\left(\frac{s^2 + s + 1}{s^2}\right) = \frac{s^2 + s + 1}{s^3}(-s^3 + s^2 + 2s + 1) \geq 0 \quad (s \leq s_0), \quad (100)$$

and hence $s^2 + s + 1 - s^2 R \geq 0$, with equality only at $s = s_0$. The denominator in b is positive and exceeds its numerator by $sR(s + 1)$, proving $b < 1$.

We will also use the following lower bound on R :

$$9T_s(5/3) = 6s^3 - 12s^2 - 2s - 3 < 0 \quad (101)$$

throughout the present interval. Indeed, on $[4/5, 1]$ the polynomial is decreasing and has value $-1151/125$ at $4/5$; on $[1, s_0]$ it has one interior minimum, while its value at 1 is -11 and its value at s_0 is smaller than its value $-95/36$ at $13/6$. Here $s_0 < 13/6$ follows by substituting $13/6$ in $s^3 - (s + 1)^2$. Thus $R > 5/3$, while $s > 4/5$, and $s^2 R > 16/15$. $\square$

Claim 4.6 (The range $u_2^{\text{det}} \leq u \leq 2$). *For every $u_2^{\text{det}} \leq u \leq 2$, FORCEDPREFIXUTE(J, u, b) satisfies*

$$\text{ALG} \leq R \cdot \text{OPT} + O_u(n).$$

Proof. We first consider a binary input in which every job is either zero ($p_i = 0$) or long ($p_i = u$). The first bn positions are the forced prefix. A long job in this prefix is processed immediately; a long job in the suffix is deferred; every zero job is processed immediately.

Fix the total number L of long jobs, and write $k = bn$ for the moment; integer rounding will be restored at the end. Their positions do not affect OPT, which knows all values and sorts by effective length. We determine the worst positions for ALG directly.

Suppose exactly q long jobs lie in the prefix and put $D = L - q$. Within the prefix, moving a long job before a zero delays that zero and every later immediate completion by the long processing time u . Within the suffix, moving a deferred long job before a zero delays the zero by one unit of test time and does not change the final queue. Thus, for fixed q , the worst order is

$$\underbrace{u, \dots, u}_q \underbrace{0, \dots, 0}_{k-q} \mid \underbrace{u, \dots, u}_D \underbrace{0, \dots, 0}_{n-k-D},$$

where the vertical bar separates the forced prefix from the suffix. The completion cost of this order is

$$C(q) = \frac{n(n+1)}{2} + D \left(n - k - \frac{D+1}{2} \right) + u \left(qn - \frac{q(q-1)}{2} + \frac{D(D+1)}{2} \right). \quad (102)$$

The first term is the unit-test baseline. The second records the positions of the tests of the D long jobs in the suffix, and the last records all delays caused by long processing. Moving one long job from the suffix into the prefix changes q to $q + 1$ and D to $D - 1$, and hence

$$C(q+1) - C(q) = (u-1)(n-L) + k - q \geq 0.$$

Consequently a worst input puts as many long jobs in the prefix as possible, then all remaining long jobs at the start of the suffix, and all zeros last.

Divide all job counts by n , and let $x = L/n$ be the fraction of long jobs. The fraction of long jobs in the prefix is therefore $a = \min\{b, x\}$. If $x > b$, the remaining fraction $d = x - b$ consists of deferred long jobs; if $x \leq b$, then $d = 0$. Ignore for the moment the $O_u(n)$ one-job and rounding terms, and write $A = \text{ALG}/n^2$ and $O = \text{OPT}/n^2$ for the leading coefficients. Direct pair counting gives

$$\begin{aligned} O(x) &= \frac{1}{2} + \frac{s}{2}x^2, \\ A(x) &= \begin{cases} \frac{1}{2} + u \left(x - \frac{x^2}{2} \right), & 0 \leq x \leq b, \\ \frac{1}{2} + u \left(b - \frac{b^2}{2} \right) + (1-b)(x-b) + \frac{s}{2}(x-b)^2, & b \leq x \leq 1. \end{cases} \end{aligned} \quad (103)$$

The term $1/2$ is the common unit-time baseline. In $O(x)$, the extra amount $s = u - 1$ is paid only by pairs of long jobs. The two cases for $A(x)$ distinguish whether all long jobs fit into the forced prefix.

Put $G(x) = A(x) - R \cdot O(x)$. We show that its maximum is zero. On $b \leq x \leq 1$, substituting the second formula for $A(x)$ from Equation (103) makes $G(x)$ a strictly concave quadratic. Its stationary point is

$$\alpha = \frac{s^2 + s + 1}{s^2 + s + 1 + sR}.$$

Moreover, $b = (s+1)\alpha - s$, and hence $\alpha - b = s(1-\alpha) > 0$. Substitution gives $G(\alpha) = 0$; after cancelling the positive denominator, this is exactly Equation (92). Since $b < \alpha$, its right derivative at b is positive. The left derivative satisfies $G'_-(b) = G'_+(b) + s(1-b) > 0$, and the first quadratic is concave, so it is increasing on $[0, b]$. Thus $G(x) \leq 0$ for every $x \in [0, 1]$, including the limiting case $b = 0$: the two pieces agree at b , the first increases up to that common value, and the second never exceeds its zero maximum.

We now reduce a general input to the binary case just analyzed. In the present range the threshold is $u - 1 \leq 1$. Thus a suffix job is immediate exactly when $p \leq u - 1$ and deferred otherwise. First increase every deferred suffix value to u . This cannot decrease ALG , and it leaves OPT unchanged because such a job already has effective length $\min\{u, 1+p\} = u$.

For the remaining coordinates, construct a random binary vector as follows, independently from job to job. Replace a prefix value p by u with probability p/u and by zero otherwise. Replace an immediate suffix value $p \leq u - 1$ by u with probability p and by zero otherwise. The latter probability is valid because $p \leq u - 1 \leq 1$. A suffix job rounded to u changes from immediate to deferred; the comparison below includes this change of action.

To compare online costs, consider a pair of positions $i < j$ in the test order. Its leading contribution to ALG is

$$g_{ij} = \begin{cases} 1 + p_i, & i \text{ is immediate,} \\ 2 + p_j, & i \text{ is deferred and } j \text{ is immediate,} \\ 2 + \min\{p_i, p_j\}, & i, j \text{ are deferred.} \end{cases} \quad (104)$$

A prefix job remains immediate, and its rounding preserves its expected processing value. If i is an immediate suffix job, then after rounding its pair contribution is one when it becomes zero and at least two when it becomes u ; its expectation is therefore at least $1 + p_i$. If i is deferred and j is an immediate suffix job, then i has already been raised to u ,

and rounding j gives expected contribution $2 + up_j \geq 2 + p_j$. Finally, a pair of deferred jobs can only become more expensive. These cases show that the expected leading online cost does not decrease.

It remains to check that the same rounding does not increase OPT. If λ' denotes the effective length after rounding, then for a prefix job and an immediate suffix job, respectively,

$$\mathbb{E}\lambda' = 1 + \frac{u-1}{u}p \leq \min\{u, 1+p\}, \quad \mathbb{E}\lambda' = 1 + (u-1)p \leq 1+p. \quad (105)$$

A deferred suffix job has effective length u both before and after the first step. Since the minimum is jointly concave,

$$\mathbb{E} \min\{\lambda'_i, \lambda'_j\} \leq \min\{\mathbb{E}\lambda'_i, \mathbb{E}\lambda'_j\} \leq \min\{\lambda_i, \lambda_j\}.$$

The offline pair identity therefore shows that expected OPT does not increase. Consequently the expected value of $\text{ALG} - R \cdot \text{OPT}$ for the binary vector is at least its value for the original input, up to $O_u(n)$ one-job terms. Some binary realization is at least as bad as this expectation, and the binary bound proves the claim. $\square$

Claim 4.7 (The range $2 \leq u \leq u_3^{\det}$). *For every $2 \leq u \leq u_3^{\det}$, $\text{FORCEDPREFIXUTE}(J, u, b)$ satisfies*

$$\text{ALG} \leq R \cdot \text{OPT} + O_u(n).$$

Proof. Here $u \geq 2$, so the algorithm uses threshold one. A prefix job is always immediate, a suffix job with $p \leq 1$ is immediate, and a suffix job with $p > 1$ is deferred. First keep this immediate/deferred pattern fixed and vary one processing time at a time in the leading excess $\text{ALG} - R \cdot \text{OPT}$. On each interval between the threshold and the offline cap u , the pair formula Equation (104) makes this excess convex in that coordinate. Hence a maximum occurs at an interval endpoint. This reduces

- a prefix value to 0 or u ;
- an immediate suffix value to 0 or 1; and
- a deferred suffix value to 1^+ or u .

Here 1^+ means a value tending to one from above, so the job remains deferred. For the last reduction, on $1 < p < u-1$ the only nonlinear term is a concave minimum multiplied by $1 - R < 0$, hence it is convex; on $[u-1, u]$ the offline effective length is already the constant u , while the online cost is nondecreasing in p . In the rest of this proof, we call a job with $p = u$ *capped*.

We next choose the worst test order for fixed numbers of these endpoint types. OPT depends only on those numbers. In ALG, swapping two neighboring positions shows the following: capped jobs in the prefix may be moved before zero jobs in the prefix; deferred jobs in the suffix may be moved before immediate jobs in the suffix; and suffix jobs of value one may be moved before zero jobs in the suffix. These are three direct comparisons, not an additional scheduling principle: substitute the processing times of the two jobs being swapped in Equation (104); all unaffected pair terms cancel and the remaining difference is nonnegative. The order among two deferred jobs is irrelevant because the last line of Equation (104) is symmetric.

Divide each class size by n . Let a be the resulting fraction of prefix jobs with $p = u$, d the fraction of suffix jobs with $p = u$, t the fraction of deferred suffix jobs with $p = 1^+$, and

m the fraction of immediate suffix jobs with $p = 1$. All other jobs have value zero. The constraints are

$$0 \leq a \leq b, \quad d, t, m \geq 0, \quad d + t + m \leq 1 - b. \quad (106)$$

As in the preceding claim, let A and O denote the leading coefficients of ALG/n^2 and OPT/n^2 . Summing Equation (104) over the ordered endpoint classes gives

$$A = \frac{1}{2} + (s+1) \left(a - \frac{a^2}{2} \right) + (1-b)(d+t+m) - \frac{m^2}{2} + \frac{s}{2}d^2, \quad (107)$$

$$O = \frac{1}{2} + \frac{(a+d+t+m)^2}{2} + \frac{s-1}{2}(a+d)^2. \quad (108)$$

The formula for O is especially transparent: every job contributes the unit baseline; all four nonzero classes have one additional unit of effective length; and the two capped classes a, d have another $s-1 = u-2$ units. Replacing an immediate value-one suffix job by a deferred value-1⁺ job does not change O in the limit and increases A . Thus a worst reduced instance has $m = 0$. Put $x = a+d$, the total fraction of capped jobs, and maximize $A - R \cdot O$ over the remaining split between a, d, t . We retain the stationary point from the preceding claim,

$$\alpha = \frac{s^2 + s + 1}{s^2 + s + 1 + sR}.$$

For fixed x , the derivative with respect to the deferred $p = 1^+$ fraction t is $1 - b - R(x+t)$. Thus the best choice is $t = (\chi - x)_+$, where

$$\chi = \frac{1-b}{R} = \frac{s(s+1)}{s^2 + s + 1 + sR}.$$

The identities

$$\alpha - \chi = \frac{1}{s^2 + s + 1 + sR}, \quad \chi - b = \frac{s^2 R - 1}{s^2 + s + 1 + sR}$$

show that $b < \chi < \alpha$ on the present range. At fixed x, t , replacing a capped suffix job by a capped prefix job leaves O unchanged and changes A at rate $s(1-x) + b - a \geq 0$. We therefore fill the prefix with capped jobs as far as the constraints permit. Altogether,

$$t = (\chi - x)_+, \quad a = \min\{b, x\}, \quad d = (x - b)_+. \quad (109)$$

It remains to optimize the single variable x . If $x \geq \chi$, then $t = 0$, so only values zero and u remain. This is exactly the binary calculation from claim 4.6; its maximum gap is zero at $x = \alpha$. If $x < \chi$, then $t = \chi - x > 0$, and substitution gives

$$A - R \cdot O = -\frac{R-1}{2} + \frac{(1-b)^2}{2R} + a(s+b-sx) - \frac{a^2}{2} + \frac{s-R(s-1)}{2}x^2. \quad (110)$$

The coefficient $s - R(s-1)$ is positive on $1 \leq s \leq s_0$. Indeed, $T_s(2) = s(s^2 - s + 1) > 0$, so for $s \leq 2$ the equation $T_s(R) = 0$ gives $R < 2$, and hence $s - R(s-1) > 2 - s \geq 0$. For $2 \leq s \leq s_0 < 13/6$,

$$16T_s(7/4) = 12s^3 - 20s^2 + s - 4 > 0; \quad (111)$$

the polynomial is 14 at $s = 2$ and has positive derivative thereafter. Thus $R < 7/4$ and $s - R(s-1) > 7/4 - 3s/4 > 1/8$.

On $0 \leq x \leq b$, one has $a = x$, and

$$A - R \cdot O = -\frac{R-1}{2} + \frac{(1-b)^2}{2R} + (s+b)x - \frac{s+1+R(s-1)}{2}x^2. \quad (112)$$

Its derivative is decreasing, and the derivative at b multiplied by the positive denominator $s^2 + s + 1 + sR$ is

$$R(1 + s^2 + Rs^2(s - 1)) > 0, \quad (113)$$

so the maximum is at b . On $b \leq x \leq \chi$, Equation (110) is convex and its maximum is at b or χ . At $x = \chi$ one has $t = 0$, so this endpoint is already covered by the binary calculation. It remains to compare the other endpoint $x = b$ with it:

$$F(b) - F(\chi) = \frac{\chi - b}{2} [2sb - (s - R(s - 1))(b + \chi)], \quad (114)$$

and direct substitution of b and χ gives

$$(s^2 + s + 1 + sR)[(s - R(s - 1))(b + \chi) - 2sb] = -s + (1 + s - s^3)R + s^2(s - 1)R^2 > 0. \quad (115)$$

To prove the remaining sign without numerical approximation, observe that T_s is increasing on positive arguments and

$$9T_s(5/3) = 6s^3 - 12s^2 - 2s - 3 < 0, \quad (116)$$

so $R > 5/3$. At $R = 5/3$, the derivative of the right-hand side of Equation (115) satisfies

$$3 \frac{\partial}{\partial R} [-s + (1 + s - s^3)R + s^2(s - 1)R^2] \Big|_{R=5/3} = 7s^3 - 10s^2 + 3s + 3 > 0. \quad (117)$$

The last polynomial is increasing for $s \geq 1$. Since the coefficient of R^2 is nonnegative, the right-hand side of Equation (115) is therefore increasing for $R \geq 5/3$. At $R = 5/3$ its value satisfies

$$9[-s + (1 + s - s^3)(5/3) + s^2(s - 1)(25/9)] = 10s^3 - 25s^2 + 6s + 15 > 0. \quad (118)$$

For the last sign, write $s = 3/2 + z$; the expression becomes $3/2 - 3z/2 + 20z^2 + 10z^3$. It is at least $3/2$ for $-1/2 \leq z \leq 0$, and for $z \geq 0$ it is larger than $3/2 - 9/320$. Hence both possible maxima on $[b, \chi]$ are nonpositive. Together with the binary case $x \geq \chi$, this proves $A - R \cdot O \leq 0$ for every reduced configuration. Restoring the one-job terms and rounding $\lfloor bn \rfloor$ contributes only $O_u(n)$. $\square$

We can now prove the lemma.

Proof of Lemma 4.4. By Claim 4.5, the displayed value of b is a valid algorithm parameter throughout $[u_2^{\det}, u_3^{\det}]$. The guarantees in Claims 4.6 and 4.7 cover this interval and give Equation (99) with $R = R_{\text{quad}}(u)$. $\square$

At $u = u_3^{\det}$, $b = 0$. For $u_3^{\det} \leq u \leq u_4^{\det}$, the zero-prefix specialization gives the ratio $1 + 1/\sqrt{u - 1}$.

Lemma 4.8 (Zero-prefix guarantee). *Fix $u_3^{\det} \leq u \leq u_4^{\det}$ and set $b = 0$ in algorithm 10. Then*

$$\text{ALG} \leq \left(1 + \frac{1}{\sqrt{u - 1}}\right) \cdot \text{OPT} + O_u(n). \quad (119)$$

Proof. Here $b = 0$, so there is no forced prefix, and the threshold is one. The coordinatewise argument from claim 4.7 reduces every input to four types: deferred jobs with $p = u$, deferred jobs with $p = 1^+$, immediate jobs with $p = 1$, and zero jobs. Divide their class sizes

by n and denote the first three resulting fractions by d, t, m , respectively; put $s = u - 1$. Thus $d, t, m \geq 0$ and $d + t + m \leq 1$. The leading online and offline coefficients are

$$A = \frac{1}{2} + d + t + m - \frac{m^2}{2} + \frac{sd^2}{2}, \quad O = \frac{1}{2} + \frac{(d + t + m)^2}{2} + \frac{(s - 1)d^2}{2}. \quad (120)$$

As before, replacing an immediate value-one job by a deferred value-1⁺ job preserves O in the limit and increases A . We may therefore take $m = 0$ and write $y = d + t$ for the total positive fraction. For fixed y , both A and O are affine functions of d^2 , so their ratio is monotone in d^2 . Its maximum is attained at one of the two extreme compositions:

$$\frac{1 + 2y}{1 + y^2} \quad (d = 0), \quad 1 + \frac{2y}{1 + sy^2} \quad (d = y). \quad (121)$$

The first case has only value-1⁺ positive jobs; the second has only positive jobs with $p = u$. For $d = y$, the desired bound is equivalent to $(\sqrt{sy} - 1)^2 \geq 0$. For $d = 0$, put $r = 1/\sqrt{s}$. The desired inequality is

$$(1 + r)y^2 - 2y + r \geq 0.$$

On the range of the lemma, $s = u - 1 \leq u_4^{\det} - 1 = \varphi^2$, and hence $r(1 + r) \geq 1$. The quadratic on the left has nonpositive discriminant and is therefore nonnegative for every y . Thus both extreme compositions are bounded by $1 + 1/\sqrt{u - 1}$, as required. Binary instances attain this value. $\square$

4.1.3 Adaptive thresholds for the final two parameter intervals

The final algorithm changes its threshold as work accumulates. We prove its guarantee on $[u_4^{\det}, u_5^{\det}]$ and its uniform plateau guarantee for $u \geq u_5^{\det}$. In the proof, a revealed value $p_i = u$ must be treated separately from a smaller value placed in the final queue, because the two have different pair contributions to the offline optimum. Throughout this subsection, $R = 1 + c$, and we retain φ , $u_5^{\det}$, and r from Equations (89) and (94).

For every $c > 0$ define the history-dependent threshold

$$a_c(y) = \begin{cases} 1, & y \leq -1, \\ 1 + \frac{1}{2} \ln\left(\frac{c + 2}{c - 2y}\right), & -1 \leq y \leq 0. \end{cases} \quad (122)$$

The following single algorithm covers both parameter regimes $u_4^{\det} \leq u \leq u_5^{\det}$ and $u \geq u_5^{\det}$. It does not use raw execution, so its schedule does not depend on u .

Algorithm 11 ADAPTIVETHRESHOLD(J, c)

Input: jobs J and parameter $c > 0$.

- 1: Fix the test order $1, \dots, n$.
 - 2: Set $N \leftarrow 0$, $N_{\text{def}} \leftarrow 0$, and $Q \leftarrow \emptyset$.
 - 3: **For** $i = 1, \dots, n$ **do**:
 - 4: Set $x \leftarrow n - i + 1$ and $y \leftarrow (N_{\text{def}} - (1 + c)N)/x$.
 - 5: Set $a \leftarrow a_c(y)$ according to Equation (122).
 - 6: Test job i for one unit and observe p_i .
 - 7: **If** $p_i \leq a$, process job i immediately.
 - 8: **Otherwise**, add job i to Q and set $N_{\text{def}} \leftarrow N_{\text{def}} + 1$.
 - 9: **If** $p_i > 0$, set $N \leftarrow N + 1$.
 - 10: Process the jobs of Q in nondecreasing order of their revealed processing times.
-

For $c \in [r, 1/\varphi]$, let $m = m(c) \in [0, 1/\varphi]$ and $u = u(c)$ be determined by the (c, m) -parametrization in Equation (95). The right-hand side of the first equation is strictly increasing in c , as is the left-hand side in m . Consequently $m(c)$ is increasing and $u(c)$ is decreasing. At the two boundary values of the parameter c , the parametrization gives

$$c = \frac{1}{\varphi} : (m, u) = \left(\frac{1}{\varphi}, u_4^{\det}\right), \quad c = r : (m, u) = (0, u_5^{\det}).$$

Theorem 4.9 (Middle and high revealing upper bounds). *For every finite $u \in [u_4^{\det}, \infty)$, the formally specified algorithm ADAPTIVETHRESHOLD from algorithm 11 satisfies*

$$\text{ALG} \leq \begin{cases} (1+c) \cdot \text{OPT} + O_u(n), & u_4^{\det} \leq u \leq u_5^{\det}, \quad c \text{ given by (95)}, \\ (1+r) \cdot \text{OPT} + O(n), & u \geq u_5^{\det}. \end{cases}$$

The constant in the last $O(n)$ is absolute and uniform over $u \in [u_5^{\det}, \infty)$. In particular these are upper bounds on the deterministic size-asymptotic competitive ratio.

We prove the theorem through six claims. The first two rewrite $\text{ALG} - R \cdot \text{OPT}$ as a sum of contributions from ordered pairs of jobs plus $O_u(n)$ one-job terms, reduce the processing times to five boundary values, and assign the resulting pair contributions to local charges ℓ_i . The next three claims construct and calibrate a nonnegative potential W such that, for every endpoint type T with state-update direction v_T ,

$$\ell_T + \nabla W \cdot v_T \leq 0.$$

Here $\nabla W \cdot v_T$ is the first-order change obtained by differentiating W in the direction of that state update. The sixth claim uses a Taylor estimate to compare this directional derivative with the actual potential change caused by one job, and then telescopes the resulting finite inequalities.

Common setup. Fix a finite raw-execution cap $u \geq u_4^{\det}$ and put $R = 1 + c$. Use ADAPTIVETHRESHOLD(J, c) from algorithm 11. When $u \geq u_5^{\det}$, set $c = r$. For the analysis, immediately before a test let x be the number of jobs not yet tested, N the number of earlier tests that revealed a positive value, and N_{def} the number of those jobs placed in the final queue. Thus

$$y = \frac{N_{\text{def}} - RN}{x} \leq 0 \tag{123}$$

and the threshold is given by Equation (122). When $u_4^{\det} \leq u \leq u_5^{\det}$, the parametrization Equation (95) gives $c \geq r$, and hence

$$a_c(y) \leq a_c(0) \leq a_r(0) = \frac{1}{r} < u - 1. \tag{124}$$

When $u \geq u_5^{\det}$, we set $c = r$, so Equation (124) holds there as well. Thus every revealed value $p_i \in [u - 1, u]$, for which the offline effective length has reached the offline cap u , is placed in the final queue.

The state y never increases. Indeed, the numerator in (123) changes by 0, $-R$, or $-c$ after a zero, a positive job processed immediately after its test, or a positive job placed in the final queue, respectively, while x decreases. This monotonicity will allow us to allocate every pair cost when its relevant endpoint becomes known.

Claim 4.10 (Reduction to five endpoint types). *Fix a decision transcript of ADAPTIVETHRESHOLD: for each tested job, fix whether its revealed value is zero or positive and whether the job is processed immediately or placed in the final queue. This also fixes the state and threshold before every subsequent test. Among all processing-time vectors consistent with this transcript, the supremum of the pairwise term Γ_n from Equation (128) is unchanged if every job is restricted to one of the five endpoint types in Equation (130). The symbols 0^+ and $a_c(y)^+$ denote one-sided limits at which the fixed transcript is preserved.*

Proof. Write

$$\bar{p}_i = \min\{p_i, u - 1\}.$$

By Equation (2) and lemma 2.1, the clairvoyant revealing optimum is SPT on jobs of effective lengths $1 + \bar{p}_i = \min\{1 + p_i, u\}$. For two jobs tested in the order $i < j$, let

$$e_{ij} = \begin{cases} (p_i - p_j)_+, & i \text{ is processed immediately,} \\ 1 + (p_j - p_i)_+, & i \text{ is placed in the final queue and } j \text{ is processed immediately,} \\ 1, & i \text{ and } j \text{ are placed in the final queue.} \end{cases} \quad (125)$$

The first line applies whenever i is processed after its test, regardless of the action taken on j . Pairwise completion-time accounting for the online schedule gives

$$\text{ALG} = \sum_i (1 + p_i) + \sum_{i < j} (1 + \min\{p_i, p_j\} + e_{ij}), \quad (126)$$

whereas the common offline pair identity Equation (4) gives

$$\text{OPT} = \sum_i (1 + \bar{p}_i) + \sum_{i < j} (1 + \min\{\bar{p}_i, \bar{p}_j\}). \quad (127)$$

Collecting the pair contributions in $\text{ALG} - R \cdot \text{OPT}$, define

$$\Gamma_n = \sum_{i < j} (e_{ij} + \min\{p_i, p_j\} - R \min\{\bar{p}_i, \bar{p}_j\}). \quad (128)$$

Equations (126)–(128) yield the exact identity

$$\text{ALG} - R \cdot \text{OPT} = \Gamma_n - c \binom{n}{2} + \sum_i [(1 + p_i) - R(1 + \bar{p}_i)]. \quad (129)$$

The final sum contains one term per job, rather than one per pair, and is nonpositive. If $p_i < u - 1$, so that the offline cap u is inactive, its i -th summand is $-c(1 + p_i)$. If $p_i \geq u - 1$, its offline effective length is u and the summand is at most $1 - cu \leq 0$. Thus it is enough to control Γ_n .

Now fix the decision transcript described in the claim. It determines all later threshold states because these states depend only on whether each earlier value was zero or positive and whether it was processed immediately or placed in the final queue. Varying a processing value inside an interval that preserves this transcript therefore does not change any later threshold. On each such interval, (128) is convex in that value, so its maximum occurs at a boundary. The resulting five endpoint types are

type	value	action
Z	0	process after the test
E	0^+	process after the test
M	$a_c(y)$	process after the test
Q	$a_c(y)^+$	place in the final queue
U	u	place in the final queue.

(130)

Here 0^+ means a positive value tending to zero, and $a_c(y)^+$ means a value tending to the current threshold from above. The types E and Q are kept distinct from Z and M , respectively, because approaching either boundary from the other side produces a different state update or action. For a job processed after its test, the positive interval has endpoints 0^+ and $a_c(y)$. If a value $p_i < u - 1$ is placed in the final queue, its interval has endpoints $a_c(y)^+$ and $u - 1$. On $[u - 1, u]$ the offline contribution is constant and the online contribution is nondecreasing, so u is worst. Thus the intermediate value $u - 1$ can be moved further to u . This proves the claimed reduction without changing any later threshold decision. $\square$

Claim 4.11 (Local pair allocation). *Assign every term of Γ_n for a pair $i < j$ to the later endpoint j . The resulting total charge is at most the sum of the local quantities ℓ_i defined below. To express them, distinguish earlier endpoints of type E from the positive endpoint types M, Q, U . Let*

$$P = \#\{\text{earlier } M, Q, U\}, \quad E = \#\{\text{earlier } E\}, \quad N_{\text{def}} = \#\{\text{earlier } Q, U\},$$

and let K be the number of earlier endpoints of type U . Let I, d be the numbers of earlier endpoints of types M, Q , respectively, and let A_I, A_D be the corresponding sums of their processing values. Thus $P = I + d + K$ and $N_{\text{def}} = d + K$.

For a current endpoint of type Z, E, M , or Q , with $a = a_c(y)$, the allocated charges are bounded by

$$\ell_Z = \ell_E = N_{\text{def}}, \quad \ell_M = N_{\text{def}} + a(x + N_{\text{def}} - RP), \quad \ell_Q = N_{\text{def}} + a(N_{\text{def}} - RP). \quad (131)$$

For a current endpoint U , representing $p = u$, direct pair counting gives the exact charge

$$\begin{aligned} \ell_U &= N_{\text{def}} + A_D + uK - R(A_I + A_D + (u - 1)K) \\ &= d - RA_I - cA_D + Kh_U, \quad h_U = 1 + u - R(u - 1). \end{aligned} \quad (132)$$

Proof. To verify these formulas, pair the current job once with each earlier endpoint type and use Equation (128). An earlier job processed after its test was already charged for the delay that its processing causes when it appeared. Hence a current endpoint U , whose value is u , contributes only $-Rp$ against such an earlier job. Against an earlier endpoint Q of value p it contributes $p - Rp = -cp$, and against an earlier endpoint U it contributes h_U . Summing over the five endpoint types gives

$$\Gamma_n \leq \sum_i \ell_i; \quad (133)$$

the difference is exactly the sum of nonnegative extra delays by which the displayed M charges overestimate the true pair contributions. Formula (132), rather than a substitution $p = u$ into ℓ_Q , is essential before the U -endpoints have been put into a prefix.

Threshold monotonicity gives $A_I \geq aI$ and $A_D \geq ad$. Comparing (132) with the hypothetical Q -charge at the same state gives the exact identity

$$\begin{aligned} \ell_U - \ell_Q &= R(aI - A_I) + c(ad - A_D) + K[1 - c(u - 1 - a)] \\ &\leq K[1 - c(u - 1 - a)]. \end{aligned} \quad (134)$$

$\square$

Claim 4.12 (Base potential for types Z, E, M, Q). *There is a nonnegative potential W_{base} for which the directional-derivative bound*

$$\ell_T + \nabla W_{\text{base}} \cdot v_T \leq 0$$

holds for every $T \in \{Z, E, M, Q\}$. The type U , representing $p = u$, is handled separately in claim 4.13.

Proof. Define

$$\eta = \frac{N_{\text{def}} - RP}{x}, \quad b = \frac{E}{x}, \quad y = \eta - Rb. \quad (135)$$

Call $-1 \leq y \leq 0$ Region I, and in this region put

$$f_c(y) = \frac{c+2}{4} + \frac{c-2y}{4} \left(\ln \frac{c-2y}{c+2} - 1 \right), \quad (136)$$

$$h_c(y) = a_c(y)(1+y),$$

$$G_c(y, b) = f_c(y) + bh_c(y) + \frac{R}{2}b^2. \quad (137)$$

Call $y \leq -1$ Region II, and in this region set instead

$$G_c(y, b) = g_c(\eta), \quad g_c(\eta) = \frac{(1+\eta)_+^2}{2R}. \quad (138)$$

The two definitions are C^1 across $y = -1$. Indeed, feasibility implies $0 \leq b \leq 1/R$ there, and both sides have value $Rb^2/2$ and derivatives

$$(G_y, G_b) = (b, Rb).$$

They are also nonnegative: $f_c(-1) = 0$, $f'_c = a_c - 1 \geq 0$, and $h_c, b, g_c \geq 0$.

Define the base potential

$$W_{\text{base}}(x, P, E, N_{\text{def}}) = xN_{\text{def}} + x^2G_c(y, b). \quad (139)$$

In the unnormalized state vector $(x, P, E, N_{\text{def}}, K)$, revealing one endpoint of type T changes the coordinates in the direction v_T shown below:

	x	P	E	N_{def}	K
v_Z	-1	0	0	0	0
v_E	-1	0	1	0	0
v_M	-1	1	0	0	0
v_Q	-1	1	0	1	0
v_U	-1	1	0	1	1.

(140)

Let $J_T = \ell_T + \nabla W_{\text{base}} \cdot v_T$. This is the local charge plus the directional derivative of the potential; we must show $J_T \leq 0$. For the moment $T \in \{Z, E, M, Q\}$. In Region I, direct differentiation and the identities

$$\begin{aligned} -2f_c + (y-R)f'_c + a_c(1+y) &= 0, \\ 1 - 2f_c + (y-c)f'_c + a_c y &= 0 \end{aligned} \quad (141)$$

give

$$\frac{J_Z}{x} = (-2f_c + yf'_c) + b(-h_c + yh'_c),$$

$$\begin{aligned}
\frac{J_E}{x} &= b[-h_c + (y - R)h'_c + R], \\
\frac{J_M}{x} &= b[-h_c + (y - R)h'_c + a_c R], \\
\frac{J_Q}{x} &= b[-h_c + (y - c)h'_c + a_c R].
\end{aligned} \tag{142}$$

All four quantities are nonpositive. For the first one,

$$-2f_c + yf'_c = a_c(c - y) - R \leq 0; \tag{143}$$

the right-hand side is zero at $y = -1$ and is nonincreasing because $a'_c(c - y) - a_c \leq 1 - a_c \leq 0$. Also $h_c, h'_c \geq 0$. For the remaining three drifts, observe that

$$h_c + (c - y)h'_c = a_c R + (c - y)(1 + y)a'_c \geq a_c R. \tag{144}$$

The Q -bracket in (142) is therefore nonpositive; the M -bracket is the Q -bracket minus h'_c , and the E -bracket is the M -bracket minus $R(a_c - 1)$.

In Region II, substituting $W = x^2 g_c(\eta)$ into the definitions of the four drifts gives

$$\begin{aligned}
J_Z/x &= J_E/x = -2g_c + \eta g'_c, \\
J_M/x &= -2g_c + (\eta - R)g'_c + 1 + \eta, \\
J_Q/x &= 1 - 2g_c + (\eta - c)g'_c + \eta.
\end{aligned} \tag{145}$$

For $-1 \leq \eta \leq 0$, substitution of $g'_c = (1 + \eta)/R$ makes these values $-(1 + \eta)/R$, $-(1 + \eta)/R$, and 0, respectively. For $\eta \leq -1$ their signs are immediate from $g_c = g'_c = 0$. Hence $J_T \leq 0$ for $T \in \{Z, E, M, Q\}$, as claimed. $\square$

Claim 4.13 (Reserve potential for jobs with $p = u$). *There is a nonnegative reserve term W_{cap} , depending on the number K of earlier U endpoints, such that the full potential $W = W_{\text{base}} + W_{\text{cap}}$ satisfies*

$$\ell_T + \nabla W \cdot v_T \leq 0 \quad (T \in \{Z, E, M, Q, U\}).$$

Proof. Define

$$Z(c) = 2 + \frac{1}{c} + \frac{1}{2} \ln\left(1 + \frac{2}{c}\right), \quad \delta = Z(c) - u. \tag{146}$$

For $u_4^{\text{det}} \leq u \leq u_5^{\text{det}}$, with (c, m) chosen by Equation (95), the definition of δ gives

$$\delta = \text{artanh } m. \tag{147}$$

For $0 \leq t \leq m$, set

$$\beta(t) = ct[\delta - \text{artanh } t], \quad \mathcal{V}(t) = \int_t^m \beta(s) ds, \tag{148}$$

and extend $\mathcal{V}$ by zero for $t \geq m$. Since $\beta(m) = 0$, this extension is C^1 . With

$$q = \frac{K}{x}, \quad \mu = \frac{q}{1 + q}, \quad \mathcal{H}(q) = (1 + q)^2 \mathcal{V}(\mu), \tag{149}$$

define the full potential

$$W = xN_{\text{def}} + x^2(G_c(y, b) + \mathcal{H}(q)). \tag{150}$$

Thus $W_{\text{cap}} = x^2 \mathcal{H}(q)$. Both terms in W are nonnegative.

For an endpoint of type Z, E, M , or Q , the value of K is fixed while x decreases, so the reserve term contributes, after division by x ,

$$L_0(q) = -2\mathcal{H}(q) + q\mathcal{H}'(q). \quad (151)$$

An endpoint U also increments K , and its contribution is $L_0 + \mathcal{H}'$. On the range $\mu \leq m$, where $\mathcal{V}$ may be nonzero, differentiation of (149) gives

$$\mathcal{H}' = 2(1+q)\mathcal{V} - \beta(\mu), \quad L_0 = -2(1+q)\mathcal{V} - q\beta(\mu) \leq 0. \quad (152)$$

Thus the reserve term can only decrease the left-hand side of the directional-derivative bound for each of Z, E, M, Q .

For a current endpoint U , feasibility $N_{\text{def}} \leq P$ and $P \geq K$ gives

$$y = \frac{N_{\text{def}} - R(P + E)}{x} \leq -\frac{cK}{x} = -cq. \quad (153)$$

While $\mu \leq m$, we have $q \leq m/(1-m) \leq \varphi$ and hence $cq \leq 1$. Monotonicity of a_c , together with (134), yields

$$\frac{\ell_U - \ell_Q}{x} \leq cq \left[\delta - \frac{1}{2} \ln(1+2q) \right] =: B(q). \quad (154)$$

The change of variables in (149) satisfies

$$\text{artanh } \mu = \frac{1}{2} \ln(1+2q), \quad (1+q)\beta(\mu) = B(q). \quad (155)$$

Consequently the extra positive charge of an endpoint of type U is cancelled exactly by the directional derivative of the reserve term:

$$L_0 + \mathcal{H}' + B(q) = 0 \quad (\mu \leq m). \quad (156)$$

When $\mu \geq m$, the reserve term vanishes. At $q_* = m/(1-m)$, the bracket in (134), evaluated at the largest possible threshold $a_c(-cq_*)$, is zero by (147) and (155). For $q \geq q_*$, (153) only decreases the threshold, so that bracket is nonpositive. The directional-derivative bound for type U therefore holds for every state and for arbitrary interleavings of type U with the other four types. $\square$

Claim 4.14 (Initial calibration). *The full potential starts with exactly the amount needed to cancel the constant pairwise term: $W_{\text{initial}} = cn^2/2$.*

Proof. Initially $y = b = q = 0$, and

$$f_c(0) = \frac{1}{2} - \frac{c}{4} \ln\left(1 + \frac{2}{c}\right). \quad (157)$$

Elementary integration in (148) gives

$$\mathcal{V}(0) = \frac{c}{2}(\delta - m). \quad (158)$$

Indeed,

$$\int_0^m t \text{artanh } t \, dt = \frac{(m^2 - 1) \text{artanh } m + m}{2}.$$

Using (95) and (147), we obtain the exact calibration

$$f_c(0) + \mathcal{H}(0) = f_c(0) + \mathcal{V}(0) = \frac{c}{2}. \quad (159)$$

Thus $W_{\text{initial}} = cn^2/2$. $\square$

Claim 4.15 (Finite transitions and telescoping). *Let z_i be the state before job i is revealed, write $W_i = W(z_i)$, and let T_i be its endpoint type. For $u_4^{\text{det}} \leq u \leq u_5^{\text{det}}$, each actual transition satisfies*

$$W_{i+1} - W_i = \nabla W(z_i) \cdot v_{T_i} + O_u(1).$$

Together with claims 4.12 and 4.13, this gives

$$\text{ALG} \leq (1 + c) \cdot \text{OPT} + O_u(n). \quad (160)$$

For every finite $u \geq u_5^{\text{det}}$, there is an absolute constant B , independent of u , such that

$$\text{ALG} \leq (1 + r) \cdot \text{OPT} + Bn. \quad (161)$$

Proof. We first verify the needed regularity on feasible states. In Region I, $y \geq -1$ implies

$$-xy = R(P + E) - N_{\text{def}} = cP + (P - N_{\text{def}}) + RE \leq x. \quad (162)$$

All terms in the middle expression are nonnegative. Hence $P/x, N_{\text{def}}/x, q \leq 1/c$ and $b \leq 1/R$. The normalized arguments of G_c in (139) therefore stay in a bounded set, and all derivatives needed for the Taylor estimate of one transition are bounded. In Region II the unnormalized base potential is simply

$$xN_{\text{def}} + \frac{(x + N_{\text{def}} - RP)_+^2}{2R}, \quad (163)$$

which has bounded one-sided second derivatives. The calculation following Equation (138) shows that the two pieces are C^1 at $y = -1$, and the positive-part square is C^1 at $\eta = -1$.

For the additional term $\mathcal{R}(x, K) = x^2 \mathcal{H}(K/x)$, whenever $0 \leq q \leq q_*$ one has bounded $\mathcal{H}''$, and

$$\mathcal{R}_{xx} = 2\mathcal{H} - 2q\mathcal{H}' + q^2\mathcal{H}'', \quad \mathcal{R}_{xK} = \mathcal{H}' - q\mathcal{H}'', \quad \mathcal{R}_{KK} = \mathcal{H}''. \quad (164)$$

For $q \geq q_*$, $\mathcal{R} = 0$; moreover $\mathcal{H}(q_*) = \mathcal{H}'(q_*) = 0$. Thus the join is C^1 with bounded one-sided Hessians. At $x = 0$, set $\mathcal{R} = 0$: for $K > 0$ it is already locally zero, while $\mathcal{R}(\theta x, \theta K) = \theta^2 \mathcal{R}(x, K)$ gives zero gradient at $(x, K) = (0, 0)$. Altogether W is continuously differentiable with a Lipschitz gradient on all states that the algorithm can reach. For $u_4^{\text{det}} \leq u \leq u_5^{\text{det}}$, its regularity constant may depend on the fixed value of u . When $c = r$ and $m = 0$, however, the additional term is identically zero and the base potential depends only on the absolute constant r ; its regularity constant is therefore uniform over all $u \geq u_5^{\text{det}}$. This includes the last transition from $x = 1$ to $x = 0$.

It follows that each transition caused by one job has Taylor remainder

$$\ell_i + W_{i+1} - W_i \leq O_u(1). \quad (165)$$

The endpoint $E = 0^+$ is interpreted one-sidedly. For a fixed sequence of endpoint types, replace all E coordinates by positive values ε . Its state trajectory is unchanged, and the exact finite pair objective, being a finite sum of affine and minimum terms, converges as $\varepsilon \downarrow 0$. Equivalently, retain an $O(\varepsilon n^2)$ error and take this limit for each fixed n . Therefore the directional-derivative bound for the formal type E transfers to actual input vectors without an asymptotic loss.

Summing (165), using $W \geq 0$, and applying (159) gives

$$\sum_i \ell_i \leq \frac{c}{2} n^2 + O_u(n). \quad (166)$$

Together with (129) and (133), this yields

$$\text{ALG} - R \cdot \text{OPT} \leq -c \binom{n}{2} + \frac{c}{2} n^2 + O_u(n) = O_u(n). \quad (167)$$

At $u = u_5^{\det}$, one has $c = r$, $m = 0$, and $Z(r) = u_5^{\det}$. The reserve term for type U is identically zero, while (159) remains valid. If u is increased above $u_5^{\det}$, the bracket controlling an endpoint of type U , $1 - r(u - 1 - a_r(y))$ decreases, and the contribution of an individual job of type U in (129) also decreases.

Here is the promised uniform finite calculation. The base potential and the set of states reachable by the algorithm contain no u , so there is one constant C_{step} , depending only on r , for which, whenever $u \geq u_5^{\det}$, every transition, whether its endpoint is U or not, satisfies

$$\ell_i + W_{i+1} - W_i \leq C_{\text{step}}.$$

Because $W_0 = rn^2/2$ and $W_n = 0$, telescoping gives

$$\sum_i \ell_i \leq \frac{r}{2} n^2 + C_{\text{step}} n. \quad (168)$$

In the final sum over individual jobs in Equation (129), every endpoint other than U has $p_i < u - 1$ and contributes $-r(1 + p_i)$. The endpoint U has $p_i = u$ and contributes at most $1 - ru \leq 0$. Hence this sum is nonpositive uniformly on $[u_5^{\det}, \infty)$. Combining Equations (129), (133) and (168) now yields

$$\text{ALG} - (1 + r) \cdot \text{OPT} \leq -r \binom{n}{2} + \frac{r}{2} n^2 + C_{\text{step}} n = \left(C_{\text{step}} + \frac{r}{2} \right) n. \quad (169)$$

Thus, with $B = C_{\text{step}} + r/2$,

$$\text{ALG} \leq (1 + r) \cdot \text{OPT} + Bn \quad \text{for every finite } u \in [u_5^{\det}, \infty), \quad (170)$$

and B is independent of u . The estimate is genuinely uniform rather than the result of sending a parameter-dependent remainder to infinity. This proves the claim. $\square$

We can now prove the theorem.

Proof of Theorem 4.9. Claims 4.10 and 4.11 reduce the competitive excess to the local charges. The potentials in claims 4.12 and 4.13 satisfy the required directional-derivative bounds, and claim 4.14 gives their initial value. Claim 4.15 then yields both displayed finite bounds.

Finally, $u \geq u_4^{\det} > 1$, so every effective offline length is at least one and $\text{OPT} \geq n(n+1)/2$. Hence the additive terms vanish in the size-asymptotic ratios, proving the final assertion. $\square$

4.2 Lower bounds

The lower bounds use three adversarial constructions. A hidden binary stopping game covers $0 < u \leq u_4^{\det}$. For $u_4^{\det} \leq u \leq u_5^{\det}$, we prepend a block of jobs with $p_j = u$ to a scaled harmonic construction. Finally, for $u \geq u_5^{\det}$, we use the obligatory-testing harmonic instances with $1 + p_j < u$ for every job. On these instances the offline cap u is inactive, and the transfer lemma preserves the obligatory-testing lower bound.

4.2.1 The binary stopping game and its piecewise lower bound

Binary instances $p_j \in \{0, u\}$ give the lower bounds 1 for $0 < u \leq u_1^{\det}$, u for $u_1^{\det} \leq u \leq u_2^{\det}$, $R_{\text{quad}}(u)$ for $u_2^{\det} \leq u \leq u_3^{\det}$, and $1 + 1/\sqrt{u-1}$ for $u_3^{\det} \leq u \leq u_4^{\det}$. The stopping point must be hidden from the algorithm; announcing it in advance gives a strictly weaker lower bound. In this subsection, a job with $p_j = 0$ is a *zero job*, and one with $p_j = u$ is a *long job*.

Lemma 4.16 (Hidden binary stopping with raw execution). *Fix $u > 1$, a target $R \leq u$, and a stopping fraction $\alpha \in (0, 1)$. Against a deterministic algorithm, there is a legal binary adversary whose leading normalized excess $2(\text{ALG} - R \cdot \text{OPT})/n^2$ at its stopping time is, up to $O_u(1/n)$,*

$$(u - R)(1 - \sigma^2) + \sigma^2 f_{u,R}(y, b), \quad (171)$$

where $0 \leq b \leq y \leq 1$ and

$$0 \leq y - \alpha < \frac{1}{n - v}. \quad (172)$$

Moreover,

$$f_{u,R}(y, b) = 1 - R + 2y + (u - 1)(1 - R)y^2 + b^2 + 2b(u - 1 - uy). \quad (173)$$

Here $\sigma \in [0, 1]$ is the fraction of jobs not already executed raw, y is the fraction of those jobs revealed to be long, and b is the fraction both revealed long and already completed. Consequently, if

$$\min_{0 \leq b \leq \alpha} f_{u,R}(\alpha, b) \geq 0,$$

then the adversary gives ratio at least $R - O_u(1/n)$.

Proof. Until the stopping time, answer every test of a fresh job with $p = u$; if the algorithm instead runs a fresh job raw, assign that job the hidden value zero. Let v be the number of raw first touches and L the number of tests, all of which have revealed long jobs. Stop immediately after the first touch for which

$$L \geq \alpha(n - v). \quad (174)$$

All untouched jobs are then assigned value zero. Only for the purpose of lower-bounding the remaining cost, we grant the algorithm this future information and allow an optimal continuation; in the actual interaction, values are still revealed only by tests. If the line is never crossed, every job was run raw: if even one job had been tested, then after the last first touch the tested fraction would be one. On the all-zero hidden instance the ratio is $u \geq R$.

At a crossing, let e of the L revealed long jobs already be complete, let $d = L - e$ be tested but still pending, and let $x = n - v - e - d$ jobs remain untouched. To obtain a lower bound on the algorithm's cost, we give it the benefit of rearranging its work into the following cheaper schedule: first the v raw completions, then the tests and processing of the e completed long jobs, then the d tests whose long jobs remain pending, then the x zero test-completions, and finally the processing of the d pending long jobs.

Here is why the rearrangement can only decrease total completion time. If a noncompleting unit test is immediately before an available length- u completion while h jobs remain, the two orders contribute $h + hu$ and $hu + (h - 1)$, respectively; moving the completion left saves one. Raw completions are available from time zero, and identical long jobs may be relabeled so that the e completed ones are tested first. After the switch, a zero

test-completion has length one and therefore precedes a pending operation of length u . Repeating only these explicit neighboring swaps produces the displayed relaxed schedule and respects every test-before-processing precedence constraint.

The exact costs of this relaxed schedule and of OPT are

$$A_{\text{ex}} = u \left(vn - \frac{v(v-1)}{2} \right) + (1+u) \left(e(n-v) - \frac{e(e-1)}{2} \right) + d(n-v-e) + x(d+x) - \frac{x(x-1)}{2} + u \frac{d(d+1)}{2}, \quad (175)$$

$$O_{\text{ex}} = \frac{n(n+1)}{2} + (u-1) \frac{(e+d)(e+d+1)}{2}. \quad (176)$$

Put $\nu = v/n$, $\delta = (v+e+d)/n$, and $\lambda = e/n$. Twice the quadratic coefficients in Equations (175) and (176) are

$$\begin{aligned} \mathcal{A} &= 1 + 2\delta(1 - u\nu) + (u-1)\delta^2 + 2\nu(\nu + u - 2) \\ &\quad + \lambda^2 + 2\lambda(\nu + u - 1 - u\delta), \\ \mathcal{O} &= 1 + (u-1)(\delta - \nu)^2. \end{aligned}$$

Now put $\sigma = 1 - \nu$, $y = (e+d)/(n-v)$, and $b = e/(n-v)$. Thus σ is the nonraw fraction of the original instance, while y and b are the long and already-completed fractions within those nonraw jobs. Direct expansion gives exactly Equation (171) after normalizing by $n^2/2$. The first-crossing overshoot in Equation (174) is less than one: a test increases $e+d-\alpha(n-v)$ by one and a raw first touch increases it by α . This proves Equation (172), and in particular $\sigma(y - \alpha) < 1/n$. The minimization in the lemma uses $b \leq \alpha$, whereas the actual value only satisfies $b \leq y$ and may exceed α by the one-job overshoot. Put $\bar{b} = \min\{b, \alpha\}$. Then $0 \leq b - \bar{b} \leq y - \alpha$. Since $f_{u,R}$ is Lipschitz on the unit square,

$$\sigma^2 f_{u,R}(y, b) \geq \sigma^2 f_{u,R}(\alpha, \bar{b}) - O_u(1/n). \quad (177)$$

Thus the stated minimum over $0 \leq \bar{b} \leq \alpha$ suffices uniformly even when $n-v$ is small. The omitted one-job terms are $O_u(n)$ in the unnormalized costs. Finally, freeze every revealed or hidden answer on its job label. Because the algorithm is deterministic and a raw action never reveals its assigned zero, rerunning it on this fixed vector reproduces the same transcript. The adaptive description therefore yields a legal fixed hard instance. $\square$

Proposition 4.17 (Binary lower curve). *Every deterministic algorithm has size-asymptotic ratio at least*

$$B(u) = \begin{cases} 1, & 0 < u \leq 1, \\ u, & u_1^{\text{det}} \leq u \leq u_2^{\text{det}}, \\ R_{\text{quad}}(u), & u_2^{\text{det}} \leq u \leq u_3^{\text{det}}, \\ 1 + \frac{1}{\sqrt{u-1}}, & u \geq u_3^{\text{det}}. \end{cases} \quad (178)$$

The lower bound uses only $p_j \in \{0, u\}$.

We optimize the stopping lemma in three steps. First we minimize over the fraction b of revealed long jobs that the algorithm has already completed. This produces two quadratic formulas, according to whether the minimizing b is positive or zero. We then locate their transition points and choose the stopping fraction α at which the relevant quadratic is largest.

Claim 4.18 (Minimization over completed long jobs). *For fixed revealed-long fraction y , minimizing Equation (173) over the completed-long fraction b gives*

$$b_*(y) = \max\{0, uy - (u - 1)\}. \quad (179)$$

The upper constraint $b \leq y$ never binds, because $uy - (u - 1) \leq y$ for $y \leq 1$. When $uy - (u - 1) > 0$, the minimum equals

$$\min_b f_{u,R}(y, b) = 1 - R - (u - 1)^2 + 2(u^2 - u + 1)y - (u^2 - u + 1 + (u - 1)R)y^2. \quad (180)$$

and its unconstrained maximum over y occurs at

$$\alpha = \frac{u^2 - u + 1}{u^2 - u + 1 + (u - 1)R}.$$

This maximum is zero exactly when

$$(1 - R - (u - 1)^2)(u^2 - u + 1 + (u - 1)R) + (u^2 - u + 1)^2 = 0. \quad (181)$$

which is equivalent to Equation (92). When $uy - (u - 1) \leq 0$, the minimum equals

$$\min_b f_{u,R}(y, b) = 1 - R + 2y + (u - 1)(1 - R)y^2. \quad (182)$$

and its unconstrained maximum is zero for

$$R = 1 + \frac{1}{\sqrt{u - 1}}, \quad \alpha = \frac{1}{\sqrt{u - 1}}. \quad (183)$$

This maximizing point indeed lies in the region where $b = 0$ if and only if $(u - 1)^3 \geq u^2$, with equality at $u = u_3^{\det}$.

Proof. The part of $f_{u,R}(y, b)$ that depends on b is $b^2 + 2b(u - 1 - uy)$, whose unconstrained minimizer is $uy - (u - 1)$. Taking the maximum with zero gives Equation (179); its upper bound follows from $uy - (u - 1) \leq y$. Substitution gives the two displayed quadratics. Differentiating each concave quadratic in y gives the stated maximizing value of α . Substitution of that value gives Equation (181) in the positive- b case and Equation (183) in the zero- b case. Expanding the former gives Equation (92); substituting the latter value of α into $u\alpha - (u - 1) \leq 0$ gives $(u - 1)^3 \geq u^2$. $\square$

Claim 4.19 (The transition from the ratio u). *For $R = u$, the maximum in Equation (180) is nonnegative exactly while*

$$[u(u - 1)]^2 - u(u - 1) - 1 \leq 0, \quad (184)$$

whose positive equality is $u = u_2^{\det}$.

Proof. Substitute $R = u$ and the maximizing value of α from claim 4.18 into Equation (180). After multiplication by its positive denominator, nonnegativity is equivalent to the displayed inequality. With $x = u(u - 1)$, its equality is $x^2 - x - 1 = 0$, so $x = \varphi$. Since $u(u - 1)$ is strictly increasing for $u > 1$, the unique positive solution is

$$u = \frac{1 + \sqrt{1 + 4\varphi}}{2} = \frac{1 + \sqrt{3 + 2\sqrt{5}}}{2} = u_2^{\det}.$$

$\square$

Claim 4.20 (Stopping fractions for the three nontrivial intervals). *For $u_1^{\det} < u \leq u_2^{\det}$, use $R = u$ and the maximizing point of Equation (180). For $u_2^{\det} \leq u \leq u_3^{\det}$, use the positive root $R_{\text{quad}}(u)$ and $\alpha = (u^2 - u + 1)/(u^2 - u + 1 + (u - 1)R_{\text{quad}}(u))$; the zero-maximum identity makes the maximum zero. The minimizing completed-long fraction b is positive in the interior and becomes zero at $u = u_3^{\det}$. For $u \geq u_3^{\det}$, use the zero- b maximizing point from Equation (183). Each choice satisfies*

$$\min_{0 \leq b \leq \alpha} f_{u,R}(\alpha, b) \geq 0.$$

Proof. For $u_1^{\det} < u \leq u_2^{\det}$, the assertion follows from claim 4.19. For $u_2^{\det} \leq u \leq u_3^{\det}$, the defining equation for $R_{\text{quad}}(u)$ is the zero-maximum identity in claim 4.18; the minimizing completed-long fraction is positive in the interior and becomes zero at $u = u_3^{\det}$. For $u \geq u_3^{\det}$, the zero- b formula and its feasibility are exactly Equation (183) and the final assertion of claim 4.18. $\square$

We can now prove the proposition.

Proof of Proposition 4.17. For $u \leq 1$, the lower bound $R_{\text{RO}}^{\det}(u) \geq 1$ is immediate. For $u > 1$, use the target ratio and stopping fraction from claim 4.20 in lemma 4.16, and let the number of jobs tend to infinity. The recording step in that lemma converts every adaptive interaction into a fixed binary instance. This proves all four formulas in Equation (178). $\square$

4.2.2 The harmonic construction for short jobs

This subsection develops the harmonic construction used for the lower bounds on $u_4^{\det} \leq u \leq u_5^{\det}$ and on $u \geq u_5^{\det}$. Its largest processing time will be below two. Consequently, whenever the construction is later used with a raw-execution cap $u > 3$, every job satisfies $1 + p_j < u$, so the offline cap u is inactive. Its notation is local to this subsection. We first record its extremal specialization and then extract the scaled form used when $u_4^{\det} \leq u \leq u_5^{\det}$. The construction has many equally large groups of positive jobs, revealed from longest to shortest, followed by one large group of zero jobs. The only design choice is the gap between two consecutive positive processing times. Choosing these gaps harmonically will make all decisions of the online algorithm disappear from the leading term.

Theorem 4.21 (Extremal harmonic core). *For every deterministic online algorithm $\mathcal{B}$ and every $\varepsilon > 0$, there are arbitrarily large fixed instances I with unit tests and rational processing times such that*

$$\text{ALG}_{\mathcal{B}}(I) \geq (R_{\text{OT}}^{\det} - \varepsilon) \cdot \text{OPT}(I).$$

We prove the theorem through five claims. They establish, in order, that the adaptive reveal order defines a fixed input, that a shortest-first relaxation lower-bounds every phase, that this relaxation gives an explicit finite accounting formula, that harmonic processing-time gaps remove all algorithm-dependent terms, and that the resulting one-variable limit is maximized at $R_{\text{OT}}^{\det}$.

Common construction. Fix an integer K , a rational number $\xi > 0$, and a rational parameter $\gamma \in (0, 1)$; one may keep $\gamma = 1/2$ throughout. For a large scaling integer H , the instance will contain H jobs of each of K positive types $0, 1, \dots, K - 1$ and ξH zero jobs. We choose H to clear all denominators.

The adversary assigns processing times according to the order in which tests are started. The first H tested jobs receive processing time p_0 , the next H receive p_1 , and so on; after the K positive blocks, every remaining job receives processing time zero. Thus

$$p_0 > p_1 > \cdots > p_{K-1} > 1,$$

so the algorithm sees the positive types from longest to shortest and learns about the zero jobs only at the end.

Claim 4.22 (The reveal order defines a fixed input). *For every deterministic algorithm, assigning each answer produced by this reveal order to the label of the tested job gives a fixed input on which the algorithm repeats the same tests, answers, and completion decisions.*

Proof. This adaptive description is only a convenient way to construct the input. After running the interaction against a fixed deterministic algorithm, assign to every job label the value revealed when that label was tested. Rerunning the algorithm on this fixed assignment produces exactly the same transcript. If the algorithm never tests or completes some job, its cost is infinite and the lower bound is immediate; otherwise every job label is recorded. Hence it is enough to lower-bound the cost under this adaptive reveal order. $\square$

Claim 4.23 (A shortest-first lower bound in each phase). *Let T_ℓ be the start of the first test in positive block ℓ , and let T_K be the start of the first test in the zero block. Phase ℓ contains the completions in $(T_\ell, T_{\ell+1}]$; a completion exactly at T_ℓ belongs to the preceding phase. In this construction, the mass of a group is its number of jobs divided by H . For group masses $q_1, \dots, q_m$ and remaining lengths $v_1 \leq \dots \leq v_m$, define*

$$\mathcal{S}((q_g, v_g)_{g=1}^m) = \sum_{g=1}^m \left(\frac{v_g q_g^2}{2} + q_g \sum_{h < g} v_h q_h \right). \quad (185)$$

If groups of size $q_g H$ finish within one phase, their relative completion cost is at least $H^2 \mathcal{S} + O(H)$. If

$$p_0 - p_{K-1} < 1. \quad (186)$$

then the nondecreasing remaining lengths in a positive phase are

$$p_{\ell-1} < p_{\ell-2} < \cdots < p_0 < 1 + p_\ell.$$

In the final phase, the order starts with the zero jobs of remaining length one and continues with the positive types from shortest to longest.

Proof. Each T_ℓ is the boundary between two nonpreemptive operations. Consequently, a previously tested job that is unfinished at T_ℓ has not started processing and still needs its entire processing time.

Suppose jobs with remaining work $v_1, \dots, v_m$ complete after some time T . If their completion order is $\pi_1, \dots, \pi_m$, then the g th relative completion time is at least $\sum_{h \leq g} v_{\pi_h}$. Summing these prefix inequalities shows that their total relative completion time is minimized by arranging the v_i in nondecreasing order. For groups of size $q_g H$, the within-group sums give $v_g q_g^2 H^2 / 2 + O(H)$, and the waiting caused by shorter groups gives the cross term in Equation (185).

In phase ℓ , an unfinished job of an older type $j < \ell$ has remaining length p_j , whereas a type- ℓ job completed in its own phase needs its test and processing, of total length $1 + p_\ell$. The inequalities in the claim follow from $p_0 - p_{K-1} < 1$. At T_K , each untested zero job has remaining length one, which proves the asserted final-phase order. $\square$

Claim 4.24 (Finite phase accounting). *For each positive type j , let*

- $s_j \in [0, 1]$ *be the mass of type- j jobs completed in their own phase;*
- $d_{j,\ell} \geq 0$ *be the mass of type- j jobs completed in the later phase $\ell > j$; and*
-

$$X_{j,\ell} = s_j + \sum_{h=j+1}^{\ell-1} d_{j,h} \leq 1 \quad (\ell > j)$$

be the mass of type j completed before phase ℓ . Thus $X_{j,\ell+1}$ is the mass completed by the end of phase ℓ .

Then the start of phase ℓ satisfies

$$\frac{T_\ell}{H} \geq \ell + \sum_{j < \ell} p_j X_{j,\ell}. \quad (187)$$

For each phase, charge its completions their common start time from Equation (187), and charge their relative completion times using Equation (185). Do the same at T_K for the remaining positive jobs and the ξH zero jobs. Expanding and collecting equal variables gives

$$\begin{aligned} \frac{\text{ALG}}{H^2} &\geq C_K + \sum_{j=0}^{K-1} \left(\frac{s_j^2}{2} - \theta_j s_j \right) \\ &\quad + \sum_{j < \ell} d_{j,\ell} \left[\ell - j - \theta_j - \sum_{j < k < \ell} (p_j - p_k) X_{k,\ell+1} \right] - O(1/H), \end{aligned} \quad (188)$$

where K, ξ, γ are fixed while $H \rightarrow \infty$, and

$$\theta_j = 1 - \xi(p_j - 1) - \sum_{k > j} (p_j - p_k - 1), \quad (189)$$

$$C_K = \frac{\xi^2}{2} + 2\xi K + K^2 + P_K, \quad P_K = \sum_{j=0}^{K-1} \left(j + \frac{1}{2} \right) p_j. \quad (190)$$

Proof. At time T_ℓ , the machine has already performed the tests of the ℓH jobs in earlier positive blocks and has processed every earlier job that has completed. This gives Equation (187).

For each phase, charge its completions their common start time from Equation (187), and charge their relative completion times using claim 4.23. In phase ℓ , the masses in the shortest-first relaxation are

$$d_{\ell-1,\ell}, d_{\ell-2,\ell}, \dots, d_{0,\ell}, s_\ell$$

with respective lengths

$$p_{\ell-1}, p_{\ell-2}, \dots, p_0, 1 + p_\ell.$$

Multiply their total mass by the right-hand side of Equation (187) and add the functional Equation (185). In the final phase use masses

$$\xi, 1 - X_{K-1,K}, \dots, 1 - X_{0,K}$$

and lengths $1, p_{K-1}, \dots, p_0$. The constant terms sum to C_K . The terms containing only s_j are $s_j^2/2 - \theta_j s_j$; the coefficient of each $d_{j,\ell}$ is precisely the bracket in Equation (188). Thus the displayed identity is just the phasewise shortest-first bound with the phase start times retained. No optimality assumption about the online schedule is hidden in it. $\square$

The meaning of the expression is useful. Completing an s_j fraction of a type in its own phase creates a quadratic completion cost but saves later waiting cost, producing $s_j^2/2 - \theta_j s_j$. A completion postponed to a later positive phase is represented by $d_{j,\ell}$ and its bracket. We next choose the processing times so that every such bracket is nonnegative and all θ_j are equal.

Claim 4.25 (Harmonic gaps remove the algorithm-dependent terms). *Assume Equation (186) and choose the processing times by*

$$p_{K-1} = 1 + \frac{\gamma}{\xi}, \quad p_i = p_{i+1} + \frac{1}{\xi + K - 1 - i} \quad (0 \leq i < K - 1). \quad (191)$$

Then every θ_j in Equation (189) equals

$$\theta_j = 1 - \gamma, \quad (192)$$

every coefficient of $d_{j,\ell}$ in Equation (188) is nonnegative, and

$$\liminf_{H \rightarrow \infty} \frac{\text{ALG}}{H^2} \geq Q_K := C_K - \frac{K}{2}(1 - \gamma)^2. \quad (193)$$

The corresponding offline coefficient is

$$D_K = \frac{\xi^2}{2} + \xi K + \frac{K^2}{2} + P_K, \quad \lim_{H \rightarrow \infty} \frac{\text{OPT}}{H^2} = D_K. \quad (194)$$

Thus Q_K/D_K is a lower bound for the limiting ratio of the finite- K construction.

Proof. Measured upward from the shortest type, the consecutive gaps in Equation (191) are

$$\frac{1}{\xi + 1}, \frac{1}{\xi + 2}, \dots, \frac{1}{\xi + K - 1}.$$

If $m = K - 1 - j$, then

$$p_j - 1 = \frac{\gamma}{\xi} + \sum_{h=1}^m \frac{1}{\xi + h},$$

$$\sum_{k>j} (p_j - p_k) = \sum_{h=1}^m \frac{h}{\xi + h}.$$

Adding the first line multiplied by ξ to the second line makes each denominator cancel:

$$\xi(p_j - 1) + \sum_{k>j} (p_j - p_k) = \gamma + \sum_{h=1}^m \frac{\xi + h}{\xi + h} = \gamma + m. \quad (195)$$

Substitution in Equation (189) gives Equation (192). This cancellation explains the harmonic increments: the factor ξ from the zero block and the multiplicity h from the h shorter positive types add up to the denominator $\xi + h$.

The delayed-completion terms are now harmless as well. Since $X_{k,\ell+1} \leq 1$, their brackets are at least

$$\ell - j - (1 - \gamma) - \sum_{j < k < \ell} (p_j - p_k)$$

$$= \gamma + \sum_{j < k < \ell} (1 - (p_j - p_k)) > 0,$$

where the final inequality follows from Equation (186). We may therefore drop every term containing $d_{j,\ell}$. The remaining minimization is not a multivariable optimization problem: complete one square for each j ,

$$\frac{s_j^2}{2} - (1 - \gamma)s_j = \frac{1}{2}(s_j - (1 - \gamma))^2 - \frac{1}{2}(1 - \gamma)^2.$$

Because $1 - \gamma \in (0, 1)$, this minimum is feasible for $s_j \in [0, 1]$, which proves Equation (193).

Knowing all processing times, the optimum runs the zero jobs first and the positive types from shortest to longest. Equivalently, the pairwise contribution of two jobs is $1 + \min\{p_i, p_j\}$. The four terms respectively account for zero-zero pairs, zero-positive pairs, the unit-test part of positive-positive pairs, and the processing part of positive-positive pairs. This gives Equation (194) and completes the proof. $\square$

Claim 4.26 (Limit and one-variable maximization). *Fix $0 < \alpha < e - 1$ and set $\xi = K/\alpha$. As $K \rightarrow \infty$, the processing times in Equation (191) satisfy Equation (186), and*

$$\lim_{K \rightarrow \infty} \frac{Q_K}{D_K} = R(\alpha) := 1 + \frac{\alpha + \alpha^2/2}{\frac{1}{2} + \frac{\alpha}{2} + \frac{\alpha^2}{4} + \frac{1}{2}(1 + \alpha)^2 \ln(1 + \alpha)}. \quad (196)$$

The function $R(\alpha)$ has a unique maximum at

$$\alpha_\star = \sqrt{u_5^{\det}} - 1 < e - 1, \quad R(\alpha_\star) = R_{\text{OT}}^{\det}. \quad (197)$$

Proof. The total span is

$$p_0 - p_{K-1} = \sum_{h=1}^{K-1} \frac{1}{\xi + h} < \ln\left(1 + \frac{K}{\xi}\right) = \ln(1 + \alpha) < 1,$$

which verifies Equation (186).

The processing contribution P_K has the exact form

$$P_K = \frac{K^2}{2} \left(1 + \frac{\gamma}{\xi}\right) + \frac{1}{2} \sum_{h=1}^{K-1} \frac{(K - h)^2}{\xi + h}. \quad (198)$$

Indeed, the common baseline $1 + \gamma/\xi$ contributes the first term, and swapping the order of summation shows that the increment $1/(\xi + h)$ receives total weight $(K - h)^2/2$. After division by ξ^2 , the sum becomes a Riemann integral:

$$\begin{aligned} \frac{P_K}{\xi^2} &\rightarrow I(\alpha) = \frac{\alpha^2}{2} + \frac{1}{2} \int_0^\alpha \frac{(\alpha - t)^2}{1 + t} dt \\ &= \int_0^\alpha (\alpha - t)(1 + \ln(1 + t)) dt \\ &= \frac{1}{2}(1 + \alpha)^2 \ln(1 + \alpha) - \frac{\alpha}{2} - \frac{\alpha^2}{4}. \end{aligned} \quad (199)$$

The second integral form follows either by integration by parts or by summing the limiting processing-time curve $1 + \ln(1 + t)$ against the remaining positive mass $\alpha - t$. The subtraction in Equation (193) is only $O(K)$, whereas $C_K, D_K = \Theta(K^2)$. Since $C_K - D_K = \xi K + K^2/2$, we obtain Equation (196).

Put $z = (1 + \alpha)^2$. In terms of z , the limiting ratio is

$$\widehat{R}(z) = 1 + \frac{2(z - 1)}{1 + z + z \ln z}. \quad (200)$$

The derivative of the fraction has the sign of

$$3 - z + \ln z.$$

For $z > 1$ this expression is strictly decreasing, is positive at $z = 1$, and tends to $-\infty$. Hence it has one zero $u_5^{\det}$, characterized by

$$u_5^{\det} - 3 = \ln u_5^{\det}.$$

The derivative is positive before $u_5^{\det}$ and negative afterward, so this is the global maximum. At the root,

$$1 + u_5^{\det} + u_5^{\det} \ln u_5^{\det} = (u_5^{\det} - 1)^2,$$

and therefore

$$\widehat{R}(u_5^{\det}) = 1 + \frac{2}{u_5^{\det} - 1} = R_{\text{OT}}^{\det}.$$

Moreover, $\sqrt{u_5^{\det}} - 1 = 1.12255\dots < e - 1$, so the condition $\alpha < e - 1$ holds at the maximizer. This proves Equation (197). $\square$

We can now prove the theorem.

Proof of Theorem 4.21. By claim 4.26, choose a rational α sufficiently close to $\alpha_\star = \sqrt{u_5^{\det}} - 1$, and choose any rational $\gamma \in (0, 1)$. For a sufficiently large K , set $\xi = K/\alpha$ and use Equation (191); then $Q_K/D_K > R_{\text{OT}}^{\det} - \varepsilon/2$. All parameters are rational. Let H tend to infinity through multiples of their common denominators. Claims 4.24 and 4.25 make the actual ratio exceed $R_{\text{OT}}^{\det} - \varepsilon$ for all sufficiently large H . Finally, freeze the adaptively revealed answers on their job labels as in claim 4.22. The resulting fixed input produces the same transcript for the given deterministic algorithm. $\square$

The lower bound for $u_4^{\det} \leq u \leq u_5^{\det}$ needs the same construction at an arbitrary total scale and at a prescribed point of its limiting curve. The next lemma packages this rescaled form so that no phase accounting has to be repeated later.

Lemma 4.27 (Scaled harmonic core). *Fix $s > 0$ and a parameter $t \in [1, e)$. Against every deterministic obligatory-testing algorithm, there are scales $M_\nu \rightarrow \infty$ and finite rational multilevel instances with $sM_\nu + o(M_\nu)$ jobs such that*

$$\frac{\text{OPT}}{M_\nu^2} \rightarrow O_0 = \frac{s^2}{4}(1 + t^{-2} + 2 \ln t), \quad \liminf_{\nu \rightarrow \infty} \frac{\text{ALG}}{M_\nu^2} \geq A_0^{\text{lb}} := O_0 + \frac{s^2}{2}(1 - t^{-2}). \quad (201)$$

Their total positive processing time, divided by M_ν , converges to

$$\frac{1}{M_\nu} \sum_{j: p_j > 0} p_j \rightarrow L = s \ln t, \quad (202)$$

and their largest processing time converges to $1 + \ln t < 2$. Freezing each revealed answer on its job label, as in claim 4.22, converts the adaptive description into a fixed instance.

Proof. The case $t = 1$ is obtained by continuity from instances with vanishing positive mass, so suppose $1 < t < e$. In the construction from the proof of Theorem 4.21, set $\alpha = t - 1 = K/\xi$. Before rescaling, the number of jobs divided by H is $K + \xi = \xi t$. Multiply every normalized group size by the common factor $s/(\xi t)$.

If every normalized group size is multiplied by a common factor, each term in the phase accounting, harmonic cancellation, and completion of squares in Equations (188), (193) and (195) is multiplied by the square of that factor. The Riemann-sum limit Equation (199) substituted into Equation (194) therefore gives

$$O_0 = \frac{s^2}{4}(1 + t^{-2} + 2 \ln t).$$

Likewise $C_K - D_K = \xi K + K^2/2$, while the $O(K)$ loss from completing the square in Equation (193) disappears after division by the quadratic scale. After the same rescaling this gives the guaranteed lower benchmark

$$A_0^{\text{lb}} - O_0 = \frac{s^2}{2}(1 - t^{-2}).$$

Finally, summing the limiting processing-time curve $1 + \ln(1 + x)$ over the positive mass gives $L = s \ln t$, and Equation (191) gives $p_0 \rightarrow 1 + \ln t$. Choose successively finer rational approximations, clear their denominators, and apply the recording argument from claim 4.22: assign each revealed answer to its job label and rerun the deterministic algorithm on the resulting fixed vector. This produces arbitrarily large fixed rational instances with the stated limit and liminf bounds. $\square$

4.2.3 Combining jobs with $p = u$ and a harmonic block

For $u_4^{\text{det}} < u < u_5^{\text{det}}$, neither the binary family nor the pure harmonic family is tight. The adversary first reveals a prescribed fraction of jobs with $p = u$. It then reveals the scaled harmonic instance from lemma 4.27, chosen so that every later job satisfies $1 + p \leq u$. Two lemmas show that neither raw execution nor processing one of the initial jobs early can avoid the resulting lower bound. The later part is exactly the construction from lemma 4.27, so its accounting need not be repeated here.

Lemma 4.28 (Deferring jobs with $p = u$). *Prepend a fraction m of jobs with $p = u$ to a later block occupying the remaining fraction $s = 1 - m$ and having total normalized processing time L . Suppose every job in the later block satisfies $1 + p \leq u$ and*

$$s(u - 1) - L - m \geq 0. \tag{203}$$

Then processing any initial job with $p = u$ before all later jobs are complete cannot decrease the leading online cost.

Proof. Consider an initial job with $p = u$ that is completed after all initial jobs have been tested but before a later job of value p . The pair contributes $1 + u$ to the online cost. If we postpone the initial job until after the later job, the same pair contributes $2 + p$. The postponement lowers the cost by $u - 1 - p \geq 0$, so the postponed schedule is a valid lower bound on the algorithm's actual cost.

It remains to consider jobs completed while the initial group of $p = u$ jobs is still being tested. Let q be the fraction of the full instance formed by these completed jobs. Their pairs with all later jobs add $q[s(u - 1) - L]$. Their largest possible saving against not-yet-tested jobs with $p = u$ is $qm - q^2/2$. Relative to leaving every such job pending, the change is therefore at least

$$q[s(u - 1) - L - m] + \frac{q^2}{2} \geq 0. \tag{204}$$

The omitted one-job terms are lower order. $\square$

Lemma 4.29 (Accounting for the initial $p = u$ jobs). *Let O_0 , A_0^{lb} , and L be the offline coefficient, guaranteed online coefficient, and total normalized positive processing time of the later block, where every job satisfies $1 + p \leq u$. After postponing all initial jobs with $p = u$, the corresponding coefficients O and A^{lb} of the combined instance satisfy*

$$A^{\text{lb}} = A_0^{\text{lb}} + m(1 - m) + m(1 + L) + \frac{u}{2}m^2, \quad (205)$$

$$O = O_0 + m(1 - m + L) + \frac{u}{2}m^2. \quad (206)$$

In particular, $A^{\text{lb}} - O = (A_0^{\text{lb}} - O_0) + m$.

Proof. A pair consisting of one initial job with $p = u$ and one later job of value p contributes $2 + p$ online and $1 + p$ offline. Summed over the initial fraction m and later fraction $1 - m$, these give the cross terms involving $m(1 - m)$ and mL . A pair of initial jobs with $p = u$ contributes $2 + u$ online and u offline, which gives the terms proportional to m^2 . Adding the coefficients of the later block yields the two displayed formulas. Subtracting the two identities gives the final assertion. $\square$

Lemma 4.30 (Quota adversary allowing raw execution). *Fix $m \in (0, 1)$ and a later block satisfying the hypotheses of lemma 4.28, in particular $1 + p \leq u$ for each of its jobs. Against every deterministic revealing algorithm there are fixed instances whose asymptotic excess is at least the combined coefficient $A^{\text{lb}} - R \cdot O$ obtained from the initial $p = u$ jobs and the later block in Equations (205) and (206) for every target $R \leq u$.*

Proof. In the initial phase, a tested fresh job receives $p = u$, while a job run raw is assigned hidden value zero. If v raw jobs and ℓ tested jobs with $p = u$ have been touched, stop at the first crossing

$$\frac{\ell}{n - v} \geq m. \quad (207)$$

With $S = n - v$, the overshoot is less than one, so $\ell = mS + O(1)$. If no crossing occurs, no test can have occurred: after all first touches, any positive ℓ would make the ratio in Equation (207) equal one. Thus every job was run raw and the all-zero hidden instance has ratio $u \geq R$.

At a crossing, scale the prescribed later instance to the remaining $S - \ell$ first touches. The crossing exceeds the target by fewer than one job; absorb this error and integer rounding into the zero group. Fix an order of the job types in this later block. A test reveals the next value in that order; a raw action assigns the next value to the touched job but does not reveal it to the algorithm.

Move every initial raw-zero completion to the start. It has the same length u as processing a known job with $p = u$, and crossing a unit test only lowers completion cost. With

$$Z_v = vn - \frac{v(v - 1)}{2}, \quad (208)$$

this prefix contributes uZ_v online and Z_v offline. Subtract these known contributions from the comparison. The remaining S labels consist, up to one-job rounding, of a fraction m of jobs with $p = u$ and the prescribed later block satisfying $1 + p \leq u$.

To compare with the harmonic lower-bound construction, run a virtual copy of the revealing algorithm. If this virtual algorithm later runs a fresh job of assigned value p raw, the obligatory-testing simulation instead tests and immediately processes that job in time $1 + p \leq u$, hides the answer from the virtual copy, and idles until the virtual raw block of length u ends. Inductively every simulated obligatory completion is no later than the

corresponding virtual revealing completion. Since $1 + p \leq u$ on the later block, its offline lengths are $1 + p$ in both the revealing and obligatory models. Write ALG_{hyb} and OPT_{hyb} for the online and offline costs of this physical obligatory-testing simulation. Then

$$\text{ALG}_{\text{RO}} - R \cdot \text{OPT}_{\text{RO}} \geq (u - R)Z_v + (\text{ALG}_{\text{hyb}} - R \cdot \text{OPT}_{\text{hyb}}) - O_u(n). \quad (209)$$

The first term is nonnegative. Finally, record every assigned value on its job label. Replaying this fixed vector produces the same transcript because the values learned only by the physical simulation were hidden from the virtual algorithm. This proves legality. $\square$

Proposition 4.31 (Lower bound for $u_4^{\text{det}} \leq u \leq u_5^{\text{det}}$). *For every $u_4^{\text{det}} \leq u \leq u_5^{\text{det}}$, every deterministic revealing algorithm has size-asymptotic competitive ratio at least $1 + c(u)$. Equivalently,*

$$R_{\text{RO}}^{\text{det}}(u) \geq 1 + c(u). \quad (210)$$

The construction has two continuous parameters: m is the fraction of initial jobs with $p = u$, and t selects the scaled harmonic block that follows. We first write its online/offline ratio, then solve the first-order conditions for a worst parameter pair. The remaining two claims verify that the pair is feasible and can be approximated by fixed finite instances.

Claim 4.32 (Ratio of the combined construction). *For $m \in [0, 1)$ and $t \in [1, e)$, apply lemma 4.27 with $s = 1 - m$ and prepend a fraction m of jobs with $p = u$. The resulting guaranteed lower ratio is*

$$\mathcal{R}_u(m, t) = 1 + \frac{m + \frac{(1-m)^2}{2}(1 - t^{-2})}{\frac{(1-m)^2}{4}(1 + t^{-2} + 2 \ln t) + m(1 - m)(1 + \ln t) + \frac{u}{2}m^2}. \quad (211)$$

Proof. Substitute $s = 1 - m$, $L = (1 - m) \ln t$, and the coefficients from Equation (201) into Equations (205) and (206), and divide the online coefficient by the offline coefficient. Collecting the excess over one gives exactly Equation (211). $\square$

Claim 4.33 ((c, m) -parametrization of a stationary point). *Suppose the ratio in Equation (211) has an interior stationary point with value $1 + c$. Combining this value equation with the two vanishing partial derivatives gives*

$$\ln t = u - 2 - \frac{1}{c}, \quad t^2 = \frac{1 - m}{1 + m} \left(1 + \frac{2}{c}\right). \quad (212)$$

Eliminating t gives precisely Equation (95). Conversely, those equations make $\mathcal{R}_u(m, t) = 1 + c$.

Proof. Write Equation (211) as $1 + N/D$. At a stationary point of value $1 + c$, the three equations are

$$N = cD, \quad \partial_m N = c \partial_m D, \quad \partial_t N = c \partial_t D.$$

The t -derivative equation is

$$(1 - m)^2 t^{-3} = c \left[\frac{(1 - m)^2}{2} (t^{-1} - t^{-3}) + \frac{m(1 - m)}{t} \right],$$

which rearranges to the second identity in Equation (212). Substituting it into the remaining two equations and simplifying gives the first identity. Eliminating t from these identities gives Equation (95). Reversing the substitutions verifies both the value and the two vanishing derivatives. $\square$

Claim 4.34 (Feasibility and uniqueness of the parameters). *For each $u \in [u_4^{\det}, u_5^{\det}]$, the equations in Equations (95) and (212) determine a unique triple (c, m, t) . As c decreases from $1/\varphi$ to r , this triple runs continuously from*

$$(u, m, t) = \left(u_4^{\det}, \frac{1}{\varphi}, 1\right) \quad \text{to} \quad (u, m, t) = (u_5^{\det}, 0, \sqrt{u_5^{\det}}). \quad (213)$$

It satisfies the deferral condition Equation (203); moreover, the finite harmonic block can be chosen with $1 + p < u$ for every one of its jobs.

Proof. The parameter relations give $u \geq u_4^{\det}$, $\ln t < 1$, and $m \leq 1/\varphi$. Therefore

$$(1 - m)(u - 1 - \ln t) > (1 - m)\varphi \geq \frac{1}{\varphi} \geq m, \quad (214)$$

so Equation (203) holds; also $1 + p < 3 < u$ for every later positive value in a sufficiently fine finite construction.

In the first equation of Equation (95), the left-hand side $\operatorname{artanh} m - m$ has derivative $m^2/(1 - m^2) > 0$, while the right-hand side has derivative $2/[c^2(c + 2)] > 0$. Hence it determines a unique $m = m(c)$, which is strictly increasing. Consequently

$$u(c) = 1 + \frac{2}{c} - m(c) \quad (215)$$

is strictly decreasing. At $c = r$ both sides of the first equation vanish, so $m = 0$; at $c = 1/\varphi$, direct substitution gives $m = 1/\varphi$. Together with t from Equation (212), the boundary value $c = r$ gives $(u, m, t) = (u_5^{\det}, 0, \sqrt{u_5^{\det}})$, while $c = 1/\varphi$ gives $(u, m, t) = (u_4^{\det}, 1/\varphi, 1)$, as recorded in Equation (213). Consequently every $u \in [u_4^{\det}, u_5^{\det}]$ has exactly one parameter pair. $\square$

Claim 4.35 (Approximation by fixed finite instances). *Fix $u_4^{\det} < u < u_5^{\det}$ and the associated parameters (c, m, t) from claim 4.34. Against every deterministic revealing algorithm there is a sequence of fixed finite instances I_H such that*

$$\liminf_{H \rightarrow \infty} \frac{\operatorname{ALG}(I_H) - (1 + c) \cdot \operatorname{OPT}(I_H)}{|I_H|^2} \geq 0. \quad (216)$$

Proof. To pass to fixed finite instances, first approximate (m, t) rationally while retaining the strict inequalities $1 + p < u$, then choose a sufficiently fine harmonic family, and finally multiply all rational block sizes by a common integer H . For a fixed number of positive levels, the error of fewer than one job at the quota crossing, the terms contributed by individual jobs, and rounding are $O_u(H)$, whereas the displayed pair costs are $\Theta(H^2)$. Let $H \rightarrow \infty$ first, then increase the harmonic depth, and finally let the rational approximation converge to (m, t) . Together with lemma 4.30, this proves Equation (216). $\square$

We can now prove the proposition.

Proof of Proposition 4.31. For $u_4^{\det} < u < u_5^{\det}$, claims 4.32 and 4.33 identify the target ratio as $1 + c(u)$, claim 4.34 verifies the hypotheses of the construction, and claim 4.35 supplies fixed instances approaching that ratio. The bounds at $u = u_4^{\det}$ and $u = u_5^{\det}$ follow by taking limits from the open interval. $\square$

4.2.4 Transferring the lower bound when raw execution is no shorter

The harmonic construction uses only values with $1 + p_j \leq u$, for which the offline cap u is inactive. The next lemma shows that any revealing algorithm with raw-execution cap u induces an obligatory-testing algorithm on these inputs without changing either the transcript comparison or the offline optimum.

Lemma 4.36 (Transfer from jobs with $1 + p \leq u$). *Let $u \geq 1$, and let $L_{\text{OT}}(P)$ denote the deterministic size-asymptotic value in obligatory testing, restricted to $0 \leq p_j \leq P$. Then*

$$R_{\text{RO}}^{\text{det}}(u) \geq L_{\text{OT}}(u - 1). \quad (217)$$

In particular,

$$R_{\text{RO}}^{\text{det}}(u) \geq R_{\text{OT}}^{\text{det}} \left(u \geq 1 + \frac{1}{r} = 2.752620747896442 \dots \right). \quad (218)$$

Proof. Given a revealing algorithm with raw-execution cap u , construct an obligatory-testing algorithm on instances with $p_j \leq u - 1$ by running a virtual copy of the revealing algorithm. Copy each test and processing action. When the virtual algorithm runs an untouched job raw for u time, physically test and immediately process it in $1 + p_j \leq u$ time, hide the answer from the virtual copy, and wait for the virtual clock. Every physical completion is no later than its virtual completion. Moreover, $\min\{u, 1 + p_j\} = 1 + p_j$, so the revealing and obligatory optima coincide. Thus we obtain the following direct comparison. For every revealing algorithm $\mathcal{A}$ with raw-execution cap u , the simulation produces an obligatory-testing algorithm $\mathcal{B}$ and, pointwise on every instance with $p_j \leq u - 1$,

$$\text{ALG}_{\mathcal{A}} \geq \text{ALG}_{\mathcal{B}}, \quad \text{OPT}_u = \text{OPT}_{\infty}.$$

Consequently the size-asymptotic worst-case ratio of $\mathcal{A}$ is at least the corresponding ratio of $\mathcal{B}$ restricted to $p_j \leq u - 1$, which is at least $L_{\text{OT}}(u - 1)$ by its definition. Taking the infimum over $\mathcal{A}$ proves Equation (217).

The harmonic family attaining $R_{\text{OT}}^{\text{det}}$ has limiting largest processing time

$$1 + \ln(1 + y_*) = 1 + \frac{1}{2} \ln u_5^{\text{det}} = \frac{u_5^{\text{det}} - 1}{2} = \frac{1}{r}, \quad y_* = \sqrt{u_5^{\text{det}}} - 1. \quad (219)$$

Choosing finite constructions whose largest processing time approaches $1/r$ from below includes the equality case $u = 1 + 1/r$ and proves Equation (218). $\square$

4.3 Assembly of the curve

It remains to verify the shape and joins of the six formulas in Equation (96) and to pair each upper bound with its matching lower bound. We do this explicitly before transferring the uniform plateau witnesses to obligatory testing.

4.3.1 Shape of the candidate curve

Before matching the bounds, we verify the shape of the six formulas in Equation (96). The only nontransparent monotonicity is that of $R_{\text{quad}}(u)$ on $[u_2^{\text{det}}, u_3^{\text{det}}]$; the formulas on $[u_3^{\text{det}}, u_4^{\text{det}}]$ and $[u_4^{\text{det}}, u_5^{\text{det}}]$ follow directly from the reciprocal formula and the (c, m) -parametrization in Equation (95).

Lemma 4.37 (Monotonicity of the deterministic curve). *The function $R_{\text{quad}}(u)$ is strictly decreasing on $[u_2^{\text{det}}, u_3^{\text{det}}]$. The function $1 + 1/\sqrt{u-1}$ is strictly decreasing on $[u_3^{\text{det}}, u_4^{\text{det}}]$, and $1 + c(u)$ is strictly decreasing on $[u_4^{\text{det}}, u_5^{\text{det}}]$. Consequently $R_{\text{RO}}^{\text{det}}(u)$ from Equation (96) attains its global maximum only at $u = u_2^{\text{det}}$.*

Proof. Implicit differentiation of $T_s(R_{\text{quad}}(u)) = 0$, with $s = u - 1$, gives

$$R'_{\text{quad}}(u) = -\frac{R^2 + (3s^2 + 2s)R - 3(s+1)^2}{2sR + s^3 + s^2 + 1}. \quad (220)$$

Using $T_s(R) = 0$, its numerator factors as

$$\frac{2s^3 + s^2 - 1}{s} \left(R - \frac{2s^3 + 3s^2 - 2}{2s^3 + s^2 - 1} \right).$$

On the relevant interval $s \geq u_2^{\text{det}} - 1 > 2/3$, the prefactor is positive, while the fraction in parentheses is at most $5/3$ because $4s^3 - 4s^2 + 1 > 0$. The explicit inequality (101) gives $R > 5/3$, and hence $R'_{\text{quad}}(u) < 0$. The function $1 + 1/\sqrt{u-1}$ is visibly decreasing. In the (c, m) -parametrization on $[u_4^{\text{det}}, u_5^{\text{det}}]$, $m(c)$ is increasing and $u(c) = 1 + 2/c - m(c)$ is decreasing, so $1 + c$ decreases with u . Thus the formula $R_{\text{RO}}^{\text{det}}(u) = u$ increases up to $u = u_2^{\text{det}}$; the formulas $R_{\text{quad}}(u)$, $1 + 1/\sqrt{u-1}$, and $1 + c(u)$ then decrease until $u = u_5^{\text{det}}$; and $R_{\text{RO}}^{\text{det}}(u) = R_{\text{OT}}^{\text{det}}$ is constant for $u \geq u_5^{\text{det}}$. This proves the global-maximum assertion. $\square$

4.3.2 Exact joins and completion of the proof

We next check that adjacent formulas agree at every transition point. Once this consistency is recorded, the theorem follows by pairing the upper and lower result already proved for each interval.

Lemma 4.38 (Exact joins). *The six formulas in Equation (96) agree at all five transition points. More explicitly, the formulas 1 and u agree at $u_1^{\text{det}} = 1$, and*

$$\begin{aligned} R_{\text{quad}}(u_2^{\text{det}}) &= u_2^{\text{det}}, \\ R_{\text{quad}}(u_3^{\text{det}}) &= 1 + \frac{1}{\sqrt{s_0}} = 1.682327803828019\dots, \\ 1 + \frac{1}{\sqrt{u_4^{\text{det}} - 1}} &= \varphi = 1 + c(u_4^{\text{det}}), \\ 1 + c(u_5^{\text{det}}) &= 1 + r = R_{\text{OT}}^{\text{det}}. \end{aligned} \quad (221)$$

Proof. The equality at $u_1^{\text{det}} = 1$ is immediate. Substitution of the defining equations for u_2^{det} and $s_0 = u_3^{\text{det}} - 1$ into T_s gives the first two displayed identities; uniqueness of the positive root identifies the values with R_{quad} . Since $u_4^{\text{det}} - 1 = \varphi + 1 = \varphi^2$, the formula $1 + 1/\sqrt{u-1}$ equals $1 + 1/\varphi = \varphi$ at $u = u_4^{\text{det}}$. The identities $1 + c(u_4^{\text{det}}) = \varphi$ and $1 + c(u_5^{\text{det}}) = 1 + r$ follow from the parameter triples at $u = u_4^{\text{det}}$ and $u = u_5^{\text{det}}$ in Equation (213). $\square$

Proof of Theorem 4.1. For the upper bounds, lemma 4.3 covers $0 < u \leq u_2^{\text{det}}$, lemma 4.4 covers $u_2^{\text{det}} \leq u \leq u_3^{\text{det}}$, and lemma 4.8 covers $u_3^{\text{det}} \leq u \leq u_4^{\text{det}}$. The guarantees on $u_4^{\text{det}} \leq u \leq u_5^{\text{det}}$ and $u \geq u_5^{\text{det}}$ follow from Theorem 4.9.

For the matching lower bounds, proposition 4.17 covers $0 < u \leq u_4^{\text{det}}$, proposition 4.31 covers $u_4^{\text{det}} \leq u \leq u_5^{\text{det}}$, and lemma 4.36 and Theorem 4.21 cover $u \geq u_5^{\text{det}}$ with instances satisfying $1 + p_j < u$, so the offline cap u is inactive. The exact joins are given by lemma 4.38,

and the claimed shape follows from lemma 4.37. The finite upper bounds have the stated $O_u(n)$ remainders, while Theorem 4.9 gives one absolute $O(n)$ remainder on the plateau. This proves every assertion of the theorem. $\square$

Remark 4.39 (Representative values for $u_4^{\text{det}} \leq u \leq u_5^{\text{det}}$). For orientation, some values for $u \in [u_4^{\text{det}}, u_5^{\text{det}}]$ are

u	$R_{\text{RO}}^{\text{det}}(u)$	m	t
3.800000	1.598851588	0.539725635	1.138984621
4.000000	1.583283633	0.428863572	1.330517832
4.071679486	1.579204806	0.381330539	1.412236296
4.300000	1.571692026	0.198387083	1.734651379

Remark 4.40 (Asymptotic scope). All statements above concern the fixed- u size-asymptotic ratio. For $0 < u < u_5^{\text{det}}$, the upper bounds have additive error $O_u(n)$. For $u \in [u_5^{\text{det}}, \infty)$ the same algorithm has one uniform $O(n)$ error. The unbounded obligatory endpoint is not included in the statement of the curve; it is derived in Section 4.4.

4.4 The obligatory endpoint

The obligatory model is the unbounded endpoint of revealing optimization, but the passage to that endpoint is not a formal substitution $u = \infty$: both the algorithm and the additive error must be uniform in u . The finite curve has just established exactly those two properties, so the endpoint now follows by a short transfer.

Proof of Theorem 4.2. For the upper bound, use ADAPTIVETHRESHOLD with $c = r$. This algorithm never runs a job raw and is independent of u . Given a finite obligatory instance, choose any finite

$$u > \max\{u_5^{\text{det}}, 1 + \max_j p_j\}.$$

Its execution is identical in the revealing model with raw-execution cap u , and every offline effective length is $\min\{u, 1 + p_j\} = 1 + p_j$. Thus the revealing and obligatory optima coincide. The constant B in Equation (170) is independent of u , so that inequality also bounds the obligatory instance by $\text{ALG} \leq R_{\text{OT}}^{\text{det}} \cdot \text{OPT} + Bn$.

For the lower bound, regard an arbitrary obligatory algorithm $\mathcal{B}$ as a revealing algorithm with the fixed raw-execution cap $u = u_5^{\text{det}}$ that simply never uses raw execution. The plateau lower bound in Theorem 4.1 may be witnessed by instances with $p_j < u_5^{\text{det}} - 1$, so the offline cap $u = u_5^{\text{det}}$ is inactive. On those instances the online transcript is unchanged and $\min\{u_5^{\text{det}}, 1 + p_j\} = 1 + p_j$, so again the two offline optima coincide. The revealing lower inequality at $u = u_5^{\text{det}}$ therefore gives the claimed obligatory lower bound for arbitrarily large instances. $\square$

5 Randomized revealing optimization: the exact four-piece curve

Theorem 3.14 identified the optimal leading online cost for every fixed input in the revealing-optimization model. We now compare that cost with the clairvoyant optimum and maximize over all inputs. The raw-execution cap u is fixed, but the algorithm is not told the entries

of the input vector. This section proves Theorem 5.1, the rigorous randomized half of the informal curve theorem Theorem 1.6. Its three transition points are

$$u_1^{\text{rand}} = 1, \quad u_2^{\text{rand}} = 5.048917339522306\dots, \quad u_3^{\text{rand}} = \frac{25}{4}. \quad (222)$$

Here $u_2^{\text{rand}} > 1$ is the unique root of $u^3 - 6u^2 + 5u - 1 = 0$ above one.

Theorem 5.1 (Randomized revealing-optimization curve). *For every fixed $u > 0$, the optimal randomized size-asymptotic competitive ratio against an oblivious adversary is*

$$R_{\text{RO}}^{\text{rand}}(u) = \begin{cases} 1, & 0 < u \leq u_1^{\text{rand}}, \\ \frac{u^3}{u^2 + (u-1)^3}, & u_1^{\text{rand}} \leq u \leq u_2^{\text{rand}}, \\ 1 + \frac{1}{2(\sqrt{u}-1)}, & u_2^{\text{rand}} \leq u \leq u_3^{\text{rand}}, \\ \frac{4}{3}, & u \geq u_3^{\text{rand}}. \end{cases} \quad (223)$$

More precisely, there are randomized nonanticipating algorithms $\mathcal{A}_{n,u}$ and numbers $\varepsilon_u(n) \rightarrow 0$ such that every vector $p \in [0, u]^n$ satisfies

$$\mathbb{E} \text{ALG}_{\mathcal{A}_{n,u}}(p) \leq R_{\text{RO}}^{\text{rand}}(u) \cdot \text{OPT}_u(p) + \varepsilon_u(n)n^2. \quad (224)$$

Conversely, for every randomized online algorithm $\mathcal{A}$ and every $\varepsilon > 0$, there are arbitrarily large n and a fixed vector $p \in [0, u]^n$ for which

$$\mathbb{E} \text{ALG}_{\mathcal{A}}(p) \geq (R_{\text{RO}}^{\text{rand}}(u) - \varepsilon) \cdot \text{OPT}_u(p). \quad (225)$$

The curve is continuous and has global maximum

$$\max_{u>0} R_{\text{RO}}^{\text{rand}}(u) = \frac{27 + 6\sqrt{3}}{23} = 1.625752384583185\dots, \quad (226)$$

attained at $u = (3 + \sqrt{3})/2$.

Theorem 3.14 identifies the best asymptotic online cost on each input. Here we compare that value with the clairvoyant SPT cost and maximize over all empirical distributions. Replacing an initial part of the survival function by its average reduces this optimization to distributions supported on two values. Two one-variable maximizations give the four formulas in Equation (223). A sublinear sample removes advance knowledge of the input, and shuffled binary vectors give matching lower bounds.

5.1 Offline and stationary fluid coefficients

Assume initially that $u > 1$ and write

$$s = u - 1. \quad (227)$$

Let D be a finite distribution on $[0, u]$ and use the survival function

$$S_D(t) = D((t, u]), \quad S_D(t) = 0 \quad (t > u). \quad (228)$$

The convention at point masses does not affect any integral below. Since a clairvoyant job of length p has effective length $1 + \min\{s, p\}$, write a nonnegative value x as $\int_0^\infty \mathbf{1}_{t < x} dt$ in the symmetric pair identity Equation (5). This gives

$$\begin{aligned}\Omega_{\text{RO},u}(D) &:= \frac{1}{2} \iint \min\{u, 1+p, 1+q\} dD(p)dD(q) \\ &= \frac{1}{2} \left(1 + \int_0^s S_D(t)^2 dt \right),\end{aligned}\tag{229}$$

$$\text{OPT}_u(p) = n^2 \Omega_{\text{RO},u}(D[p]) + \frac{1}{2} \sum_i \min\{u, 1+p_i\}\tag{230}$$

for every empirical distribution $D[p]$.

We next specialize the maximum-density module lemma 3.1 to a form suited to the offline cap u . If its threshold prefix has mass a and first moment ℓ , put

$$\tau_D = \frac{1 + \ell}{a}.\tag{231}$$

The common threshold equation and the layer-cake identity give

$$\int (\tau_D - p)^+ dD(p) = 1, \quad \int_0^{\tau_D} S_D(t) dt = \tau_D - 1.\tag{232}$$

The stationary algorithm tests all jobs in a uniform random order. It immediately processes a positive job if its revealed value lies in the selected threshold prefix, and after the last test it processes all remaining jobs in SPT order. The following formula is the stationary pair identity specialized to the finite offline cap u .

Lemma 5.2 (Stationary survival formula). *The doubled leading coefficient of the stationary algorithm is*

$$\mathcal{T}_u(D) = \begin{cases} \tau_D + \int_{\tau_D}^u S_D(t)^2 dt, & \tau_D \leq u, \\ \tau_D, & \tau_D > u. \end{cases}\tag{233}$$

Consequently the better of the raw and stationary algorithms has leading ratio

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s S_D(t)^2 dt}.\tag{234}$$

Proof. Let $h = 1 - a$ be the residual mass. Direct pair accounting gives the leading coefficient

$$(1 + \ell) \left(1 - \frac{a}{2} \right) + \text{SPT}(\nu).\tag{235}$$

If $\tau_D \leq u$, the threshold property gives

$$\text{SPT}(\nu) = \frac{\tau_D h^2}{2} + \frac{1}{2} \int_{\tau_D}^u S_D(t)^2 dt, \quad (1 + \ell) \left(1 - \frac{a}{2} \right) = \frac{\tau_D}{2} (1 - h^2).$$

Their sum is one half of the first line of (233). If $\tau_D > u$, the whole distribution is in the prefix and (235) equals $\tau_D/2$. Raw execution has leading coefficient $u/2$, and division by (229) proves (234). $\square$

This is where the instance-optimal theorem enters the curve calculation. In Equation (57), $q = 0$ is the raw algorithm and $q = 1$ is precisely the stationary algorithm using the selected threshold prefix. Hence

$$2\Phi_{\text{RO},u}(D) \leq \min\{u, \mathcal{T}_u(D)\}, \quad (236)$$

and Equation (234) is an upper bound on the instance-specific ratio $\Phi_{\text{RO},u}(D)/\Omega_{\text{RO},u}(D)$. Interior values of q can strictly improve individual inputs; the point of the next two subsections is that the two choices $q = 0$ and $q = 1$ already give the sharp value after maximizing over all D . The binary lower bounds at the end apply to every algorithm, not only to algorithms restricted to $q \in \{0, 1\}$.

5.2 Flattening the survival function

We now maximize (234) over arbitrary D . Suppose first that $1 \leq \tau_D \leq u$ and put

$$y = \frac{\tau_D - 1}{\tau_D}. \quad (237)$$

By (232), y is the average of S_D on $[0, \tau_D]$. Since S_D is nonincreasing, its average on every initial subinterval is at least y . Thus Cauchy–Schwarz gives

$$\int_0^a S_D(t)^2 dt \geq ay^2 \quad (0 \leq a \leq \tau_D), \quad (238)$$

and $S_D(t) \leq y$ almost everywhere for $t > \tau_D$.

Replace S_D on $(0, \tau_D)$ by the constant y and leave its tail unchanged. This remains a nonincreasing survival function, preserves (232) and $\mathcal{T}_u(D)$, and by (238) can only decrease the offline coefficient. Hence it can only increase the ratio. We may therefore assume

$$S_D(t) = y \quad (0 < t < \tau_D). \quad (239)$$

If $u - 1 \leq \tau_D \leq u$, the offline integral lies entirely in the flat part, and $S_D(t)^2 \leq y^2$ on the remaining tail. It follows that

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s S_D(t)^2 dt} \leq B_u(\tau_D) := \frac{\tau_D + (u - \tau_D)y^2}{1 + (u - 1)y^2}. \quad (240)$$

The numerator displayed on the right is at most u , so the raw minimum is inactive.

Now suppose $\tau_D < u - 1$. Put

$$L = u - 1 - \tau_D, \quad E = \int_{\tau_D}^{u-1} S_D(t)^2 dt, \quad G = \int_{u-1}^u S_D(t)^2 dt. \quad (241)$$

Monotonicity gives

$$0 \leq E \leq Ly^2, \quad G \leq E/L. \quad (242)$$

Consequently

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s S_D(t)^2 dt} \leq \frac{\tau_D + \frac{u - \tau_D}{u - 1 - \tau_D} E}{1 + \tau_D y^2 + E}. \quad (243)$$

For fixed τ_D this is linear-fractional in E , so its derivative has a constant sign. Its maximum on $[0, Ly^2]$ therefore occurs at $E = 0$ or $E = Ly^2$. Substitution of these two values gives, respectively,

$$A(\tau_D) := \frac{\tau_D}{1 + \tau_D y^2} = \frac{\tau_D^2}{\tau_D^2 - \tau_D + 1}, \quad (244)$$

$$B_u(\tau_D) = \frac{\tau_D + (u - \tau_D)y^2}{1 + (u - 1)y^2}. \quad (245)$$

They are attained by rectangular survival functions: respectively a binary distribution on $\{0, \tau_D\}$ and one on $\{0, u\}$.

It remains to cover $\tau_D > u$. Equation (232) and the zero extension of S_D give $\int_0^u S_D(t) dt = \tau_D - 1 \geq u - 1$. The average of a nonincreasing function on $[0, u - 1]$ is no smaller than its average on $[0, u]$, and hence

$$\int_0^{u-1} S_D(t)^2 dt \geq (u - 1) \left(\frac{\tau_D - 1}{u} \right)^2 \geq \frac{(u - 1)^3}{u^2}. \quad (246)$$

Here the raw algorithm is chosen, and its ratio is at most $B_u(u)$. We have proved the reduction

$$\sup_D \frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s S_D(t)^2 dt} \leq \max \left\{ \sup_{1 \leq \tau \leq u-1} A(\tau), \sup_{1 \leq \tau \leq u} B_u(\tau) \right\}, \quad (247)$$

where the first supremum is omitted for $u < 2$. In fact equality holds in (247), because the binary distributions on $\{0, \tau_D\}$ and $\{0, u\}$ identified before the reduction attain the A and B_u families, respectively.

5.3 The two scalar maximizations

The first family satisfies

$$A'(\tau) = \frac{\tau(2 - \tau)}{(\tau^2 - \tau + 1)^2}. \quad (248)$$

Thus its maximum is attained at $\tau = \min\{2, u - 1\}$ and equals $4/3$ as soon as $u \geq 3$.

For the second family, substitution of (237) gives

$$B_u(\tau) = \frac{\tau^2 u + 2\tau^2 - 2\tau u - \tau + u}{\tau^2 u - 2\tau u + 2\tau + u - 1}. \quad (249)$$

The sign of its derivative is the sign of

$$g_u(\tau) = (2\tau - 1)^2 - u(\tau - 1)^2. \quad (250)$$

For $u \leq 4$ this is positive on $[1, \infty)$. For $u > 4$, its unique root above one is

$$\tau_*(u) = \frac{u - 2 + \sqrt{u}}{u - 4} = \frac{\sqrt{u} - 1}{\sqrt{u} - 2}, \quad (251)$$

and B_u increases before this point and decreases afterwards. The identity $\tau_*(u) = u$ is equivalent to

$$u^3 - 6u^2 + 5u - 1 = 0. \quad (252)$$

The cubic is strictly increasing beyond its last critical point and changes sign there exactly once; its unique root above one is u_2^{rand} from (222). Hence

$$\max_{1 \leq \tau \leq u} B_u(\tau) = B_u(u) = \frac{u^3}{u^3 - 2u^2 + 3u - 1} = \frac{u^3}{u^2 + (u - 1)^3}, \quad u \leq u_2^{\text{rand}}, \quad (253)$$

$$= B_u(\tau_*(u)) = 1 + \frac{1}{2(\sqrt{u} - 1)}, \quad u \geq u_2^{\text{rand}}. \quad (254)$$

It remains to compare the candidate A with the values obtained from B_u . On $2 \leq u \leq 3$, clearing the positive denominators in $B_u(u) - A(u - 1)$ and putting $v = u - 2$ leaves

$$v^4 + 3v^3 + 3v^2 + 3v + 3 > 0.$$

On $3 \leq u \leq u_2^{\text{rand}}$, one has $B_u(u) > 4/3$. On $u_2^{\text{rand}} \leq u \leq u_3^{\text{rand}}$, (254) is at least $4/3$, with equality exactly at $u = u_3^{\text{rand}}$; afterwards $A(2) = 4/3$ is larger. Combining these observations with (247) proves the announced upper bound

$$\frac{\min\{u, \mathcal{T}_u(D)\}}{1 + \int_0^s S_D(t)^2 dt} \leq R_{\text{RO}}^{\text{rand}}(u) \quad (255)$$

for every finite distribution D .

The junction statements now follow directly. Equations (252) to (254) give continuity at u_2^{rand} , and the formula $1 + 1/[2(\sqrt{u} - 1)]$ equals $4/3$ at $u = u_3^{\text{rand}}$. Differentiating (253) shows that its only interior maximum above one occurs at $u = (3 + \sqrt{3})/2$; substitution gives (226).

5.4 Worst-case upper bound from instance optimality

No second learning argument is needed. Apply the universal unannounced algorithm of Theorem 3.14 to $D = D[p]$. The endpoint comparison Equation (236) and the scalar bound Equation (255) give

$$\Phi_{\text{RO},u}(D[p]) \leq R_{\text{RO}}^{\text{rand}}(u) \cdot \Omega_{\text{RO},u}(D[p]).$$

The exact empirical identity Equation (230) has a nonnegative linear correction. Therefore the instance-optimal upper bound immediately yields

$$\mathbb{E} \text{ALG}_{\mathcal{A}_{n,u}} \leq R_{\text{RO}}^{\text{rand}}(u) \cdot \text{OPT}_u + o_u(n^2), \quad (256)$$

which is (224). For $0 < u \leq 1$, raw execution is exactly clairvoyantly optimal, so no learning is required.

5.5 Binary oblivious lower bounds

It remains to show that no adaptive algorithm improves the resulting worst-case upper bound. We first record an envelope for shuffled binary vectors.

Lemma 5.3 (Binary-input envelope at the offline cap u). *Fix $u > 1$, $x \in [0, 1]$, and put $z = 1 - x$. Let $p^{(n)}$ have $xn + O(1)$ coordinates of value u and $zn + O(1)$ zero coordinates. For every randomized nonanticipating algorithm, even an announced one,*

$$\frac{\mathbb{E} \text{ALG}}{n^2} \geq \begin{cases} \frac{u}{2} - o_u(1), & x \geq (u - 1)/u, \\ \frac{1 + x + ux^2}{2} - o_u(1), & x \leq (u - 1)/u. \end{cases} \quad (257)$$

Every labeling has

$$\frac{\text{OPT}_u}{n^2} = \frac{1 + (u - 1)x^2}{2} + O_u(1/n). \quad (258)$$

Proof. First fix a deterministic algorithm. If it does not complete all jobs, the claim is immediate. Otherwise place the individual jobs uniformly on the public labels, treating jobs of the same value as distinct. Reveal a job's value in the analysis only when the algorithm first touches its label. Conditional on all earlier reveals, the unrevealed jobs

remain uniformly permuted, and the algorithm has already made its test/raw choice before the next value is revealed.

Take $\delta_n = n^{-1/4}$. The predictable-permutation bound lemma 2.7, applied to the zero indicator and the test selector, gives one event of probability $1 - o(1)$ on which every prefix of the order in which labels are first touched that ends with at least $\delta_n n$ untouched labels and contains t tests satisfies

$$|N_0(t) - zt| \leq \Delta_n, \quad \Delta_n = O(\delta_n^{-1} \sqrt{n \ln n}) = o(n). \quad (259)$$

Here $N_0(t)$ is the number of tested zeros. The event is simultaneous, so we do not condition on the algorithm's adaptive stopping time.

Edit the suffix by running raw every still-untouched job that the original schedule would later test, deleting its later known-processing operation. Only $\delta_n n + O(1)$ jobs change, so the edit changes total completion cost by $o_u(n^2)$. Let q be the final tested fraction of the edited schedule. On (259), every prefix of its completion curve is dominated, up to $o(1)$ vertical error, by the divisible items

item	completion capacity	work per completion
tests that reveal zero jobs	zq	$1/z$
processing of tested $p = u$ jobs	xq	u
raw jobs	$1 - q$	u

When $z = 0$, the first row is simply omitted. Indeed, t tests complete at most $zt + \Delta_n$ zeros and expose at most $xt + \Delta_n$ jobs with $p = u$; raw execution and processing of a known $p = u$ job both use u work per completion. Ordering these three divisible blocks by work per completion, as in lemma 3.2, therefore upper-bounds completed mass at every work value. Integrating the remaining mass gives a lower bound on total completion time.

If $z \leq 1/u$, every displayed item costs at least u per completion, so the normalized area is at least $u/2 - o_u(1)$. If $z \geq 1/u$, the tests that reveal zero jobs come first. For fixed q the resulting remaining-mass area is

$$F_z(q) = q - \frac{zq^2}{2} + \frac{u}{2}(1 - zq)^2, \quad F'_z(q) = (1 - zq)(1 - uz). \quad (260)$$

It is minimized at $q = 1$, giving $F_z(1) = (1 + x + ux^2)/2$. Restoring the edited final $\delta_n n + O(1)$ jobs, the $o(1)$ error in completed mass, and the sampling event's failure probability together contribute only $o_u(n^2)$.

The argument holds for each deterministic algorithm. For a randomized algorithm, fix its seed and then average over the seed. The private shuffle is already part of the input convention. Finally, the effective lengths of the two job types are 1 and u , so (258) follows from lemma 2.1. $\square$

We now select the binary masses. If $u_1^{\text{rand}} < u \leq u_2^{\text{rand}}$, take

$$x = \frac{u - 1}{u}. \quad (261)$$

This is the boundary $z = 1/u$ in lemma 5.3, and its online and offline coefficients have ratio

$$\frac{u}{1 + (u - 1)((u - 1)/u)^2} = \frac{u^3}{u^2 + (u - 1)^3}. \quad (262)$$

If $u_2^{\text{rand}} \leq u \leq u_3^{\text{rand}}$, take

$$x = \frac{1}{\sqrt{u} - 1}. \quad (263)$$

The definition of u_2^{rand} is precisely the point from which $x \leq (u - 1)/u$. Since $(u - 1)x^2 = 1 + 2x$, the second line of (257) divided by (258) becomes

$$1 + \frac{x}{2} = 1 + \frac{1}{2(\sqrt{u} - 1)}. \quad (264)$$

Irrational masses are approximated by integer job counts divided by n ; the rounding error and the one-job terms omitted from the leading pair coefficient are $O_u(n)$.

For $u \geq u_3^{\text{rand}}$, the lower bound $4/3$ uses a fixed bounded distribution supported on $\{0, 2\}$ instead of the binary distribution on $\{0, u\}$ used for smaller u .

Lemma 5.4 (Balanced zero-two lower bound). *For every $u \geq u_3^{\text{rand}}$ and every randomized algorithm, arbitrarily large even n admit a vector with $n/2$ zero coordinates and $n/2$ coordinates of value two such that*

$$\mathbb{E} \text{ALG} \geq n^2 - o_u(n^2), \quad \text{OPT}_u = \frac{3}{4}n^2 + O(n). \quad (265)$$

Proof. Expose values when their labels are first touched, apply the predictable-permutation event Equation (259) simultaneously to every prefix of that order, and replace tests among the final $o(n)$ untouched jobs by raw execution as in the proof of lemma 5.3. For a realized tested fraction q , the three divisible blocks now have capacities and costs

item	completion capacity	work per completion
tests that reveal zero jobs	$q/2$	2
processing of tested jobs with $p = 2$	$q/2$	2
raw jobs	$1 - q$	u

Every item costs at least two per completion. Thus the completion envelope lies below $\min\{1, t/2\}$ independently of q , whose remaining-mass area is one. This proves the online estimate after the $o_u(n^2)$ repairs and averaging over the algorithm seed. Since $u > 3$, the two effective lengths are one and three; their equally weighted divisible SPT cost gives offline coefficient $3/4$. $\square$

Equations (262) and (264), together with lemma 5.4, match (256) on $1 < u \leq u_2^{\text{rand}}$, $u_2^{\text{rand}} \leq u \leq u_3^{\text{rand}}$, and $u \geq u_3^{\text{rand}}$, respectively. For $0 < u \leq 1$, raw execution is both an online and a clairvoyant optimum. This completes the proof of the exact curve in Theorem 5.1.

5.6 The randomized obligatory endpoint

We now pass from the randomized curve with a finite raw-execution cap u to the unbounded obligatory endpoint. The adversary fixes a vector $p \in [0, \infty)^n$ before the algorithm draws its random bits; the input convention then privately shuffles its coordinates. The entries are not announced, and no upper bound on the processing times is assumed. The zero-two lower bound at the finite raw-execution cap $u = u_3^{\text{rand}}$ transfers directly. The upper bound needs one additional safeguard: the algorithm groups every sampled value above a threshold into one category and switches to a safe test-all schedule when the sample does not identify a dense enough early group. Letting this threshold grow slowly also makes the same

algorithm instance-optimal on every bounded class. This subsection proves Theorem 5.5, the rigorous randomized part of the informal obligatory-testing result Theorem 1.1 from the introduction.

Theorem 5.5 (Randomized obligatory testing). *There are a universal constant C and a randomized nonanticipating algorithm $\mathcal{A}_n^*$, depending only on n , such that every fixed vector $p \in [0, \infty)^n$ satisfies*

$$\mathbb{E} \text{ALG}_{\mathcal{A}_n^*}(p) \leq \frac{4}{3} \cdot \text{OPT}(p) + Cn^{19/10} \leq \left(\frac{4}{3} + 2Cn^{-1/10} \right) \cdot \text{OPT}(p). \quad (266)$$

A fixed-cutoff companion $\mathcal{A}_n^{(32)}$, using $B = 32$ and the more conservative fallback criterion $\hat{\kappa} < 2/B$, satisfies the sharper explicit estimate

$$\mathbb{E} \text{ALG}_{\mathcal{A}_n^{(32)}}(p) \leq \frac{4}{3} \cdot \text{OPT}(p) + 20378n^{7/4}. \quad (267)$$

Conversely, for every randomized algorithm $\mathcal{A}$ and every $\varepsilon > 0$, arbitrarily large even n admit a vector with $n/2$ zero coordinates and $n/2$ coordinates of value two such that

$$\mathbb{E} \text{ALG}_{\mathcal{A}}(p) \geq \left(\frac{4}{3} - \varepsilon \right) \cdot \text{OPT}(p), \quad \text{OPT}(p) = \frac{3}{4}n^2 + n. \quad (268)$$

Hence the exact randomized size-asymptotic competitive ratio against an oblivious adversary is $R_{\text{OT}}^{\text{rand}} = 4/3$.

When the multiset of values is revealed in advance, the algorithmic core is the prefix of the shortest processing times that maximizes completions per unit of testing and processing work. Testing jobs in a uniformly random order gives an exact pair identity whose nonnegative terms prove the factor $4/3$. The universal algorithm learns an approximately separated prefix from a sublinear sample; the threshold and the safe test-all schedule described below protect it from unbounded values missed by the sample. The matching lower bound is inherited from the zero–two lower bound at the finite raw-execution cap $u = u_3^{\text{rand}}$. We first prove the uniform upper bound and then give the short transfer.

5.6.1 Normalization of the offline pair identity

For a distribution D on $[0, \infty)$, define the leading offline coefficient

$$\Omega_{\text{OT}}(D) := \frac{1}{2} + \text{SPT}(D). \quad (269)$$

For a fixed vector p , the finite SPT identity gives

$$\Omega_{\text{OT}}(D[p]) = \frac{1}{2} + \frac{\text{SPT}([n])}{n^2} - \frac{1}{2n^2} \sum_i p_i, \quad \Delta_{\text{OT}}(p) = \frac{n + \sum_i p_i}{2}. \quad (270)$$

Combining Equation (5) with the finite SPT identity, with $\lambda_i = 1 + p_i$, gives

$$\text{OPT}(p) = n^2 \Omega_{\text{OT}}(D[p]) + \Delta_{\text{OT}}(p). \quad (271)$$

The linear correction $\Delta_{\text{OT}}(p)$ will be kept rather than hidden: this makes the analysis valid for unbounded processing times.

5.6.2 Stationary prefixes

For a fixed input p , let $E \subseteq [n]$ contain every zero job and suppose that every processing time in E is at most every processing time outside E . Put $T = [n] \setminus E$ and

$$e = |E|, \quad a = \frac{e}{n}, \quad h = 1 - a, \quad \ell_X = \frac{1}{n} \sum_{i \in X} p_i \quad (X \subseteq [n]). \quad (272)$$

We call E a *processing-time prefix*. Since every job in E is no longer than every job in T , the SPT pair formula gives

$$\text{SPT}([n]) = \text{SPT}(E) + n^2 h \ell_E + \text{SPT}(T). \quad (273)$$

Consider the stationary algorithm that tests the jobs in a uniformly random order, processes each positive E -job immediately, and after the last test processes T in SPT order. Here *stationary* means that E and this immediate-or-deferred rule are fixed before the random order is exposed. We write its leading coefficient as

$$\text{STAT}(E) := (1 + \ell_E) \left(1 - \frac{a}{2}\right) + \frac{\text{SPT}(T)}{n^2} - \frac{\ell_T}{2n}. \quad (274)$$

This notation has a direct exact interpretation. Let $W = n + \sum_{i \in E} p_i$ and $W_E = e + \sum_{i \in E} p_i$. Every test or immediate processing block precedes a fixed E -job with probability $1/2$, so the E -jobs contribute $(eW + W_E)/2$ in expectation. Every T -job then sees the common offset W , and the final phase contributes $(n - e)W + \text{SPT}(T)$. Adding these quantities gives

$$\mathbb{E} \text{ALG}_{\text{stat}}(p, E) = n^2 \text{STAT}(E) + \frac{e + \sum_i p_i}{2}. \quad (275)$$

For a nonempty prefix define its completion density by

$$\kappa(E) = \frac{a}{1 + \ell_E}. \quad (276)$$

Choose a maximum-density prefix and put

$$\tau = \frac{1 + \ell_E}{a} = \frac{1}{\kappa(E)}. \quad (277)$$

All zeros belong to every maximizer: adding a zero increases the numerator without increasing the denominator. For an occurrence of length p outside E , adding it does not decrease $a/(1 + \ell_E)$ exactly when $p \leq (1 + \ell_E)/a = \tau$; the analogous removal calculation applies to an occurrence inside E . We may therefore choose a maximizer with

$$\tau = \tau_{D[p]}, \quad p_i \leq \tau \ (i \in E), \quad p_i \geq \tau \ (i \notin E), \quad (278)$$

where tied occurrences may be assigned to either side.

The next identity is the algebraic source of the constant $4/3$.

Lemma 5.6 (Exact identity and robust bound). *For every ordered prefix split with $a > 0$ and $\tau = (1 + \ell_E)/a$,*

$$\begin{aligned} & 2\tau(4\Omega_{\text{OT}}(D[p]) - 3\text{STAT}(E)) \\ &= (\tau - 2)^2 + 4 \left[\tau \left(\frac{2\text{SPT}(E)}{n^2} - \frac{\ell_E}{n} \right) - \ell_E^2 \right] \\ &+ \tau \left(\frac{2\text{SPT}(T)}{n^2} - \frac{\ell_T}{n} - \tau h^2 \right). \end{aligned} \quad (279)$$

Consequently:

- (i) if values in E are at most τ and values in T are at least τ , then $\text{STAT}(E) \leq 4\Omega_{\text{OT}}(D[p])/3$;
- (ii) if values in E are at most $\tau + s$ and values in T are at least $\tau - s$, where $s \geq 0$, then

$$\text{STAT}(E) \leq \frac{4}{3}\Omega_{\text{OT}}(D[p]) + \frac{2}{3}s. \quad (280)$$

Proof. Expansion using $\ell_E = a\tau - 1$, $h = 1 - a$, and Equation (273) gives Equation (279). Under exact separation, $xy \leq \tau \min\{x, y\}$ for $x, y \in E$, hence

$$\ell_E^2 \leq \tau \left(\frac{2 \text{SPT}(E)}{n^2} - \frac{\ell_E}{n} \right).$$

For $x, y \in T$, one has $\min\{x, y\} \geq \tau$, hence

$$\frac{2 \text{SPT}(T)}{n^2} - \frac{\ell_T}{n} \geq \tau h^2.$$

This proves (i).

Under the relaxed inequalities,

$$\tau \left(\frac{2 \text{SPT}(E)}{n^2} - \frac{\ell_E}{n} \right) - \ell_E^2 \geq -s \left(\frac{2 \text{SPT}(E)}{n^2} - \frac{\ell_E}{n} \right),$$

and

$$\frac{2 \text{SPT}(T)}{n^2} - \frac{\ell_T}{n} - \tau h^2 \geq -sh^2.$$

Moreover $2 \text{SPT}(E)/n^2 - \ell_E/n \leq a\ell_E \leq a^2\tau$. Substitution into Equation (279) yields

$$\text{STAT}(E) \leq \frac{4}{3}\Omega_{\text{OT}}(D[p]) + \frac{2}{3}sa^2 + \frac{1}{6}sh^2 \leq \frac{4}{3}\Omega_{\text{OT}}(D[p]) + \frac{2}{3}s,$$

because $(2/3)a^2 + (1/6)(1 - a)^2 \leq 2/3$ on $[0, 1]$. $\square$

Combining the exact identity with Equations (271) and (275) already gives a finite statement useful below.

Corollary 5.7 (Announced multiset). *If the multiset is announced, the stationary algorithm for a maximum-density prefix satisfies*

$$\mathbb{E} \text{ALG}_{\text{stat}}(p, E) \leq \frac{4}{3} \cdot \text{OPT}(p).$$

Indeed, the correction in Equation (275) is at most $\Delta_{\text{OT}}(p)$. Equality in the quadratic coefficient of Equation (279) forces $\tau = 2$, zero mass $1/2$, and all positive mass at two. This predicts the lower-bound instance before any probabilistic argument is used.

We also need an estimate whose error is independent of the values outside the early set. Direct expansion gives

$$\text{STAT}(E) - \Omega_{\text{OT}}(D[p]) = \frac{1}{2} \left(h + a\ell_E - \frac{2 \text{SPT}(E)}{n^2} + \frac{\ell_E}{n} \right). \quad (281)$$

If all values in E are at most $L \geq 1$, then $\ell_E \leq aL$ and therefore

$$\text{STAT}(E) \leq \Omega_{\text{OT}}(D[p]) + \frac{L}{2}. \quad (282)$$

5.6.3 Learning the prefix from a sublinear sample

For the main algorithm, set the slowly growing cutoff

$$B = B_n = 32 + n^{1/20}. \quad (283)$$

For an integer d , let $\eta = B/d$. Keep zero as a separate category, partition $(0, B]$ into the consecutive bins $((b-1)\eta, b\eta]$, and add one overflow category for values above B . A finite positive value is rounded upward to the endpoint $q_b = b\eta$.

Let S be a uniformly random subset of k labels. If μ_b and $\hat{\mu}_b$ are the population and sample fractions of category b , put

$$\Delta = \sum_b |\hat{\mu}_b - \mu_b|. \quad (284)$$

Lemma 5.8 (Histogram error). *For $1 \leq k < n$,*

$$\mathbb{E}\Delta \leq \sqrt{\frac{d+2}{k}}.$$

Proof. This is Equation (12), with the zero bin and the overflow bin included among the $d+2$ categories. $\square$

For a finite category prefix J , define

$$\hat{a}(J) = \sum_{b \in J} \hat{\mu}_b, \quad \hat{\ell}(J) = \sum_{b \in J} q_b \hat{\mu}_b, \quad \hat{\kappa}(J) = \frac{\hat{a}(J)}{1 + \hat{\ell}(J)}.$$

Let $\hat{\kappa}$ be the maximum and $\hat{\tau} = 1/\hat{\kappa}$. Starting with a maximizing category prefix, also include every finite category whose endpoint is at most $\hat{\tau}$. Thus the algorithm uses

$$\hat{J} = \{\text{finite categories } b : q_b \leq \hat{\tau}\}. \quad (285)$$

Lemma 5.9 (Closing a maximum-density prefix at its threshold). *The prefix $\hat{J}$ from Equation (285) has the same maximum sample density. Every category in it has endpoint at most $\hat{\tau}$, and every finite category outside it has endpoint greater than $\hat{\tau}$, including categories with zero sample count.*

Proof. For every positive-mass sample category, adding it preserves or improves the density exactly when its endpoint is at most $\hat{\tau}$; removing it preserves or improves the density under the reverse inequality. Empty categories do not change the sample statistics, and adding mass at equality preserves the identity $1 + \hat{\ell} = \hat{a}\hat{\tau}$. Adding all categories up to the threshold therefore preserves optimality and makes the two pointwise assertions tautological. $\square$

For large n , set

$$k = \lfloor n^{3/4} \rfloor, \quad d = \lfloor n^{1/4} \rfloor, \quad \eta = B/d. \quad (286)$$

The algorithm draws a uniform permutation and first tests its first k jobs, leaving positive sample jobs pending. It computes $\hat{J}$ according to Equation (285). If

$$\hat{\kappa} < \frac{1}{B}, \quad (287)$$

it enters *fallback mode*: test every remaining job and then process all positive jobs in SPT order. Otherwise it enters *learned mode*: first process the positive sampled jobs whose categories lie in $\hat{J}$; then test the remaining jobs in random order and immediately process each positive job whose category lies in $\hat{J}$; finally process all other positive tested jobs in SPT order. The overflow category never belongs to $\hat{J}$.

5.6.4 Analysis on good and bad samples

Put

$$\gamma = \frac{1}{B(B+1)}, \quad G = \{\Delta \leq \gamma\}. \quad (288)$$

Markov's inequality, recalled in Section 2.5, and lemma 5.8 give

$$\mathbb{P}(G^c) \leq B(B+1) \sqrt{\frac{d+2}{k}}. \quad (289)$$

Assume that G holds and the algorithm uses the learned schedule. Let E contain all jobs whose categories belong to $\hat{J}$, and let $a, \ell_E, \tau = (1 + \ell_E)/a$ be its true statistics. Since $\hat{\tau} \leq B$ and $\hat{a} = (1 + \hat{\ell})/\hat{\tau} \geq 1/B$, we have $a \geq 1/(B+1)$. For this fixed category set,

$$|a - \hat{a}| \leq \Delta, \quad |\ell_E - \hat{\ell}| \leq B\Delta + \eta.$$

Consequently

$$|\tau - \hat{\tau}| \leq 2B(B+1)\Delta + (B+1)\eta. \quad (290)$$

Every value in E is at most $\hat{\tau}$, while every value outside E is greater than $\hat{\tau} - \eta$. Thus the robust bound applies with

$$s = 2B(B+1)\Delta + (B+2)\eta. \quad (291)$$

The algorithm tests the sample before all other jobs, so its order is not exactly the stationary schedule in which all jobs share one uniform order. The difference is only lower order. Conditional on the unordered sample, put $e_S = |E \cap S|$, $e_T = |E \setminus S|$, and

$$W_S = k + \sum_{i \in E \cap S} p_i, \quad W_T = n - k + \sum_{i \in E \setminus S} p_i.$$

Comparing the two independent random internal orders with one fully random order, the difference in the cost contributed by jobs in E is

$$\frac{e_T W_S - e_S W_T}{2} \leq \frac{1}{2} n k (1 + B).$$

Postponing the positive jobs in $E \cap S$ until all sample tests finish adds at most $k^2(1+B)$. Jobs outside E are processed identically at the end because every zero belongs to E . Therefore, on event G when the learned schedule is used,

$$\mathbb{E}[\text{ALG} \mid S] \leq \frac{4}{3} \cdot \text{OPT} + n^2 \left[\frac{4}{3} B(B+1)\Delta + \frac{2}{3} (B+2)\eta \right] + (1+B)(nk/2 + k^2). \quad (292)$$

On event G when the fallback schedule is used, comparison of sample and population densities shows that the reciprocal of the true maximum density satisfies $\tau_{D[p]} \geq B/4 \geq 8$. The exact threshold inequalities imply

$$\frac{2 \text{SPT}([n])}{n^2} - \frac{1}{n^2} \sum_i p_i \geq \frac{(\tau_{D[p]} - 1)^2}{\tau_{D[p]}}, \quad \Omega_{\text{OT}}(D[p]) \geq \frac{\tau_{D[p]} - 1 + 1/\tau_{D[p]}}{2} \geq \frac{57}{16}. \quad (293)$$

For every test order, test-all-then-SPT satisfies

$$\text{ALG}_{\text{fb}} \leq \text{OPT} + \frac{n(n-1)}{2}. \quad (294)$$

Indeed, if there are z zeros, moving their completions from the first z SPT positions to their test positions increases their sum of completion times by at most $zn - z(z+1)/2$. Delaying the remaining $n - z$ positive jobs until all tests are finished adds at most $(n - z)(n - z - 1)/2$; the two bounds sum to $n(n - 1)/2$. Combining Equations (293) and (294) gives $\text{ALG}_{\text{fb}} < \frac{4}{3} \cdot \text{OPT}$.

Because processing times are unbounded, the estimate on G^c must control excess over $\frac{4}{3} \cdot \text{OPT}$, not raw cost. Under the learned schedule every value in E is at most B , so Equation (282) bounds the stationary excess by $(B/2)n^2$; under the fallback schedule Equation (294) bounds it by $n^2/2$. Averaging Equation (292) and these bounds on G^c , and using Equation (289), yields the uniform estimate

$$\mathbb{E} \text{ALG} \leq \frac{4}{3} \cdot \text{OPT} + n^2 \left[\frac{B(B+1)(3B+8)}{6} \sqrt{\frac{d+2}{k}} + \frac{2B(B+2)}{3d} \right] + (1+B)(nk/2 + k^2). \quad (295)$$

For $n \geq 12^4$, the floors in Equation (286) give $\sqrt{(d+2)/k} \leq 2n^{-1/4}$, $1/d \leq 2n^{-1/4}$, and $k \leq n^{3/4}$. Since $B_n = O(n^{1/20})$, the three error terms in Equation (295) are respectively $O(n^{19/10})$, $O(n^{37/20})$, and $O(n^{9/5})$. Enlarging one universal constant covers smaller n and proves the first inequality in Equation (266). The bound $\text{OPT} \geq n^2/2$ gives its multiplicative form.

For completeness, the fixed-cutoff companion uses $B = 32$ and the more conservative fallback criterion $\hat{\kappa} < 2/B$. Substituting these fixed-cutoff choices into the error calculation gives

$$\mathbb{E} \text{ALG} \leq \frac{4}{3} \cdot \text{OPT} + n^2 \left[9472 \sqrt{\frac{d+2}{k}} + \frac{704}{d} \right] + 17(nk/2 + k^2).$$

The estimates $k \leq n^{3/4}$, $1/d \leq 2n^{-1/4}$, and $\sqrt{(d+2)/k} \leq 2n^{-1/4}$ give $\mathbb{E} \text{ALG} \leq \frac{4}{3} \cdot \text{OPT} + 20378n^{7/4}$, with the same deliberately loose constant covering smaller n . This proves Equation (267).

5.6.5 Transfer of the zero–two lower bound

Fix the finite raw-execution cap $u = u_3^{\text{rand}}$. Any randomized obligatory algorithm can be viewed as a revealing-optimization algorithm at this value of u that simply never uses raw execution. Apply lemma 5.4. Its fixed hard input has only values zero and two, so every revealing effective length is

$$\min\{u_3^{\text{rand}}, 1 + p_j\} = 1 + p_j.$$

The online transcript and the offline optimum are therefore exactly the same in the revealing model with raw-execution cap $u = u_3^{\text{rand}}$ and in the obligatory model. For even n , the common offline value is

$$\text{OPT} = \frac{3}{4}n^2 + n,$$

whereas the selected labeling has expected online cost $n^2 - o(n^2)$. Its ratio tends to $4/3$, which proves Equation (268). Together with Equation (266), this completes the proof of Theorem 5.5.

6 Blind optimization: speedup without revelation

We finally separate the two benefits that were bundled together in the revealing-optimization model. Raw execution still takes the known time u , and one unit of optimization still

reduces the processing time to $p_i \in [0, u]$. The difference is that optimization does not reveal p_i . An optimized job must be started blindly and its duration becomes known only when it finishes. We call this the *blind-optimization* model $\text{BO}(u)$.

We study deterministic worst-case algorithms, and randomized algorithms against an oblivious adversary. The deterministic optimum has a particularly simple form: run everything raw up to $u = 2$, and optimize everything after that point. To prove that no adaptive mixture improves these two algorithms, the adversary assigns value zero to a raw job and value u to an optimized job, and stops after a prescribed fraction has been optimized. Randomization is still essential for learning the input: a private random sample is representative after the initial shuffle, whereas an adversary can make a deterministic exploration history atypical.

This small change removes the threshold decision entirely. It also shows that the bounded randomized ratio in the revealing model is genuinely an information effect.

The section first proves that deterministic instance optimality is impossible and determines the exact deterministic worst-case curve. It then proves two closely related randomized statements. The first, Theorem 6.5, is the rigorous version of the informal instance-optimality theorem Theorem 1.5. The second, Theorem 6.6, is the rigorous version of Theorem 1.4 and maximizes that instance-specific value relative to the clairvoyant optimum. Keeping the results together avoids repeating the same reduction to contiguous optimization-and-processing blocks.

For a vector p , put

$$\mu_p = \frac{1}{n} \sum_{i=1}^n p_i$$

and define

$$\Phi_{\text{BO},u}(p) = \frac{1}{2} \min\{u, 1 + \mu_p\}. \quad (296)$$

The two terms have an immediate meaning. Running every job raw uses u units per completion. Optimizing and then blindly processing every job uses $1 + \mu_p$ units per completion on average.

Randomization is necessary for attaining this benchmark on every input.

Theorem 6.1 (No deterministic instance-optimal algorithm). *For every unannounced deterministic nonanticipating algorithm $\mathcal{A}$ and every even n , there are a fixed labeled input $I \in \{0, 2\}^n$ and an announced randomized algorithm $\mathcal{A}'$ such that*

$$\text{ALG}_{\mathcal{A}}(I) \geq \mathbb{E} \text{ALG}_{\mathcal{A}'}(I) + \frac{1}{8}n^2. \quad (297)$$

Consequently no deterministic algorithm is asymptotically instance-optimal in $\text{BO}(2)$.

Proof. Run $\mathcal{A}$ against the following deterministic revelation rule. Give a job value zero if $\mathcal{A}$ runs it raw, and value two if $\mathcal{A}$ optimizes it. Optimization reveals nothing, and the value two is seen only when the optimized job is subsequently processed. Since $\mathcal{A}$ is deterministic, recording the resulting value of every label produces a fixed labeled input with the same transcript.

Let m be the number of optimized jobs on this input. Every raw job consumes two units before completion, whereas every optimized job consumes three. Irrespective of interleaving, the completion times are therefore bounded below by the shortest-first schedule of $n - m$ blocks of length two and m blocks of length three. Hence

$$\text{ALG}_{\mathcal{A}}(I) \geq n^2 + n + \frac{m^2 + m}{2}. \quad (298)$$

The multiset has $n - m$ zeros and m twos. If $m \leq n/2$, let $\mathcal{A}'$ privately permute the jobs and optimize and process them one at a time in that order. Every position sees the average block length $(n + 2m)/n$, so

$$\mathbb{E} \text{ALG}_{\mathcal{A}'}(I) = \frac{(n + 1)(n + 2m)}{2}. \quad (299)$$

Subtracting this from Equation (298) gives $(n - m)(n - m + 1)/2 \geq n^2/8$.

If $m > n/2$, let $\mathcal{A}'$ run every job raw. Its cost is $n(n + 1)$, and the difference in Equation (298) is $(m^2 + m)/2 > n^2/8$. This proves Equation (297). $\square$

An optimization operation reveals no information. We will repeatedly use the following normal form.

Lemma 6.2 (Contiguous-block normal form for optimized jobs). *For every nonanticipating algorithm $\mathcal{A}$ in $\text{BO}(u)$ there is a nonanticipating algorithm $\mathcal{A}^{\text{nf}}$ which completes every optimized job as one contiguous block of length $1 + p_i$ and satisfies*

$$\text{ALG}_{\mathcal{A}^{\text{nf}}}(I) \leq \text{ALG}_{\mathcal{A}}(I) \quad (300)$$

on every input I . If $\mathcal{A}$ is deterministic, then so is $\mathcal{A}^{\text{nf}}$.

Proof. Maintain a virtual execution of $\mathcal{A}$. When the virtual algorithm optimizes a job, record that fact and advance the virtual clock by one unit, but perform no physical work. Immediately before the virtual algorithm later processes that job, perform its postponed optimization physically and then process it. Observations that occur earlier in physical time are buffered until the corresponding virtual completion. Thus the virtual algorithm sees exactly its original history and makes the same decisions. Moving a unit of optimization to the right leaves the optimized job and every later job no later than before, while jobs completed in between move one unit earlier. Applying this to every separated optimization proves (300). $\square$

6.1 The deterministic worst-case curve

For $u > 1$ define

$$G_{\text{BO}}(u) := \frac{\sqrt{4u^3 - 4u + 1} - 1}{2(u - 1)}. \quad (301)$$

The two deterministic algorithms needed below are RAW, which runs every job for time u , and OPTIMIZEALL, which fixes an arbitrary label order and completes every job by one contiguous optimization-and-processing block of length $1 + p_i$.

Theorem 6.3 (Exact deterministic blind-optimization curve). *The optimal deterministic size-asymptotic competitive ratio in $\text{BO}(u)$ is*

$$R_{\text{BO}}^{\text{det}}(u) = \begin{cases} 1, & 0 < u \leq 1, \\ u, & 1 \leq u \leq 2, \\ G_{\text{BO}}(u), & u \geq 2. \end{cases} \quad (302)$$

The algorithm RAW is optimal for $u \leq 2$, and OPTIMIZEALL is optimal for $u \geq 2$; both are optimal at $u = 2$. In particular,

$$R_{\text{BO}}^{\text{det}}(u) = \sqrt{u} + o(1) \quad (u \rightarrow \infty). \quad (303)$$

We first prove the adaptive lower bound. Its stopping rule is the direct analogue of lemma 4.16 for optimization without revelation.

Lemma 6.4 (Hidden binary stopping for optimized jobs). *Fix $u > 1$, a target $R \leq u$, and $\alpha \in (0, 1)$. Against every deterministic algorithm there is a fixed binary input with $p_j \in \{0, u\}$ on which*

$$\text{ALG} - R \cdot \text{OPT} \geq \frac{n^2}{2} \left((u - R)(1 - \sigma^2) + \sigma^2 f_{u,R}^{\text{BO}}(y) \right) - O_u(n), \quad (304)$$

where $\sigma \in [0, 1]$, $0 \leq y - \alpha < 1/(\sigma n)$ when $\sigma > 0$, and

$$f_{u,R}^{\text{BO}}(y) = 1 - R + 2uy - (u + R(u - 1))y^2. \quad (305)$$

Consequently, if $f_{u,R}^{\text{BO}}(\alpha) \geq 0$, then every deterministic algorithm has size-asymptotic ratio at least R .

Proof. By lemma 6.2, it is enough to lower-bound an algorithm that optimizes and processes each such job in one contiguous block. Until the stopping condition below is met, give value zero to every job completed raw and value u to every optimized job. If v raw jobs and L optimized jobs with $p = u$ have been completed, stop at the first completion for which $L > 0$ and

$$L \geq \alpha(n - v). \quad (306)$$

Give value zero to all remaining jobs. The answers are deterministic functions of the transcript and therefore define one fixed labeled binary input that reproduces the same execution.

If the algorithm fails to complete all jobs, its ratio is infinite. Otherwise, if the line is never crossed, then $L = 0$ after all jobs are completed: otherwise $n - v = L$ at the end and $L \geq \alpha L$. Thus the algorithm ran every job raw on the all-zero input and has ratio $u \geq R$.

Suppose the line is crossed. Put $x = n - v - L$. For a lower bound, discard idle time, put the already completed raw jobs before the already completed optimized jobs with $p = u$, reveal the future, and let all x remaining zero jobs be optimized. These exchanges only favor the algorithm: among the operations already completed, shortest first is optimal, while no untouched future job could have completed before the crossing. The resulting exact cost and the clairvoyant optimum are

$$A_{\text{ex}} = u \left(vn - \frac{v(v-1)}{2} \right) + (u+1) \left(L(n-v) - \frac{L(L-1)}{2} \right) + \frac{x(x+1)}{2}, \quad (307)$$

$$O_{\text{ex}} = \frac{n(n+1)}{2} + (u-1) \frac{L(L+1)}{2}. \quad (308)$$

Here the offline optimum first optimizes and processes the $n - L$ zero jobs, using one unit per job, and then runs the L jobs with $p = u$ raw.

Set

$$\sigma = \frac{n-v}{n}, \quad y = \frac{L}{n-v}. \quad (309)$$

Twice the quadratic coefficients of Equations (307) and (308) are

$$\begin{aligned} \mathcal{A} &= u(1 - \sigma^2) + \sigma^2(1 + 2uy - uy^2), \\ \mathcal{O} &= (1 - \sigma^2) + \sigma^2(1 + (u-1)y^2). \end{aligned}$$

Subtracting $R \cdot \mathcal{O}$ from $\mathcal{A}$ gives exactly the bracket in (304). Immediately before the crossing, $L - \alpha(n - v) < 0$. An optimized completion increases this expression by one and a raw completion by α , so

$$0 \leq y - \alpha < \frac{1}{n - v} = \frac{1}{\sigma n}. \quad (310)$$

The quadratic $f_{u,R}^{\text{BO}}$ is uniformly Lipschitz on $[0, 1]$ for fixed u and $R \leq u$. Hence $\sigma^2 f_{u,R}^{\text{BO}}(y) \geq \sigma^2 f_{u,R}^{\text{BO}}(\alpha) - O_u(1/n)$. The exact terms contributed by individual jobs add another $O_u(n)$ before normalization. If $u - R$ and $f_{u,R}^{\text{BO}}(\alpha)$ are nonnegative, this proves the claimed lower bound. $\square$

Proof of Theorem 6.3. If $u \leq 1$, raw execution is an offline-optimal action for every job, so the ratio is one. If $1 < u \leq 2$, use

$$R = u, \quad \alpha = \frac{1}{u}. \quad (311)$$

Then $f_{u,u}^{\text{BO}}(1/u) = 2 - u \geq 0$, and lemma 6.4 gives the lower bound u . Conversely, every offline effective length is at least one, whereas RAW has cost $un(n+1)/2$, so its ratio is at most u .

Now let $u \geq 2$ and put $R = G_{\text{BO}}(u)$. The definition (301) is equivalently the positive solution of

$$(R - 1)(u + R(u - 1)) = u^2. \quad (312)$$

Moreover $R \leq u$, with equality only at $u = 2$. Choose

$$\alpha = \frac{u}{u + R(u - 1)}. \quad (313)$$

This is the maximizer of Equation (305), and (312) gives

$$f_{u,R}^{\text{BO}}(\alpha) = 1 - R + \frac{u^2}{u + R(u - 1)} = 0. \quad (314)$$

Thus lemma 6.4 proves the lower bound R .

It remains to analyze OPTIMIZEALL . Put $a(p) = 1 + p$. For a fixed multiset, its worst label order has the $a(p_i)$ in nonincreasing order. Direct pair accounting gives

$$\text{ALG}_{\text{OPTIMIZEALL}} \leq \frac{1}{2} \sum_{i,j} \max\{a(p_i), a(p_j)\} + \frac{1}{2} \sum_i a(p_i), \quad (315)$$

$$\text{OPT} = \frac{1}{2} \sum_{i,j} \min\{u, 1 + p_i, 1 + p_j\} + \frac{1}{2} \sum_i \min\{u, 1 + p_i\}. \quad (316)$$

Write $S_D(t) = D((t, u])$ for $0 \leq t \leq u$. Expressing a nonnegative value x as $\int_0^\infty \mathbf{1}_{t < x} dt$ yields

$$\iint \max\{a(p), a(q)\} dD(p) dD(q) = 1 + \int_0^u (2S_D(t) - S_D(t)^2) dt, \quad (317)$$

$$\iint \min\{u, 1 + p, 1 + q\} dD(p) dD(q) = 1 + \int_0^{u-1} S_D(t)^2 dt. \quad (318)$$

Set

$$b = \frac{1}{u-1} \int_0^{u-1} S_D(t) dt. \quad (319)$$

Cauchy–Schwarz bounds the square integral on $[0, u-1]$ from below by $(u-1)b^2$. Since S_D is nonincreasing,

$$T := \int_{u-1}^u S_D(t) dt \leq S_D(u-1) \leq b. \quad (320)$$

The function $h \mapsto 2h - h^2$ is increasing and concave on $[0, 1]$. Cauchy–Schwarz on the prefix and Jensen’s inequality on the unit tail therefore give

$$\iint \max\{a(p), a(q)\} dD(p)dD(q) \leq 1 + 2ub - ub^2, \quad (321)$$

$$\iint \min\{u, 1+p, 1+q\} dD(p)dD(q) \geq 1 + (u-1)b^2. \quad (322)$$

Consequently their ratio is at most

$$g_u(b) = \frac{1 + 2ub - ub^2}{1 + (u-1)b^2}. \quad (323)$$

The derivative of g_u changes sign exactly once on $[0, 1]$. At its maximum, if $R = g_u(b)$, the stationary equation $g'_u(b) = 0$ says

$$u(1-b) = R(u-1)b, \quad b = \frac{u}{u + R(u-1)}. \quad (324)$$

Substitution into $g_u(b) = R$ gives (312). That equation has the unique positive solution $G_{\text{BO}}(u)$; hence

$$\max_{0 \leq b \leq 1} g_u(b) = G_{\text{BO}}(u). \quad (325)$$

All survival-function inequalities above are tight for a binary multiset with a b -fraction of values u and the remaining values zero, labeled so that the optimized jobs with $p = u$ occur first. Equations (315)–(323) prove $\text{ALG}_{\text{OPTIMIZEALL}} \leq R \cdot \text{OPT} + O_u(n)$. This matches the lower bound and proves (302). The asymptotic formula (303) follows directly from (301). $\square$

Theorem 6.5 (Blind-optimization instance optimality). *For every fixed $u < \infty$ there exist a randomized nonanticipating algorithm $\mathcal{A}_{\text{BO}}$, depending only on n, u , and a sequence $\varepsilon_u(n) \rightarrow 0$ such that every vector $p \in [0, u]^n$ satisfies*

$$\mathbb{E} \text{ALG}_{\mathcal{A}_{\text{BO}}}(p) \leq n^2 \Phi_{\text{BO},u}(p) + \varepsilon_u(n)n^2. \quad (326)$$

Conversely, every $p \in [0, u]^n$ and every randomized nonanticipating algorithm $\mathcal{A}$, even an announced one, satisfy

$$\mathbb{E} \text{ALG}_{\mathcal{A}}(p) \geq n^2 \Phi_{\text{BO},u}(p) - \varepsilon_u(n)n^2. \quad (327)$$

Thus one algorithm is asymptotically optimal, at the n^2 scale, for every input vector simultaneously. Unlike $\Phi_{\text{RO},u}$ and Φ_{BE} , $\Phi_{\text{BO},u}$ depends only on the empirical mean.

The proof is simpler than in the revealing models. Because optimization reveals nothing, it can be moved next to the processing of its job. Every job is then completed either by raw execution of known length u or by one contiguous optimization-and-processing operation whose average length is $1 + \mu_p$. A sampling bound that holds for every adaptive prefix rules out an adaptive advantage, while a sample estimates the one number μ_p . We first prove this instance-optimal statement and then maximize its value relative to the clairvoyant optimum.

6.2 Proof of randomized instance optimality

By the input convention, the coordinates of p are shuffled uniformly before the algorithm starts. The lower half simply says that no announced algorithm beats $n^2\Phi_{\text{BO},u}(p) - o_u(n^2)$ in expectation.

By lemma 6.2, we may regard every optimized job as one indivisible operation of hidden length $1 + p_i$. Raw execution has the known length u . Either action completes exactly one job.

The announced lower bound. Fix the private seed of an algorithm and place the announced occurrences uniformly behind the labels. We may assume that the algorithm completes every job. Treat equal-valued jobs as distinct and reveal their values in the analysis in first-touch order. By deferred decisions this order is a uniform permutation, and the decision whether to optimize the next job uses only the values revealed by earlier optimized jobs. Running a job raw does not reveal its value to the algorithm; exposing it afterwards only for the analysis does not affect the algorithm's decisions.

Fix $\delta > 0$ and ignore the last δn first touches for the moment. The predictable-permutation estimate lemma 2.7, applied to the processing times and the indicator of the decision to optimize, gives one event of probability $1 - o(1)$ on which every earlier prefix containing tn completed optimized jobs has total hidden processing work

$$\sum_{i \text{ optimized in the prefix}} p_i = \mu_p tn + o_\delta(n). \quad (328)$$

The event is simultaneous over all prefixes, so a stopping fraction chosen adaptively by the algorithm may be substituted only after the event has been fixed. Editing the last δn jobs to either mode changes total completion cost by $O_u(\delta n^2)$.

For one error bound that applies directly at every n , take $\delta_n = n^{-1/4}$. The bounded selected-work estimate used in Section 3.7 has deviation

$$O_u\left((1 + \delta_n^{-1})\sqrt{n \ln(n+2)}\right) = O_u\left(n^{3/4}\sqrt{\ln(n+2)}\right)$$

and failure probability $o(1)$, simultaneously over all relevant prefixes. After division by n , both this deviation and the modification of the last $\delta_n n$ jobs contribute $O_u(n^{-1/4}\sqrt{\ln(n+2)})$ to the normalized completion-area bound.

Put $a = 1 + \mu_p$. If the final optimized fraction is q , the best possible order of the two divisible block types puts the smaller work per completion first. When $a \leq u$ its area is

$$F_{a,u}(q) = a\left(q - \frac{q^2}{2}\right) + \frac{u(1-q)^2}{2}, \quad F'_{a,u}(q) = (a-u)(1-q). \quad (329)$$

It is minimized at $q = 1$, with value $a/2$. When $a \geq u$, jobs run raw go first and the area is

$$F_{a,u}(q) = \frac{u(1-q^2)}{2} + \frac{aq^2}{2} = \frac{u}{2} + \frac{a-u}{2}q^2, \quad (330)$$

which is minimized at $q = 0$, with value $u/2$. The shortest-first argument of lemma 3.2 proves that no interleaving has smaller area. Equations (328)–(330), with the uniform choice $\delta = \delta_n$, give

$$\frac{1}{n^2} \mathbb{E} \text{ALG}_{\mathcal{A}}(p) \geq \frac{1}{2} \min\{u, 1 + \mu_p\} - o_u(1).$$

The estimate is uniform in the deterministic algorithm. Averaging it over the algorithm seed proves (327). Together with the upper bound below, one may take $\varepsilon_u(n) = O_u(n^{-1/4}\sqrt{\ln(n+2)})$ after enlarging its constant.

One algorithm for all input vectors. Take $k = \lceil n^{2/3} \rceil$ uniformly random sample jobs. Optimize each sample job and process it immediately, thereby observing its duration, and let $\hat{\mu}$ be their sample mean. If $1 + \hat{\mu} < u$, optimize and immediately process every remaining job in fresh random order. Otherwise run every remaining job raw.

The sample contributes at most $O_u(nk) = o_u(n^2)$ beyond the main schedule. The variance estimate Equation (11), after scaling from $[0, 1]$ to $[0, u]$, gives

$$\mathbb{E}|\hat{\mu} - \mu_p| = O_u(k^{-1/2} + k/n) = o_u(1).$$

Choosing the wrong mode can cost at most $O(n^2)|\hat{\mu} - \mu_p|$, while choosing the correct mode has leading cost $n^2\Phi_{\text{BO},u}(p)$. This proves (326). In particular, the universal algorithm needs to learn only one number, not a histogram or a threshold.

6.3 The exact worst-case curve

The clairvoyant scheduler knows p_i , uses the effective length $\min\{u, 1 + p_i\}$ from Equation (2), and orders these blocks shortest first. For a distribution D on $[0, u]$, define its leading offline coefficient by

$$\Omega_{\text{BO},u}(D) := \frac{1}{2} \iint \min\{u, 1 + p, 1 + q\} dD(p) dD(q). \quad (331)$$

Consequently the instance-specific ratio from Theorem 6.5 is

$$\alpha_{\text{BO},u}(D) = \frac{\min\{u, 1 + \int p dD(p)\}}{\iint \min\{u, 1 + p, 1 + q\} dD(p) dD(q)}. \quad (332)$$

Let $u_{\text{BO}} > 1$ denote the unique root greater than one of

$$u^3 - 2u^2 - u + 1 = 0, \quad u_{\text{BO}} = 2.2469796037 \dots \quad (333)$$

Theorem 6.6 (Exact blind-optimization curve). *The optimal randomized size-asymptotic competitive ratio in $\text{BO}(u)$ against an oblivious adversary is*

$$R_{\text{BO}}^{\text{rand}}(u) = \begin{cases} 1, & 0 < u \leq 1, \\ \frac{u^3}{u^2 + (u-1)^3}, & 1 \leq u \leq u_{\text{BO}}, \\ \frac{1}{2} \left(1 + \sqrt{\frac{u^2 + u - 1}{u - 1}} \right), & u \geq u_{\text{BO}}. \end{cases} \quad (334)$$

The formulas $u^3/[u^2 + (u-1)^3]$ and $\frac{1}{2}(1 + \sqrt{(u^2 + u - 1)/(u - 1)})$ agree at u_{BO} . Moreover,

$$R_{\text{BO}}^{\text{rand}}(u) = \frac{\sqrt{u}}{2} + \frac{1}{2} + o(1) \quad (u \rightarrow \infty). \quad (335)$$

Proof. The claim is trivial for $u \leq 1$, because raw execution is no longer than the unit optimization operation. Assume $u > 1$ and let

$$S_D(t) = D((t, u]), \quad 0 \leq t \leq u.$$

Writing each nonnegative minimum as an integral of its level indicators in (331) gives

$$2\Omega_{\text{BO},u}(D) = 1 + \int_0^{u-1} S_D(t)^2 dt. \quad (336)$$

Put $S = \int_0^{u-1} S_D(t) dt$. Since S_D is nonincreasing,

$$\int_{u-1}^u S_D(t) dt \leq S_D(u-1) \leq \frac{S}{u-1}.$$

Therefore, writing $\mu = \int p dD(p) = \int_0^u S_D(t) dt$,

$$S \geq \frac{u-1}{u} \mu. \quad (337)$$

Cauchy–Schwarz now yields

$$\int_0^{u-1} S_D(t)^2 dt \geq \frac{S^2}{u-1} \geq (u-1) \left(\frac{\mu}{u} \right)^2. \quad (338)$$

Set $b = \mu/u$. Equations (332)–(338) imply

$$\alpha_{\text{BO},u}(D) \leq f_u(b) := \frac{\min\{u, 1+ub\}}{1+(u-1)b^2}. \quad (339)$$

Equality is attained by the binary distribution

$$D(\{u\}) = b, \quad D(\{0\}) = 1-b. \quad (340)$$

Indeed, its survival function is constant b on $[0, u)$, so both inequalities in (338) are equalities.

It remains to maximize the scalar function f_u . Put

$$b_0 = \frac{u-1}{u}.$$

For $b \geq b_0$ the numerator is u and f_u decreases, so this region is maximized at b_0 . For $b \leq b_0$, differentiation gives the unique unconstrained maximizer

$$b_*(u) = \frac{u}{u-1 + \sqrt{(u-1)(u^2+u-1)}}. \quad (341)$$

The equality $b_*(u) = b_0$ is equivalent to $u^3 - 2u^2 - u + 1 = 0$. Hence b_0 is optimal up to u_{BO} and $b_*(u)$ is optimal afterwards. Substitution gives

$$f_u(b_0) = \frac{u^3}{u^2 + (u-1)^3}, \quad f_u(b_*) = \frac{1}{2} \left(1 + \sqrt{\frac{u^2+u-1}{u-1}} \right).$$

This proves (334). The shuffled binary vectors from (340) give the lower bounds under the input convention. Together with the universal upper algorithm of Theorem 6.5, they prove both sides of the competitive-ratio claim. If the maximizing mass b is irrational, finite vectors use $\lfloor bn \rfloor$ copies of u ; continuity gives the same asymptotic ratio. Finally, $(u^2+u-1)/(u-1) = u+2+1/(u-1)$ gives (335). $\square$

The hard distribution explains the separation. Most jobs have optimized length zero, while a carefully chosen small fraction have length u . Blind optimization is attractive on average, so the online algorithm optimizes the jobs. It cannot recognize a job with $p = u$ before committing to it, and processing such a job early delays many zeros. The clairvoyant optimum optimizes the zeros, runs the jobs with $p = u$ raw, and places them last. In the revealing model, by contrast, the same jobs can be deferred after their tests. The gap between the two curves therefore measures the scheduling value of the revealed information itself.

Acknowledgements

An initial improvement over the then-current deterministic lower bound for obligatory testing was obtained with the Bolzano open-source multi-agent research system [GHK⁺26]. That preliminary result motivated the present project. Most of the subsequent conjecture generation, proof development, counterexample search, exposition, and Lean formalization was carried out with substantial assistance from OpenAI’s GPT-5.6 through interactive Codex sessions. The author selected the research directions, reviewed and revised the arguments, and takes responsibility for the claims in this paper.

A Auditing the main statements in Lean

The accompanying Lean 4 development [dMU21] uses mathlib [The20] and is pinned to version 4.30.0. This appendix is organized as a reader-facing audit of the formal statements. Each item first states, in ordinary mathematical English, what one would like to verify and points to its informal counterparts in Sections 1.2 and 1.4 to 1.6. It then displays the decisive part of a public Lean signature and explains how to read it.

The excerpts suppress namespace prefixes, routine positivity hypotheses, and local parameter definitions, but not the quantified input, algorithm, completion premise, cost comparison, or error term. The exact export name is shown in every excerpt, so that the complete statement can be found by text search in the accompanying source. Randomized Lean statements represent private randomness by a nonempty finite type of seeds and `uniformAverage`; permutations, samples, and tie-breaking coins are all included in such seeds.

A.1 The common asymptotic meaning

All competitive-ratio statements below use the following two predicates.

```
def SizeAsymptoticUpper (alg opt : forall n, Family n -> Real)
  (c : Real) : Prop :=
  forall eps, 0 < eps -> exists N, forall n, N <= n ->
    forall I, alg n I <= (c + eps) * opt n I

def FixedSizeLowerBound (cap : Cap) (ratio : Real) : Prop :=
  forall strategies : forall n, Online.Strategy n,
    forall eps, 0 < eps -> forall N,
      exists n, N <= n /\
        exists I : FixedInput cap n, exists fuel,
          I.defeats (strategies n) fuel (ratio - eps)
```

Here `I.defeats` means that the finite run has settled and either leaves a job incomplete or satisfies $(R - \varepsilon) \text{OPT}(I) \leq \text{ALG}(I)$. Thus the two predicates are exactly the upper and lower quantifier orders used for the size-asymptotic ratios in the paper; in particular, the lower input is fixed after the algorithm has been chosen.

A.2 Obligatory testing: exact competitive ratios

Theorem A.1 (Lean-audited obligatory-testing ratios). *Let $z_* > 1$ be the unique solution of $z_* - 3 = \ln z_*$ and put*

$$R_* = 1 + \frac{2}{z_* - 1} = 1.570574\dots$$

In obligatory testing, the optimal deterministic size-asymptotic competitive ratio is R_* . Against an oblivious adversary, the optimal randomized ratio is $4/3$. This is the formal counterpart of Theorem 1.1.

For the deterministic statement, the public record contains both directions and an actual completing online strategy.

```
structure ObligatoryExactRatioConclusion (ratio : Real) : Prop where
  lower : LowerBound.FixedSizeLowerBound .infinite ratio
  upper :
    exists strategy : forall n, Online.Strategy n,
      UpperBound.CompletesByAnalysisFuel .infinite strategy /\
      SizeAsymptoticUpper
        (UpperBound.strategyCost .infinite strategy)
        (UpperBound.fixedOfflineCost .infinite) ratio

theorem exact_ratio_main :
  (forall {u : Real}, 0 < u ->
    FiniteExactRatioConclusion u (exactCurve u)) /\
    ObligatoryExactRatioConclusion RStar
```

The definitions in `SchedulingPaper/Constants.lean` identify `RStar` with R_* , while the second conjunct supplies the matching lower and upper bounds. The completion field rules out a certificate that only analyzes a partial transcript.

For the randomized statement, Lean exposes a finite upper error and a fixed binary input for the lower bound.

```
theorem unknownMultiset_operational_expectedCost_le_20378
  ... (p : Fin (k + rest) -> Real) (hp : forall i, 0 <= p i) :
  uniformAverage
    (physicalSampledRunCost (k + rest) k d eta h_eta p) <=
    4 / 3 * finiteObligatoryOPT p + 20378 * R ^ 7

theorem oblivious_iid_binary_lower_online_actualOPT
  {n : Nat} (hn : 0 < n)
  (strategy : Seeds -> Online.Strategy n) (fuel : Nat)
  (hdone : forall seed input job, ... = .done) :
  exists input : BinaryInput n,
    (4 * n / (3 * n + 5)) *
      finiteObligatoryOPT (iidBinaryProcessingTime input) <=
      uniformAverage fun seed =>
        Online.runCompletionCost .infinite ...
```

The upper theorem uses the literal fixed-cutoff sample-learn-fallback algorithm $\mathcal{A}_n^{(32)}$ from Theorem 5.5, with $d = \lfloor n^{1/4} \rfloor$, $k = \lfloor n^{3/4} \rfloor$, and cutoff 32. With this checked parameter choice, $R^7 = o(n^2)$. Thus Lean verifies the exact competitive ratio. This is the fixed-cutoff branch now displayed explicitly in algorithm 2, not its growing-cutoff companion. In the lower theorem the selected $\{0, 2\}$ input is outside the seed average, and $4n/(3n + 5) \rightarrow 4/3$.

A.3 Obligatory-testing instance optimality

Theorem A.2 (Lean-audited obligatory instance comparison). *Fix $L < \infty$. There is one randomized nonanticipating algorithm, whose parameters depend on n but not on the input or on L , such that for every $p \in [0, L]^n$ and every completing announced algorithm $\mathcal{B}$ with a finite private seed,*

$$\mathbb{E} \text{ALG}_{\mathcal{A}^*}(p) \leq \mathbb{E} \text{ALG}_{\mathcal{B}}(p) + o_L(n^2).$$

The assignment of the entries of p to public labels is hidden by a private uniform permutation. This is the finite-seed version of Theorem 1.2.

The final Lean comparison is already between the physical learned run and an arbitrary family of completing competitors.

```

theorem exists_fixedPlacement_universalGrowingPolicy_concrete_rate
  {Seeds : Type*} [Fintype Seeds] [Nonempty Seeds]
  ... {L : Real} (hL : 0 < L)
  (p : Fin (k + r) -> Real)
  (hp0 : forall job, 0 <= p job)
  (hpL : forall job, p job <= L)
  (competitor : Seeds -> CompletePolicy p)
  (htest : forall seed, FirstTouchesAreTests (competitor seed)) :
  exists placement,
    uniformAverage (physicalGrowingRunCost ... p) <=
      (k + r : Real) ^ 2 *
        (uniformAverage (fun seed =>
          normalizedCost p (competitor seed) placement) +
          2 * concreteUniversalGrowingError L n)

theorem concreteUniversalGrowingError_tendsto_zero (L : Real) :
  Tendsto (concreteUniversalGrowingError L) atTop (nhds 0)

```

`FirstTouchesAreTests` is exactly the obligatory-testing restriction. The conclusion chooses one hidden placement before averaging over the competitor's seed, while the learned algorithm averages over its own private sample and order. The second theorem turns the displayed normalized error into $o_L(1)$.

The checked universal growing-cutoff family is structurally the same sample-learn-fallback algorithm but uses a deliberately slower convenient parameter sequence: if $m = \lfloor n^{1/32} \rfloor$, it has cutoff $m - 1$, $m^2 - 1$ positive grid intervals, mesh $1/m$, and pilot size of order $n^{7/8}$. The development does not currently identify this literal family with the fixed-cutoff family used above or prove the uniform unbounded $4/3$ estimate for this growing family. The manuscript separately proves its sharper $n^{3/4}$ -pilot, $32 + n^{1/20}$ -cutoff version. Hence the exact ratio and the bounded instance comparison are both checked, but their realization by one definitionally identical Lean strategy is not part of the audit.

A.4 Why the obligatory instance assumptions are necessary

Lean also checks both limitations stated after Theorem 1.2. Every completing deterministic policy has a fixed balanced zero-two input on which its literal cost is at least $9n^2/8$. Without a fixed input bound, every finite-seed randomized policy has a fixed input in $[0, n^2]^n$ on which it loses at least $n^2/8$ against the corresponding announced comparison schedule.

```

theorem deterministic_balanced_cost_lower
  (m : Nat) (strategy : Strategy (2 * m)) (fuel : Nat)
  (hcomplete : forall processing, ... -> CompletesAll processing strategy fuel) :
  exists processing,
    (forall job, processing job = 0 \/\ processing job = 2) /\
    (Finset.univ.filter fun job => processing job = 2).card = m /\
    9 * ((2 * m : Nat) : Real) ^ 2 / 8 <=
      runCompletionCost .infinite processing ...

theorem finiteSeed_exists_operational_quadratic_gap
  {n : Nat} (hn : 7 <= n)

```

```

(strategy : Seeds -> Strategy n) (fuel : Nat)
(hdoneHigh : ...) (hdoneZero : ...) :
n ^ 2 / 8 <= averageCostOnAllHigh - announcedAllHighCost \ /
exists label,
  n ^ 2 / 8 < averageCostWithOneZero label - announcedOneZeroUpper

```

The first input is frozen after the deterministic strategy is fixed. In the second theorem the exceptional zero label, when that branch is selected, is outside the private-seed average. These are the two operational quantifier orders required by Theorem 3.18.

A.5 Blind execution

Theorem A.3 (Lean-audited blind-execution statements). *For every fixed $L < \infty$, one randomized nonanticipating algorithm is instance-optimal on every $p \in [0, L]^n$ up to $o_L(n^2)$, even against an announced completing competitor. If no common upper bound on processing times is imposed, no randomized algorithm has a finite asymptotic competitive ratio. The bounded statement is Theorem 1.3; the preceding paragraph in Section 1.4 states the unbounded limitation.*

The bounded comparison is a direct per-input inequality.

```

theorem boundedUniform_blindPilot_concrete_rate
  {n : Nat} ... {L : Real} (hL : 0 < L)
  (p : Fin n -> Real)
  (hp0 : forall job, 0 <= p job)
  (hpL : forall job, p job <= L)
  (competitor : CompletePolicy p) :
  uniformAverage (fun pilotOrder =>
    uniformAverage (blindPilotLearnedCost G pilot pilotOrder)) /
    (n : Real) ^ 2 <=
    uniformAverage (normalizedCost p competitor) +
    7830 * (L + 1) ^ 2 / concreteUnknownParameter n

```

The left side is the learned four-block algorithm; the right side is the private-shuffle cost of an arbitrary announced competitor. The checked parameter tends to infinity, so the last term tends to zero. The finite-seed corollary `boundedUniform_blindPilot_finiteSeed_concrete_rate` performs the pointwise conditioning and averaging in Lean itself.

For the unbounded part, Lean states the contradiction entirely in terms of literal completion costs.

```

theorem no_finite_ratio_at_quadratic_scale_operational
  {n : Nat} (hn : 0 < n)
  (strategy : Seeds -> Strategy n)
  (hcomplete : forall seed, CompletesAllNonnegative (strategy seed))
  (R eta : Real) ...
  (hzeroCost : average zeroCost <= eta * n ^ 2)
  (hcompetitive : forall label,
    average (oneLongCost (n ^ 2) label) <=
      R * n ^ 2 + eta * n ^ 2) : False

```

Thus a low cost on the all-zero input and an R -competitive cost on every input with one length- n^2 job cannot hold simultaneously for large n .

A.6 Blind optimization: the two worst-case curves

Theorem A.4 (Lean-audited blind-optimization curves). *For every fixed $u > 0$, the exact deterministic ratio is*

$$R_{\text{BO}}^{\text{det}}(u) = \begin{cases} 1, & u \leq 1, \\ u, & 1 \leq u \leq 2, \\ \frac{\sqrt{4u^3 - 4u + 1} - 1}{2(u - 1)}, & u \geq 2, \end{cases}$$

and the exact randomized ratio is

$$R_{\text{BO}}^{\text{rand}}(u) = \begin{cases} 1, & u \leq 1, \\ \frac{u^3}{u^2 + (u - 1)^3}, & 1 \leq u \leq u_{\text{BO}}, \\ \frac{1}{2} \left(1 + \sqrt{\frac{u^2 + u - 1}{u - 1}} \right), & u \geq u_{\text{BO}}, \end{cases}$$

where $u_{\text{BO}} > 1$ is the unique root of $u^3 - 2u^2 - u + 1 = 0$. This expands the two curves summarized in Theorem 1.4.

The Lean definitions contain these formulas literally.

```
def deterministicCurve (u : Real) : Real :=
  if u <= 1 then 1
  else if u <= 2 then u
  else (sqrt (4 * u ^ 3 - 4 * u + 1) - 1) / (2 * (u - 1))

def randomizedCurve (u : Real) : Real :=
  if u <= 1 then 1
  else if u <= randomizedTransition then
    u ^ 3 / (u ^ 2 + (u - 1) ^ 3)
  else
    (1 + sqrt ((u ^ 2 + u - 1) / (u - 1))) / 2
```

For the deterministic curve, one exported theorem gives both a completing upper strategy and a fixed input against every completing strategy, with linear remainders.

```
theorem deterministicCurve_operational_exact (hu : 0 < u) :
  (exists strategy : forall n, Online.Strategy n,
    (forall n, Online.CompletesAll u (strategy n)) /\
    forall n processing, admissible u processing ->
      runCost u processing (strategy n) n <=
        deterministicCurve u * offlineCost u processing +
        (1 + u) * n / 2) /\
  (forall n, 0 < n -> forall strategy,
    Online.CompletesAll u strategy ->
    exists processing, admissible u processing /\
      runCost u processing strategy n >=
        deterministicCurve u * offlineCost u processing -
        (2 * u ^ 2 + u) * n)
```

For randomization, the literal pilot algorithm supplies the upper direction, and finite Yao selects a fixed binary input for the lower direction.

```
theorem operationalPilot_expectedCost_le_curve_normalized ... :
  uniformAverage (fun sampleOrder =>
```

```

conditionalExpectedPilotRunCost ... sampleOrder) <=
  randomizedCurve u * offlineCost u processing +
  pilotNormalizedError u n pilotSize * n ^ 2

theorem exists_binary_mass_operational_ratio_tendsto_curve (hu : 1 < u) :
  exists b in Set.Icc (0 : Real) 1,
    Tendsto (fun n => binaryFiniteYaoRatio n u b)
      atTop (nhds (randomizedCurve u)) /\
  forall n Seeds, 0 < n -> forall strategy : Seeds -> Strategy n,
    (forall seed, CompletesAll u (strategy seed)) ->
      exists input : BinaryInput n,
        binaryFiniteYaoRatio n u b * offlineCost u input <=
          uniformAverage fun seed => runCost u input (strategy seed) n

```

The theorem `pilotNormalizedError_tendsto_zero` verifies that any growing sublinear pilot makes the upper error vanish.

A.7 Blind-optimization instance optimality

Theorem A.5 (Blind-optimization instance statement). *For fixed $u < \infty$, a sample-then-commit randomized algorithm satisfies, for every $p \in [0, u]^n$,*

$$\mathbb{E} \text{ALG}_{\mathcal{A}^*}(p) \leq \frac{n(n+1)}{2} \min\{u, 1 + \mu_p\} + o_u(n^2), \quad \mu_p = \frac{1}{n} \sum_i p_i.$$

Conversely, every completing algorithm with a finite private seed has expected cost at least

$$\frac{n^2}{2} \min\{u, 1 + \mu_p\} - nu\sqrt{2n}.$$

Hence both sides have the same leading value and Lean verifies the full instance-optimal comparison in Theorem 1.5.

Lean checks both the operational sample-then-commit construction and the arbitrary-policy lower benchmark:

```

theorem conditionalExpectedPilotRunCost_eq ... :
  conditionalExpectedPilotRunCost ... = conditionalExpectedPilotCost ...

theorem pilotPolicy_expectedCost_le ... :
  uniformAverage (fun sampleOrder =>
    conditionalExpectedPilotCost ... sampleOrder) <=
    n * (n + 1) / 2 * min u (1 + populationMean processing) +
    (1 + u) * pilotSize * n +
    n * (n + 1) / 2 * (u / sqrt pilotSize)

theorem finiteSeed_expectedCost_ge_instanceBenchmark_nsq
  {Seeds : Type*} [Fintype Seeds] [Nonempty Seeds]
  ... (p : Fin n -> Real)
  (hp : forall job, p job in Set.Icc 0 u)
  (strategy : Seeds -> Online.Strategy n)
  (hcompletes : forall seed placement,
    Online.Completes (placedProcessing p placement) (strategy seed)) :
  n ^ 2 / 2 * min u (1 + populationMean p) -
  n * u * sqrt (2 * n) <=
  uniformAverage fun seed =>
    uniformAverage fun placement =>
      Online.runCost u (placedProcessing p placement)
        (strategy seed) n

```



```

theorem instanceBenchmark_normalizedError_tendsto_zero (u : Real) :
  Tendsto (fun n : Nat => u * sqrt (2 / (n : Real))) atTop (nhds 0)

```

The upper errors are $o_u(n^2)$ for any growing sublinear pilot. The lower error is explicit; the third export proves that after division by n^2 it is $u\sqrt{2/n} = o_u(1)$. Its proof compiles a literal completing online strategy into a causal first-touch trace, applies a uniform-permutation first-moment bound at every position, and averages directly over the algorithm's private seed. Thus no conditioning step is left to prose.

A.8 Revealing optimization: exact competitive-ratio curves

Theorem A.6 (Lean-audited revealing-optimization curves). *For every $u > 0$, the deterministic ratio is the six-piece function $R_{\text{RO}}^{\text{det}}(u)$ displayed in Equation (96). The randomized ratio is*

$$R_{\text{RO}}^{\text{rand}}(u) = \begin{cases} 1, & 0 < u \leq 1, \\ \frac{u^3}{u^2 + (u-1)^3}, & 1 \leq u \leq u_2, \\ 1 + \frac{1}{2(\sqrt{u}-1)}, & u_2 \leq u \leq 25/4, \\ 4/3, & u \geq 25/4, \end{cases}$$

where $u_2 > 1$ is the unique root of $u^3 - 6u^2 + 5u - 1 = 0$. These are the curves summarized in Theorem 1.6.

The deterministic curve is the first conjunct of the same top-level theorem used for the obligatory endpoint:

```

theorem exact_ratio_main :
  (forall {u : Real}, 0 < u ->
    FiniteExactRatioConclusion u (exactCurve u)) /\
    ObligatoryExactRatioConclusion RStar

def exactCurve (u : Real) : Real :=
  if u <= 1 then 1
  else if u <= uDiamond then u
  else if u <= uZero then rhoI u
  else if u <= goldenRatio + 2 then reciprocalBranch u
  else if u <= zStar then mixedFiniteCurve u
  else RStar

```

Here `FiniteExactRatioConclusion` has the same lower field, completing strategy, and `SizeAsymptoticUpper` field displayed at the start of this appendix. The branch functions and all five joins are proved in `SchedulingPaper/ExactCurve.lean`.

For the randomized curve, the scalar definition and the two attaining binary families are explicit:

```

def randomizedCurve (u : Real) : Real :=
  if u <= 1 then 1
  else if u <= transition then boundaryRatio u
  else if u <= 25 / 4 then interiorRatio u
  else 4 / 3

theorem binaryFamilies_le_curve (hu : 1 < u) :
  (forall tau, 1 <= tau -> tau <= u ->
    familyB u tau <= randomizedCurve u) /\

```

```

(forall tau, 1 <= tau -> tau <= u - 1 ->
  familyA tau <= randomizedCurve u)

theorem binaryFamilies_attain_curve (hu : 1 < u) :
  (exists tau in Set.Icc 1 u,
    familyB u tau = randomizedCurve u) \ /
  (exists tau in Set.Icc 1 (u - 1),
    familyA tau = randomizedCurve u)

```

The operational Yao theorem `exists_operational_binary_family_ratio_tendsto_curve` then selects one fixed labeled binary input outside every finite seed average. The theorem `randomizedRevealingCurveExact` is one public package containing this lower theorem, the announced operational upper theorem, the checked announced-to-unannounced transfer, and vanishing of the transfer error. The transfer itself is exported as `compiledExpectedNormalizedCost_le_curve_of_competitor`: given a finite-seed announced policy satisfying the curve bound on every placement, it returns the corresponding direct bound for the literal unannounced pilot learner.

A.9 Revealing-optimization instance optimality

Theorem A.7 (Lean-audited revealing instance comparison). *Fix $u < \infty$. There is one randomized nonanticipating algorithm such that for every $p \in [0, u]^n$ and every completing announced algorithm $\mathcal{B}$ with a finite private seed,*

$$\mathbb{E} \text{ALG}_{\mathcal{A}^*}(p) \leq \mathbb{E} \text{ALG}_{\mathcal{B}}(p) + o_u(n^2).$$

This is the finite-seed version of Theorem 1.7.

The final public theorem compares the literal learned strategy directly with the arbitrary competitor on one hidden placement.

```

theorem exists_fixedPlacement_compiled_le_randomizedCompetitor_concrete_rate
  {Seeds : Type*} [Fintype Seeds] [Nonempty Seeds]
  ... {u : Real} (hu : 0 < u)
  (p : Fin n -> Real)
  (hp0 : forall job, 0 <= p job)
  (hpu : forall job, p job <= u)
  (competitor : Seeds -> CompletePolicy p) :
  exists placement,
    compiledExpectedNormalizedCost grid pilot u <=
      (uniformAverage fun seed =>
        normalizedRawCost u p (competitor seed) placement) +
        concreteRevealingComparisonError u n

theorem concreteRevealingComparisonError_tendsto_zero (u : Real) :
  Tendsto (concreteRevealingComparisonError u) atTop (nhds 0)

```

The left side is an actual fixed-fuel transcript-only strategy averaged over independent pilot and main permutations. The competitor may adapt among tests, processing, and raw execution and is given the input multiset. The placement is chosen before its seed is averaged, and the second theorem supplies the required $o_u(1)$ normalized error.

The concrete checked pilot has size $\lfloor n/m^2 \rfloor = \Theta(n^{7/8})$, where $m = \lfloor n^{1/16} \rfloor$. This is now the parameter choice displayed in the introductory pseudocode as well.

A.10 Certified numerical summaries

The curve definitions remain exact, but the short decimal summaries in Section 1 are no longer justified only by external calculation. The file `SchedulingPaper/NumericalBounds.lean` exports, among others,

```
theorem RStar_decimal_bounds :  
  (15705 : Real) / 10000 < RStar /\ RStar < 15706 / 10000  
  
theorem uDiamond_decimal_bounds :  
  (1866 : Real) / 1000 < uDiamond /\ uDiamond < 1867 / 1000  
  
theorem uZero_decimal_bounds :  
  (3147 : Real) / 1000 < uZero /\ uZero < 3148 / 1000  
  
theorem RevealingOptimization.transition_decimal_bounds :  
  (5048917 : Real) / 1000000 < transition /\  
    transition < 5048918 / 1000000  
  
theorem RevealingOptimization.globalMaximum_decimal_bounds :  
  (1625 : Real) / 1000 < globalMaximum /\  
    globalMaximum < 1626 / 1000
```

The algebraic intervals are discharged by exact polynomial arithmetic. For R_* , Lean brackets $\log q$ by a finite, rational atanh series, proves $4.5051 < z_* < 4.5056$, and then derives the displayed interval for R_* .

A.11 Proof boundary and reproduction

The root module `SchedulingPaper.lean` imports the developments used by all exports above. The checked project contains no `sorry`, `admit`, or project-defined `axiom`. It uses `mathlib`'s classical real analysis and noncomputable choices of exact real parameters; it proves mathematical properties of the algorithms but does not extract a floating-point implementation. The randomized lower statements shown here quantify over nonempty finite seed types.

All mathematical directions of the Section 1 headline theorems now have Lean exports. The checked obligatory 4/3 and bounded instance-optimal upper bounds currently use different concrete parameter families, as recorded above. Lean verifies the exact symbolic constants, branch transition equations, and the rational decimal brackets listed above. Longer printed digit strings remain readability aids unless a correspondingly narrow interval is explicitly exported.

From the repository root, the complete development is checked with

```
lake build
```

and each displayed export can be located by searching for its theorem name.

References

- [ABC17] Peyman Afshani, J  r  my Barbay, and Timothy M. Chan. Instance-optimal geometric algorithms. *Journal of the ACM*, 64(1):3:1–3:38, 2017. doi:10.1145/3046673.
- [AE21] Susanne Albers and Alexander Eckl. Explorable uncertainty in scheduling with non-uniform testing times. In Christos Kaklamanis and Asaf Levin, editors,

- Approximation and Online Algorithms: 18th International Workshop, WAOA 2020, Revised Selected Papers*, volume 12806 of *Lecture Notes in Computer Science*, pages 127–142. Springer, 2021. doi:10.1007/978-3-030-80879-2_9.
- [Azu67] Kazuoki Azuma. Weighted sums of certain dependent random variables. *Tohoku Mathematical Journal*, 19(3):357–367, 1967. doi:10.2748/tmj/1178243286.
- [BS26] Felix Buld and Andreas S. Schulz. Scheduling with testing: Competitive algorithms for minimizing the total weighted completion time in the adversarial model, 2026. arXiv:2606.25166.
- [Che52] Herman Chernoff. A measure of asymptotic efficiency for tests of a hypothesis based on the sum of observations. *The Annals of Mathematical Statistics*, 23(4):493–507, 1952. doi:10.1214/aoms/1177729330.
- [DDN⁺22] Fanny Dufossé, Christoph Dürr, Noël Nadal, Denis Trystram, and Óscar C. Vásquez. Scheduling with a processing time oracle. *Applied Mathematical Modelling*, 104:701–720, 2022. doi:10.1016/j.apm.2021.12.020.
- [DEL24] Konstantinos Dogeas, Thomas Erlebach, and Ya-Chun Liang. Scheduling with obligatory tests. In Timothy Chan, Johannes Fischer, John Iacono, and Grzegorz Herman, editors, *32nd Annual European Symposium on Algorithms (ESA 2024)*, volume 308 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 48:1–48:14, Dagstuhl, Germany, 2024. Schloss Dagstuhl – Leibniz-Zentrum für Informatik. doi:10.4230/LIPIcs.ESA.2024.48.
- [DEMM20] Christoph Dürr, Thomas Erlebach, Nicole Megow, and Julie Meißner. An adversarial model for scheduling with testing. *Algorithmica*, 82(12):3630–3675, 2020. doi:10.1007/s00453-020-00742-2.
- [DKL⁺23] Christoph Damerius, Peter Kling, Minming Li, Chenyang Xu, and Ruilong Zhang. Scheduling with a limited testing budget: Tight results for the offline and oblivious settings. In *31st Annual European Symposium on Algorithms (ESA 2023)*, volume 274 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 38:1–38:15. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2023. doi:10.4230/LIPIcs.ESA.2023.38.
- [dMU21] Leonardo de Moura and Sebastian Ullrich. The Lean 4 theorem prover and programming language. In *Automated Deduction – CADE 28*, volume 12699 of *Lecture Notes in Computer Science*, pages 625–635. Springer, 2021. doi:10.1007/978-3-030-79876-5_37.
- [Doo53] Joseph L. Doob. *Stochastic Processes*. Wiley, New York, 1953.
- [GCH24] Mingyang Gong, Zhi-Zhong Chen, and Kunitaru Hayashi. Approximation algorithms for multiprocessor scheduling with testing to minimize the total job completion time. *Algorithmica*, 86(5):1400–1427, 2024. doi:10.1007/s00453-023-01198-w.
- [GHK⁺26] Jan Grebík, Pavel Hubáček, Martin Koucký, Matěj Kripner, Václav Rozhoň, Robert Šámal, and Adrián Zámečník. Bolzano: Case studies in LLM-assisted mathematical research, 2026. URL: <https://arxiv.org/abs/2604.16989>, arXiv:2604.16989.

- [Hoe63] Wassily Hoeffding. Probability inequalities for sums of bounded random variables. *Journal of the American Statistical Association*, 58(301):13–30, 1963. doi:10.1080/01621459.1963.10500830.
- [Kah91] Simon Kahan. A model for data in motion. In *Proceedings of the 23rd Annual ACM Symposium on Theory of Computing (STOC 1991)*, pages 265–277. Association for Computing Machinery, 1991. doi:10.1145/103418.103449.
- [KLL⁺26] Bob Krekelberg, Alison Hsiang-Hsuan Liu, Fu-Hong Liu, Prudence W. H. Wong, and Xiao-Ou Zhang. The power of amortization on minimizing total completion time with explorable uncertainty, 2026. Version 2, revised 8 May 2026. arXiv:2503.13357.
- [LL26] Kao-Chuan Liang and Ya-Chun Liang. The completion-threshold framework for obligatory-test scheduling on multiple machines, 2026. Version 2, revised 14 July 2026. arXiv:2606.02029.
- [LLWZ23] Alison Hsiang-Hsuan Liu, Fu-Hong Liu, Prudence W. H. Wong, and Xiao-Ou Zhang. The power of amortization on scheduling with explorable uncertainty. In Jaroslaw Byrka and Andreas Wiese, editors, *Approximation and Online Algorithms: 21st International Workshop, WAOA 2023*, volume 14297 of *Lecture Notes in Computer Science*, pages 90–103. Springer, 2023. doi:10.1007/978-3-031-49815-2_7.
- [LMS19] Retsef Levi, Thomas L. Magnanti, and Yaron Shaposhnik. Scheduling with testing. *Management Science*, 65(2):776–793, 2019. doi:10.1287/mnsc.2017.2973.
- [LMS24] Retsef Levi, Thomas L. Magnanti, and Yaron Shaposhnik. Scheduling with testing of heterogeneous jobs. *Management Science*, 70(5):2934–2953, 2024. doi:10.1287/mnsc.2023.4833.
- [MPT94] Rajeev Motwani, Steven J. Phillips, and Eric Torng. Nonclairvoyant scheduling. *Theoretical Computer Science*, 130(1):17–47, 1994. doi:10.1016/0304-3975(94)90151-1.
- [NRTT24] Shyam Narayanan, Václav Rozhoň, Jakub Tětek, and Mikkel Thorup. Instance-optimality in I/O-efficient sampling and sequential estimation. In *65th IEEE Annual Symposium on Foundations of Computer Science (FOCS 2024)*, pages 658–688. IEEE, 2024. doi:10.1109/FOCS61266.2024.00048.
- [The20] The mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020)*, pages 367–381. Association for Computing Machinery, 2020. doi:10.1145/3372885.3373824.
- [Yao77] Andrew Chi-Chih Yao. Probabilistic computations: Toward a unified measure of complexity. In *18th Annual Symposium on Foundations of Computer Science*, pages 222–227. IEEE, 1977. doi:10.1109/SFCS.1977.24.