

Exact Deterministic Size-Asymptotic Ratios for Single-Machine Scheduling with Uniform Tests

Bolzano team

August 2026

Abstract

We determine the exact deterministic size-asymptotic competitive ratio for a single-machine family with unit tests, total completion time, and a common upper limit $u \in (0, \infty]$. For finite u , a job may be executed raw for u time or tested for one unit and then processed for its revealed time. The endpoint $u = \infty$ is obligatory testing: raw execution is unavailable. Its exact value is

$$R_{\text{obl}} = 1 + r = 1.570574096649395\dots, \quad r = \frac{2}{z_\star - 1}, \quad z_\star - 3 = \log z_\star.$$

For finite u , the exact curve $V(u)$ has six explicit regimes: 1, u , an algebraic root, $1 + 1/\sqrt{u-1}$, a one-parameter transcendental interpolation, and the plateau R_{obl} . In fact, $V(u) = V(\infty) = R_{\text{obl}}$ for every $u \geq z_\star$. The global maximum is

$$\sup_{u>0} V(u) = \frac{1 + \sqrt{3 + 2\sqrt{5}}}{2} = 1.866760399173862\dots$$

We provide matching lower bounds and formally specified online algorithms. The smaller- u regimes use binary stopping games and two simple policies. The obligatory endpoint and the two large- u regimes share one harmonic lower-bound core and one four-endpoint bank potential. Finite u adds one capped endpoint; a reserve pays for it on the transition interval and becomes unnecessary at $z_\star$. The upper estimates are $\text{ALG} \leq V(u) \text{OPT} + O_u(n)$, with one absolute $O(n)$ term simultaneously for all $u \in [z_\star, \infty]$. All equalities are exact in the size-asymptotic sense.

1 Introduction and proof intuition

Scheduling with testing separates information acquisition from productive work. A test reveals the unknown processing time of a job, but the test itself occupies the machine and delays every unfinished job. This makes a locally cheap experiment globally expensive: one unit spent testing near the beginning of a schedule can be counted in the completion time of almost every job.

We study one machine, unit tests, nonpreemptive operations, and total completion time. All jobs are initially available. For a finite common upper limit u , a job can either be executed raw for u time units, or be tested for one unit and then processed for its revealed time $p \in [0, u]$. It is useful to extend this family by one endpoint:

$$u = \infty \quad \Longleftrightarrow \quad \text{raw execution is unavailable and every job must be tested.} \quad (1)$$

Thus obligatory testing is not a separate phenomenon; it is the cap-free endpoint of the common-upper family. This viewpoint exposes which parts of the proofs are shared and which difficulty is created by a finite cap.

For a clairvoyant scheduler, a job has effective length

$$q_u(p) = \min\{u, 1 + p\}, \quad q_\infty(p) = 1 + p. \quad (2)$$

The optimum simply orders these effective lengths from shortest to longest. The online scheduler does not know whether a test will save almost u time or reveal a job so long that the optimum would have skipped the test. The entire problem is the cost of making this distinction too late.

1.1 The exact answers

The obligatory endpoint had a gap between the best known deterministic lower and upper bounds [DEL24, LL26]. The finite common-upper problem likewise had no exact answer for arbitrary revealed values [DEMM20]. We close both gaps in the natural size-asymptotic convention.

Let $z_\star > 1$ be the unique solution of

$$z_\star - 3 = \log z_\star, \quad (3)$$

where all logarithms are natural, and put

$$R_\star = 1 + \frac{2}{z_\star - 1} = 1.570574096649395\dots, \quad z_\star = 4.505241495792883\dots \quad (4)$$

Theorem 1.1 (Obligatory endpoint). *The following matching statements hold for $u = \infty$.*

- (i) *For every deterministic online algorithm $\mathcal{B}$ and every $\varepsilon > 0$, there are arbitrarily large instances I for which*

$$\text{ALG}_{\mathcal{B}}(I) \geq (R_\star - \varepsilon) \text{OPT}(I).$$

- (ii) *There is one deterministic algorithm $\mathcal{A}$ and one absolute constant C such that every n -job instance satisfies*

$$\text{ALG}_{\mathcal{A}}(I) \leq R_\star \text{OPT}(I) + Cn.$$

Consequently the optimal deterministic size-asymptotic value at $u = \infty$ is exactly $R_\star$.

For the finite curve, let $\varphi = (1 + \sqrt{5})/2$ and define

$$u_\diamond = \frac{1 + \sqrt{3 + 2\sqrt{5}}}{2}, \quad u_0 = 1 + s_0, \quad s_0^3 = (s_0 + 1)^2,$$

where $s_0 > 1$. For $s = u - 1$, let $\rho_I(u)$ be the positive root of

$$s\rho^2 + (s^3 + s^2 + 1)\rho - (s^2 + s + 1)(s + 2) = 0.$$

On $\varphi + 2 \leq u \leq z_\star$, define $C_{\text{mix}}(u) = 1 + c$ by the unique pair (c, m) satisfying

$$\text{artanh } m - m = 1 - \frac{1}{c} + \frac{1}{2} \log\left(1 + \frac{2}{c}\right), \quad u = 1 + \frac{2}{c} - m, \quad (5)$$

with $2/(z_\star - 1) \leq c \leq 1/\varphi$ and $0 \leq m \leq 1/\varphi$.




Figure 1: The exact deterministic size-asymptotic ratio for finite common upper limit u . The curve stabilizes at the obligatory endpoint value; dotted lines mark its five regime changes.

Theorem 1.2 (Finite common-upper curve). *For every fixed $u > 0$, the optimal deterministic size-asymptotic competitive ratio is*

$$V(u) = \begin{cases} 1, & 0 < u \leq 1, \\ u, & 1 \leq u \leq u_\diamond, \\ \rho_I(u), & u_\diamond \leq u \leq u_0, \\ 1 + 1/\sqrt{u-1}, & u_0 \leq u \leq \varphi + 2, \\ C_{\text{mix}}(u), & \varphi + 2 \leq u \leq z_\star, \\ R_\star, & u \geq z_\star. \end{cases} \quad (6)$$

Every lower branch holds against arbitrary deterministic algorithms on arbitrarily large fixed instances. Every upper branch is attained up to $O_u(n)$. On the finite part of the last branch, one algorithm and one absolute $O(n)$ constant work simultaneously for all $u \geq z_\star$. Together with Theorem 1.1, the same witness covers the endpoint $u = \infty$.

The finite curve is continuous, rises from one to its global maximum

$$V(u_\diamond) = u_\diamond = 1.866760399173862\dots,$$

and then decreases to the obligatory value. In fact it does more than converge:

$$V(u) = V(\infty) = R_\star \quad \text{for every } u \geq z_\star. \quad (7)$$

This is equality of size-asymptotic game values, not equality of the finite problems. At any finite u , the optimum may still save a test on a job with $p \geq u - 1$.

1.2 Why total completion time leads to pair accounting

The most useful way to think about total completion time is pairwise. Every operation of job i contributes once to C_i , and it contributes once more for each other job that remains

unfinished while that operation is executed. Equivalently, the objective consists of diagonal job terms and one delay term for every unordered pair of jobs. The offline pair term is simply the shorter effective length. The online pair term records whether an earlier revealed job was processed immediately or left waiting.

This representation explains both sides of our proof. A lower-bound adversary tries to make many pairs wait in the wrong order. An upper-bound algorithm tries to ensure that every expensive local decision can be charged to a corresponding offline pair. Since there are $\Theta(n^2)$ pairs but only n diagonal terms, controlling the quadratic pair coefficient gives the exact size-asymptotic ratio; bounded rounding errors of $O(1)$ per job become only $O(n)$.

1.3 Lower-bound intuition: hide the short jobs until the end

At the obligatory endpoint, the adversary reveals equal blocks of positive jobs from longer to shorter values and reveals a large zero block last. The algorithm is forced to choose between two risks. Processing a revealed job early may delay many still-hidden zeros. Deferring it may leave a large tail that delays everything completed later. The adaptive revelation is frozen after the interaction, so a standard replay argument produces one fixed instance for the chosen deterministic algorithm.

The processing times are separated by harmonic gaps. This choice is not a guess about the final constant. The zero mass and the number of shorter positive blocks give two multiplicities; their sum exactly cancels the denominator of each harmonic gap. After this cancellation, every possible postponement made by the algorithm has a nonnegative coefficient and every own-phase choice contributes the same completed square. All algorithmic decisions disappear from the leading term. What remains is a one-variable ratio describing the relative positive and zero masses, and its stationary equation is (3).

For finite u , two additional adversarial pictures appear. When u is small or moderate, binary values $p \in \{0, u\}$ already suffice: the adversary watches how much work the algorithm completes before revealing whether the next jobs are zeros or caps. For the transition to the obligatory value, the hard instance is a capped prefix followed by a scaled copy of the harmonic core. The prefix exploits the optimum's ability to skip tests; the core exploits uncertainty in the test order. As u grows, the optimal cap mass decreases continuously to zero, leaving exactly the obligatory construction.

1.4 Upper-bound intuition: three policies and one common bank

The exact upper curve uses three simple algorithmic shapes. Their formal pseudocode appears later; here we describe why they are natural.

approximate range	upper-bound policy at a glance
$u \lesssim 1.87$	Do not test: execute every job raw.
$1.87 \lesssim u \lesssim 3.15$	Test all jobs; always finish an initial fraction, then use a short/long threshold. The initial fraction shrinks as u grows.
$3.15 \lesssim u \lesssim 3.62$	Test all jobs; finish values at most one immediately and sort the rest in a final tail.
$3.62 \lesssim u \lesssim 4.51$	Use the history-dependent threshold; the proof adds a reserve for capped outcomes.
$u \gtrsim 4.51$, including ∞	Use the same adaptive threshold, now with no cap reserve.

Raw policy. When u is very small, information costs at least as much as it can save. The algorithm therefore runs every job raw. Slightly beyond that trivial range, raw execution need no longer be clairvoyantly optimal for each job, but comparing its length u with the optimum’s effective lengths already gives the first nontrivial minimax branch.

Forced-prefix threshold policy. In the next regimes the algorithm tests every job. During an initial fraction of the test order it processes every revealed value immediately. After that prefix it processes only sufficiently short outcomes and sends the rest to a shortest-processing-time tail. The forced prefix guarantees that some jobs finish before too many tests accumulate; the suffix threshold protects the schedule from long revealed jobs. As u increases, the useful prefix shrinks continuously to zero. The algebraic branch is exactly the point where the worst binary mixture is tangent to the target ratio; after the prefix vanishes, balancing a capped tail against a threshold-one tail gives the square-root branch.

Adaptive-threshold policy. For large u , a fixed prefix is too rigid. The algorithm again tests jobs in a fixed order, but its immediate-processing threshold reacts to the observed history. It remembers only the number N of positive outcomes and the number D of those outcomes that were deferred. A normalized balance of the form

$$\frac{D - (1 + c)N}{\text{number of untested jobs}}$$

measures how much future delay has already been promised to the tail. Near a neutral balance the algorithm can afford to process moderately long jobs immediately. Every positive outcome drives the balance downward, while a deferral refunds one unit because future tests will also delay that tail job. When the normalized balance becomes sufficiently negative, the threshold falls toward one. All deferred jobs are sorted only after the tests finish.

The analysis first fixes the symbolic immediate/deferred path. Convexity then reduces every ordinary revealed value to one of four endpoints: a true zero, a positive value tending to zero, the immediate side of the live threshold, or its deferred side. A nonnegative bank potential pays the excess pair cost at each endpoint. The logarithmic threshold is forced by making the immediate and deferred boundary equations balance simultaneously; the obligatory constant is forced by calibrating the bank’s initial value.

Finite u creates exactly one new endpoint: a capped value $p = u$, for which the optimum can use raw execution and avoid the test. On the middle bridge this endpoint can temporarily overdraw the ordinary bank, so we add a reserve whose balance depends only on the observed cap fraction. At the end of the bridge the reserve becomes zero. With $r = R_\star - 1$, the worst residual charge of the cap endpoint is at most $2 - r(u - 1)$. The right-hand side is nonpositive precisely when $u \geq 1 + 2/r = z_\star$. Thus beyond $z_\star$ the obligatory four-endpoint bank already pays for the fifth endpoint. The plateau in Equation (7) is a proof-level fact, not merely a numerical coincidence.

1.5 Why the infinite endpoint needs a uniform proof

For each fixed processing-time vector, the finite- u offline lengths $q_u(p)$ eventually equal $1 + p$. Nevertheless, a competitive-ratio proof cannot blindly send u to infinity: the worst instance may choose $p \approx u$, the algorithm may depend on u , and the hidden constant in an $O_u(n)$ estimate may diverge. We avoid every such interchange. The lower bound uses one bounded harmonic core, which embeds in all sufficiently large finite caps. The

high- u upper bound uses one u -independent adaptive algorithm and one absolute linear remainder. Consequently the endpoint $u = \infty$ and the finite plateau are certified by the same witnesses.

The linear remainder is lower order because $\text{OPT} \geq n(n+1)/2$ at $u = \infty$ and, for fixed u , the analogous finite bound is also quadratic. It implies an endpoint coefficient $R_\star + \varepsilon$ with a fixed additive constant for every $\varepsilon > 0$. We do *not* prove the stronger statement $\text{ALG} \leq R_\star \text{OPT} + B$ with one fixed B , nor the unrelaxed inequality $\text{ALG} \leq R_\star \text{OPT}$ on every finite instance.

1.6 How to read the proofs

After the literature review and common model, Section 4 begins with the cap-free endpoint. Its lower subsection gives the harmonic adversary and a reusable scaled-core lemma. Its upper subsection formally states the adaptive algorithm and proves the ordinary four-endpoint bank. These are matching and may still be read independently.

Section 5 then treats finite caps. The binary stopping game and the two simple policies prove the first four branches. The capped-prefix lower construction reuses the harmonic core, while the middle/high upper bound reuses the ordinary bank and analyzes only the new cap endpoint and its reserve. At $u = z_\star$ both cap corrections vanish. Appendices A and B contain an exact rational checkpoint for the cap-free lower construction and an independent cap-bubbling certificate.

2 Related work

Scheduling with testing belongs to the broader framework of optimization with explorable uncertainty, where uncertain data can be queried at a cost; the query model goes back at least to Kahan [Kah91]. In the stochastic scheduling model of Levi, Magnanti, and Shaposhnik, testing diagnoses random job attributes and the scheduler trades the information gained against the delay caused by diagnosis [LMS19]. Their later work treats heterogeneous distributions and job priorities [LMS24]. Those policies exploit distributional information. The present paper instead uses worst-case processing times and competitive analysis.

Dürr, Erlebach, Megow, and Meißner introduced the adversarial model now usually called scheduling with testing [DEMM20]. A job has a known upper bound and may either be executed untested for that bound or tested for one unit to reveal a possibly smaller processing time. For total completion time on one machine they proved a deterministic lower bound of $1.8546\dots$ and an upper bound of 2. When all upper bounds are the same, they retained the lower bound and obtained the asymptotic upper bound $1.9338\dots$; on the further restriction $p \in \{0, u\}$, their UTE algorithm achieves $1.8668\dots$. Our finite-cap theorem closes the common-upper gap for arbitrary $p \in [0, u]$, determines the dependence on the fixed parameter u , and shows that the UTE constant is exactly the maximum of the resulting curve.

A related binary-information model is the processing-time-oracle problem of Dufossé et al. [DDN⁺22]. There, every job has one of two known possible processing times and an untested execution lasts for the true processing time; optimal adaptive and nonadaptive two-phase policies can be computed. This differs from our raw mode, whose duration is the declared upper bound regardless of the hidden value. Other variants alter a different part of the model. Nonuniform test times were introduced by Albers and Eckl [AE21]; the competitive guarantees for that setting were subsequently sharpened by Liu et al. [LLWZ23]

and, most recently, by Krekelberg et al. [KLL⁺26]. Damerius et al. [DKL⁺23] place job-dependent testing costs under an external budget, rather than charging test time directly to every unfinished job. Multiprocessor extensions for total completion time were studied by Gong, Chen, and Hayashi [GCH24]. Buld and Schulz [BS26] recently obtained the first constant guarantees for weighted total completion time in the adversarial optional-testing model. These variants do not specialize to the common-upper single-machine game solved here, but they delimit which assumptions make an exact curve possible.

Dogeas, Erlebach, and Liang introduced obligatory-test scheduling for total completion time [DEL24]. For arbitrary test times, their 1-SORT algorithm is 1.861-competitive. For uniform unit tests, their fixed-threshold algorithm SIDLE processes a sufficiently short revealed job immediately and sends longer jobs to a shortest-processing-time tail. They proved the upper bound $1.58451\dots$, exhibited matching instances for their analysis of SIDLE, and gave a $\sqrt{2}$ deterministic lower bound. Liang and Liang then developed a completion-threshold framework on identical machines; their dyadic construction improves the single-machine lower bound to $3/2$ [LL26]. Our harmonic multilevel construction raises that lower bound to R_* , while the adaptive-threshold algorithm lowers the upper bound to the same value. Unlike SIDLE, its threshold depends on both the number of positive outcomes observed and the number deferred.

Our pair representation is related to the delay accounting of Dogeas et al. [DEL24], but the certificate is different. Once the immediate/deferred decisions are fixed, coordinatewise convexity reduces the obligatory upper bound to four boundary outcomes. A state-dependent bank then certifies those outcomes. The distinction between 0 and 0^+ forces an extra state variable; in the finite-cap model, the endpoint $p = u$ creates a fifth outcome because the offline optimum may skip the test. The reserve in Section 5.6 pays precisely for this cap endpoint.

For context, if processing times are revealed only when jobs finish and no tests are available, one obtains the classical nonclairvoyant model of Motwani, Phillips, and Torng [MPT94]. Lindermayr et al. [LSSS26] instead consider online arrivals, preemption, and total flow time when a job’s operations are revealed successively. For their uniform two-operation specialization, operation-level SRPT is 2-competitive and this is tight. That result is not directly comparable with ours because release dates, preemption, and the flow-time benchmark all differ. Finally, the offline shortest-processing-time rule used below is the unweighted special case of Smith’s rule [Smi56].

3 Unified model and preliminaries

There are n jobs, all available at time zero, and one machine. Operations are nonpreemptive, and at most one operation can run at a time. A unit test of job j reveals its processing time $p_j \geq 0$.

The family is indexed by $u \in (0, \infty]$. For finite u , all jobs share this known upper limit, $p_j \in [0, u]$, and either of two modes completes a job: execute it untested for time u , or test it for one unit and subsequently process it for time p_j . An untested execution reveals no value and needs no later test. At the endpoint $u = \infty$, processing times are arbitrary finite nonnegative numbers and raw execution is unavailable. Thus every job consists of its test and a later, not necessarily immediate, processing operation of length p_j . This is exactly obligatory testing.

The completion time C_j is the completion time of the processing operation; in tested mode, when $p_j = 0$, this is the instant immediately after the test. The objective is $\sum_j C_j$. The number n and the parameter u are known to the online algorithm.

We write $\text{ALG}_{\mathcal{A}}(I)$ for the objective value of an online algorithm $\mathcal{A}$ on an instance I , and $\text{OPT}(I)$ for the value of an offline optimum that knows all processing times. We omit subscripts and arguments when they are clear.

3.1 One offline optimum for the whole family

Define the clairvoyant effective length by

$$q_u(p) := \begin{cases} \min\{u, 1 + p\}, & u < \infty, \\ 1 + p, & u = \infty. \end{cases} \quad (8)$$

Lemma 3.1 (Unified offline order). *Let $q_i = q_u(p_i)$ and let $q_{(1)} \leq \dots \leq q_{(n)}$. For every $u \in (0, \infty]$, an offline optimum completes the jobs in nondecreasing order of q_i and has value*

$$\text{OPT}_u = \sum_{k=1}^n \sum_{i=1}^k q_{(i)}, \quad (9)$$

$$= \sum_i q_i + \sum_{i < j} \min\{q_i, q_j\}. \quad (10)$$

For finite u , it uses raw mode when $u \leq 1 + p_j$ and tested mode otherwise. At $u = \infty$, it tests every job and processes the jobs in nondecreasing p_j order.

Proof. Let $C_{[1]} \leq \dots \leq C_{[n]}$ be the completion times of any feasible schedule. Completing job j consumes at least q_j units: for finite u this is the shorter of its two modes, and for $u = \infty$ it is its compulsory test-process length. By time $C_{[k]}$ the machine has completed some k jobs, so

$$C_{[k]} \geq \sum_{i=1}^k q_{(i)}.$$

The schedule that takes the shorter available mode for every job and executes the resulting blocks in nondecreasing q_i order attains every inequality. Summing and expanding the prefix sums gives both formulas. $\square$

At the cap-free endpoint, the pair formula specializes to

$$\text{OPT}_{\infty} = \sum_{k=1}^n \left(k + \sum_{i=1}^k p_{(i)} \right), \quad (11)$$

$$= \sum_i (1 + p_i) + \sum_{i < j} (1 + \min\{p_i, p_j\}). \quad (12)$$

In particular,

$$\text{OPT}_{\infty} \geq \frac{n(n+1)}{2}. \quad (13)$$

For finite u , $q_j \geq \min\{u, 1\}$ gives

$$\text{OPT}_u \geq \min\{u, 1\} \frac{n(n+1)}{2}. \quad (14)$$

3.2 Competitive-ratio conventions

The ordinary competitive ratio of $\mathcal{A}$ is

$$\text{CR}(\mathcal{A}) = \sup_I \frac{\text{ALG}_{\mathcal{A}}(I)}{\text{OPT}(I)}.$$

For the result of this paper, the relevant size-asymptotic ratio is

$$\text{CR}_{\infty}(\mathcal{A}) = \limsup_{n \rightarrow \infty} \sup_{|I|=n} \frac{\text{ALG}_{\mathcal{A}}(I)}{\text{OPT}(I)}. \quad (15)$$

The additive-constant convention calls a coefficient c admissible for $\mathcal{A}$ if there exists $B < \infty$ such that

$$\text{ALG}_{\mathcal{A}}(I) \leq c \text{OPT}(I) + B \quad \text{for every } I.$$

We write

$$\text{CR}_{\text{add}}(\mathcal{A}) = \inf\{c : c \text{ is admissible for } \mathcal{A}\}. \quad (16)$$

The optimal deterministic values in the two conventions are obtained by taking the infimum of $\text{CR}_{\infty}(\mathcal{A})$ and $\text{CR}_{\text{add}}(\mathcal{A})$, respectively, over deterministic algorithms. For any fixed u , an estimate $\text{ALG} \leq c \text{OPT} + O_u(n)$ fixes Equation (15) at most c and, by Equations (13) and (14), makes every coefficient $c + \varepsilon$ admissible. It need not make the endpoint coefficient c admissible.

Remark 3.2 (No hidden interchange of limits). The notation $u = \infty$ identifies one model exactly. We never infer an endpoint upper bound merely by taking $u \rightarrow \infty$ in an estimate whose algorithm or additive term may depend on u . The endpoint estimate is proved directly in Section 4.2; Section 5.6 then verifies that the same algorithm and certificate work uniformly for every finite $u \geq z_{\star}$.

3.3 Test-order adversaries

The lower bound will assign processing times according to the order in which tests finish. This is only a proof device.

Lemma 3.3 (Replay). *Fix a deterministic online algorithm. Suppose all untested jobs have the same known parameters, and an adversary assigns processing times deterministically from the transcript and test-completion order. The resulting interaction is reproduced by a fixed input instance.*

Proof. Run the interaction once and record the value revealed for every tested job label. Assign those values to the corresponding labels in advance. An induction over operation completions shows that the deterministic algorithm chooses the same tests, processing operations, and idle intervals on the fixed instance. If the algorithm fails to test or complete some job forever, its competitive ratio is already infinite; otherwise every relevant label is recorded. $\square$

4 The cap-free endpoint $u = \infty$

This section contains the two estimates needed for the cap-free endpoint theorem and isolates the harmonic and four-endpoint cores reused later. The first subsection applies to an arbitrary deterministic online algorithm; the second analyzes one explicit algorithm. Each restates the local notation it uses.

4.1 Harmonic core and matching lower bound

This subsection proves the lower bound in one self-contained calculation. Its notation is local to this subsection. The construction has many equally large groups of positive jobs, revealed from longest to shortest, followed by one large group of zero jobs. The only design choice is the gap between two consecutive positive processing times. Choosing these gaps harmonically will make all decisions of the online algorithm disappear from the leading term.

Theorem 4.1 (Lower bound). *For every deterministic online algorithm $\mathcal{B}$ and every $\varepsilon > 0$, there are arbitrarily large fixed instances I with unit tests and rational processing times such that*

$$\text{ALG}_{\mathcal{B}}(I) \geq (R_{\star} - \varepsilon) \text{OPT}(I).$$

The revelation experiment. Fix an integer K , a rational number $\xi > 0$, and a rational slack $\gamma \in (0, 1)$; one may keep $\gamma = 1/2$ throughout. For a large scaling integer H , the instance will contain H jobs of each of K positive types $0, 1, \dots, K-1$ and ξH zero jobs. We choose H to clear all denominators.

The adversary assigns processing times according to the order in which tests are started. The first H tested jobs receive processing time p_0 , the next H receive p_1 , and so on; after the K positive blocks, every remaining job receives processing time zero. Thus

$$p_0 > p_1 > \dots > p_{K-1} > 1,$$

so the algorithm sees the positive types from longest to shortest and learns about the zero jobs only at the end.

This adaptive description is only a convenient way to construct the input. After running the interaction against a fixed deterministic algorithm, assign to every job label the value revealed when that label was tested. Replaying the algorithm on this fixed assignment produces exactly the same transcript. If the algorithm never tests or completes some job, its cost is infinite and the lower bound is immediate; otherwise every job label is recorded. Hence it is enough to lower-bound the cost in the revelation experiment.

Why each phase can be relaxed to shortest processing time. Let T_ℓ be the start of the first test in positive block ℓ , and let T_K be the start of the first test in the zero block. Phase ℓ contains the completions in $(T_\ell, T_{\ell+1}]$; a completion exactly at T_ℓ belongs to the preceding phase. Each T_ℓ is the boundary between two nonpreemptive operations. Consequently, a previously tested job that is unfinished at T_ℓ has not started processing and still needs its entire processing time.

We repeatedly use the following elementary observation. Suppose jobs with remaining work $v_1, \dots, v_m$ complete after some time T . If their completion order is $q_1, \dots, q_m$, then the r th relative completion time is at least $\sum_{h \leq r} v_{q_h}$. Summing these prefix inequalities shows that their total relative completion time is minimized by arranging the v_i in nondecreasing order. This is merely a lower bound; the online algorithm need not actually use that order.

For normalized group masses $q_1, \dots, q_m$ and remaining lengths $v_1 \leq \dots \leq v_m$, this shortest-first bound has leading coefficient

$$\mathcal{S}((q_g, v_g)_{g=1}^m) = \sum_{g=1}^m \left(\frac{v_g q_g^2}{2} + q_g \sum_{h < g} v_h q_h \right). \quad (17)$$

More precisely, the corresponding contribution for groups of size $q_g H$ is $H^2 \mathcal{S} + O(H)$. The quadratic term is simply the familiar sum $v_g(1 + \dots + q_g H)$ inside one group; the cross term is the waiting caused by all shorter groups.

We shall ensure that

$$p_0 - p_{K-1} < 1. \quad (18)$$

In phase ℓ , an unfinished job of an older type $j < \ell$ therefore has remaining length p_j , whereas a type- ℓ job completed in its own phase needs both its test and processing, of total length $1 + p_\ell$. Their order in the shortest-first relaxation is

$$p_{\ell-1} < p_{\ell-2} < \dots < p_0 < 1 + p_\ell.$$

At T_K , an untested zero job has remaining length 1, so the relaxed tail starts with all zero jobs and then uses the positive types from shortest to longest. These are the only two places where Equation (18) is used.

The finite accounting identity. We now record the online algorithm's possible behavior. Normalize all job counts by H . Let

- $s_j \in [0, 1]$ be the mass of type- j jobs completed in their own phase;
- $d_{j,\ell} \geq 0$ be the mass of type- j jobs completed in the later phase $\ell > j$; and
-

$$X_{j,\ell} = s_j + \sum_{r=j+1}^{\ell-1} d_{j,r} \leq 1 \quad (\ell > j)$$

be the mass of type j completed before phase ℓ . Thus $X_{j,\ell+1}$ is the mass completed by the end of phase ℓ .

At time T_ℓ , the machine has already performed the tests of the ℓH jobs in earlier positive blocks and has processed every earlier job that has completed. Thus

$$\frac{T_\ell}{H} \geq \ell + \sum_{j < \ell} p_j X_{j,\ell}. \quad (19)$$

For each phase, charge its completions their common start time from Equation (19), and charge their relative completion times using Equation (17). Do the same at T_K for the remaining positive jobs and the ξH zero jobs. Expanding and collecting equal variables gives

$$\begin{aligned} \frac{\text{ALG}}{H^2} &\geq C_K + \sum_{j=0}^{K-1} \left(\frac{s_j^2}{2} - \theta_j s_j \right) \\ &\quad + \sum_{j < \ell} d_{j,\ell} \left[\ell - j - \theta_j - \sum_{j < k < \ell} (p_j - p_k) X_{k,\ell+1} \right] - O(1/H), \end{aligned} \quad (20)$$

where K, ξ, γ are fixed while $H \rightarrow \infty$, and

$$\theta_j = 1 - \xi(p_j - 1) - \sum_{k > j} (p_j - p_k - 1), \quad (21)$$

$$C_K = \frac{\xi^2}{2} + 2\xi K + K^2 + P_K, \quad P_K = \sum_{j=0}^{K-1} \left(j + \frac{1}{2} \right) p_j. \quad (22)$$

For completeness, here is how to audit Equation (20) without guessing its origin. In phase ℓ , the masses in the shortest-first relaxation are

$$d_{\ell-1,\ell}, d_{\ell-2,\ell}, \dots, d_{0,\ell}, s_\ell$$

with respective lengths

$$p_{\ell-1}, p_{\ell-2}, \dots, p_0, 1 + p_\ell.$$

Multiply their total mass by the right-hand side of Equation (19) and add the functional Equation (17). In the final phase use masses

$$\xi, 1 - X_{K-1,K}, \dots, 1 - X_{0,K}$$

and lengths $1, p_{K-1}, \dots, p_0$. The constant terms sum to C_K . The terms containing only s_j are $s_j^2/2 - \theta_j s_j$; the coefficient of each $d_{j,\ell}$ is precisely the bracket in Equation (20). Thus the displayed identity is just the phasewise shortest-first bound with the phase start times retained. No optimality assumption about the online schedule is hidden in it.

The meaning of the expression is useful. Completing an s_j fraction of a type in its own phase creates a quadratic completion cost but saves later waiting cost, producing $s_j^2/2 - \theta_j s_j$. A completion postponed to a later positive phase is represented by $d_{j,\ell}$ and its bracket. We next choose the processing times so that every such bracket is nonnegative and all θ_j are equal.

The harmonic choice and its telescoping cancellation. Set

$$p_{K-1} = 1 + \frac{\gamma}{\xi}, \quad p_i = p_{i+1} + \frac{1}{\xi + K - 1 - i} \quad (0 \leq i < K - 1). \quad (23)$$

In other words, measured upward from the shortest type, the consecutive gaps are

$$\frac{1}{\xi + 1}, \frac{1}{\xi + 2}, \dots, \frac{1}{\xi + K - 1}.$$

This is the key design choice. If $m = K - 1 - j$, then

$$p_j - 1 = \frac{\gamma}{\xi} + \sum_{r=1}^m \frac{1}{\xi + r},$$

$$\sum_{k>j} (p_j - p_k) = \sum_{r=1}^m \frac{r}{\xi + r}.$$

Adding the first line multiplied by ξ to the second line makes each denominator cancel:

$$\xi(p_j - 1) + \sum_{k>j} (p_j - p_k) = \gamma + \sum_{r=1}^m \frac{\xi + r}{\xi + r} = \gamma + m. \quad (24)$$

Substitution in Equation (21) gives the same value for every type,

$$\theta_j = 1 - \gamma. \quad (25)$$

This is why the increments in Equation (23) are harmonic: the factor ξ coming from the zero block and the multiplicity r coming from the r shorter positive types add up to the denominator $\xi + r$.

The delayed-completion terms are now harmless as well. Since $X_{k,\ell+1} \leq 1$, their brackets are at least

$$\begin{aligned} \ell - j - (1 - \gamma) - \sum_{j < k < \ell} (p_j - p_k) \\ = \gamma + \sum_{j < k < \ell} (1 - (p_j - p_k)) > 0, \end{aligned}$$

where the final inequality follows from Equation (18). We may therefore drop every term containing $d_{j,\ell}$. The remaining minimization is not a multivariable optimization problem: complete one square for each j ,

$$\frac{s_j^2}{2} - (1 - \gamma)s_j = \frac{1}{2}(s_j - (1 - \gamma))^2 - \frac{1}{2}(1 - \gamma)^2.$$

Because $1 - \gamma \in (0, 1)$, this minimum is feasible for $s_j \in [0, 1]$. Hence every online algorithm obeys

$$\liminf_{H \rightarrow \infty} \frac{\text{ALG}}{H^2} \geq Q_K := C_K - \frac{K}{2}(1 - \gamma)^2. \quad (26)$$

It remains to compute the offline benchmark. Knowing all processing times, the optimum runs the zero jobs first and the positive types from shortest to longest. Equivalently, the pairwise contribution of two jobs is $1 + \min\{p_i, p_j\}$. Its leading coefficient is

$$D_K = \frac{\xi^2}{2} + \xi K + \frac{K^2}{2} + P_K, \quad \lim_{H \rightarrow \infty} \frac{\text{OPT}}{H^2} = D_K. \quad (27)$$

The four terms respectively account for zero-zero pairs, zero-positive pairs, the unit-test part of positive-positive pairs, and the processing part of positive-positive pairs. Thus the finite- K lower-bound certificate is simply Q_K/D_K .

Passing to a smooth curve. Write

$$\alpha = \frac{K}{\xi}$$

for the ratio of positive mass to zero mass, and keep α fixed while $K \rightarrow \infty$. We restrict to $0 < \alpha < e - 1$. Then

$$p_0 - p_{K-1} = \sum_{r=1}^{K-1} \frac{1}{\xi + r} < \log\left(1 + \frac{K}{\xi}\right) = \log(1 + \alpha) < 1,$$

which verifies Equation (18).

The processing contribution P_K has the exact form

$$P_K = \frac{K^2}{2} \left(1 + \frac{\gamma}{\xi}\right) + \frac{1}{2} \sum_{h=1}^{K-1} \frac{(K-h)^2}{\xi + h}. \quad (28)$$

Indeed, the common baseline $1 + \gamma/\xi$ contributes the first term, and swapping the order of summation shows that the increment $1/(\xi + h)$ receives total weight $(K-h)^2/2$. After division by ξ^2 , the sum becomes a Riemann integral:

$$\frac{P_K}{\xi^2} \rightarrow I(\alpha) = \frac{\alpha^2}{2} + \frac{1}{2} \int_0^\alpha \frac{(\alpha - t)^2}{1 + t} dt$$

$$\begin{aligned}
&= \int_0^\alpha (\alpha - t)(1 + \log(1 + t)) dt \\
&= \frac{1}{2}(1 + \alpha)^2 \log(1 + \alpha) - \frac{\alpha}{2} - \frac{\alpha^2}{4}.
\end{aligned} \tag{29}$$

The second integral form follows either by integration by parts or by summing the limiting processing-time curve $1 + \log(1 + t)$ against the remaining positive mass $\alpha - t$. The subtraction in Equation (26) is only $O(K)$, whereas $C_K, D_K = \Theta(K^2)$. Since $C_K - D_K = \xi K + K^2/2$, we obtain

$$R(\alpha) := \lim_{K \rightarrow \infty} \frac{Q_K}{D_K} = 1 + \frac{\alpha + \alpha^2/2}{\frac{1}{2} + \frac{\alpha}{2} + \frac{\alpha^2}{4} + \frac{1}{2}(1 + \alpha)^2 \log(1 + \alpha)}. \tag{30}$$

This last one-variable maximization is where the constant in the theorem comes from. Put $z = (1 + \alpha)^2$. Then

$$R(z) = 1 + \frac{2(z - 1)}{1 + z + z \log z}. \tag{31}$$

The derivative of the fraction has the sign of

$$3 - z + \log z.$$

For $z > 1$ this expression is strictly decreasing, is positive at $z = 1$, and tends to $-\infty$. Hence it has one zero z_* , characterized by

$$z_* - 3 = \log z_*.$$

The derivative is positive before z_* and negative afterward, so this is the global maximum. At the root,

$$1 + z_* + z_* \log z_* = (z_* - 1)^2,$$

and therefore

$$R(z_*) = 1 + \frac{2}{z_* - 1} = R_*.$$

Moreover, $\sqrt{z_*} - 1 = 1.12255\dots < e - 1$, so the maximizer lies in the range where Equation (18) holds.

Proof of Theorem 4.1. Choose a rational α sufficiently close to $\alpha_* = \sqrt{z_*} - 1$, and choose any rational $\gamma \in (0, 1)$. For a sufficiently large K , set $\xi = K/\alpha$ and use Equation (23); then $Q_K/D_K > R_* - \varepsilon/2$. All parameters are rational. Let H tend to infinity through multiples of their common denominators. Equations (26) and (27) make the actual ratio exceed $R_* - \varepsilon$ for all sufficiently large H . Finally, the replay argument at the beginning of the proof converts the revelation experiment into a fixed instance for the given deterministic algorithm. $\square$

Lemma 4.2 (Scaled harmonic core). *Fix a total normalized mass $s > 0$ and a parameter $t \in [1, e)$. Against every deterministic algorithm for the cap-free endpoint, there are scales $M_\nu \rightarrow \infty$ and finite rational multilevel instances with $sM_\nu + o(M_\nu)$ jobs such that*

$$\frac{\text{OPT}}{M_\nu^2} \rightarrow O_0 = \frac{s^2}{4}(1 + t^{-2} + 2 \log t), \quad \liminf_{\nu \rightarrow \infty} \frac{\text{ALG}}{M_\nu^2} \geq A_0^{\text{lb}} := O_0 + \frac{s^2}{2}(1 - t^{-2}). \tag{32}$$

Their normalized total positive processing mass converges to

$$\frac{1}{M_\nu} \sum_{j: p_j > 0} p_j \rightarrow L = s \log t, \tag{33}$$

and their largest processing time converges to $1 + \log t < 2$. The adaptive revelation is converted by replay into a fixed instance.

Proof. The case $t = 1$ is obtained by continuity from instances with vanishing positive mass, so suppose $1 < t < e$. In the construction above set $\alpha = t - 1 = K/\xi$. Before rescaling, the total normalized mass is $K + \xi = \xi t$. Give every unit of normalized mass the common factor $s/(\xi t)$.

The phase accounting, harmonic cancellation, and completion of squares in Equations (20), (24) and (26) are homogeneous of degree two in the job masses. The Riemann-sum limit Equation (29) substituted into Equation (27) therefore gives

$$O_0 = \frac{s^2}{4}(1 + t^{-2} + 2 \log t).$$

Likewise $C_K - D_K = \xi K + K^2/2$, while the $O(K)$ square-completion loss in Equation (26) disappears after division by the quadratic scale. After the same rescaling this gives the guaranteed lower benchmark

$$A_0^{\text{lb}} - O_0 = \frac{s^2}{2}(1 - t^{-2}).$$

Finally, summing the limiting processing-time curve $1 + \log(1 + x)$ over the positive mass gives $L = s \log t$, and Equation (23) gives $p_0 \rightarrow 1 + \log t$. Rational approximation, clearing denominators, and the replay argument used in the theorem produce a diagonal sequence of arbitrarily large fixed rational instances with the stated limit and liminf bounds. $\square$

4.2 Upper bound: a self-contained proof

We prove the following statement from scratch in this subsection.

Theorem 4.3 (Upper bound). *Let $z > 1$ be the solution of $z - 3 = \log z$, and set*

$$R = 1 + \rho, \quad \rho = \frac{2}{z - 1} = 0.5705740966 \dots$$

There is a deterministic online algorithm and an absolute constant C such that every instance with n jobs satisfies

$$\text{ALG} \leq R \text{OPT} + Cn.$$

The proof has three ingredients. First, pairwise accounting gives an exact formula for the cost of a schedule. Second, a simple convexity argument says that it is enough to consider four boundary outcomes for every test. Third, we give a nonnegative *bank balance* W : whenever a boundary outcome creates positive excess cost, the balance drops by at least that amount, up to a constant rounding loss. Thus the proof below uses derivatives only as shorthand for the first-order change of this bank balance; no optimization background is needed.

The algorithm. For a parameter $c > 0$, the formal procedure $\text{ADAPTIVETHRESHOLD}(c)$ is given in algorithm 1. Immediately before a test, let

- x be the number of jobs not yet tested, including the job about to be tested;
- N be the number of earlier tests that revealed a strictly positive processing time; and
- D be the number of these N jobs that the algorithm deferred.

Define the normalized state

$$y = \frac{D - RN}{x}. \quad (34)$$

Because $D \leq N$ and $R > 1$, we always have $y \leq 0$. For later reuse define the parameterized threshold

$$a_c(y) = \begin{cases} 1, & y \leq -1, \\ 1 + \frac{1}{2} \log\left(\frac{c+2}{c-2y}\right), & -1 \leq y \leq 0. \end{cases} \quad (35)$$

The theorem uses $c = \rho$; throughout this subsection we abbreviate $A(y) = a_\rho(y)$ and $R = 1 + \rho$. The defining equation for $z = (\rho + 2)/\rho$ says

$$\log \frac{\rho + 2}{\rho} = \frac{2}{\rho} - 2. \quad (36)$$

Consequently, A is continuous at -1 , increases from $A(-1) = 1$ to $A(0) = 1/\rho = 1.7526207478\dots$, and

$$A'(y) = \frac{1}{\rho - 2y} \quad (-1 < y < 0). \quad (37)$$

Algorithm 1 ADAPTIVETHRESHOLD(J, c)

Input: jobs J and parameter $c > 0$.

- 1: Fix an arbitrary test order $j_1, \dots, j_n$.
 - 2: Set $N \leftarrow 0$, $D \leftarrow 0$, and $Q \leftarrow \emptyset$.
 - 3: **For** $i = 1, \dots, n$ **do**:
 - 4: Set $x \leftarrow n - i + 1$ and $y \leftarrow (D - (1 + c)N)/x$.
 - 5: Set $a \leftarrow a_c(y)$ according to Equation (35).
 - 6: Test j_i for one unit and observe p_{j_i} .
 - 7: **If** $p_{j_i} \leq a$, process j_i immediately.
 - 8: **Otherwise**, add j_i to Q and set $D \leftarrow D + 1$.
 - 9: **If** $p_{j_i} > 0$, set $N \leftarrow N + 1$.
 - 10: Process the jobs of Q in nondecreasing order of their revealed processing times.
-

For $c = \rho$, the state in the pseudocode is exactly (34). A genuine zero increments neither counter and completes at the end of its test. The procedure never uses raw execution; the same pseudocode will therefore apply to the large finite- u regimes.

The numerator $D - RN$ in (34) is best viewed as a balance, not as an estimate of a processing time. Every revealed positive job reduces the balance by R ; deferring it returns one unit, because every later unit test will also delay that deferred job. Dividing by x spreads this balance over the tests that remain. A state close to 0 can afford a threshold larger than one. Once the normalized balance falls below -1 , the proof can only support the conservative threshold one, so A becomes flat.

We will later derive (35), rather than asking the reader to guess why this particular logarithm appears.

Pairwise accounting. For two jobs $i < j$ in test order, charge to their unordered pair the duration of operations of either job performed while the other job is unfinished. If job i is processed immediately, this charge in the algorithm is $1 + p_i$. If i is deferred and j is

immediate, it is $2 + p_j$. If both are deferred, it is $2 + \min\{p_i, p_j\}$. Hence

$$\text{ALG} = \sum_i (1 + p_i) + \sum_{i < j} g_{ij}, \quad g_{ij} = \begin{cases} 1 + p_i, & i \text{ immediate,} \\ 2 + p_j, & i \text{ deferred, } j \text{ immediate,} \\ 2 + \min\{p_i, p_j\}, & i, j \text{ deferred.} \end{cases} \quad (38)$$

The offline optimum tests and immediately processes jobs in nondecreasing order of p_i , and therefore

$$\text{OPT} = \sum_i (1 + p_i) + \sum_{i < j} (1 + \min\{p_i, p_j\}). \quad (39)$$

For completeness, the latter claim follows because the k th completed job cannot finish before the machine has performed k tests and the processing of the k shortest jobs. The shortest-first schedule attains all these lower bounds simultaneously.

We shall bound the excess

$$F = \text{ALG} - R \text{OPT}.$$

Why four boundary outcomes suffice. Fix only the symbolic decisions made along a run: for each job, specify whether its value is zero, positive and immediate, or positive and deferred. This fixes all later counters and thresholds. The allowed values of a coordinate are then, respectively,

$$\{0\}, \quad (0, A(y)], \quad (A(y), \infty).$$

If all processing times except p_i are fixed inside such a symbolic run, (38)–(39) express F as an affine function of p_i plus negative multiples of $\min\{p_i, h\}$. Since $p \mapsto \min\{p, h\}$ is concave, F is convex in p_i . A convex function on $(0, A(y)]$ has its supremum at one of the two ends. On $(A(y), \infty)$, the eventual slope is $1 - R = -\rho < 0$; convexity then implies that every earlier slope is also negative, so the supremum is at the lower end.

It is therefore enough to consider the following four formal outcomes.

type	limiting value	algorithmic decision	counter update
Z	0	immediate	none
E	0^+	immediate	$N \leftarrow N + 1$
I	$A(y)$	immediate	$N \leftarrow N + 1$
Q	$A(y)^+$	deferred	$N \leftarrow N + 1, D \leftarrow D + 1$

Here 0^+ is not a new processing time. It means that a positive immediate value tends to zero *after its symbolic decision has been fixed*. Although its processing contribution vanishes, it still increments N and therefore changes all future thresholds. This is why 0 and 0^+ cannot be identified. Similarly, $A(y)^+$ approaches the threshold from the deferred side. Applying the one-coordinate argument successively shows that a uniform bound on all formal four-type words implies the same bound on all genuine instances. Because F is a finite sum of affine and minimum terms on a fixed symbolic path, the simultaneous one-sided limit at every formal vertex exists and is independent of the rates at which its open coordinates approach their endpoints.

One more elementary property will simplify every minimum in the pairwise formulas. Put $u = D - RN \leq 0$. After an outcome of type Z, E or I, or Q, respectively, the pair (u, x) changes as follows:

$$(u, x - 1), \quad (u - R, x - 1), \quad (u - R, x - 1), \quad (u - \rho, x - 1).$$

Whenever $x \geq 2$, the new quotient is at most u/x . Thus the states, and hence the thresholds, seen by successive tests never increase. When $x = 1$ there is no next test and therefore no next threshold to compare. The nonzero boundary values $A(y)$ consequently arrive in nonincreasing order. In particular, the minimum of the current nonzero boundary value and any earlier one is always the current value.

The proof state and the local excess. The algorithm needs only (x, N, D) . The analysis splits N into two kinds. Among earlier formal outcomes, let

- S be the number of *substantive* positive outcomes I, Q;
- e be the number of infinitesimal outcomes E; and
- d be the number of deferred outcomes Q.

Thus $N = S + e$ and $D = d$. Define

$$\eta = \frac{d - RS}{x}, \quad b = \frac{e}{x}, \quad y = \eta - Rb. \quad (40)$$

We have $d \leq S$, hence $\eta \leq 0$. The variable b records exactly the part of the algorithmic state that is invisible to limiting processing-cost minima: an infinitesimal value affects N , but its minimum with every other processing time tends to zero.

Consider the current boundary outcome at a pre-state (x, S, e, d) , and write $a = A(y)$. Pairwise accounting gives

$$F = -\rho \frac{n(n+1)}{2} + \sum_i \tilde{\ell}_i, \quad (41)$$

where the exact contribution of the current outcome is

$$\tilde{\ell}_Z = \tilde{\ell}_E = d, \quad (42)$$

$$\tilde{\ell}_I = d + a[x + d - RS - R], \quad (43)$$

$$\tilde{\ell}_Q = d + a[d - RS - \rho]. \quad (44)$$

Here is a quick derivation. The current unit test gives one extra unit of algorithmic pair cost for each earlier deferred job, producing d . Its own test and its pairs with all earlier jobs give a universal excess $-\rho(i+1)$ at step i ; summing this term produces the first term in (41). For an I outcome, the current processing operation contributes $a(x+d)$ to the algorithm and $Ra(S+1)$ to the scaled optimum. For a Q outcome, the corresponding quantities are $a(d+1)$ and $Ra(S+1)$. Monotonicity of the thresholds justifies using the current a in every earlier processing-time minimum.

For the potential calculation it is convenient to discard the two favorable diagonal terms $-Ra$ and $-\rho a$. Define the relaxed local rewards

$$\ell_Z = \ell_E = d, \quad (45)$$

$$\ell_I = d + a(x + d - RS) = d + ax(1 + \eta), \quad (46)$$

$$\ell_Q = d + a(d - RS) = d + ax\eta. \quad (47)$$

Then $\tilde{\ell}_c \leq \ell_c$ for all four types, and

$$F \leq -\rho \frac{n(n+1)}{2} + \sum_i \ell_i. \quad (48)$$

How the logarithmic threshold is found. Before giving the full bank balance, it is useful to see where the formula comes from. Ignore 0^+ outcomes for a moment, so $b = 0$, and look in the range $-1 \leq y \leq 0$. Suppose the nonlinear part of the balance has the form $x^2 f(y)$. Asking the immediate boundary I and the deferred boundary Q to be exactly paid by the first-order decrease of this balance gives

$$-2f + (y - R)f' + A(1 + y) = 0, \quad (49)$$

$$1 - 2f + (y - \rho)f' + Ay = 0. \quad (50)$$

Subtracting the equations gives $f' = A - 1$. Differentiating either equation then gives

$$1 + (2y - \rho)A'(y) = 0.$$

With the natural joining condition $A(-1) = 1$, this is exactly (35)–(37).

The universal negative term in (48) has leading part $-\rho n^2/2$. The initial balance will be $n^2 f(0)$, so exact cancellation requires $f(0) = \rho/2$. Since

$$f(y) = \int_{-1}^y (A(t) - 1) dt, \quad (51)$$

direct integration gives

$$f(0) = \frac{1}{2} - \frac{\rho}{4} \log\left(\frac{\rho + 2}{\rho}\right).$$

Thus $f(0) = \rho/2$ is equivalent to (36), or, after setting $z = (\rho + 2)/\rho$, to $z - 3 = \log z$. Thus both the threshold and the constant R are forced by two local balance equations and the initial accounting; they are not the output of an unexplained numerical optimization.

The bank balance. For $-1 \leq y \leq 0$, put

$$f(y) = \int_{-1}^y (A(t) - 1) dt, \quad h(y) = A(y)(1 + y). \quad (52)$$

The term xd in the balance below pays for the fact that each of the x remaining tests will delay each of the d already deferred jobs. The $x^2 f(y)$ term handles substantive outcomes. Finally, the terms involving b repair the discontinuity between a genuine zero and 0^+ .

Define

$$W(x, S, e, d) = xd + x^2 G(y, b), \quad (53)$$

using two formulas for G :

$$G(y, b) = \begin{cases} f(y) + bh(y) + \frac{R}{2}b^2, & y \geq -1, \\ g(\eta), & g(\eta) = \frac{(1 + \eta)_+^2}{2R}, \quad y \leq -1, \end{cases} \quad \eta = y + Rb. \quad (54)$$

The first formula is used while the threshold is changing. The second is used once the threshold has saturated at one. In that flat region the limiting processing-cost terms depend on the substantive balance η , not on the number of 0^+ outcomes separately, which explains why g contains no b .

This piecewise definition is smooth enough for a first-order argument. At $y = -1$, feasibility gives $0 \leq b \leq 1/R$ and $\eta = -1 + Rb$. Since

$$f(-1) = h(-1) = 0, \quad h'(-1) = 1,$$

the first formula in (54) has value $Rb^2/2$ and derivatives $G_y = b$, $G_b = Rb$. The second formula has exactly the same value and derivatives. This also checks the derivative in the raw x coordinate: at fixed (S, e, d) ,

$$\partial_x [x^2 G(y, b)] = x(2G - yG_y - bG_b),$$

which agrees on the two sides because G , G_y , and G_b agree. The remaining raw derivatives are linear combinations of G_y and G_b . The internal join of $(1 + \eta)_+^2$ at $\eta = -1$ also has a continuous first derivative. Hence W is continuously differentiable. Moreover, $W \geq 0$: in the first region $f \geq 0$, $h \geq 0$, and $b, d \geq 0$; in the second this is immediate from (54).

The choice $h = A(1 + y)$ also has a direct interpretation. At $b = 0$, it makes the first infinitesimal outcome 0^+ exactly neutral in the local balance. The quadratic correction $Rb^2/2$ then maintains the balance for all b and is precisely what makes the two regions meet smoothly.

Four local balance checks. We now verify that every formal outcome can be paid from the bank. In raw coordinate order (x, S, e, d) , the four unit-step directions are

$$v_Z = (-1, 0, 0, 0), \quad v_E = (-1, 0, 1, 0), \quad v_I = (-1, 1, 0, 0), \quad v_Q = (-1, 1, 0, 1). \quad (55)$$

For an endpoint type $q \in \{Z, E, I, Q\}$, define its first-order balance

$$J_q = \ell_q + \nabla W \cdot v_q. \quad (56)$$

The inequality $J_q \leq 0$ simply says: local reward plus first-order change in the bank balance is nonpositive.

We need only a few one-variable identities. On $-1 \leq y \leq 0$,

$$f' = A - 1, \quad f(-1) = 0, \quad f(0) = \frac{\rho}{2}, \quad (57)$$

$$-2f + (y - R)f' + A(1 + y) = 0, \quad (58)$$

$$1 - 2f + (y - \rho)f' + Ay = 0, \quad (59)$$

$$-2f + yf' = A(\rho - y) - R \leq 0, \quad (60)$$

$$h + (\rho - y)h' = AR + (\rho - y)(1 + y)A' \geq AR. \quad (61)$$

The first line follows from the definition and (36). For (58)–(59), differentiate the left-hand sides and use $A' = 1/(\rho - 2y)$; both derivatives vanish, and both expressions equal zero at $y = -1$. Equation (60) follows from (58). Its right-hand side is zero at -1 and decreases afterward, because

$$\frac{d}{dy} [A(y)(\rho - y)] = \frac{\rho - y}{\rho - 2y} - A(y) \leq 0.$$

Indeed, $y \leq 0$ gives $(\rho - y)/(\rho - 2y) \leq 1$, while $A(y) \geq 1$. Finally, (61) is obtained by differentiating $h = A(1 + y)$. Notice also that $h, h' \geq 0$.

In the changing-threshold region $y \geq -1$, direct differentiation of (53) first gives, for an arbitrary $G(y, b)$,

$$J_Z/x = -2G + yG_y + bG_b, \quad (62)$$

$$J_E/x = -2G + (y - R)G_y + (b + 1)G_b, \quad (63)$$

$$J_I/x = -2G + (y - R)G_y + bG_b + A(1 + y + Rb), \quad (64)$$

$$J_Q/x = 1 - 2G + (y - \rho)G_y + bG_b + A(y + Rb). \quad (65)$$

Substituting $G = f + bh + Rb^2/2$ and using (58)–(59) reduces the four expressions to

$$J_Z/x = (-2f + yf') + b(-h + yh'), \quad (66)$$

$$J_E/x = b[-h + (y - R)h' + R], \quad (67)$$

$$J_I/x = b[-h + (y - R)h' + AR], \quad (68)$$

$$J_Q/x = b[-h + (y - \rho)h' + AR]. \quad (69)$$

Now the signs are transparent. The Z line is nonpositive by (60), $h, h' \geq 0$, and $y \leq 0$. The Q line is nonpositive by (61). The I line is no larger because $y - R < y - \rho$ and $h' \geq 0$; the E line is no larger again because $A \geq 1$. Thus all four checks hold in the changing-threshold region.

In the flat region $y \leq -1$, we have $A = 1$ and $G = g(\eta)$. The four checks become

$$J_Z/x = J_E/x = -2g + \eta g', \quad (70)$$

$$J_I/x = -2g + (\eta - R)g' + 1 + \eta, \quad (71)$$

$$J_Q/x = 1 - 2g + (\eta - \rho)g' + \eta. \quad (72)$$

If $\eta \leq -1$, then $g = g' = 0$, and these expressions are 0, $1 + \eta$, and $1 + \eta$. If $-1 \leq \eta \leq 0$, then $g' = (1 + \eta)/R$, and they are, respectively,

$$-\frac{1 + \eta}{R}, \quad -\frac{1 + \eta}{R}, \quad 0.$$

They are therefore nonpositive in both cases. We have proved

$$J_Z, J_E, J_I, J_Q \leq 0 \quad (73)$$

at every feasible state.

From infinitesimal checks to unit steps. Equation (73) compares a local reward with the *derivative* of W , whereas one job changes the integer counters by one. This loses only a constant per job. More precisely, there is an absolute constant C_0 such that every admissible transition satisfies

$$W(s + v_q) - W(s) \leq \nabla W(s) \cdot v_q + C_0. \quad (74)$$

We include the bounded-curvature argument because it is what makes the $O(n)$ term uniform over all instances.

Suppose first that $x \geq 2$ and linearly interpolate the unit transition. The numerators of both y and η have the form $u - \kappa t$, where $u \leq 0$ and $\kappa \geq 0$, while the denominator is $x - t$. Their derivatives are therefore nonpositive, so the segment crosses each joining surface at most once. In the changing-threshold region, $y \geq -1$ implies

$$R(S + e) - d = \rho S + (S - d) + Re \leq x.$$

Consequently,

$$0 \leq \frac{S}{x} \leq \frac{1}{\rho}, \quad 0 \leq \frac{e}{x} \leq \frac{1}{R}, \quad 0 \leq \frac{d}{x} \leq \frac{1}{\rho}.$$

After normalizing by x , all reachable states in this region thus lie in a fixed compact set. The balance W is homogeneous of degree two, so its second derivatives in the raw

coordinates are homogeneous of degree zero and are uniformly bounded on this normalized set.

In the flat region the formula is even simpler:

$$W = xd + \frac{(x + d - RS)_+^2}{2R}. \quad (75)$$

Its gradient is globally Lipschitz. Because the two regional formulas have the same first derivative at their interface, an interpolated unit step has a uniformly bounded second derivative except at finitely many harmless joining points. More explicitly, let $\phi(t) = W(s + tv_q)$. The preceding bounds and the C^1 gluing make ϕ' Lipschitz with one uniform constant L along the whole segment, so

$$|\phi(1) - \phi(0) - \phi'(0)| \leq \int_0^1 |\phi'(t) - \phi'(0)| dt \leq \int_0^1 Lt dt = \frac{L}{2}.$$

This proves (74) for $x \geq 2$. When $x = 1$, the changing-threshold pre-states lie in the same compact normalized set, while (75) applies globally in the flat region. At $x = 0$ we define $W = 0$; this agrees with (75), since terminal feasibility gives $d \leq S$ and hence $d - RS \leq 0$.

Combining (56)–(74) gives the actual one-step bank inequality

$$\ell_i + W(s_{i+1}) - W(s_i) \leq C_0. \quad (76)$$

Telescoping the account. Initially $S = e = d = 0$, $x = n$, and $y = 0$. Therefore

$$W(s_0) = n^2 f(0) = \frac{\rho}{2} n^2. \quad (77)$$

The terminal balance is zero. Summing (76) over all n jobs yields

$$\sum_i \ell_i \leq \frac{\rho}{2} n^2 + C_0 n.$$

Substitution into (48) gives

$$\text{ALG} - R \text{OPT} = F \leq -\rho \frac{n(n+1)}{2} + \frac{\rho}{2} n^2 + C_0 n = \left(C_0 - \frac{\rho}{2}\right) n.$$

The constant is independent of the four-type word. Taking $C = \max\{0, C_0 - \rho/2\}$, the boundary reduction therefore extends the same estimate to every vector of nonnegative processing times, proving Theorem 4.3.

Finally, $\text{OPT} \geq n(n+1)/2$, so the additive $O(n)$ term is lower order and

$$\frac{\text{ALG}}{\text{OPT}} \leq R + O(1/n).$$

This proves the desired size-asymptotic upper bound. Notice exactly where the remaining finite-instance loss enters: it is only the uniform Taylor remainder in (74) (together with the deliberately discarded favorable diagonal terms), not any unproved limiting step in the endpoint reduction. In particular, this argument proves a linear additive term; it does not prove one fixed additive constant at the endpoint R .

Corollary 4.4 (Exact asymptotic values). *The optimal deterministic size-asymptotic value and the optimal deterministic additive-constant infimum both equal $R_\star$. Moreover, for every $\varepsilon > 0$, the Adaptive-Threshold algorithm satisfies*

$$\text{ALG}(I) \leq (R_\star + \varepsilon) \text{OPT}(I) + B_\varepsilon \quad (78)$$

for all instances, with B_ε independent of I .

Proof. The upper bound and Equation (13) imply

$$\frac{\text{ALG}}{\text{OPT}} \leq R_\star + \frac{2C}{n+1},$$

so the size-asymptotic ratio is at most $R_\star$. The lower bound is Theorem 4.1. For the additive statement, Young's inequality and Equation (13) give

$$Cn \leq \frac{\varepsilon}{2}n^2 + \frac{C^2}{2\varepsilon} \leq \varepsilon \text{OPT} + \frac{C^2}{2\varepsilon}.$$

Thus one can take $B_\varepsilon = C^2/(2\varepsilon)$, proving that every coefficient above $R_\star$ is admissible for this algorithm. Conversely, fix $c < R_\star$. By Theorem 4.1, every deterministic algorithm has a sequence of instances with $\text{OPT} \rightarrow \infty$ and $\text{ALG} / \text{OPT} > c + \delta$ for some $\delta > 0$. The inequality $\text{ALG} \leq c \text{OPT} + B$ would then force $\delta \leq B / \text{OPT} \rightarrow 0$, a contradiction. This proves the additive-constant claim. $\square$

5 Finite caps: the exact curve

We now move from the cap-free endpoint to finite u . Raw execution becomes available, all jobs have the same known upper limit, and a test reveals $p_j \in [0, u]$. Thus the offline effective length of job j is

$$q_j = \min\{u, 1 + p_j\}, \quad (79)$$

and the offline optimum schedules the q_j in nondecreasing order. We write $V(u)$ for the optimal deterministic size-asymptotic ratio at this fixed common upper limit.

The answer has six branches. We first define their transition points. Put

$$\varphi = \frac{1 + \sqrt{5}}{2}, \quad u_\diamond = \frac{1 + \sqrt{3 + 2\sqrt{5}}}{2} = 1.866760399173862\dots \quad (80)$$

Let $s_0 > 1$ be the root of

$$s_0^3 = (s_0 + 1)^2, \quad u_0 = 1 + s_0 = 3.147899035704787\dots \quad (81)$$

For $s = u - 1$, put

$$T_s(z) = sz^2 + (s^3 + s^2 + 1)z - (s^2 + s + 1)(s + 2), \quad (82)$$

and let $\rho_I(u)$ be the positive root of

$$s\rho^2 + (s^3 + s^2 + 1)\rho - (s^2 + s + 1)(s + 2) = 0. \quad (83)$$

Denote the left-hand side by $T_s(\rho)$. Equivalently,

$$\rho_I(u) = \frac{-1 - u + 2u^2 - u^3 + \sqrt{u^6 - 4u^5 + 10u^4 - 6u^3 - 3u^2 + 6u - 3}}{2(u - 1)}. \quad (84)$$

At the other end, let $z_\star > 1$ solve $z_\star - 3 = \log z_\star$ and define

$$r = \frac{2}{z_\star - 1}, \quad R_{\text{obl}} = 1 + r = 1.570574096649395\dots, \quad z_\star = 4.505241495792883\dots \quad (85)$$

For $\varphi + 2 \leq u \leq z_\star$, define $C_{\text{mix}}(u) = 1 + c$ by the unique solution (c, m) of

$$\text{artanh } m - m = 1 - \frac{1}{c} + \frac{1}{2} \log\left(1 + \frac{2}{c}\right), \quad u = 1 + \frac{2}{c} - m, \quad (86)$$

with $r \leq c \leq 1/\varphi$ and $0 \leq m \leq 1/\varphi$.

Theorem 5.1 (Exact common-upper curve). *For every fixed $u > 0$, the optimal deterministic size-asymptotic ratio in the optional-testing model is*

$$V(u) = \begin{cases} 1, & 0 < u \leq 1, \\ u, & 1 \leq u \leq u_\diamond, \\ \rho_I(u), & u_\diamond \leq u \leq u_0, \\ 1 + \frac{1}{\sqrt{u-1}}, & u_0 \leq u \leq \varphi + 2, \\ C_{\text{mix}}(u), & \varphi + 2 \leq u \leq z_*, \\ R_{\text{obl}}, & u \geq z_*. \end{cases} \quad (87)$$

For every branch there is an explicit deterministic algorithm satisfying

$$\text{ALG} \leq V(u) \text{OPT} + O_u(n).$$

On finite $u \geq z_*$ the remainder is one absolute $O(n)$ term, and Theorem 1.1 supplies the same algorithm at $u = \infty$. Conversely, for every deterministic algorithm $\mathcal{A}$ and every $\epsilon > 0$, there are arbitrarily large fixed instances I such that

$$\text{ALG}_{\mathcal{A}}(I) \geq (V(u) - \epsilon) \text{OPT}(I).$$

The upper-bound witness can be read from the following policy map. The algorithms themselves are specified formally in algorithms 1 to 3.

range	policy	parameter choice
$0 < u \leq u_\diamond$	RAW	none
$u_\diamond < u \leq u_0$	FORCEDPREFIXUTE	$b = b(u)$ from (104)
$u_0 < u \leq \varphi + 2$	FORCEDPREFIXUTE	$b = 0$
$\varphi + 2 < u < z_*$	ADAPTIVETHRESHOLD	$c = c(u)$ from (86)
$z_* \leq u \leq \infty$	ADAPTIVETHRESHOLD	$c = r$

The global-maximum assertion follows directly from the displayed formulas. Only the algebraic branch needs a check. Implicit differentiation of $T_s(\rho_I(u)) = 0$, with $s = u - 1$, gives

$$\rho'_I(u) = -\frac{\rho^2 + (3s^2 + 2s)\rho - 3(s+1)^2}{2s\rho + s^3 + s^2 + 1}. \quad (88)$$

Using $T_s(\rho) = 0$, its numerator factors as

$$\frac{2s^3 + s^2 - 1}{s} \left(\rho - \frac{2s^3 + 3s^2 - 2}{2s^3 + s^2 - 1} \right).$$

On the relevant interval $s \geq u_\diamond - 1 > 2/3$, the prefactor is positive, while the fraction in parentheses is at most $5/3$ because $4s^3 - 4s^2 + 1 > 0$. The sign certificate (108) gives $\rho > 5/3$. Hence $\rho'_I(u) < 0$. The reciprocal branch and the mixed parametrization are also decreasing, so V attains its global maximum only at $u = u_\diamond$.

The lower bounds, including the fact that the middle construction is safe against raw execution, are proved next. We then give self-contained upper bounds through $u = \varphi + 2$. The parameterized upper bound for the last two branches is proved separately in Section 5.6.

5.1 The binary stopping game and its lower envelope

The first four branches already occur on binary instances $p_j \in \{0, u\}$. The stopping point must be hidden from the algorithm; announcing it in advance gives a strictly weaker lower bound.

Lemma 5.2 (Raw-safe hidden stopping). *Fix $u > 1$, a target $\rho \leq u$, and a stopping fraction $\alpha \in (0, 1)$. Against a deterministic algorithm, there is a legal binary adversary whose leading excess at its first crossing is*

$$(u - \rho)(1 - \sigma^2) + \sigma^2 f_{u,\rho}(y, b), \quad (89)$$

where $0 \leq b \leq y \leq 1$ and

$$0 \leq y - \alpha < \frac{1}{n - v}. \quad (90)$$

Moreover,

$$f_{u,\rho}(y, b) = 1 - \rho + 2y + (u - 1)(1 - \rho)y^2 + b^2 + 2b(u - 1 - uy). \quad (91)$$

Here $\sigma \in [0, 1]$ is the fraction of jobs not already executed raw. Consequently, if

$$\min_{0 \leq b \leq \alpha} f_{u,\rho}(\alpha, b) \geq 0,$$

then the adversary gives ratio at least $\rho - O_u(1/n)$.

Proof. Start in a long phase. A fresh job that is tested receives $p = u$, while a fresh job run raw is assigned the hidden value zero. If v jobs have been run raw and L tested-long jobs have been seen, stop at the first touch for which

$$L \geq \alpha(n - v). \quad (92)$$

All untouched jobs are then assigned value zero. Only for the purpose of lower-bounding the remaining cost, we grant the algorithm this future information and allow an optimal continuation; in the actual interaction, values are still revealed only by tests. If the line is never crossed, every job was run raw: if even one job had been tested, then after the last first touch the tested fraction would be one. On the all-zero hidden instance the ratio is $u \geq \rho$.

At a crossing, let e of the L long jobs already be complete, let $d = L - e$ be pending, and let $x = n - v - e - d$ jobs remain untouched. The following favorable canonical schedule lower-bounds the algorithm: run the v raw jobs; test and immediately process the e completed long jobs; perform the d noncompleting long tests; test the x zero jobs; and finally process the d pending long jobs. Indeed, if a noncompleting unit test is immediately before an available length- u completion while h jobs remain, the two orders contribute $h + hu$ and $hu + (h - 1)$, respectively; moving the completion left saves one. Raw completions are available from time zero, and identical long jobs may be relabelled so that the e completed ones are tested first. After the switch, a zero test-completion has length one and therefore precedes a pending operation of length u . These exchanges respect precedence.

The exact costs of this relaxed schedule and of OPT are

$$A_{\text{ex}} = u \left(vn - \frac{v(v-1)}{2} \right) + (1+u) \left(e(n-v) - \frac{e(e-1)}{2} \right)$$

$$+ d(n - v - e) + x(d + x) - \frac{x(x - 1)}{2} + u \frac{d(d + 1)}{2}, \quad (93)$$

$$O_{\text{ex}} = \frac{n(n + 1)}{2} + (u - 1) \frac{(e + d)(e + d + 1)}{2}. \quad (94)$$

Put $\nu = v/n$, $\delta = (v + e + d)/n$, and $\lambda = e/n$. Twice the quadratic coefficients in Equations (93) and (94) are

$$\begin{aligned} \mathcal{A} &= 1 + 2\delta(1 - u\nu) + (u - 1)\delta^2 + 2\nu(\nu + u - 2) \\ &\quad + \lambda^2 + 2\lambda(\nu + u - 1 - u\delta), \\ \mathcal{O} &= 1 + (u - 1)(\delta - \nu)^2. \end{aligned}$$

Now put $\sigma = 1 - \nu$, $y = (e + d)/(n - v)$, and $b = e/(n - v)$. Direct expansion gives exactly Equation (89) after normalizing by $n^2/2$. The first-crossing overshoot in Equation (92) is less than one: a test increases $e + d - \alpha(n - v)$ by one and a raw first touch increases it by α . This proves Equation (90), and in particular $\sigma(y - \alpha) < 1/n$. There is one small domain issue: the actual $b \leq y$ can exceed α . Put $\bar{b} = \min\{b, \alpha\}$. Then $0 \leq b - \bar{b} \leq y - \alpha$. Since $f_{u,\rho}$ is Lipschitz on the unit square,

$$\sigma^2 f_{u,\rho}(y, b) \geq \sigma^2 f_{u,\rho}(\alpha, \bar{b}) - O_u(1/n). \quad (95)$$

Thus the stated minimum over $0 \leq \bar{b} \leq \alpha$ suffices uniformly even when $n - v$ is small. The discarded diagonal terms are $O_u(n)$ in the unnormalized costs. Finally, the adaptive transcript defines a fixed instance by replay: every tested answer was fixed when revealed, and every raw job's zero value stayed hidden. This proves the lemma. $\square$

Minimizing Equation (91) over the algorithm's completion choice gives

$$b_*(y) = \max\{0, uy - (u - 1)\}. \quad (96)$$

The upper constraint $b \leq y$ never binds, because $uy - (u - 1) \leq y$ for $y \leq 1$. There are therefore two elementary quadratic branches. On the positive branch, put

$$A_u = u^2 - u + 1, \quad D_{u,\rho} = A_u + (u - 1)\rho. \quad (97)$$

Then

$$\min_b f_{u,\rho}(y, b) = 1 - \rho - (u - 1)^2 + 2A_u y - D_{u,\rho} y^2. \quad (98)$$

Its maximizer is $\alpha = A_u/D_{u,\rho}$, and tangency to zero is

$$(1 - \rho - (u - 1)^2)D_{u,\rho} + A_u^2 = 0. \quad (99)$$

Expanding Equation (99) gives Equation (83). On the zero- b branch,

$$\min_b f_{u,\rho}(y, b) = 1 - \rho + 2y + (u - 1)(1 - \rho)y^2, \quad (100)$$

and tangency gives

$$\rho = 1 + \frac{1}{\sqrt{u - 1}}, \quad \alpha = \frac{1}{\sqrt{u - 1}}. \quad (101)$$

The latter point belongs to the zero- b branch iff $(u - 1)^3 \geq u^2$, with equality at $u = u_0$.

For $\rho = u$, the maximum in Equation (98) is nonnegative exactly while

$$[u(u - 1)]^2 - u(u - 1) - 1 \leq 0, \quad (102)$$

whose positive equality is $u = u_\diamond$. We have proved the following binary lower envelope.

Proposition 5.3 (Binary lower curve). *Every deterministic algorithm has size-asymptotic ratio at least*

$$B(u) = \begin{cases} 1, & 0 < u \leq 1, \\ u, & 1 \leq u \leq u_\diamond, \\ \rho_I(u), & u_\diamond \leq u \leq u_0, \\ 1 + \frac{1}{\sqrt{u-1}}, & u \geq u_0. \end{cases} \quad (103)$$

The lower bound uses only $p_j \in \{0, u\}$.

Proof. For $1 < u \leq u_\diamond$, use $\rho = u$ and the maximizing point of Equation (98). For $u_\diamond \leq u \leq u_0$, use the positive root $\rho_I(u)$ and $\alpha = A_u/D_{u,\rho_I(u)}$; the tangency identity makes the maximum zero. The minimizing completion mass is positive in the interior and becomes zero at $u = u_0$. For $u \geq u_0$, use the zero- b tangency point from Equation (101). The branch conditions are precisely the two inequalities checked above. Finally, $V(u) \geq 1$ is trivial for $u \leq 1$. In each nontrivial case Lemma 5.2 supplies fixed binary instances by replay. $\square$

5.2 Matching upper bounds through the binary regime

The first policy is formalized in algorithm 2.

Algorithm 2 RAW(J, u)

Input: jobs J and a finite common upper limit u .

- 1: Fix an arbitrary order $j_1, \dots, j_n$.
 - 2: **For** $i = 1, \dots, n$, execute j_i untested for u time.
-

For $u \leq u_\diamond$, use RAW. It is optimal when $u \leq 1$; when $u > 1$, every effective length in Equation (79) is at least one, so the ratio is at most u .

For the next interval, let $\rho = \rho_I(u)$, $s = u - 1$, and define

$$K = s^2 + s + 1, \quad D = K + s\rho, \quad b = \frac{K - s^2\rho}{D}, \quad \chi = \frac{1 - b}{\rho}, \quad \alpha = \frac{K}{D}. \quad (104)$$

The defining equation for ρ gives

$$0 \leq b < \chi < \alpha < 1, \quad \chi = \frac{s(s+1)}{D}, \quad b = (s+1)\alpha - s; \quad (105)$$

For completeness, these inequalities follow without numerical root isolation. Since T_s is increasing on positive arguments,

$$T_s\left(\frac{K}{s^2}\right) = \frac{K}{s^3}(-s^3 + s^2 + 2s + 1) \geq 0 \quad (s \leq s_0), \quad (106)$$

and hence $b = (K - s^2\rho)/D \geq 0$, with equality only at $s = s_0$. Moreover,

$$\alpha - \chi = \frac{1}{D}, \quad \chi - b = \frac{s^2\rho - 1}{D} > 0. \quad (107)$$

For the last sign,

$$9T_s(5/3) = f(s) := 6s^3 - 12s^2 - 2s - 3 < 0. \quad (108)$$

throughout the present interval. Indeed, on $[4/5, 1]$ the polynomial is decreasing and $f(4/5) = -1151/125$; on $[1, s_0]$ it has one interior minimum, while $f(1) = -11$ and $f(s_0) < f(13/6) = -95/36$. Here $s_0 < 13/6$ follows by substituting $13/6$ in $s^3 - (s+1)^2$. Thus $\rho > 5/3$, while $s > 4/5$, and $s^2\rho > 16/15$. Finally $K < D$ gives $\alpha < 1$. In particular, $b = 0$ occurs only at $u = u_0$. The remaining two elementary branches use the single procedure in algorithm 3.

Algorithm 3 FORCEDPREFIXUTE(J, u, b)

- Input:** jobs J , finite $u > 1$, and $b \in [0, 1]$.
- 1: Fix a test order $j_1, \dots, j_n$; set $k \leftarrow \lfloor bn \rfloor$, $\tau \leftarrow \min\{1, u - 1\}$, and $Q \leftarrow \emptyset$.
 - 2: **For** $i = 1, \dots, n$ **do**:
 - 3: Test j_i for one unit and observe p_{j_i} .
 - 4: **If** $i \leq k$ or $p_{j_i} \leq \tau$, process j_i immediately.
 - 5: **Otherwise**, add j_i to Q .
 - 6: Process the jobs of Q in nondecreasing revealed processing time.
-

On $u_\diamond \leq u \leq u_0$, use FORCEDPREFIXUTE with b from Equation (104). Rounding bn changes the cost by $O_u(n)$.

Here is the complete binary calculation. Adjacent exchanges place as many long jobs as possible in the forced prefix and put all suffix-long jobs before suffix zeros. If x is the total long mass, then the prefix-long mass is $a = \min\{b, x\}$ and the deferred-long mass is $d = (x - b)_+$. After division by n^2 , the leading excess $G(x) = A - \rho O$ is

$$G(x) = \begin{cases} \frac{1-\rho}{2} + u \left(x - \frac{x^2}{2} \right) - \frac{\rho s}{2} x^2, & 0 \leq x \leq b, \\ \frac{1-\rho}{2} + u \left(b - \frac{b^2}{2} \right) + (1-b)(x-b) + \frac{s}{2}(x-b)^2 - \frac{\rho s}{2} x^2, & b \leq x \leq 1. \end{cases} \quad (109)$$

The second quadratic is strictly concave, has its stationary point at $x = \alpha$, and satisfies $G(\alpha) = 0$; after substituting Equation (104), the latter identity is exactly Equation (83). Since $b < \alpha$, its right derivative at b is positive. The left derivative satisfies $G'_-(b) = G'_+(b) + s(1-b) > 0$, and the first quadratic is concave, so it is increasing on $[0, b]$. Thus $G(x) \leq 0$ for every $x \in [0, 1]$, including the limiting case $b = 0$. This proves the binary upper bound without a hidden local-versus-global step.

For $u_\diamond \leq u \leq 2$, arbitrary values reduce to binary values by independent Bernoulli extremalization, in the following order. First raise *every* suffix-deferred value to u . This cannot lower ALG, while its offline effective length was already u . Only after this deterministic step, replace a prefix value p by u with probability p/u and by zero otherwise, and replace a suffix-immediate value $p \leq u - 1$ by u with probability p and by zero otherwise. Notice that the u realization of a suffix-immediate job becomes deferred, so status is not preserved. Nevertheless the online check is pairwise. For jobs i before j , the leading pair contribution is

$$g_{ij} = \begin{cases} 1 + p_i, & i \text{ is immediate,} \\ 2 + p_j, & i \text{ is deferred and } j \text{ is immediate,} \\ 2 + \min\{p_i, p_j\}, & i, j \text{ are deferred.} \end{cases} \quad (110)$$

A rounded prefix job stays immediate and preserves its expected value. If i is suffix-immediate, its rounded contribution is one when it becomes zero and at least two when it

becomes u , hence its expectation is at least $1 + p_i$. If i is deferred and j is suffix-immediate, then i has been raised to u and the expected pair contribution is $2 + up_j \geq 2 + p_j$; a deferred-deferred pair can only increase. Thus expected ALG does not decrease, up to its $O_u(n)$ diagonal terms.

The expected effective lengths are, respectively,

$$1 + \frac{u-1}{u}p \leq q(p), \quad 1 + (u-1)p \leq 1 + p, \quad (111)$$

and u for a deferred value. Since the minimum is jointly concave, $\mathbb{E} \min(q'_i, q'_j) \leq \min(q_i, q_j)$. Thus some binary realization has competitive excess at least that of the original instance.

It remains to justify arbitrary values for $2 \leq u \leq u_0$. The following quadratic lemma records the entire endpoint calculation.

Lemma 5.4 (UTE endpoint game). *For $2 \leq u \leq u_0$, UTE with the parameters Equation (104) satisfies*

$$\text{ALG} \leq \rho_I(u) \text{OPT} + O_u(n). \quad (112)$$

Proof. Fix the symbolic prefix/immediate/deferred decisions. Convexity in one coordinate reduces a prefix job to $p \in \{0, u\}$, a suffix-immediate job to $p \in \{0, 1\}$, and a suffix-deferred job to $p \in \{1^+, u\}$. For the last claim, on $1 < p < u-1$ the deferred-deferred term has coefficient $1 - \rho < 0$ multiplying a concave minimum; on $[u-1, u]$, OPT is flat and ALG is nondecreasing. Adjacent exchanges give the canonical endpoint order.

Normalize counts by n . Let a be prefix- u mass, d suffix- u mass, t suffix- 1^+ mass, and m suffix-immediate-one mass. Their constraints are

$$0 \leq a \leq b, \quad d, t, m \geq 0, \quad d + t + m \leq 1 - b. \quad (113)$$

Pairwise accounting using Equation (110) gives

$$A = \frac{1}{2} + (s+1) \left(a - \frac{a^2}{2} \right) + (1-b)(d+t+m) - \frac{m^2}{2} + \frac{s}{2}d^2, \quad (114)$$

$$O = \frac{1}{2} + \frac{(a+d+t+m)^2}{2} + \frac{s-1}{2}(a+d)^2. \quad (115)$$

Moving m to t does not change O and increases A , so $m = 0$. Put $x = a + d$ and write $F(x) = A - \rho O$ after optimizing the other variables. The derivative of the gap in t is $1 - b - \rho(x+t)$, so its maximizing value is $(\chi - x)_+$. At fixed x, t , moving mass from d to a leaves O unchanged and changes A at rate $s(1-x) + b - a \geq 0$. Hence

$$t = (\chi - x)_+, \quad a = \min\{b, x\}, \quad d = (x - b)_+. \quad (116)$$

For $x \geq \chi$, the gap $A - \rho O$ is the binary concave quadratic already discussed; it is tangent to zero at $x = \alpha$. For $x < \chi$, put

$$C_0 = -\frac{\rho-1}{2} + \frac{(1-b)^2}{2\rho}, \quad \gamma = s - \rho(s-1). \quad (117)$$

Substitution gives

$$A - \rho O = C_0 + a(s+b-sx) - \frac{a^2}{2} + \frac{\gamma x^2}{2}. \quad (118)$$

One has $\gamma > 0$ on $1 \leq s \leq s_0$. Indeed, $T_s(2) = s(s^2 - s + 1) > 0$, so for $s \leq 2$ the equation $T_s(\rho) = 0$ gives $\rho < 2$, and hence $\gamma > 2 - s \geq 0$. For $2 \leq s \leq s_0 < 13/6$,

$$16T_s(7/4) = 12s^3 - 20s^2 + s - 4 > 0; \quad (119)$$

the polynomial is 14 at $s = 2$ and has positive derivative thereafter. Thus $\rho < 7/4$ and $\gamma > 7/4 - 3s/4 > 1/8$.

On $0 \leq x \leq b$, one has $a = x$, and

$$A - \rho O = C_0 + (s + b)x - \frac{s + 1 + \rho(s - 1)}{2}x^2. \quad (120)$$

Its derivative is decreasing, and the derivative at b multiplied by D is

$$\rho(1 + s^2 + \rho s^2(s - 1)) > 0, \quad (121)$$

so the maximum is at b . On $b \leq x \leq \chi$, Equation (118) is convex and its maximum is at b or χ . The endpoint χ lies on the binary branch. Finally,

$$F(b) - F(\chi) = \frac{\chi - b}{2} [2sb - \gamma(b + \chi)], \quad (122)$$

and direct use of Equation (104) gives

$$D[\gamma(b + \chi) - 2sb] = -s + (1 + s - s^3)\rho + s^2(s - 1)\rho^2 > 0. \quad (123)$$

For a rational sign certificate, T_s is increasing on positive arguments and

$$9T_s(5/3) = 6s^3 - 12s^2 - 2s - 3 < 0, \quad (124)$$

so $\rho > 5/3$. If the right-hand side of Equation (123) is denoted by $q_s(\rho)$, then

$$3 \partial_\rho q_s(5/3) = 7s^3 - 10s^2 + 3s + 3 > 0. \quad (125)$$

The last polynomial is increasing for $s \geq 1$, so q_s is increasing for $\rho \geq 5/3$. Finally,

$$9q_s(5/3) = 10s^3 - 25s^2 + 6s + 15 > 0. \quad (126)$$

For the last sign, write $s = 3/2 + z$; the expression becomes $3/2 - 3z/2 + 20z^2 + 10z^3$. It is at least $3/2$ for $-1/2 \leq z \leq 0$, and for $z \geq 0$ it is larger than $3/2 - 9/320$. These inequalities prove that every endpoint gap is nonpositive. Diagonals and prefix rounding contribute $O_u(n)$. $\square$

At $u = u_0$, $b = 0$. The same zero-prefix algorithm remains exact through $u = \varphi + 2$.

Lemma 5.5 (Zero-prefix extension). *For $u_0 \leq u \leq \varphi + 2$, FORCEDPREFIXUTE($J, u, 0$) from algorithm 3 is $(1 + 1/\sqrt{u - 1})$ -competitive up to $O_u(n)$.*

Proof. The same coordinatewise reduction leaves capped-deferred mass d , boundary-deferred mass t , immediate-one mass m , and zeros. With $s = u - 1$, these masses satisfy $d, t, m \geq 0$ and $d + t + m \leq 1$, and the leading costs are

$$A = \frac{1}{2} + d + t + m - \frac{m^2}{2} + \frac{sd^2}{2}, \quad O = \frac{1}{2} + \frac{(d + t + m)^2}{2} + \frac{(s - 1)d^2}{2}. \quad (127)$$

Move m to t and put $y = d + t$. For fixed y , the ratio is fractional-linear in d^2 , so its maximum is on one of two faces:

$$\frac{1 + 2y}{1 + y^2} \leq \varphi \quad (d = 0), \quad 1 + \frac{2y}{1 + sy^2} \leq 1 + \frac{1}{\sqrt{s}} \quad (d = y). \quad (128)$$

The first one-variable function is maximized at $y = 1/\varphi$ and the second at $y = 1/\sqrt{s}$, which gives the two displayed constants. The second value dominates the first exactly when $u \leq \varphi + 2$. Binary instances attain it. $\square$

5.3 The unrestricted mixed lower bound

For $\varphi + 2 < u < z_*$, neither the binary family nor the pure cap-free family is tight. The correct adversary puts a capped block before a scaled harmonic core. Two lemmas ensure that optional raw execution and early processing of caps cannot escape it.

The cap-free construction needed below is exactly the scaled harmonic core from Lemma 4.2; no second phase-accounting proof is needed here. We now prepend capped jobs to that reusable core.

Lemma 5.6 (Cap deferral). *Prepend capped mass m with value $p = u$ to a later cap-free block of mass $s = 1 - m$ and total processing mass L . Suppose every later value satisfies $1 + p \leq u$ and*

$$s(u - 1) - L - m \geq 0. \quad (129)$$

Then processing any capped job before all later jobs are complete cannot decrease the leading online cost.

Proof. A cap completed after its test block but before a later job of value p has mutual cost $1 + u$; leaving it pending until after that job gives $2 + p$. The exchange saves $u - 1 - p \geq 0$. If cap mass q is instead completed inside the cap-test block, its pairs with all later jobs add $q[s(u - 1) - L]$. Its largest possible saving against not-yet-tested caps is $qm - q^2/2$. Relative to leaving every cap pending, the change is therefore at least

$$q[s(u - 1) - L - m] + \frac{q^2}{2} \geq 0. \quad (130)$$

The finite diagonal discrepancy is lower order. $\square$

After cap deferral, pairwise accounting gives the offline coefficient O and the guaranteed online lower benchmark A^{lb} :

$$A^{\text{lb}} = A_0^{\text{lb}} + m(1 - m) + m(1 + L) + \frac{u}{2}m^2, \quad (131)$$

$$O = O_0 + m(1 - m + L) + \frac{u}{2}m^2. \quad (132)$$

Indeed, a cap-later pair contributes $2 + p$ online and $1 + p$ offline, whereas a cap-cap pair contributes $2 + u$ online and u offline; collecting these pairs yields the two cross terms and the cap squares above. In particular, $A^{\text{lb}} - O = (A_0^{\text{lb}} - O_0) + m$.

Lemma 5.7 (Raw-safe quota and replay). *Fix $m \in (0, 1)$ and a later cap-free block satisfying the hypotheses of Lemma 5.6. Against every deterministic optional algorithm there are fixed instances whose asymptotic excess is at least the cap-free hybrid benchmark $A^{\text{lb}} - CO$ from Equations (131) and (132) for every target $C \leq u$.*

Proof. In the initial phase, a tested fresh job receives $p = u$, while a job run raw is assigned hidden value zero. If v raw jobs and ℓ tested caps have been touched, stop at the first crossing

$$\frac{\ell}{n - v} \geq m. \quad (133)$$

With $S = n - v$, the overshoot is less than one, so $\ell = mS + O(1)$. If no crossing occurs, no test can have occurred: after all first touches, any positive ℓ would make the ratio in Equation (133) equal one. Thus every job was run raw and the all-zero hidden instance has ratio $u \geq C$.

At a crossing, scale the prescribed later instance to the remaining $S - \ell$ first touches, absorbing the one-job cap overshoot and integer rounding into its zero group. A later test reveals the next value in this fixed type stream; a later raw action consumes the next value but keeps it hidden from the optional algorithm.

Move every initial raw-zero completion to the start. It has the same length u as cap processing, and crossing a unit test only lowers completion cost. With

$$Z_v = vn - \frac{v(v-1)}{2}, \quad (134)$$

this prefix contributes uZ_v online and Z_v offline. Delete it. The remaining S labels consist, up to one-job rounding, of capped fraction m and the prescribed later cap-free block.

If the optional algorithm later runs a fresh assigned value p raw, a physical cap-free simulation tests and immediately processes that job in time $1+p \leq u$, suppresses the answer from the virtual optional algorithm, and idles until its virtual u -block ends. Inductively every physical completion is no later than its virtual optional completion. Since the finite-cap and cap-free offline lengths are both $1+p$ on the later block,

$$\text{ALG}_{\text{opt}} - C \text{OPT}_{\text{opt}} \geq (u - C)Z_v + (\text{ALG}_{\text{hyb}} - C \text{OPT}_{\text{hyb}}) - O_u(n). \quad (135)$$

The first term is nonnegative. Finally, record every assigned value on its job label. Replaying this fixed vector produces the same transcript because the values learned only by the physical simulation were hidden from the virtual algorithm. This proves legality. $\square$

Apply Lemma 4.2 with $s = 1 - m$. Substitution into Equations (131) and (132) gives the guaranteed two-parameter lower ratio

$$\mathcal{R}_u(m, t) = 1 + \frac{m + \frac{(1-m)^2}{2}(1-t^{-2})}{\frac{(1-m)^2}{4}(1+t^{-2} + 2\log t) + m(1-m)(1+\log t) + \frac{u}{2}m^2}. \quad (136)$$

At an interior stationary point with value $1 + c$, differentiation and the zero-gap equation give

$$\log t = u - 2 - \frac{1}{c}, \quad t^2 = \frac{1-m}{1+m} \left(1 + \frac{2}{c}\right). \quad (137)$$

Eliminating t gives precisely Equation (86). Conversely, those equations make $\mathcal{R}_u(m, t) = 1 + c$.

The branch runs from

$$(u, m, t) = \left(\varphi + 2, \frac{1}{\varphi}, 1\right) \quad \text{to} \quad (u, m, t) = (z_*, 0, \sqrt{z_*}). \quad (138)$$

It satisfies $u \geq \varphi + 2$, $\log t < 1$, and $m \leq 1/\varphi$. Therefore

$$(1-m)(u-1-\log t) > (1-m)\varphi \geq \frac{1}{\varphi} \geq m, \quad (139)$$

so Equation (129) holds; also $1+p < 3 < u$ for every later positive value in a sufficiently fine finite construction. The functions

$$g(c) = 1 - \frac{1}{c} + \frac{1}{2} \log \left(1 + \frac{2}{c}\right), \quad h(m) = \text{artanh } m - m \quad (140)$$

have $g'(c) = 2/[c^2(c+2)] > 0$ and $h'(m) = m^2/(1-m^2) > 0$. Hence $m(c) = h^{-1}(g(c))$ is strictly increasing, while

$$u(c) = 1 + \frac{2}{c} - m(c) \quad (141)$$

is strictly decreasing. The identities $g(r) = h(0) = 0$ and $g(1/\varphi) = h(1/\varphi)$ give the two endpoints in Equation (138); consequently every $u \in [\varphi + 2, z_*]$ has exactly one parameter pair.

To pass to fixed finite instances, first approximate (m, t) rationally while retaining the strict cap inequalities, then choose a sufficiently fine harmonic family, and finally multiply all rational block sizes by a common integer H . For fixed harmonic depth, quota overshoot, diagonals, and rounding are $O_u(H)$ whereas the displayed costs are $\Theta(H^2)$. Send first H , then the harmonic depth, and finally the rational approximation to their limits. Together with Lemma 5.7, this proves

$$V(u) \geq C_{\text{mix}}(u) \quad (\varphi + 2 \leq u \leq z_*). \quad (142)$$

The construction above applies directly in the open interval. At $u = \varphi + 2$ and $u = z_*$ the same statement follows by approaching from the interior, so the endpoints are included.

5.4 The large-upper-limit lower bound by cap-free transfer

Lemma 5.8 (Bounded cap-free transfer). *Let $u \geq 1$, and let $V_\infty(P)$ denote the deterministic size-asymptotic value at the cap-free endpoint, restricted to $0 \leq p_j \leq P$. Then*

$$V(u) \geq V_\infty(u - 1). \quad (143)$$

In particular,

$$V(u) \geq R_{\text{obl}} \quad \left(u \geq 1 + \frac{1}{r} = 2.752620747896442 \dots \right). \quad (144)$$

Proof. Given a finite-cap algorithm, construct a cap-free algorithm on instances with $p_j \leq u - 1$ by running a virtual copy of the finite-cap algorithm. Copy ordinary tests and processing actions. When the virtual algorithm runs an untouched job raw for u time, physically test and immediately process it in $1 + p_j \leq u$ time, hide the answer from the virtual copy, and wait for the virtual clock. Every physical completion is no later than its virtual completion. Moreover, $\min\{u, 1 + p_j\} = 1 + p_j$, so finite-cap and cap-free optima coincide. Thus we obtain the following game-value comparison. For every finite-cap algorithm $\mathcal{A}$, the simulation produces a cap-free algorithm $\mathcal{B}$ and, pointwise on every instance with $p_j \leq u - 1$,

$$\text{ALG}_{\mathcal{A}} \geq \text{ALG}_{\mathcal{B}}, \quad \text{OPT}_u = \text{OPT}_\infty.$$

Consequently

$$\text{CR}_\infty(\mathcal{A}) \geq \text{CR}_\infty(\mathcal{B}; p \leq u - 1) \geq V_\infty(u - 1).$$

Taking the infimum over $\mathcal{A}$ proves Equation (143).

The harmonic cap-free family attaining R_{obl} has limiting largest processing time

$$1 + \log(1 + y_*) = 1 + \frac{1}{2} \log z_* = \frac{z_* - 1}{2} = \frac{1}{r}, \quad y_* = \sqrt{z_*} - 1. \quad (145)$$

Approximation from below also gives the endpoint, proving Equation (144). $\square$

5.5 Completion of the curve and consistency at the joins

The lower-bound side of Theorem 5.1 is now complete: Proposition 5.3 supplies the first four branches, Equation (142) the fifth, and Equation (144) the last. The raw algorithm, Lemmas 5.4 and 5.5, supplies the matching upper bounds through $\varphi + 2$. The remaining upper bounds are proved in Section 5.6.

For clarity, all five joins are exact:

$$\begin{aligned}\rho_I(u_\phi) &= u_\phi, \\ \rho_I(u_0) &= 1 + \frac{1}{\sqrt{s_0}} = 1.682327803828019\dots, \\ 1 + \frac{1}{\sqrt{(\varphi + 2) - 1}} &= \varphi = C_{\text{mix}}(\varphi + 2), \\ C_{\text{mix}}(z_*) &= 1 + r = R_{\text{obl}}.\end{aligned}$$

At $u = 1$, both initial branches equal one. Representative values of the new mixed branch are

u	$V(u)$	m	t
3.800000	1.598851588	0.539725635	1.138984621
4.000000	1.583283633	0.428863572	1.330517832
4.071679486	1.579204806	0.381330539	1.412236296
4.300000	1.571692026	0.198387083	1.734651379

Remark 5.9 (Asymptotic scope). All statements above concern the fixed- u size-asymptotic ratio. Below the plateau, the upper bounds have additive error $O_u(n)$. On $u \in [z_*, \infty]$ the same algorithm has one uniform $O(n)$ error. We do not claim a finite-instance endpoint inequality with one additive constant.

The next subsection, Section 5.6, proves the matching middle- and high- u upper bounds and thereby completes the proof of Theorem 5.1.

5.6 Matching upper bound for the common-upper model

We now prove the upper bounds on the two ranges on which capped outcomes matter. Besides completing the upper-bound side of the common-upper curve, the proof records explicitly why an outcome on the optional cap cannot be treated as an ordinary deferred outcome. Throughout this subsection,

$$C = 1 + c, \quad \varphi = \frac{1 + \sqrt{5}}{2},$$

and $z_* > 1$ is the solution of $z_* - 3 = \log z_*$. We also set

$$r = \frac{2}{z_* - 1}.$$

For $c \in [r, 1/\varphi]$, let $m = m(c) \in [0, 1/\varphi]$ be determined by

$$\operatorname{artanh} m - m = 1 - \frac{1}{c} + \frac{1}{2} \log \left(1 + \frac{2}{c} \right), \quad u = 1 + \frac{2}{c} - m. \quad (146)$$

The right-hand side of the first equation is strictly increasing in c , as is the left-hand side in m . Consequently $m(c)$ is increasing and $u(c)$ is decreasing. The endpoints are

$$(c, m, u) = \left(\frac{1}{\varphi}, \frac{1}{\varphi}, \varphi + 2 \right) \quad \text{and} \quad (c, m, u) = (r, 0, z_*).$$

Theorem 5.10. *For every $u \in [\varphi + 2, \infty]$, the formally specified algorithm `ADAPTIVETHRESHOLD` from algorithm 1 satisfies*

$$\text{ALG} \leq \begin{cases} (1 + c) \text{OPT} + O_u(n), & \varphi + 2 \leq u \leq z_*, \quad c \text{ given by (146)}, \\ (1 + r) \text{OPT} + O(n), & u \geq z_*. \end{cases}$$

The constant in the last $O(n)$ is absolute and uniform over the extended interval $u \in [z_, \infty]$. In particular these are upper bounds on the deterministic size-asymptotic competitive ratio.*

Proof.

At $u = \infty$ this is Theorem 4.3, with the same algorithm and the same four-endpoint bank. For the finite-cap calculations below, fix $u < \infty$.

The algorithm. Use `ADAPTIVETHRESHOLD`(J, c) from algorithm 1; on the high interval use $c = r$. For the analysis, immediately before a test let x be the number of jobs not yet tested, N the number of earlier strictly positive outcomes, and D the number of those outcomes deferred to the final tail. Thus

$$y = \frac{D - CN}{x} \leq 0 \tag{147}$$

and the threshold is

$$a_c(y) = \begin{cases} 1, & y \leq -1, \\ 1 + \frac{1}{2} \log\left(\frac{c+2}{c-2y}\right), & -1 \leq y \leq 0. \end{cases} \tag{148}$$

This is the same threshold as Equation (35). On the mixed interval $c \geq r$, and hence

$$a_c(y) \leq a_c(0) \leq a_r(0) = \frac{1}{r} < u - 1.$$

The same inequality holds on the high interval, where we use $c = r$. Thus every outcome in the capped region $[u - 1, u]$ is deferred.

The state y never increases. Indeed, the numerator in (147) changes by 0, $-C$, or $-c$ after a zero, a positive immediate outcome, or a positive deferred outcome, respectively, while x decreases. This monotonicity will allow us to allocate every pair cost when its relevant endpoint becomes known.

Exact pair objective and the five endpoints. Write

$$\bar{p}_i = \min\{p_i, u - 1\}.$$

The clairvoyant optional optimum is SPT on jobs of effective lengths $1 + \bar{p}_i = \min\{1 + p_i, u\}$. For two jobs tested in the order $i < j$, let

$$e_{ij} = \begin{cases} (p_i - p_j)_+, & i \text{ is processed immediately,} \\ 1 + (p_j - p_i)_+, & i \text{ is deferred and } j \text{ is immediate,} \\ 1, & i \text{ and } j \text{ are deferred.} \end{cases} \tag{149}$$

The first line applies whenever i is immediate, regardless of the action taken on j . Direct completion-time accounting gives

$$\text{ALG} = \sum_i (1 + p_i) + \sum_{i < j} (1 + \min\{p_i, p_j\} + e_{ij}), \tag{150}$$

$$\text{OPT} = \sum_i (1 + \bar{p}_i) + \sum_{i < j} (1 + \min\{\bar{p}_i, \bar{p}_j\}). \quad (151)$$

It is therefore natural to introduce

$$\Gamma_n = \sum_{i < j} (e_{ij} + \min\{p_i, p_j\} - C \min\{\bar{p}_i, \bar{p}_j\}). \quad (152)$$

Equations (150)–(152) yield the exact identity

$$\text{ALG} - C \text{OPT} = \Gamma_n - c \binom{n}{2} + \sum_i [(1 + p_i) - C(1 + \bar{p}_i)]. \quad (153)$$

The final sum is nonpositive. Below the cap its i -th summand is $-c(1 + p_i)$; on the cap it is at most $1 - cu \leq 0$. Thus it is enough to control Γ_n .

Fix a symbolic immediate/deferred path. Coordinatewise convexity in (152) reduces a worst outcome vector to the five formal endpoint types

type	outcome	action
Z	0	immediate
E	0^+	immediate
M	$a_c(y)$	immediate
Q	$a_c(y)^+$	deferred
U	u	deferred and capped.

(154)

For an immediate coordinate the two positive endpoints are 0^+ and the live threshold; the genuine zero is kept separate because it does not enter N . A deferred coordinate below $u - 1$ moves to the live threshold or to $u - 1$. On $[u - 1, u]$, the offline contribution is flat and the algorithmic contribution is nondecreasing, so the latter endpoint moves to u . The symbolic categories, and therefore all later threshold states, remain unchanged during each coordinate move. This is the only endpoint reduction used below.

A local pair allocation. Separate the earlier E -outcomes from the substantive outcomes M, Q, U . Let

$$P = \#\{\text{earlier } M, Q, U\}, \quad E = \#\{\text{earlier } E\}, \quad D = \#\{\text{earlier } Q, U\},$$

and let K be the number of earlier U -outcomes. Among the ordinary substantive outcomes, let I, d be the numbers of M, Q -outcomes and let A_I, A_D be the sums of their processing values. Thus $P = I + d + K$ and $D = d + K$.

At a current ordinary endpoint, with $a = a_c(y)$, allocate the relaxed local charges

$$\ell_Z = \ell_E = D, \quad \ell_M = D + a(x + D - CP), \quad \ell_Q = D + a(D - CP). \quad (155)$$

For a capped endpoint the exact charge is instead

$$\begin{aligned} \ell_U &= D + A_D + uK - C(A_I + A_D + (u - 1)K) \\ &= d - CA_I - cA_D + Kh_U, \quad h_U = 1 + u - C(u - 1). \end{aligned} \quad (156)$$

These formulae follow by checking the finitely many ordered endpoint pairs. In particular, an earlier immediate outcome already allocated its online processing delay when it appeared, so a current cap contributes only $-Cp$ against that outcome. Against an earlier ordinary

deferred outcome it contributes $p - Cp = -cp$, and against an earlier cap it contributes h_U . The same check shows

$$\Gamma_n \leq \sum_i \ell_i; \quad (157)$$

the difference is exactly the sum of the nonnegative own-processing terms allocated by the M -endpoints. Formula (156), rather than a substitution $p = u$ into ℓ_Q , is essential before caps have been put into a prefix.

Threshold monotonicity gives $A_I \geq aI$ and $A_D \geq ad$. Comparing (156) with the hypothetical Q -charge at the same state gives the exact identity

$$\begin{aligned} \ell_U - \ell_Q &= C(aI - A_I) + c(ad - A_D) + K[1 - c(u - 1 - a)] \\ &\leq K[1 - c(u - 1 - a)]. \end{aligned} \quad (158)$$

The reusable four-endpoint bank. The next potential is the parameterized form of the cap-free bank from Section 4.2. The states Z, E, M, Q are the same four ordinary endpoints; only their target coefficient changes from r to c . At $c = r$ the formulas below coincide term for term with the obligatory certificate. Keeping the parameter visible lets us add the finite-cap reserve without proving a second ordinary endpoint theorem. Define

$$\eta = \frac{D - CP}{x}, \quad b = \frac{E}{x}, \quad y = \eta - Cb. \quad (159)$$

For $-1 \leq y \leq 0$, put

$$f_c(y) = \frac{c+2}{4} + \frac{c-2y}{4} \left(\log \frac{c-2y}{c+2} - 1 \right), \quad (160)$$

$$h_c(y) = a_c(y)(1+y),$$

$$G_c(y, b) = f_c(y) + bh_c(y) + \frac{C}{2}b^2. \quad (161)$$

For $y \leq -1$, set instead

$$G_c(y, b) = g_c(\eta), \quad g_c(\eta) = \frac{(1+\eta)_+^2}{2C}. \quad (162)$$

The two definitions are C^1 across $y = -1$. Indeed, feasibility implies $0 \leq b \leq 1/C$ there, and both sides have value $Cb^2/2$ and derivatives

$$(G_y, G_b) = (b, Cb).$$

They are also nonnegative: $f_c(-1) = 0$, $f'_c = a_c - 1 \geq 0$, and $h_c, b, g_c \geq 0$.

Use the base potential

$$W_{\text{base}}(x, P, E, D) = xD + x^2 G_c(y, b). \quad (163)$$

In the raw state (x, P, E, D, K) , the endpoint directions are

	x	P	E	D	K
v_Z	-1	0	0	0	0
v_E	-1	0	1	0	0
v_M	-1	1	0	0	0
v_Q	-1	1	0	1	0
v_U	-1	1	0	1	1.

(164)

Let $J_T = \ell_T + \nabla W_{\text{base}} \cdot v_T$, where for the moment $T \in \{Z, E, M, Q\}$. In Region I, direct differentiation and the identities

$$\begin{aligned} -2f_c + (y - C)f'_c + a_c(1 + y) &= 0, \\ 1 - 2f_c + (y - c)f'_c + a_c y &= 0 \end{aligned} \quad (165)$$

give

$$\begin{aligned} \frac{J_Z}{x} &= (-2f_c + yf'_c) + b(-h_c + yh'_c), \\ \frac{J_E}{x} &= b[-h_c + (y - C)h'_c + C], \\ \frac{J_M}{x} &= b[-h_c + (y - C)h'_c + a_c C], \\ \frac{J_Q}{x} &= b[-h_c + (y - c)h'_c + a_c C]. \end{aligned} \quad (166)$$

All four quantities are nonpositive. For the first one,

$$-2f_c + yf'_c = a_c(c - y) - C \leq 0; \quad (167)$$

the right-hand side is zero at $y = -1$ and is nonincreasing because $a'_c(c - y) - a_c \leq 1 - a_c \leq 0$. Also $h_c, h'_c \geq 0$. For the remaining three drifts, observe that

$$h_c + (c - y)h'_c = a_c C + (c - y)(1 + y)a'_c \geq a_c C. \quad (168)$$

The Q -bracket in (166) is therefore nonpositive; the M -bracket is the Q -bracket minus h'_c , and the E -bracket is the M -bracket minus $C(a_c - 1)$.

In Region II, the same calculation gives

$$\begin{aligned} J_Z/x = J_E/x &= -2g_c + \eta g'_c, \\ J_M/x &= -2g_c + (\eta - C)g'_c + 1 + \eta, \\ J_Q/x &= 1 - 2g_c + (\eta - c)g'_c + \eta. \end{aligned} \quad (169)$$

For $-1 \leq \eta \leq 0$, substitution of $g'_c = (1 + \eta)/C$ makes these values $-(1 + \eta)/C$, $-(1 + \eta)/C$, and 0, respectively. For $\eta \leq -1$ their signs are immediate from $g_c = g'_c = 0$. Hence the base bank pays every ordinary endpoint.

The cap reserve. Define

$$Z(c) = 2 + \frac{1}{c} + \frac{1}{2} \log \left(1 + \frac{2}{c} \right), \quad \delta = Z(c) - u. \quad (170)$$

On the mixed curve, (146) says precisely that

$$\delta = \text{artanh } m. \quad (171)$$

For $0 \leq t \leq m$, set

$$\beta(t) = ct[\delta - \text{artanh } t], \quad \mathcal{V}(t) = \int_t^m \beta(s) ds, \quad (172)$$

and extend $\mathcal{V}$ by zero for $t \geq m$. Since $\beta(m) = 0$, this extension is C^1 . With

$$q = \frac{K}{x}, \quad \mu = \frac{q}{1 + q}, \quad \mathcal{H}(q) = (1 + q)^2 \mathcal{V}(\mu), \quad (173)$$

define the full potential

$$W = xD + x^2(G_c(y, b) + \mathcal{H}(q)). \quad (174)$$

It is nonnegative.

For an ordinary endpoint, K is fixed while x decreases, so the reserve contributes, after division by x ,

$$L_0(q) = -2\mathcal{H}(q) + q\mathcal{H}'(q). \quad (175)$$

A cap also increments K , and its contribution is $L_0 + \mathcal{H}'$. On the active range $\mu \leq m$, differentiation of (173) gives

$$\mathcal{H}' = 2(1+q)\mathcal{V} - \beta(\mu), \quad L_0 = -2(1+q)\mathcal{V} - q\beta(\mu) \leq 0. \quad (176)$$

Thus the reserve can only improve every ordinary drift.

For a cap, feasibility $D \leq P$ and $P \geq K$ gives

$$y = \frac{D - C(P + E)}{x} \leq -\frac{cK}{x} = -cq. \quad (177)$$

While $\mu \leq m$, we have $q \leq m/(1-m) \leq \varphi$ and hence $cq \leq 1$. Monotonicity of a_c , together with (158), yields

$$\frac{\ell_U - \ell_Q}{x} \leq cq \left[\delta - \frac{1}{2} \log(1+2q) \right] =: B(q). \quad (178)$$

The change of variables in (173) satisfies

$$\operatorname{artanh} \mu = \frac{1}{2} \log(1+2q), \quad (1+q)\beta(\mu) = B(q). \quad (179)$$

Consequently the cap residual is cancelled exactly:

$$L_0 + \mathcal{H}' + B(q) = 0 \quad (\mu \leq m). \quad (180)$$

When $\mu \geq m$, the reserve vanishes. At $q_* = m/(1-m)$, the bracket in (158), evaluated at the largest possible threshold $a_c(-cq_*)$, is zero by (171) and (179). For $q \geq q_*$, (177) only decreases the threshold, so that bracket is nonpositive. The fifth Hamiltonian inequality therefore holds for every state and for arbitrary interleavings of capped and ordinary outcomes.

Initial calibration. Initially $y = b = q = 0$, and

$$f_c(0) = \frac{1}{2} - \frac{c}{4} \log\left(1 + \frac{2}{c}\right). \quad (181)$$

Elementary integration in (172) gives

$$\mathcal{V}(0) = \frac{c}{2}(\delta - m). \quad (182)$$

Indeed,

$$\int_0^m t \operatorname{artanh} t \, dt = \frac{(m^2 - 1) \operatorname{artanh} m + m}{2}.$$

Using (146) and (171), we obtain the exact calibration

$$f_c(0) + \mathcal{H}(0) = f_c(0) + \mathcal{V}(0) = \frac{c}{2}. \quad (183)$$

Thus $W_{\text{initial}} = cn^2/2$.

Finite jobs, and uniformity on the plateau. We spell out the regularity needed to pass from the fluid Hamiltonian to unit job transitions. It is important here to restrict attention to feasible states. In Region I, $y \geq -1$ implies

$$-xy = C(P + E) - D = cP + (P - D) + CE \leq x. \quad (184)$$

All terms in the middle expression are nonnegative. Hence $P/x, D/x, q \leq 1/c$ and $b \leq 1/C$, so every normalized derivative of the Region-I perspective in (163) is bounded on the feasible cone. In Region II the raw base bank is simply

$$xD + \frac{(x + D - CP)_+^2}{2C}, \quad (185)$$

which has bounded one-sided Hessians. The two pieces are C^1 at $y = -1$, as verified above, and the positive-part square is C^1 at $\eta = -1$.

For the reserve $R(x, K) = x^2 \mathcal{H}(K/x)$, on its active cone $0 \leq q \leq q_*$ one has bounded $\mathcal{H}''$, and

$$R_{xx} = 2\mathcal{H} - 2q\mathcal{H}' + q^2\mathcal{H}'', \quad R_{xK} = \mathcal{H}' - q\mathcal{H}'', \quad R_{KK} = \mathcal{H}''. \quad (186)$$

For $q \geq q_*$, $R = 0$; moreover $\mathcal{H}(q_*) = \mathcal{H}'(q_*) = 0$. Thus the join is C^1 with bounded one-sided Hessians. At $x = 0$, set $R = 0$: for $K > 0$ it is already locally zero, while at $(x, K) = (0, 0)$ two-homogeneity gives zero gradient. Altogether W is $C^{1,1}$ on the feasible state cone. On the mixed interval its regularity constant may depend on the fixed value of u . When $c = r$ and $m = 0$, however, the reserve is identically zero and the base bank depends only on the absolute constant r ; its regularity constant is therefore uniform over all $u \geq z_*$. This includes the last transition from $x = 1$ to $x = 0$.

It follows that a unit transition has the Taylor remainder

$$\ell_i + W_{i+1} - W_i \leq O_u(1). \quad (187)$$

The endpoint $E = 0^+$ is interpreted one-sidedly. For a fixed symbolic word, replace all E -coordinates by positive values ϵ . Its state trajectory is unchanged, and the exact finite pair objective, being a finite sum of affine and minimum terms, converges as $\epsilon \downarrow 0$. Equivalently, retain an $O(\epsilon n^2)$ error and take this limit for each fixed n . Therefore the formal E -Hamiltonian transfers to genuine outcome vectors without an asymptotic loss.

Summing (187), using $W \geq 0$, and applying (183) gives

$$\sum_i \ell_i \leq \frac{c}{2} n^2 + O_u(n). \quad (188)$$

Together with (153) and (157), this yields

$$\text{ALG} - C \text{OPT} \leq -c \binom{n}{2} + \frac{c}{2} n^2 + O_u(n) = O_u(n). \quad (189)$$

At $u = z_*$, one has $c = r$, $m = 0$, and $Z(r) = z_*$. The cap reserve is identically zero, while (183) remains valid. If u is increased above z_* , the only adverse fifth-endpoint bracket $1 - r(u - 1 - a_r(y))$ decreases, and the capped diagonal term in (153) also decreases.

Here is the promised uniform finite calculation. The base bank and its feasible cone contain no u , so there is one constant κ , depending only on r , for which every ordinary or capped transition with $u \geq z_*$ satisfies

$$\ell_i + W_{i+1} - W_i \leq \kappa.$$

Because $W_0 = rn^2/2$ and $W_n = 0$, telescoping gives

$$\sum_i \ell_i \leq \frac{r}{2}n^2 + \kappa n. \quad (190)$$

In the diagonal term of Equation (153), a value below the cap contributes $-r(1 + p_i)$. A capped value contributes at most $1 - ru \leq 0$. With the convention $\bar{p}_i = p_i$ at $u = \infty$, the first formula applies there and the capped endpoint is absent. Hence the diagonal sum is nonpositive uniformly on $[z_*, \infty]$. Combining Equations (153), (157) and (190) now yields

$$\text{ALG} - (1 + r) \text{OPT} \leq -r \binom{n}{2} + \frac{r}{2}n^2 + \kappa n = \left(\kappa + \frac{r}{2}\right)n. \quad (191)$$

Thus, with $B = \kappa + r/2$,

$$\text{ALG} \leq (1 + r) \text{OPT} + Bn \quad \text{for every } u \in [z_*, \infty], \quad (192)$$

and B is independent of u . The estimate is genuinely uniform rather than the result of sending a parameter-dependent remainder to infinity. This proves Theorem 5.10. Finally, because $u > 1$ on the present range, every effective offline length is at least one and $\text{OPT} \geq n(n + 1)/2$. The additive term in (189) vanishes in the size-asymptotic ratio. $\square$

6 Discussion and open questions

The curve identifies the leading quadratic constants of the deterministic games for every $u \in (0, \infty]$. Its scope is deliberately precise: one machine, all release dates zero, unit tests, nonpreemptive operations, total completion time, and one common upper limit when the cap is finite. Every upper proof loses at most a constant on each of the n discrete transitions. This loss is asymptotically negligible, but it is not a fixed additive constant over the whole instance. On $u \in [z_*, \infty]$ the per-transition constant is uniform in u .

Conjecture 6.1 (Endpoint additive competitiveness). There exist a deterministic online algorithm and a constant $B < \infty$ such that every instance at the cap-free endpoint satisfies

$$\text{ALG}(I) \leq R_\star \text{OPT}(I) + B.$$

The present Adaptive-Threshold algorithm is the natural first candidate. In Equations (43) and (44) the exact local contributions contain favorable negative terms that are intentionally discarded before the bank-account calculation. A discrete potential that retains those terms might prove Conjecture 6.1; alternatively, a finite-instance obstruction might show that a different policy is needed. The current lower bound does not distinguish these possibilities. In particular, our result neither proves nor claims the stronger ordinary inequality with $B = 0$.

The analogous endpoint-additive question is open for each nontrivial finite-cap branch as well: our claims concern CR_∞ and the infimum of admissible additive coefficients. Also open are heterogeneous upper limits, nonuniform test times in the exact-ratio regime, randomization, and more than one machine. The common-upper curve shows why a direct extension is unlikely to have one universal threshold: already in the uniform case the optimal policy changes form at five analytically distinct junctions.

The lower construction also suggests a structural principle for future lower bounds: hard instances reveal many decreasing positive levels, with continuum increment $dp = ds/(1 + s)$, and postpone zero jobs to the end. Our argument proves that this pattern is sufficient. It remains open whether every asymptotically extremal construction must have this form.

7 Conclusion

The family has one common cap-free core. At $u = \infty$, matching harmonic and adaptive-threshold arguments meet at

$$R_\star = 1 + \frac{2}{z_\star - 1} = 1.570574096649395\dots, \quad z_\star - 3 = \log z_\star.$$

For finite common upper limit u , binary stopping games, forced-prefix and zero-prefix UTE policies, a capped-harmonic mixture, and a cap-reserve extension of the same adaptive threshold meet on all six pieces of (6). In particular, the worst common-upper value is $u_\diamond = 1.866760399173862\dots$, while every $u \geq z_\star$ has exactly the cap-free value. Proof-theoretically, finite u adds one capped endpoint to the ordinary four-endpoint bank, and the reserve used to pay it vanishes at $z_\star$. The principal remaining questions are finite and nonuniform: endpoint additive constants, different upper limits or test times, and randomized algorithms.

A Finite rational checkpoint for the cap-free lower bound

The lower bound at $u = \infty$ takes two limits: first the scaling $H \rightarrow \infty$, and then the number K of positive levels tends to infinity. This appendix is not needed for that proof. It gives one exact finite instance showing that the leading-term calculation already predicts the true ratio accurately.

Take

$$K = 1000, \quad \alpha = \frac{9}{8}, \quad \gamma = \frac{679}{1000}, \quad \xi = \frac{K}{\alpha} = \frac{8000}{9}, \quad H = 9000.$$

There are H jobs at each positive level and ξH zero jobs. All multiplicities are integral, and all processing times defined by Equation (23) are rational.

Using exact rational arithmetic in Equations (22) and (26) to (28) gives the leading-coefficient comparison

$$\frac{Q_K}{D_K} = 1.5705287442294085\dots > 1.5705.$$

The displayed decimal is only for orientation; all comparisons below are between integers or rational numbers.

For the offline schedule, retaining the terms that are linear in H gives

$$\text{OPT}_H = H^2 D_K + H L_{\text{off}}, \tag{193}$$

where

$$\begin{aligned} L_{\text{off}} &= \frac{1}{2} \left(\xi + \sum_{i=0}^{K-1} (1 + p_i) \right) \\ &= \frac{1}{2} \left(\xi + 2K + \frac{K\gamma}{\xi} + \sum_{s=1}^{K-1} \frac{K-s}{\xi+s} \right) = 1656.440800628232\dots \end{aligned} \tag{194}$$

Again, the middle expression is the exact rational value used in the comparison.

The discrete shortest-prefix calculation behind Equation (20) also retains favorable diagonal terms that vanish after division by H^2 . Before dropping them it gives

$$\text{ALG}_H \geq H^2 Q_K + \frac{H}{2} \left(\xi + \sum_{i=0}^{K-1} p_i + \sum_{i=0}^{K-1} s_i \right) \geq H^2 Q_K.$$

Here the s_i 's are the nonnegative completion fractions from the lower proof. Exact rational arithmetic verifies

$$2000H^2Q_K > 3141(H^2D_K + HL_{\text{off}}). \quad (195)$$

Consequently, the fixed instance obtained by the replay argument in Section 4.1 satisfies

$$\frac{\text{ALG}_H}{\text{OPT}_H} \geq \frac{H^2Q_K}{H^2D_K + HL_{\text{off}}} = 1.5705286254871234\dots > \frac{3141}{2000} = 1.5705.$$

Thus the checkpoint includes the finite- H correction; it is not merely a comparison of the two leading coefficients.

B Independent certificate by bubbling caps to a prefix

For completeness, we give a second proof of the quadratic estimate. This argument is also a useful audit of the fifth-endpoint calculation: it never inserts $p = u$ into an ordinary deferred formula before the caps have actually been moved.

Work first in normalized fluid mass. Immediately before an adjacent pair XU , let E_0 be the preceding E -mass, P the preceding substantive mass, D its deferred submass, x the remaining mass, and w the mass of U -endpoints strictly after the pair. Put

$$y = \frac{D - C(P + E_0)}{x}, \quad \eta = \frac{D - CP}{x} = y + C\frac{E_0}{x} \leq 0. \quad (196)$$

Direct differentiation of the exact pair objective (152), including the mutual pair and all future caps, gives the two nontrivial commutators

$$UM - MU = 1 + \left(1 + \eta - \frac{Cw}{x}\right) (y - c)a'_c(y), \quad (197)$$

$$UQ - QU = \left(\eta - \frac{cw}{x}\right) (y - c)a'_c(y). \quad (198)$$

Here $UX - XU$ means “new objective after moving U left” minus the old objective. The second quantity is nonnegative. For the first, if $B = 1 + \eta - Cw/x \leq 0$, its second term is nonnegative. If $B > 0$, then $B \leq 1$, and on $-1 < y \leq 0$

$$\frac{(c - y)B}{c - 2y} \leq 1.$$

Thus (197) is nonnegative as well. Below $y = -1$, the threshold is flat. Direct evaluation also gives

$$UZ - ZU = UE - EU = 1. \quad (199)$$

The suffix terms involving w are crucial: they make every adjacent exchange valid even while other caps remain to the right. Repeated swaps therefore put all caps into one prefix.

After that prefix there is no future cap. Let e, b, d be the preceding E -mass, substantive mass, and deferred substantive mass, and define

$$y = \frac{d - C(e + b)}{x}, \quad v = \frac{e}{x}, \quad h = d - Cb, \quad k = x + d - Cb.$$

The four ordinary commutators are

$$ME - EM = a_c + a'_c(C - y)\frac{k}{x}, \quad QE - EQ = 1 - a'_c(C - y)(-y - Cv),$$

$$MZ - ZM = a_c - a'_c y \frac{k}{x}, \quad QZ - ZQ = 1 - a'_c y \frac{h}{x}. \quad (200)$$

They are nonnegative. Indeed, in the varying region $h/x = y + Cv \in [y, 0]$ and $k/x \in [0, 1]$. With $t = -y \in [0, 1]$,

$$a'_c(C - y)t = \frac{t(C + t)}{c + 2t} \leq 1, \quad a'_c t^2 = \frac{t^2}{c + 2t} \leq 1; \quad (201)$$

the first inequality is equivalent to $(t - 1)(t + c) \leq 0$. In the flat region $a'_c = 0$. Hence Z, E can be moved to the tail, and a worst word has the form

$$U^\mu \{M, Q\}^* \{Z, E\}^*. \quad (202)$$

On the middle block use the simpler potential

$$\widehat{W}(x, N, D) = xD + x^2 f_c \left(\frac{D - CN}{x} \right).$$

The identity

$$2f_c - yf'_c = \max\{0, -Cf'_c, -Cf'_c + a_c(1 + y), 1 - cf'_c + a_c y\} \quad (203)$$

is a direct rewriting of (165); it pays the entire ordinary continuation. If the cap prefix has mass μ , then after it $x = 1 - \mu$, $N = D = \mu$, and $y = -c\mu/(1 - \mu)$. Its internal cap-pair reward plus its worst continuation is therefore

$$F_{c,u}(\mu) = \frac{2 - c(u - 1)}{2} \mu^2 + \mu(1 - \mu) + (1 - \mu)^2 f_c \left(-\frac{c\mu}{1 - \mu} \right). \quad (204)$$

For $0 < \mu < 1/C$, differentiation gives

$$F'_{c,u}(\mu) = c\mu [Z(c) - u - \operatorname{artanh} \mu]. \quad (205)$$

On the mixed curve the unique maximizer in this range is therefore $\mu = m$, and

$$F_{c,u}(m) = f_c(0) + \frac{c}{2} (\operatorname{artanh} m - m) = \frac{c}{2}. \quad (206)$$

Here $m \leq 1/C$, since $c, m \leq 1/\varphi$. On the flat part $\mu \geq 1/C$,

$$F_{c,u}(\mu) = \mu - \frac{c(u - 1)}{2} \mu^2. \quad (207)$$

Its derivative at $1/C$ is

$$\frac{c(1 + m) - 1}{C} \leq 0,$$

and only decreases thereafter. Thus $F_{c,u}(\mu) \leq c/2$ on the whole unit interval. At $c = r, m = 0, u = z_*$, the same calculation is maximized at $\mu = 0$, and increasing u only lowers (204).

To make this independent exchange proof finite, fix $\varepsilon > 0$ and first replace the last $\varepsilon n + O(1)$ outcomes by genuine zeros. Earlier states and actions are unchanged, and only $O(\varepsilon n^2 + n)$ pair terms are affected. All cap and ordinary exchanges in the retained prefix then occur with $x \geq \varepsilon$; their displayed fluid commutators have uniform Taylor remainders there. More precisely, after assigning mass $1/n$ to every job, a normalized adjacent exchange has remainder $O_{u,\varepsilon}(n^{-3})$; there are at most n^2 exchanges. At the join $y = -1$, split an

exchange at the join and use the two one-sided formulae, whose values agree. The artificial zeros already form the terminal block, so no unubbled suffix cap invalidates (200). The normalized one-job remainder in the continuation-potential calculation is $O_{u,\varepsilon}(n^{-2})$. For fixed ε , first let $n \rightarrow \infty$, and then let $\varepsilon \downarrow 0$. Equations (203)–(206) give

$$\Gamma_n \leq c \binom{n}{2} + o(n^2),$$

which is an independent size-asymptotic certificate for the two upper bounds above.

C Machine-checked main theorem and its correspondence

This appendix records the public theorem exported by the Lean 4 development [dMU21] associated with this manuscript and explains how its formal predicates correspond to the mathematical statements in Theorems 1.1 and 1.2. The development uses the mathlib library [The20] and is pinned to Lean and mathlib version 4.30.0. Its root module imports `SchedulingPaper.MainTheorems`, whose final theorem is reproduced below. Apart from the typographic rendering of Unicode symbols, this is a verbatim excerpt.

```
theorem exact_ratio_main
  : (∀ {u : ℝ}, 0 < u →
    FiniteExactRatioConclusion u (exactCurve u)) ∧
    ObligatoryExactRatioConclusion RStar := by
  constructor
  · intro u hu
    exact finite_exact_ratio hu
  · exact obligatory_exact_ratio
```

The two conclusion records occurring in the theorem are:

```
structure FiniteExactRatioConclusion (u ratio : ℝ) : Prop where
  lower : LowerBound.FixedSizeLowerBound (.finite u) ratio
  upper :
    ∃ strategy : ∀ n, Online.Strategy n,
      UpperBound.CompletesByAnalysisFuel (.finite u) strategy ∧
      SizeAsymptoticUpper
        (UpperBound.strategyCost (.finite u) strategy)
        (UpperBound.fixedOfflineCost (.finite u)) ratio

structure ObligatoryExactRatioConclusion (ratio : ℝ) : Prop where
  lower : LowerBound.FixedSizeLowerBound .infinite ratio
  upper :
    ∃ strategy : ∀ n, Online.Strategy n,
      UpperBound.CompletesByAnalysisFuel .infinite strategy ∧
      SizeAsymptoticUpper
        (UpperBound.strategyCost .infinite strategy)
        (UpperBound.fixedOfflineCost .infinite) ratio
```

C.1 Objects on the two sides

The following table gives the direct dictionary between the formal objects and the notation of the manuscript.

Lean object	Manuscript object	Meaning
<code>Cap.finite u</code>	finite common upper limit u	processing times lie in $[0, u]$; testing and raw execution are available
<code>Cap.infinite</code>	obligatory testing	processing times are nonnegative and raw execution is unavailable
<code>exactCurve u</code>	$V(u)$	the six branches in Equation (6), including all five verified joins
<code>RStar</code>	$R_\star$	$1 + 2/(z_\star - 1)$, where $z_\star > 1$ is the unique solution of $z_\star - 3 = \log z_\star$
<code>fixedOfflineCost</code>	OPT	the clairvoyant shortest-first cost of the effective lengths $q_u(p)$
<code>strategyCost</code>	ALG	the total completion cost of the operational online transcript

A formal strategy on n jobs is a deterministic function from the public transcript to its next action. A test publishes the revealed processing time; raw execution publishes no hidden value. Thus the strategy has exactly the information available to an online algorithm in the manuscript. The formal action type uses the standard no-idling normalization appropriate when all jobs are released at time zero. Processing times and comparisons are exact real numbers.

C.2 Expansion of the lower and upper fields

For a cap and a candidate value R , `FixedSizeLowerBound` expands to the following statement. For every size-indexed family of deterministic strategies $\mathcal{B}_n$, every $\varepsilon > 0$, and every N , there are an $n \geq N$, a genuine fixed admissible n -job input I , and a finite settled execution such that either the execution leaves a job incomplete or

$$(R - \varepsilon) \text{OPT}(I) \leq \text{ALG}_{\mathcal{B}_n}(I). \quad (208)$$

An incomplete settled run represents infinite cost. In particular, for algorithms that complete all inputs, (208) is the usual lower-bound sequence on arbitrarily large fixed instances. The lower constructions are first described by adaptive revelation oracles, but the formal replay theorem freezes every such interaction into the fixed input quantified here; this is the machine-checked counterpart of Lemma 3.3.

The predicate `SizeAsymptoticUpper` is the epsilon formulation

$$\forall \varepsilon > 0 \exists N \forall n \geq N \forall I, \quad \text{ALG}_{\mathcal{A}_n}(I) \leq (R + \varepsilon) \text{OPT}(I). \quad (209)$$

The accompanying completion field proves that every job is done after the common analysis fuel $2n + 1$, rather than merely assigning a cost to a partial trace. Consequently, Equations (208) and (209) are the two matching operational statements that characterize the optimal value in Equation (15). The Lean theorem packages these two directions; it does not introduce a separate real-valued infimum or `limsup` object.

Before projecting to `SizeAsymptoticUpper`, the checked upper proof constructs a stronger `FixedUpperCertificate`: for the chosen cap it contains a constant $C \geq 0$ such that, for every n and every fixed input,

$$\text{ALG}_{\mathcal{A}_n}(I) \leq R \text{OPT}(I) + Cn. \quad (210)$$

The quadratic lower bounds Equations (13) and (14) then imply (209). For $u \geq z_\star$, the separately verified plateau certificate uses the same strategy and the same C , independent of u , including the obligatory endpoint. This uniformity is stronger than the quantifier order displayed by `exact_ratio_main`, which only exposes one strategy after a cap has been fixed.

C.3 Correspondence with the two main theorems

The first conjunct of `exact_ratio_main` quantifies over every finite $u > 0$. Substituting $R = \text{exactCurve } u = V(u)$, its lower field gives the fixed-instance lower direction of Theorem 1.2, while its upper field supplies a completing strategy and the matching upper direction. The definition of `exactCurve` is literally the six-way case split in Equation (6); the Lean development also proves its joins, continuity, plateau, and global maximum at u_∞ .

The second conjunct substitutes $R = \text{RStar} = R_\star$ in the obligatory model. Its lower field is Theorem 1.1(i), and its completing upper strategy together with (210) is Theorem 1.1(ii). The common formal infrastructure also proves that the offline benchmark is the shortest-first schedule of effective lengths and that the transcript cost is exactly its suffix-weighted pair-accounting expression.

C.4 Proof boundary

The root Lean module imports the complete lower- and upper-bound assemblies. The public theorem has no operational interface hypotheses, and the development contains no `sorry`, `admit`, or project-defined `axiom`. It does use `mathlib`’s classical real analysis and noncomputable choices of exact real parameters. Thus it certifies deterministic mathematical strategy terms, not an extracted floating-point implementation.

The theorem recorded here is specifically the size-asymptotic result. It does not assert the pointwise inequality $\text{ALG} \leq R \text{OPT}$ without a remainder, nor admissibility of the endpoint coefficient with one fixed additive constant. The formal library separately derives additive admissibility of every coefficient $R + \varepsilon$ from (210). The machine-checked mixed-regime upper bound follows the cap-reserve potential proof; the bubbling argument in Appendix B remains an independent manuscript cross-check.

References

- [AE21] Susanne Albers and Alexander Eckl. Explorable uncertainty in scheduling with non-uniform testing times. In Christos Kaklamanis and Asaf Levin, editors, *Approximation and Online Algorithms: 18th International Workshop, WAOA 2020, Revised Selected Papers*, volume 12806 of *Lecture Notes in Computer Science*, pages 127–142. Springer, 2021. doi:10.1007/978-3-030-80879-2_9.
- [BS26] Felix Buld and Andreas S. Schulz. Scheduling with testing: Competitive algorithms for minimizing the total weighted completion time in the adversarial model, 2026. arXiv:2606.25166.
- [DDN⁺22] Fanny Dufossé, Christoph Dürr, Noël Nadal, Denis Trystram, and Óscar C. Vázquez. Scheduling with a processing time oracle. *Applied Mathematical Modelling*, 104:701–720, 2022. doi:10.1016/j.apm.2021.12.020.
- [DEL24] Konstantinos Dogeas, Thomas Erlebach, and Ya-Chun Liang. Scheduling with obligatory tests. In Timothy Chan, Johannes Fischer, John Iacono, and Grzegorz Herman, editors, *32nd Annual European Symposium on Algorithms (ESA 2024)*, volume 308 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 48:1–48:14, Dagstuhl, Germany, 2024. Schloss Dagstuhl – Leibniz-Zentrum für Informatik. doi:10.4230/LIPIcs.ESA.2024.48.

- [DEMM20] Christoph Dürr, Thomas Erlebach, Nicole Megow, and Julie Meißner. An adversarial model for scheduling with testing. *Algorithmica*, 82(12):3630–3675, 2020. doi:10.1007/s00453-020-00742-2.
- [DKL⁺23] Christoph Damerius, Peter Kling, Minming Li, Chenyang Xu, and Ruilong Zhang. Scheduling with a limited testing budget: Tight results for the offline and oblivious settings. In *31st Annual European Symposium on Algorithms (ESA 2023)*, volume 274 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 38:1–38:15. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2023. doi:10.4230/LIPIcs.ESA.2023.38.
- [dMU21] Leonardo de Moura and Sebastian Ullrich. The Lean 4 theorem prover and programming language. In *Automated Deduction – CADE 28*, volume 12699 of *Lecture Notes in Computer Science*, pages 625–635. Springer, 2021. doi:10.1007/978-3-030-79876-5_37.
- [GCH24] Mingyang Gong, Zhi-Zhong Chen, and Kuniteru Hayashi. Approximation algorithms for multiprocessor scheduling with testing to minimize the total job completion time. *Algorithmica*, 86(5):1400–1427, 2024. doi:10.1007/s00453-023-01198-w.
- [Kah91] Simon Kahan. A model for data in motion. In *Proceedings of the 23rd Annual ACM Symposium on Theory of Computing (STOC 1991)*, pages 265–277. Association for Computing Machinery, 1991. doi:10.1145/103418.103449.
- [KLL⁺26] Bob Krekelberg, Alison Hsiang-Hsuan Liu, Fu-Hong Liu, Prudence W. H. Wong, and Xiao-Ou Zhang. The power of amortization on minimizing total completion time with explorable uncertainty, 2026. Version 2, revised 8 May 2026. arXiv:2503.13357.
- [LL26] Kao-Chuan Liang and Ya-Chun Liang. Hardness of obligatory-test scheduling on multiple machines, 2026. Version 2, revised 14 July 2026. arXiv:2606.02029.
- [LLWZ23] Alison Hsiang-Hsuan Liu, Fu-Hong Liu, Prudence W. H. Wong, and Xiao-Ou Zhang. The power of amortization on scheduling with explorable uncertainty. In Jaroslaw Byrka and Andreas Wiese, editors, *Approximation and Online Algorithms: 21st International Workshop, WAOA 2023*, volume 14297 of *Lecture Notes in Computer Science*, pages 90–103. Springer, 2023. doi:10.1007/978-3-031-49815-2_7.
- [LMS19] Retsef Levi, Thomas L. Magnanti, and Yaron Shaposhnik. Scheduling with testing. *Management Science*, 65(2):776–793, 2019. doi:10.1287/mnsc.2017.2973.
- [LMS24] Retsef Levi, Thomas L. Magnanti, and Yaron Shaposhnik. Scheduling with testing of heterogeneous jobs. *Management Science*, 70(5):2934–2953, 2024. doi:10.1287/mnsc.2023.4833.
- [LSSS26] Alexander Lindermayr, Guido Schäfer, Jens Schlöter, and Leen Stougie. Online flow time minimization with gradually revealed jobs, 2026. arXiv:2602.12716.

- [MPT94] Rajeev Motwani, Steven J. Phillips, and Eric Torng. Nonclairvoyant scheduling. *Theoretical Computer Science*, 130(1):17–47, 1994. doi:10.1016/0304-3975(94)90151-1.
- [Smi56] Wayne E. Smith. Various optimizers for single-stage production. *Naval Research Logistics Quarterly*, 3(1–2):59–66, 1956. doi:10.1002/nav.3800030106.
- [The20] The mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020)*, pages 367–381. Association for Computing Machinery, 2020. doi:10.1145/3372885.3373824.