

Cutaneous Adverse Drug Eruption Associated with SynthoStatin Therapy: A Detailed Clinical Case

Robert Martinez, MD; Sarah Jenkins, MD; Department of Dermatology, United States

ABSTRACT

We describe a case of severe erythematous maculopapular rash in a 71-year-old male patient following initiation of SynthoStatin 20mg oral daily for hypercholesterolemia. Symptoms commenced 10 days into therapy. Dechallenge resulted in complete resolution.

CASE PRESENTATION

A 71-year-old male patient (Weight: 75 kg, past history of hypertension) presented to our outpatient clinic with an extensive erythematous maculopapular rash covering his torso and upper extremities. Ten days prior to presentation, the patient had been prescribed oral SynthoStatin 20mg once daily. Physical examination revealed confluent erythematous macules without mucosal involvement. SynthoStatin was promptly discontinued on 14 Aug 2026. Within 7 days of cessation, the rash resolved completely without scarring (Outcome: Recovered). The reporting physician was Dr. Martinez, Dermatologist.

DISCUSSION

Statin-induced cutaneous adverse reactions are well-documented yet underreported pharmacovigilance signals. In this patient, prompt dechallenge confirmed causality without rechallenge.

REFERENCES

1. Smith AB, et al. Cutaneous reactions to lipid-lowering agents. J Dermatol Sci. 2023;45:101-108.
2. Dupont H, et al. Adverse drug reactions in pharmacovigilance databases. Lancet Pharm. 2024;12:45-52.
3. Williams R. Clinical dermatology handbook, 5th ed. London: MedPress; 2022.