

LECTURE 2

SIMULTANEOUS EQUATIONS I: DEFINITION, IDENTIFICATION, INDIRECT LS,
SINGLE-EQUATION GMM

Definition

We consider the following *system of equations*:

$$\Gamma Y_i = B Z_i + U_i, \quad (1)$$

$$E [Z_i U_i^\top] = 0, \quad (2)$$

where Y_i is an m -vector of *endogenous* variables:

$$Y_i = \begin{pmatrix} y_{1i} \\ \vdots \\ y_{mi} \end{pmatrix},$$

Z_i is an l -vector of *exogenous* variables, and U_i is an m -vector of errors:

$$U_i = \begin{pmatrix} u_{1i} \\ \vdots \\ u_{mi} \end{pmatrix}.$$

The unknown parameters are given by the $m \times m$ matrix Γ and the $m \times l$ matrix B . The econometrician observes the data $\{(Y_i^\top, Z_i^\top) : i = 1, \dots, n\}$.

We have m equations determining m endogenous variables. The first equation is given by

$$\sum_{j=1}^m \Gamma_{1j} y_{ji} = \sum_{j=1}^l B_{1j} Z_{ji} + u_{1i}.$$

Multiplying all Γ_{1j} 's and B_{1j} 's by a nonzero constant would affect the variance of the unobservable error term u_{1i} ; however, it would not change the relationship between the observable variables, that is, Γ and B can be identified only up to scale. Hence, we must introduce some normalization to the system. We will assume that $\Gamma_{jj} = 1$ for all $j = 1, \dots, m$. Later, this will allow us to treat y_{ji} as the “dependent” variable in equation j .

Next, we assume that Γ^{-1} exists, since, otherwise, there is no unique expression for Y_i in terms of the exogenous variables and shocks. Let us rewrite (1) as

$$\begin{aligned} Y_i &= \Gamma^{-1} B Z_i + \Gamma^{-1} U_i, \text{ or} \\ Y_i &= \Pi Z_i + V_i, \text{ where} \\ \Pi &= \Gamma^{-1} B, \\ V_i &= \Gamma^{-1} U_i. \end{aligned} \quad (3)$$

The system of equations in (3) is called the *reduced form* as opposed to *structural* equations in (1). Due to (2), we have that $E [Z_i V_i^\top] = 0$, and, consequently, the matrix of the reduced-form parameters Π can be estimated consistently by OLS. The reduced form is useful, for example, for forecasting Y from Z ; however, we are usually interested in estimating economic relations that are given by the structural equations. The structural parameters cannot be estimated directly by OLS, since each of the structural equations has Y 's on the right-hand side, and in general,

$$\begin{aligned} E [Y_i U_i^\top] &= \Pi E [Z_i U_i^\top] + \Gamma^{-1} E [U_i U_i^\top] \\ &= \Gamma^{-1} \text{Var}(U_i) \\ &\neq 0. \end{aligned}$$

Hence, the first question is whether it is possible to recover the structural parameters Γ from Π .

A structural equation is said to be *identified* if its coefficients are *uniquely* determined by the elements of Π . The structural equation is not identified when the structural parameters cannot be recovered *uniquely* from Π (we know that there is a solution for Γ and B because by construction $\Pi = \Gamma^{-1}B$). Simple counting shows that there are $m^2 - m + ml$ unknown parameters in Γ and B , while there are only ml reduced-form coefficients in Π . Thus, without imposing additional restrictions on the structural parameters, the system or its parts cannot be identified. The most common type of restriction is a zero restriction, or exclusion of variables from equations. It is possible that some equations are identified while others are not, and, therefore, identification should be considered equation by equation.

Without loss of generality, let us consider the first equation. Suppose that the first equation has m_1 endogenous and l_1 exogenous variables included on the right-hand side. Write

$$\Gamma = \begin{pmatrix} 1 & -\gamma_1^\top & 0^\top \\ \Gamma_1 & \Gamma_2 & \Gamma_3 \end{pmatrix},$$

where, in the first row, γ_1 is an m_1 -vector with all elements different from zero, and 0 is actually an $(m - m_1 - 1)$ -vector of zeros;

$$B = \begin{pmatrix} \beta_1^\top & 0^\top \\ B_1 & B_2 \end{pmatrix},$$

where, in the first row, β_1 is an l_1 -vector with all elements different from zero, and 0 is an $(l - l_1)$ -vector of zeros. Accordingly, let us partition Π such that $\Gamma\Pi = B$:

$$\begin{pmatrix} 1 & -\gamma_1^\top & 0^\top \\ \Gamma_1 & \Gamma_2 & \Gamma_3 \end{pmatrix} \begin{pmatrix} \pi_1 & \pi_2 \\ \Pi_1 & \Pi_2 \\ \Pi_3 & \Pi_4 \end{pmatrix} = \begin{pmatrix} \beta_1^\top & 0^\top \\ B_1 & B_2 \end{pmatrix}, \text{ and} \quad (4)$$

$$\pi_1 - \gamma_1^\top \Pi_1 = \beta_1^\top, \quad (4)$$

$$\pi_2 - \gamma_1^\top \Pi_2 = 0^\top, \quad (5)$$

where π_1 is $1 \times l_1$, π_2 is $1 \times (l - l_1)$, Π_1 is $m_1 \times l_1$, and Π_2 is $m_1 \times (l - l_1)$. The expression in (4) says that we can find β_1 if we know γ_1 . Next, (5) is a system of $l - l_1$ equations with m_1 unknowns (γ_1). To solve for γ_1 , there should be at least as many equations as unknowns, which gives us the following *order condition*:

$$l - l_1 \geq m_1,$$

that is, the number of exogenous variables excluded from the equation should be at least as large as the number of endogenous variables included on the right-hand side. The order condition is only necessary but not sufficient. Given the fact that there is a solution (because Π is determined by Γ and B), the necessary and sufficient condition for there to be a unique solution for γ_1 is the following *rank condition*:

$$\text{rank}(\Pi_2) = m_1$$

(see the appendix for a proof). Thus, the equation is *not identified* when $l - l_1 < m_1$ or the rank condition fails. It is *overidentified* when $l - l_1 > m_1$, and the rank condition is met. The equation is *exactly identified* when $l - l_1 = m_1$, and the rank condition is met. In the exactly identified case, we have

$$\gamma_1^\top = \pi_2 \Pi_2^{-1}. \quad (6)$$

Indirect LS

Since Π can be estimated consistently by OLS, we can use (6) to estimate the structural parameters of an exactly identified equation. The indirect LS estimator of γ_1 is given by

$$\hat{\gamma}_1^{\text{LS}} = \left(\hat{\Pi}_2^\top \right)^{-1} \hat{\pi}_2^\top,$$

where $\hat{\pi}_2$ and $\hat{\Pi}_2$ are the LS estimators of π_2 and Π_2 , respectively.

Let us write

$$Y_i = \begin{pmatrix} y_{1i} & Y_{1,i}^\top & Y_{1,i}^{*\top} \end{pmatrix}^\top,$$

where $Y_{1,i}$ is the m_1 -vector of the endogenous variables included on the right-hand side in equation 1, and $Y_{1,i}^*$ is the $(m - m_1 - 1)$ -vector of excluded endogenous variables. Similarly, write

$$Z_i = \begin{pmatrix} Z_{1,i}^\top & Z_{1,i}^{*\top} \end{pmatrix}^\top,$$

where $Z_{1,i}$ and $Z_{1,i}^*$ are the vectors of included and excluded exogenous variables, respectively (for the first equation). Further, write

$$\begin{pmatrix} y_{1i} \\ Y_{1,i} \\ Y_{1,i}^* \end{pmatrix} = \begin{pmatrix} \pi_1 & \pi_2 \\ \Pi_1 & \Pi_2 \\ \Pi_3 & \Pi_4 \end{pmatrix} \begin{pmatrix} Z_{1,i} \\ Z_{1,i}^* \end{pmatrix} + \begin{pmatrix} v_{1i} \\ V_{1,i} \\ V_{1,i}^* \end{pmatrix}.$$

Here, π_2 is the vector of reduced-form coefficients of $Z_{1,i}^*$ in the y_1 equation, and Π_2 gives the coefficients of $Z_{1,i}^*$ in the Y_1 equation. We have

$$\begin{aligned} \hat{\pi}_2^\top &= (Z_1^{*\top} M_1 Z_1^*)^{-1} Z_1^{*\top} M_1 y_1, \\ \hat{\Pi}_2^\top &= (Z_1^{*\top} M_1 Z_1^*)^{-1} Z_1^{*\top} M_1 Y_1, \end{aligned}$$

where

$$\begin{aligned} y_1 &= \begin{pmatrix} y_{11} \\ \vdots \\ y_{1n} \end{pmatrix}, \\ Y_1 &= \begin{pmatrix} Y_{1,1}^\top \\ \vdots \\ Y_{1,n}^\top \end{pmatrix}, \\ Z_1^* &= \begin{pmatrix} Z_{1,1}^{*\top} \\ \vdots \\ Z_{1,n}^{*\top} \end{pmatrix}, \\ Z_1 &= \begin{pmatrix} Z_{1,1}^\top \\ \vdots \\ Z_{1,n}^\top \end{pmatrix}, \\ M_1 &= I_n - Z_1 (Z_1^\top Z_1)^{-1} Z_1^\top. \end{aligned}$$

Hence,

$$\hat{\gamma}_1^{\text{ILS}} = (Z_1^{*\top} M_1 Y_1)^{-1} Z_1^{*\top} M_1 y_1.$$

This is simply the IV estimator from the regression of y_1 on Y_1 and Z_1 , where Z_1^* is used as instruments for Y_1 . Indeed, write the IV estimated equation as

$$y_1 = Y_1 \hat{\gamma}_1^{\text{IV}} + Z_1 \hat{\beta}_1^{\text{IV}} + \hat{u}_1,$$

where $Z_1^\top \hat{u}_1 = 0$ and $Z_1^{*\top} \hat{u}_1 = 0$. Then,

$$\begin{aligned} \hat{\gamma}_1^{\text{ILS}} &= (Z_1^{*\top} M_1 Y_1)^{-1} Z_1^{*\top} M_1 (Y_1 \hat{\gamma}_1^{\text{IV}} + Z_1 \hat{\beta}_1^{\text{IV}} + \hat{u}_1) \\ &= \hat{\gamma}_1^{\text{IV}} + (Z_1^{*\top} M_1 Y_1)^{-1} Z_1^{*\top} M_1 \hat{u}_1, \end{aligned}$$

and

$$\begin{aligned} Z_1^{*\top} M_1 \hat{u}_1 &= Z_1^{*\top} \hat{u}_1 - Z_1^{*\top} Z_1 (Z_1^\top Z_1)^{-1} Z_1^\top \hat{u}_1 \\ &= 0. \end{aligned}$$

The indirect LS estimator is unique only if the equation is exactly identified. In the case of an overidentified equation, one could choose some m_1 restrictions out of $l - l_1$ in (5) to construct the indirect LS estimator. However, a better approach is to use GMM estimation.

Identification and GMM estimation of a single equation

Consider the case where the first equation is identified, that is, $l - l_1 \geq m_1$, and Π_2 has rank m_1 (the rank condition is satisfied). Let us rewrite the first equation again as

$$\begin{aligned} y_{1i} &= Y_{1,i}^\top \gamma_1 + Z_{1,i}^\top \beta_1 + u_{1i} \\ &= X_{1,i}^\top \delta_1 + u_{1i}, \end{aligned} \quad (7)$$

where

$$\begin{aligned} X_{1,i} &= \begin{pmatrix} Y_{1,i}^\top & Z_{1,i}^\top \end{pmatrix}^\top, \\ \delta_1 &= \begin{pmatrix} \gamma_1^\top & \beta_1^\top \end{pmatrix}^\top. \end{aligned}$$

The equation can be estimated by GMM provided that the rank condition for GMM estimation is satisfied, that is, $\text{rank}(\mathbb{E}[Z_i X_{1,i}^\top]) = m_1 + l_1$. For this to hold, the necessary condition is that the number of instruments in Z_i must exceed the number of regressors in $X_{1,i}$: $l \geq m_1 + l_1$ or $l - l_1 \geq m_1$, which gives us back the order condition. Next, we will see that $\text{rank}(\mathbb{E}[Z_i X_{1,i}^\top]) = m_1 + l_1$ is equivalent to $\text{rank}(\Pi_2) = m_1$.

$$\begin{aligned} \mathbb{E}[Z_i X_{1,i}^\top] &= \mathbb{E}\left[\begin{pmatrix} Z_{1,i} \\ Z_{1,i}^* \end{pmatrix} \begin{pmatrix} Y_{1,i}^\top & Z_{1,i}^\top \end{pmatrix}\right] \\ &= \mathbb{E}\left[\begin{pmatrix} Z_{1,i} \\ Z_{1,i}^* \end{pmatrix} \begin{pmatrix} (\Pi_1 Z_{1,i} + \Pi_2 Z_{1,i}^* + V_{1,i})^\top & Z_{1,i}^\top \end{pmatrix}\right] \\ &= \begin{pmatrix} \mathbb{E}[Z_{1,i} Z_{1,i}^\top] \Pi_1^\top + \mathbb{E}[Z_{1,i} Z_{1,i}^{*\top}] \Pi_2^\top & \mathbb{E}[Z_{1,i} Z_{1,i}^\top] \\ \mathbb{E}[Z_{1,i}^* Z_{1,i}^\top] \Pi_1^\top + \mathbb{E}[Z_{1,i}^* Z_{1,i}^{*\top}] \Pi_2^\top & \mathbb{E}[Z_{1,i}^* Z_{1,i}^\top] \end{pmatrix}. \end{aligned}$$

Suppose this $l \times (m_1 + l_1)$ matrix does not have full column rank. This is the case if and only if there exists some nonzero $(m_1 + l_1)$ -vector $\theta = \begin{pmatrix} \theta_1^\top & \theta_2^\top \end{pmatrix}^\top$ such that $\mathbb{E}[Z_i X_{1,i}^\top] \theta = 0$:

$$\begin{aligned} 0 &= \begin{pmatrix} \mathbb{E}[Z_{1,i} Z_{1,i}^\top] \Pi_1^\top + \mathbb{E}[Z_{1,i} Z_{1,i}^{*\top}] \Pi_2^\top & \mathbb{E}[Z_{1,i} Z_{1,i}^\top] \\ \mathbb{E}[Z_{1,i}^* Z_{1,i}^\top] \Pi_1^\top + \mathbb{E}[Z_{1,i}^* Z_{1,i}^{*\top}] \Pi_2^\top & \mathbb{E}[Z_{1,i}^* Z_{1,i}^\top] \end{pmatrix} \begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix} \\ &= \begin{pmatrix} \mathbb{E}[Z_{1,i} Z_{1,i}^\top] \Pi_1^\top \theta_1 + \mathbb{E}[Z_{1,i} Z_{1,i}^{*\top}] \Pi_2^\top \theta_1 + \mathbb{E}[Z_{1,i} Z_{1,i}^\top] \theta_2 \\ \mathbb{E}[Z_{1,i}^* Z_{1,i}^\top] \Pi_1^\top \theta_1 + \mathbb{E}[Z_{1,i}^* Z_{1,i}^{*\top}] \Pi_2^\top \theta_1 + \mathbb{E}[Z_{1,i}^* Z_{1,i}^\top] \theta_2 \end{pmatrix} \\ &= \begin{pmatrix} \mathbb{E}[Z_{1,i} Z_{1,i}^\top] & \mathbb{E}[Z_{1,i} Z_{1,i}^{*\top}] \\ \mathbb{E}[Z_{1,i}^* Z_{1,i}^\top] & \mathbb{E}[Z_{1,i}^* Z_{1,i}^{*\top}] \end{pmatrix} \begin{pmatrix} \Pi_1^\top \theta_1 + \theta_2 \\ \Pi_2^\top \theta_1 \end{pmatrix}. \end{aligned}$$

Assuming that $\mathbb{E}[Z_i Z_i^\top]$ is positive definite, $\mathbb{E}[Z_i X_{1,i}^\top]$ does not have full rank if and only if

$$\begin{aligned} \Pi_1^\top \theta_1 + \theta_2 &= 0, \\ \Pi_2^\top \theta_1 &= 0, \end{aligned} \quad (8)$$

that is, the m_1 columns of $\Pi_2^\top$ are linearly dependent. (θ_1 must be nonzero, since if $\theta_1 = 0$, then $0 = \Pi_1^\top \theta_1 + \theta_2 = 0 + \theta_2 = \theta_2$, which contradicts $\theta \neq 0$.) Hence, $\text{rank}(\mathbb{E}[Z_i X_{1,i}^\top]) < m_1 + l_1$ implies that $\text{rank}(\Pi_2) < m_1$. Consequently, $\text{rank}(\Pi_2) = m_1$ implies that $\text{rank}(\mathbb{E}[Z_i X_{1,i}^\top]) = m_1 + l_1$.

Next, for any $\theta = (\theta_1^\top, \theta_2^\top)^\top$ we have that

$$\left(\mathbb{E}[Z_i X_{1,i}^\top]\right) \theta = \begin{pmatrix} \mathbb{E}[Z_{1,i} Z_{1,i}^\top] & \mathbb{E}[Z_{1,i} Z_{1,i}^{*\top}] \\ \mathbb{E}[Z_{1,i}^* Z_{1,i}^\top] & \mathbb{E}[Z_{1,i}^* Z_{1,i}^{*\top}] \end{pmatrix} \begin{pmatrix} \Pi_1^\top \theta_1 + \theta_2 \\ \Pi_2^\top \theta_1 \end{pmatrix}.$$

Suppose that $\text{rank}(\Pi_2) < m_1$. Then, there exists a nonzero θ_1 such that (8) holds. Next, define $\theta_2 = -\Pi_1^\top \theta_1$. For such a choice of $\theta = (\theta_1^\top, \theta_2^\top)^\top$ we have that $\left(\mathbb{E}[Z_i X_{1,i}^\top]\right) \theta = 0$. Hence, $\text{rank}(\Pi_2) < m_1$ implies that $\text{rank}(\mathbb{E}[Z_i X_{1,i}^\top]) < m_1 + l_1$, and, therefore, $\text{rank}(\mathbb{E}[Z_i X_{1,i}^\top]) = m_1 + l_1$ implies $\text{rank}(\Pi_2) = m_1$. Thus, the two rank conditions are equivalent: $\mathbb{E}[Z_i X_{1,i}^\top]$ has full column rank $m_1 + l_1$ if and only if the rank of Π_2 is m_1 .

We have shown that the GMM rank condition on $E[Z_i X_{1,i}^\top]$ is equivalent to the rank condition on the corresponding matrix of the reduced-form parameters. Thus, provided that the equation is identified, one can use the usual GMM technique to estimate that equation:

$$\tilde{\delta}_1(A_{1n}) = \left(\sum_{i=1}^n X_{1,i} Z_i^\top (A_{1n}^\top A_{1n}) \sum_{i=1}^n Z_i X_{1,i}^\top \right)^{-1} \sum_{i=1}^n X_{1,i} Z_i^\top (A_{1n}^\top A_{1n}) \sum_{i=1}^n Z_i y_{1i}.$$

The efficient A_{1n} is such that $A_{1n}^\top A_{1n} \rightarrow_p \left(E[u_{1i}^2 Z_i Z_i^\top] \right)^{-1}$. The efficient GMM reduces to the 2SLS estimator in the homoskedastic case ($A_{1n}^\top A_{1n} = (\sum_{i=1}^n Z_i Z_i^\top)^{-1}$), which in turn is the same as the IV estimator when the equation is exactly identified.

The estimators of δ_j are asymptotically correlated across the equations, even if different equations in the system are estimated separately. Suppose that the first two equations are identified. We have

$$n^{1/2} \begin{pmatrix} \tilde{\delta}_1(A_{1n}) - \delta_1 \\ \tilde{\delta}_2(A_{2n}) - \delta_2 \end{pmatrix} = \begin{pmatrix} S_{1,n} & 0 \\ 0 & S_{2,n} \end{pmatrix} n^{-1/2} \sum_{i=1}^n \begin{pmatrix} Z_i u_{1i} \\ Z_i u_{2i} \end{pmatrix},$$

where

$$S_{j,n}(A_{jn}) = \left(n^{-1} \sum_{i=1}^n X_{j,i} Z_i^\top (A_{jn}^\top A_{jn}) n^{-1} \sum_{i=1}^n Z_i X_{j,i}^\top \right)^{-1} n^{-1} \sum_{i=1}^n X_{j,i} Z_i^\top (A_{jn}^\top A_{jn}),$$

for $j = 1, 2$. Next, under the usual assumptions

$$\begin{aligned} S_{j,n}(A_{jn}) &\rightarrow_p S_j(A_j) \\ &= (Q_j^\top A_j^\top A_j Q_j)^{-1} Q_j^\top A_j^\top A_j, \end{aligned}$$

where $Q_j = E[Z_i X_{j,i}^\top]$. Further,

$$\begin{aligned} &n^{-1/2} \sum_{i=1}^n \begin{pmatrix} Z_i u_{1i} \\ Z_i u_{2i} \end{pmatrix} \\ &= n^{-1/2} \sum_{i=1}^n \begin{pmatrix} u_{1i} \\ u_{2i} \end{pmatrix} \otimes Z_i \rightarrow_d N(0, \Omega_{1,2}), \end{aligned}$$

where $A \otimes B$ denotes the Kronecker product of A and B ,¹ and

$$\Omega_{1,2} = E \left[\begin{pmatrix} u_{1i}^2 & u_{1i} u_{2i} \\ u_{1i} u_{2i} & u_{2i}^2 \end{pmatrix} \otimes Z_i Z_i^\top \right].$$

Thus,

$$n^{1/2} \begin{pmatrix} \tilde{\delta}_1(A_{1n}) - \delta_1 \\ \tilde{\delta}_2(A_{2n}) - \delta_2 \end{pmatrix} \rightarrow_d N(0, V(A_1, A_2)),$$

where

$$V(A_1, A_2) = \begin{pmatrix} S_1(A_1) & 0 \\ 0 & S_2(A_2) \end{pmatrix} \Omega_{1,2} \begin{pmatrix} S_1(A_1) & 0 \\ 0 & S_2(A_2) \end{pmatrix}^\top.$$

The asymptotic covariance between $\tilde{\delta}_1(A_{1n})$ and $\tilde{\delta}_2(A_{2n})$ is given by

$$(Q_1^\top A_1^\top A_1 Q_1)^{-1} Q_1^\top A_1^\top A_1 E[u_{1i} u_{2i} Z_i Z_i^\top] A_2^\top A_2 Q_2 (Q_2^\top A_2^\top A_2 Q_2)^{-1}.$$

¹Suppose that A is $k \times l$ and B is $m \times n$. Then $A \otimes B$ is a $(km) \times (ln)$ matrix given by

$$A \otimes B = \begin{pmatrix} a_{11} & \cdots & a_{1l} \\ \vdots & \ddots & \vdots \\ a_{k1} & \cdots & a_{kl} \end{pmatrix} \otimes B = \begin{pmatrix} a_{11}B & \cdots & a_{1l}B \\ \vdots & \ddots & \vdots \\ a_{k1}B & \cdots & a_{kl}B \end{pmatrix}.$$

The properties of the Kronecker product include: $(A \otimes B)^\top = A^\top \otimes B^\top$, $(A \otimes B)^{-1} = A^{-1} \otimes B^{-1}$ when A and B are invertible, and $(A \otimes B)(C \otimes D) = AC \otimes BD$ for properly defined matrices C and D .

Appendix: solving a linear system of equations

This discussion follows J.R. Magnus and H. Neudecker (2007), *Matrix Differential Calculus*.

Consider a system of linear equations

$$Ax = b,$$

where A is a known $l \times m$ matrix and b is a known l -vector. We are looking for an m -vector x that solves the system. In the case of a system of simultaneous equations, $A = \Pi_2^\top$, $x = \gamma_1$, and $b = \pi_2^\top$.

A solution can be conveniently expressed using a generalized inverse of A . The generalized inverse of A is a matrix A^- such that $AA^-A = A$; it is generally not unique. The Moore-Penrose inverse of a matrix A (denoted by A^+) is a generalized inverse that satisfies the following conditions:

1. $AA^+A = A$.
2. $A^+AA^+ = A^+$.
3. A^+A is symmetric.
4. AA^+ is symmetric.

A unique A^+ always exists for any matrix A . For example, if A has full column rank, that is, $\text{rank}(A) = m$, then

$$A^+ = (A^\top A)^{-1} A^\top,$$

as can be verified from conditions 1–4.

We will establish below several results useful in characterizing a solution to $Ax = b$. First, let us consider a *homogeneous* system of linear equations

$$Ax = 0$$

(here $x = 0$ is always a solution).

Lemma 1 *A general solution to the homogeneous system of linear equations $Ax = 0$ is*

$$x = (I_m - A^+A)q,$$

where q is an arbitrary m -vector.

Proof. The vector $x = (I_m - A^+A)q$ is a solution since, due to property 1 of a Moore-Penrose inverse,

$$\begin{aligned} A(I_m - A^+A)q &= (A - AA^+A)q \\ &= (A - A)q \\ &= 0. \end{aligned}$$

Next, suppose x is a solution. Then, it has to satisfy $x = (I_m - A^+A)x$, because $(I_m - A^+A)x = x - A^+Ax = x - A^+0 = x$. Thus, any solution can be written as $x = (I_m - A^+A)q$ for some q . ■

Lemma 2 *$Ax = b$ has a solution if and only if $\text{rank}(A) = \text{rank}(\begin{bmatrix} A & b \end{bmatrix})$.*

Proof. Write $A = \begin{bmatrix} a_1 & \cdots & a_m \end{bmatrix}$, where a_i is an l -vector, $i = 1, \dots, m$. Suppose the vector $x = (x_1, \dots, x_m)^\top$ is a solution, that is, $Ax = b$. Then,

$$a_1x_1 + \cdots + a_mx_m = b,$$

and the last column in the extended matrix $\begin{bmatrix} A & b \end{bmatrix}$ is a linear combination of the columns of A . Therefore, $\text{rank}(\begin{bmatrix} A & b \end{bmatrix}) = \text{rank}(A)$.

Now, suppose that $\text{rank}(A) = \text{rank}(\begin{bmatrix} A & b \end{bmatrix})$. Since A is an $l \times m$ matrix, its rank is less than or equal to m . Because $\begin{bmatrix} A & b \end{bmatrix}$ is $l \times (m+1)$ and has rank at most m , its columns are linearly dependent, so there is a nonzero vector $(x_1, \dots, x_m, x_{m+1})^\top$ such that

$$a_1x_1 + \cdots + a_mx_m + bx_{m+1} = 0.$$

If every nonzero vector in the null space of $\begin{bmatrix} A & b \end{bmatrix}$ had $x_{m+1} = 0$, then b would not lie in the column space of A , and $\text{rank}(\begin{bmatrix} A & b \end{bmatrix}) = \text{rank}(A) + 1$, which is a contradiction. Thus, there exists a null vector with $x_{m+1} \neq 0$, and $(-x_1/x_{m+1}, \dots, -x_m/x_{m+1})^\top$ is a solution to $Ax = b$. ■

Lemma 3 $Ax = b$ has a solution if and only if $AA^+b = b$.

Proof. From the definition of A^+ , $AA^+A = A$, and $AA^+Ax = Ax$. Suppose that x is a solution. Then, $Ax = b$, and $AA^+Ax = Ax$ implies that $AA^+b = b$. Now, suppose that $AA^+b = b$. Set $\tilde{x} = A^+b$. Then, $A\tilde{x} = AA^+b = b$, and therefore $\tilde{x}$ is a solution. ■

Next, we describe a general solution to $Ax = b$, provided that it exists.

Lemma 4 If $Ax = b$ has a solution, then it takes the following general form:

$$x = A^+b + (I_m - A^+A)q,$$

where q is an arbitrary m -vector.

Proof. Since there is a solution by the assumption, $AA^+b = b$. By the property $AA^+A = A$ of a Moore-Penrose inverse, $x = A^+b + (I_m - A^+A)q$ is a solution:

$$\begin{aligned} A(A^+b + (I_m - A^+A)q) &= AA^+b + (A - AA^+A)q \\ &= b + (A - A)q \\ &= b. \end{aligned}$$

Now, suppose $\tilde{x}$ is a solution. Then, $A\tilde{x} = b = AA^+b$ or $A(\tilde{x} - A^+b) = 0$. The last equation is a homogeneous system of equations, and by Lemma 1,

$$\tilde{x} - A^+b = (I_m - A^+A)q$$

for some $q \in \mathbb{R}^m$. ■

In the case of simultaneous equations, the identification depends on whether the structural parameters can be solved *uniquely* from the reduced-form coefficients. Here is the main identification result:

Theorem 5 A system $Ax = b$, where A is $l \times m$, $x \in \mathbb{R}^m$, and $b \in \mathbb{R}^l$ has a unique solution if and only if

$$\text{rank} \left(\begin{bmatrix} A & b \end{bmatrix} \right) = \text{rank}(A) = m.$$

Proof. Existence of a solution follows from Lemma 2, so it is left to prove uniqueness. By Lemma 4, the general solution is given by

$$x = A^+b + (I_m - A^+A)q,$$

where q is arbitrary. Thus, uniqueness of the solution is equivalent to $A^+A = I_m$. Suppose that $\text{rank}(A) = m$. In this case, A has full column rank, and $A^+ = (A^\top A)^{-1} A^\top$. It follows that $A^+A = (A^\top A)^{-1} A^\top A = I_m$.

Suppose now that $A^+A = I_m$. Then, $\min(\text{rank}(A^+), \text{rank}(A)) \geq \text{rank}(I_m) = m$. Therefore, since A is $l \times m$, it has rank m . (The rank of A cannot exceed m . If $\text{rank}(A) < m$, then $\text{rank}(A^+A) < m$ and, therefore, $A^+A \neq I_m$.) ■

In the case of simultaneous equations, the reduced-form coefficients are *defined* through the structural parameters, $\Pi = \Gamma^{-1}B$, and therefore there is at least one solution to $\Gamma\Pi = B$. Hence, identification is stated as the rank condition $\text{rank}(\Pi_2) = m_1$.