

Drug Reports → Abaloparatide → Clinical Trials → NCT05073315

Study of Cobicistat-Boosted Atazanavir (ATV/co), Cobicistat-Boosted Darunavir (DRV/co) and Emtricitabine/Tenofovir Alafenamide (F/TAF) in Children With HIV

Timeline	Design	Eligibility	Trends	Safety
----------	--------	-------------	--------	--------

Trial

NCT ID	NCT05073315
Results	False
Phase	Phase 3
Expanded Access	No expanded access
Brief Title	A Comparative Study Between ABP 501 and Humira® in Participants With Moderate to Severe Plaque Psoriasis
Official Title	A Multicenter, Randomized, Double-blind Study Evaluating the Pharmacokinetics, Efficacy, Safety, and Immunogenicity of Multiple Switches Between Humira® (Adalimumab [US]) and ABP 501 Compared With Continued Use of ... More
Sponsor	Amgen (INDUSTRY)
Indication	Plaque Psoriasis
Intervention	Adalimumab, Abp 501

Timeline

Status	NCT05073315
Timeline	
Study First Post Date	2021-10-11
Last Update Post Date	2022-04-15
Start Date	2021-10-04
Primary Completion Date	2023-02-09
Completion Date	2023-02-09
Results First Post Date	—

Design

Study type	Interventional
Enrollment	414 (Anticipated)

Eligibility

Eligibility	Inclusion Criteria	Exclusion Criteria
-------------	--------------------	--------------------

Healthy male or female subjects, between the age of 18 and 55 years, both inclusive (healthy is defined as no clinically relevant abnormalities identified by a detailed medical history, full physical examination, including blood pressure and heart rate measurement, 12-lead ECG, and clinical laboratory tests prior to administration of the study drug). Subject is informed and able to understand the full nature and purpose of the study, including possible risks and side effects, and is given ample time and opportunity to read and understand this information. Subject has the ability and agrees to cooperate with the Investigator and must sign and date the written informed consent prior to performing any of the screening procedures. BMI between 18.0 and 32.0 kg/m2, both inclusive. Subject and their partner of childbearing potential must agree to use a highly effective method of contraception throughout the study and for 5 months after the administration of assigned tre...[More](#)

Go to top