

The nature and origin of cratons constrained by their surface geology

A.M. Celal Sengör^{1,2}, Nalan Lom^{3,†}, and Ali Polat⁴

¹Eurasia Institute of Earth Sciences, Istanbul Technical University, Ayazağa, 34469, Istanbul, Turkey

²Department of Geology, Faculty of Mines, Istanbul Technical University, Ayazağa, 34469, Istanbul, Turkey

³Department of Earth Sciences, Utrecht University, Vening Meineszgebouw A, Princetonlaan 8a, 3584 CB Utrecht, the Netherlands

⁴School of the Environment, University of Windsor, Ontario, N9B 3P4, Canada

To the memory of Nicholas John (Nick) Archibald (1951–2014), master of cratonic geology.

ABSTRACT

Cratons, defined by their resistance to deformation, are guardians of crustal and lithospheric material over billion-year time scales. Archean and Proterozoic rocks can be found in many places on earth, but not all of them represent cratonic areas. Some of these old terrains, inappropriately termed “cratons” by some, have been parts of mobile belts and have experienced widespread deformations in response to mantle-plume-generated thermal weakening, uplift and consequent extension and/or various plate boundary deformations well into the Phanerozoic.

It is a common misconception that cratons consist only of metamorphosed crystalline rocks at their surface, as shown by the indiscriminate designation of them by many as “shields.” Our compilation shows that this conviction is not completely true. Some recent models argue that craton formation results from crustal thickening caused by shortening and subsequent removal of the upper crust by erosion. This process would expose a high-grade metamorphic crust at the surface, but greenschist-grade metamorphic rocks and even unmetamorphosed supracrustal sedimentary rocks are widespread on some cratonic surfaces today, showing that craton formation does not require total removal of the upper crust. Instead, the granulitization of the roots of arcs may have been responsible for weighing down the collated and thickened pieces and keeping their top surfaces usually near sea level.

In this study, we review the nature and origin of cratons on four well-studied examples. The Superior Province (the Canadian Shield), the Barberton Mountain

(Kaapvaal province, South Africa), and the Yilgarn province (Western Australia) show the diversity of rocks with different origin and metamorphic degree at their surface. These fairly extensive examples are chosen because they are typical. It would have been impractical to review the entire extant cratonic surfaces on earth today. We chose the inappropriately named North China “Craton” to discuss the requirements to be classified as a craton.

INTRODUCTION

The purpose of this paper is to point out the kind of surface (or near surface) geological data that may be used to provide *binding constraints* for models proposed for the formation and persistence of cratons. As defined and elaborated on below, cratons are defined *only* by their resistance to deformation, and they may be of any age. Neither age, nor rock composition, nor morphology are parts of the definition of a craton, although cratons of different ages may have certain differences among them. They may preserve crustal and lithospheric material over billion-year time scales. Archean and Proterozoic rocks are present in many places on earth, but not all of them represent cratonic areas. Some of these old terrains, inappropriately termed “cratons,” have been parts of mobile belts and have experienced extensive later deformation. What is and what is not a craton and how they form are still subjects of controversy largely because of the somewhat loose usage of the term and because their field geology has not everywhere been carefully assessed. The purpose of this paper is to review the concept of craton and critically discuss some recent ideas (e.g., Sengör, 1999; McKenzie and Priestley, 2008, 2016; Wang et al., 2014b; McKenzie and Tribaldos, 2018; Capitanio et al., 2020; Priestley et al., 2021) concerning their formation in the light of some relevant field geological observations.

DEFINITION, HISTORICAL BACKGROUND, AND ATTRIBUTES

Definition

It is a common misconception that cratons consist only of metamorphosed crystalline rocks at their surface, as shown by the indiscriminate designation of them by many as “shields,” regardless of how extensive their non-metamorphic sedimentary cover may be (e.g., Gibbs and Barron, 1993; Hartmann and Delgado, 2001; Furnes et al., 2015). According to our present understanding, a craton consists of a piece of continental lithosphere isolated from the convecting mantle and resistant to alpinotype orogenies, i.e., those convergent plate boundary events creating intense penetrative deformation, metamorphism and being accompanied, to various degrees, by island arc-tholeiitic and calc-alkalic magmatism, within the Phanerozoic era. Cratons were also produced during the Archean and the earlier Proterozoic, but all of those are now incorporated into larger cratonic entities that formed in the late Proterozoic to earliest Cambrian interval, i.e., no longer forming independent cratons by themselves (Hoffman, 1988). *What defines a craton is thus only its strength as revealed by the absence of deformation within it while intense deformation may be going on along its margins;* neither its age, nor its composition are parts of the definition of a craton. As we explain below, areas that have commenced, yet not completed, their cratonization (“quasicratons”) for various reasons (e.g., interference from mantle plumes, as in the Saharides: Sengör et al., 2020 and 2021; repeated deformation under the influence of the events around their periphery before cratonization was completed, as in Central and Western Europe: Küpper, 1965; Sengör, 1995; Kley, 2013; or in the U.S. Rocky Mountains: Burchfiel and Davis, 1975; Snyder et al., 1976 and Conney and Reynolds, 1977; also see Sengör et al., 2018a; or in Central and Eastern Asia: Sengör

Nalan Lom  <https://orcid.org/0000-0002-5553-4937>

†Corresponding author: nalanlom@gmail.com

and Natal'in, 1996; Calais et al., 2006; Tunini et al., 2017; also see below etc.) usually resist alpinotype orogeny, but may be deformed by germanotype orogeny¹, i.e., non-penetrative block uplifts and subsidences of the type of the U.S. Rocky Mountains or the Boothia Uplift in Canada or the Harz Mountains in Germany or the Cainozoic Tien Shan in Central Asia. Such areas may be cut by major keirogens, i.e., belts of strike-slip deformation, such as the Central African Shear Zone or the Transbrasiliense lineament, but only if the craton is of late Proterozoic age. A fully developed craton cannot thus be penetrated by mechanical means; it may be completely destroyed only by heating from below by mantle plumes to remove its protective lithospheric armor (Şengör, 2001). Although some popular earth science dictionary definitions restrict the craton to the continental crust (Neuendorf et al., 2005; Allaby, 2008), today it is well known that formation and preservation of a craton is intimately linked to the composition, thermal state, and state of stress of its underlying lithosphere, although these factors are *not* a part of the *definition* of a craton, but its currently accepted *explanation*, as emphasized above (Şengör, 1999; Aulbach, 2012). Thus, in forming a craton, the histories of their crust and the

underlying subcontinental lithospheric mantle are generally coupled in space and time.

A Terminological Problem

It should also be kept in mind that when a craton is identified it must be specified to what time period the definition applies. For example, the designation Slave Craton in Canada applies only to an interval between 2.69 and 1.88 Ma, because after that date it became only an integral part of a larger cratonic entity, the nuclear Canadian/Greenland craton. That entity was then enlarged to include the Yavapai and Mazatzal orogens in the south and the Grenville orogen in the east, finally forming the entire Laurentian Craton (Suess, 1909, p. 284²), including Greenland, until the Paleocene-Eocene opening of the Davis Strait/Baffin Bay arm of the northern Atlantic Ocean: (e.g., Agranier et al., 2019). Similarly, the Dharwar Craton in India was an independent craton until it was framed by the Proterozoic orogenic belts that finally formed a part of the Gondwanian Craton by the Cambrian period. Speaking of a Slave Craton or a Dharwar Craton today would be like calling a twig a complete tree. It would be more informative and less confusing if such well-defined, smaller (and commonly older) areas in larger cratons areas are referred to as tectonic provinces—as is done by some geologists working in Canada (e.g., Heaman and Pearson, 2010)—within the present-day cratons or blocks with the necessary qualifiers, because they are not now independent cratonic structures, but parts of larger cratons. This terminological problem is not irrelevant to the discussions on the nature and origin of cratons, because Archean and Proterozoic parts of the present-day cratons have certain differences both in structure and evolutionary path. Bearing in mind the discussion presented in this section would also help to avoid such inapposite terms as the “Euxinic Craton” referring to the tiny Moesian Platform with a complex Phanerozoic deformational history (e.g., Balintoni and Ballica, 2016).

Brief History

The observation that large areas on continents have been free of any strong visible deformation since at least the beginning of the Cambrian seems to have been already made in the first half of the nineteenth century. It had been mostly attributed to the inability of the earth’s

internal heat to reach into the higher regions. First, Murchison in 1840 (Collie and Diemer, 2004) during his expedition in western Russia, then James Hall (Hall, 1842, 1858), after studying the geology of the Midwest of the USA, noticed considerable expanses of horizontal and undeformed Phanerozoic formations occupying very wide areas. Murchison thought, following a conversation with the German geologist Baron Leopold von Buch, that the absence of deformation was a consequence of the strength of the underlying “volcanic” rocks presumed to underlie those visible at the surface. It was Dana, who considered the interior of continents as “comparatively stable areas” (Dana, 1863, p. 734–735) in contrast to the oceans that subsided under the influence of the earth’s contraction and compressed the continents between them. Within continents, Dana distinguished a stable center from less stable margins.

At this early stage, before the term craton was coined, what was found surprising was the absence of surface deformation in large areas. That was the observation everyone agreed on, but its interpretations varied. While many, such as von Buch, Murchison, and Dana thought the absence of deformation was because the underlying rocks were strong and thus resistant to deformation, others, such as James Hall, thought, quite incomprehensibly, that the flat-lying sedimentary rocks were underlain by areas less resistant than mountainous areas (Hall, 1859, p. 86–87).

The more direct path leading to the invention of the term “craton” via the idea of “consolidation” supposedly creating strong areas, begins with Eduard Suess (1875). The first definition of consolidation (“Erstarrung” in Suess’ publications, which his English translator Hertha Sollas rendered as rigefaction, i.e., stiffening: Suess, 1909, p. 625) was based on the character of the ground. The more contracted or stiffened parts of the earth’s crust caused by cooling and shrinkage were thought to be more resilient against shortening deformation. Between them lie the weaker zones, which make up the geosynclines, i.e., the mother-throughs of future mountain chains, an idea borrowed from Élie de Beaumont (1828a, 1828b, 1852) and Dufrénoy and Élie de Beaumont (1848) via Dana (1847a, 1847b, 1863) and James Hall (1859). In 1883, Suess showed that on continents, mountain belts are found bordering vast areas of deformed and intruded Precambrian rocks covered by flat-lying Phanerozoic deposits that show no orogenic deformation since the Cambrian. He called such places “Tafel” (=Table), his definition of them was based strictly on the geology seen at the surface. This term became “plate-forme” in the French translation and then translated into English as “platform.” Suess’ original interpretation

¹Orogeny is the term that covers the entirety of processes related to subduction and collision of buoyant lithospheric rafts including continents (Şengör, 1990, 2020). However, under orogeny, commonly the processes that create the great orogenic belts of our world are understood generally involving penetrative deformation, metamorphism and widespread, generally calc-alkalic magmatism. Understood as such, this term alone would leave out such lithospheric and crustal structures as the U.S. Rocky Mountains, the Cainozoic Tien Shan, the Cretaceous Harz Mountains, or the Jurassic Donbass structures. For that reason, it seems helpful to use the terms alpinotype for the orogenic structural ensembles that create generally penetrative deformation, widespread regional metamorphism and calc-alkalic magmatism and germanotype for those orogenic structures that create generally non-penetrative block uplifts and basins with little to no regional metamorphism and generally alkalic magmatism. These terms were defined by Hans Stille in 1920 to emphasize the difference between the structures in the Alps and the structures in Central Europe that formed in response to the same orogenic events (Stille, 1920). He later called the U.S. Rocky Mountains “germanotyp disloziertes Felsengebirge” (=germanotype-dislocated Rocky Mountains) as opposed to the “alpinotyp disloziertes Felsengebirge” (=alpinotype-dislocated Rocky Mountains, which are now generally referred to as the Sevier belt: Stille, 1940, fig. 59 on p. 242; notice here that Stille uses the designation “Rocky Mountains” for the entire North American Cordillera as recommended by Powell (1876, p. 4–5), but no longer followed). The distinction alpinotype versus germanotype has been found useful even in extraterrestrial geology (e.g., Byrne et al., 2014; also see Şengör, 2018).

²Inspired by the “Laurentian surface” of Lawson (1890) and “Laurentian Peneplain” of Wilson (1903). For the usage of the adjective “Laurentian,” see especially Wilson’s footnote 2 on his p. 616. Suess took Wilson’s “Laurentian” and used it to denote the entire North American craton.

followed Dana's: areas of "tables" were regions where cooling had progressed the most (Suess, 1875).

Suess later changed his ideas of the nature of rigefaction and held responsible subsidence of the folded neighboring regions for the cessation of the orogenic stresses in the "table" areas. He emphasized that it was not demonstrated that the disappearance of deformation from such areas was a consequence of advanced cooling. Suess was thus inclined to ascribe the absence of deformation to absence of stresses, not to any increase in local strength of rocks. His student Leopold Kober (1921, p. 8), disagreed with this later interpretation by Suess and thought that the disappearance of deformation in certain areas was a result of the stiffening of the underlying rocks because of preceding orogenic deformation. He thus called such, what he considered *strong*, areas *Kratogen* (usually anglicized as cratogen) and divided the earth's lithosphere into three realms: cratogens, orogens and geosynclines. The term cratogen Kober derived from the two Ancient Greek words κράτος (kratos) meaning might, strength, also act of strength, and γένεσις (genesis) meaning origin, manner of birth, creation, so the term cratogen means "born strong."

The German geologist Hans Stille agreed with Kober and originally adopted his terminology, but in 1933 introduced the shorter version, *Kraton* (anglicized as craton), for what Kober had called *Kratogen* (Stille, 1936) and defined stiffened grounds as a consolidation state that no longer allows alpinotype deformation (i.e., penetrative and intense deformation, commonly accompanied by calc-alkalic magmatism and regional metamorphism) to occur but only germanotype deformation (i.e., non-penetrative deformation creating block uplifts or subsidences). According to Stille's tectonic theory, which is fully developed on the example of the structure of the Americas (Stille, 1940), the tectonic state of the ground is divided, like Kober's earlier classification, into three mobile areas occupied by orthogeosynclines, and they supposedly generated alpinotype orogens characterized by penetrative deformation. They form the large mountain belts of our planet. Partially or semi-consolidated areas ("quasicratons" in his terminology) can only localize germanotype orogeny involving non-penetrative block faulting, and fully consolidated areas allow no orogeny at all (Fig. 1).

Thus, the starting point of the creation of the concept of craton was the surface geology. In fact, the word craton is used today to designate large parts of the lithosphere that show no alpinotype orogenic deformation at least in the Phanerozoic. The terms "Tafel," "platform," and "craton" were used as their equivalents in the past, which caused a misunderstanding in the

literature, such as East European Platform and East European Craton used as equivalents by different authors. It is important to use these terms correctly, because they denote distinct entities.

As used today, a *shield* (Suess, 1883) refers to large areas of ancient, partly crystalline but invariably strongly deformed rocks of Precambrian age that are exposed (because of subsequent erosion) in cratons, and a *platform* defines regions where the structural basement rocks are covered by horizontal or gently tilted sediments (Fig. 2). An old and rarely used Scandinavian term *glint* represents the boundary line between shields and platforms. Until Suess generalized its usage outside Scandinavia, the term *glint* (often in the form of *klin*) was confined to Scandinavian and circum-Baltic usage designating a mostly limestone questa resting on the Precambrian crystalline rocks of the shield. In this paper, we follow Suess' terminology while describing the cratonic areas. When we look at the historical development of the idea reviewed above, it is remarkable how they were formulated very early on in the history of geology, yet they could not always stick, because they commonly lacked a generally agreed upon theoretical foundation in which they could be adequately explained.

Over the past few decades, extensive research on cratons revealed common characteristics even though they experienced different overprinting by younger tectonic events. We here follow Stille's classification of cratons as follows:

- No craton: Any sort of deformation possible.
- Quasicraton: Alpinotype deformation no longer possible, but germanotype deformation still possible.
- Full craton: No copeogenic³ deformation is possible, but falcogenic deformations are possible (e.g., the uparching of the Canadian and the Baltic shields after the end of glaciations).

But the full cratons also show two subtypes: the not so fully consolidated ones still show a weakened form of germanotype deformation (Marshak and Paulsen, 1996, 1997; Marshak et al., 1998), whereas full consolidation no longer allows any copeogenic deformation. Geophysical studies show that the lithospheric thicknesses not only of different cratons, but also various types of cratonization within a single cratonic area, vary (see Şengör, 1999). Archean lithosphere is usually 200–220 km thick, where-

as the Proterozoic lithosphere can reach up to 300–350 km thickness (Jordan, 1975, 1988; Artemieva, 2006). Whereas the sub-cratonic keels are thinner in the southern hemisphere cratons (e.g., South Africa, Western Australia, South America, and India), thicker keels are found in the northern hemisphere (e.g., Baltic Shield, Siberian Platform, West Africa, and possibly the Canadian Shield) (Artemieva and Mooney, 2001, 2002; Artemieva, 2019) (Fig. 3). In cratons, Archean crust is generally thinner than the Proterozoic crust (27–40 km versus 40–55 km) (Baranov and Bobrov, 2018).

Cratons have a neutral or positive buoyancy that prevents them from sinking into the mantle. Although they are underlain by thick lithosphere, their top surfaces are usually near sea level. In the more stable areas, the long-term mean rate of denudation of the cratonic surfaces has been on the order of 1.5–4 m per million years assuming a static sea level; if the sea level is rising, the denudation takes a much longer time; if it is falling, a peneplain in the Davisian sense can never form on a craton (Pitman and Golovchenko, 1991; Flowers et al., 2006; Hall et al., 2021). The present-day elevations above sea level of the world's cratons do not exceed 1000 m, except in southern Africa (Zimbabwe, Kaapvaal and their continuation in Antarctica, Grunehogna), where the recent high topography has been attributed to the dynamic uplift of the proposed the Bouvet-Karoo mantle plume (Burke, 1996; Lithgow-Bertelloni and Silver, 1998; Burke and Gunnell, 2008; Jacobs, 2009; Artemieva and Vinnik, 2016) and in the Nilgiri Hills in southernmost India, where younger transpression has thickened the crust and raised the average elevation to above 2 km and crustal thickness to 52 km (Şengör and Natal'in, 1996; Das et al., 2015; Praseeda et al., 2015). The rest of the cratons reside between 100 and 600 m elevation where they commonly exhibit low grade to non-metamorphic rocks coeval with the time of cratonization. This indicates the cratons as a whole were never sites of high elevation and had little erosive removal after they stabilized.

RECIPES FOR A CRATON

In order to produce a craton, some conditions must be met. It requires: (a) depletion of volatiles from the sub-cratonic upper mantle, which helps the subcontinental mantle to be buoyant compared with the oceanic mantle (Pollack, 1986; Carlson et al., 2005; Capitanio et al., 2020; Priestley et al., 2021; but see Wang et al., 2014b); (b) geothermal gradient must be low (Ranalli and Murphy, 1987); (c) radioactive elements must be differentiated and depleted in




Figure 1. The North American Craton showing its constituent orogens and areas of different degrees of consolidation and their correspondence with the age of last orogeny (modified from Clouzet et al., 2018, figure 1). B.H.T.—Buffalo Head Terrane; G.B.A.—Great Bear Arc; Grt. Rhy—Granite-Rhyolite Province; Kwn.R—Keweenawan Rift; L.B.A—Little Belt Arc; Med. Hat—Medicine Hat Block; T.H.O.—Trans-Hudson Orogen; WY—Wyoming; RW—reworked; coll.—collisional; Juv.—juvenile; Qu. Depos.—Quartzite deposits.

order to decrease heat production at the base of the cratonic crust (Jaupart and Mareschal, 1999; Michaut et al., 2009), and (d) continental crust of intermediate or felsic composition must be generated in a substantial volume, probably through arc-like processes (Taylor and McLennan, 1985, 1995; Şengör et al., 1993; Şengör and Natal'in, 1996; James and Fouch, 2002).

According to Şengör (1999), a craton begins forming in a typical, evolving, subduction-accretion system with its underlying depleted mantle wedge (Fig. 4A). When continental collision occurs involving such a wedge (Fig. 4B), it thickens the lithosphere and the crust by shortening, but the predominantly mafic lower part pulls down and creates a lithospheric keel as recently shown by the preferred orientation of the fast axes of

the olivine crystals in the subcontinental mantle (Priestley et al., 2021). Şengör (1999) pointed out that this does not give rise to much uplift of the surface of the crust, because of extensive granulitization of the lower crust increasing its density to $\geq 3 \text{ g/cm}^3$ (Fig. 4C). The formation of granulite-facies rocks increases the strength, and it becomes extremely difficult to destroy the cratons they underlie (Hawkesworth et al., 2010). This increase in weight prevents the rise of the surface elevations.

There is increasing evidence indicating that subduction-accretion processes were active during the earliest Archean, even perhaps in the latest Hadean (Harrison, 2009, 2020; Windley et al., 2021; Hyung and Jacobsen, 2020; Turner et al., 2020; Korenaga, 2021). Trace-element

characteristics of Archean tonalite-trondhjemite-granodiorite gneisses resemble those predicted for Phanerozoic subduction-derived granitoids and thus are consistent with the growth of Archean continental crust at convergent plate margins (Arndt, 2013; Windley et al., 2021). In addition, the presence of ophiolites, but mainly ophirags, in many Eoarchean to Neoarchean Archean greenstone belts (Kusky, 1989; Armstrong et al., 1990; Polat et al., 2002, 2008; Hofmann and Kusky, 2004; Smithies et al., 2005; Mukhopadhyay et al., 2012; Turner et al., 2014; Furnes et al., 2015; Grosch and Slama, 2017) and the lithospheric structure derived from S-wave receiver functions suggest cratonic lithosphere was formed by arc accretion (Miller and Eaton, 2010). De Wit et al. (1992) favored the




Figure 2. Cratons of the world. The dark gray color indicates the entire craton, i.e., the shield(s) plus the surrounding platforms. The light gray color indicates areas listed as cratons in the literature but do not meet the criteria. AU—Australian Craton; BC—Brazilian Craton; IC—Indian Craton; NAC—North American Craton; SC—Siberian Craton; SSC—Sub-Saharan Craton; Y—Yangtze Craton; WAC—West African Craton. The constituent older cratons commonly mentioned in the literature, that are now no longer independent cratons, but simply provinces within the larger cratonic entities: (1) Dharwar; (2) Bastar; (3) Singhbhum; (4) Aravalli; (5) Bundlkhand; (6) North China (this has never been a craton as we discuss herein); (7) Aldan; (8) Anabar; (9) Sarmatia; (10) Karelia; (11) Kola; (12) Nain; (13) Superior Province; (14) Hearne; (15) Sask; (16) Rae; (17) Slave; (18) Wyoming; (19) Guyana; (20) São Luis; (21) Amazonia; (22) São Francisco; (23) Rio de la Plata; (24) Congo; (25) Tanzania; (26) Zimbabwe; (27) Kaapvaal; (26 and 27) Kalahari; (28) Madagascar; (29) Pilbara; (30) Yilgarn; (31) North Australia; (32) Gawler; (33) Vestfold; (34) Napier; (35) Grunehogna.

subduction process over plume origin based on the Archean age metamorphosed peridotite and basalt (eclogite) xenoliths from Kaapvaal province (Shirey et al., 2004). Nothing else but subduction seems capable not only in placing oceanic crust on to the overriding plate or carrying material to great depths, but also in the ability to bring them back to the surface in an orderly fashion, which no diapiric process, as defended by Hamilton (2007, 2019)⁴, for example, would be capable of doing. Hydrated mineral inclusions

in the Hadean Jack Hills zircons, Western Australia, indicate their host protoliths were exposed at the earth's surface in the presence of water (shales, probably implying subduction-accretion material around sialic “rockbergs” perhaps not too dissimilar to “subduction zones” seen in present-day lava lakes: see Harrison, 2009) and then transported to a depth to form parenting orthogneiss (Mojzsis et al., 2001; Harrison 2009). A recent review of the geochemistry of zircons from Jack Hills reveals important petrogenetic information that implies the initiation of subduction zone processes as early as 3.6 Ga (Ackerson et al., 2021).

Kusky (1993) suggested that Archean lithospheric mantle formed beneath arcs by accretion of oceanic lithosphere and trapped subarc mantle wedge. Using major element and Re-Os isotope data from cratonic mantle xenoliths, Servali and Korenaga (2018) proposed that Archean cratonic

mantle was derived from highly depleted oceanic lithosphere that originated beneath mid-ocean ridges or similar spreading settings. Higher ambient mantle temperatures in the Archean likely caused higher degrees of partial melting in the oceans (but not in the continents: Burke and Kidd, 1978), resulting in highly depleted oceanic residual lithospheric mantle that had olivines with high forsterite (92–94) contents (Herzberg, 2018). If Servali and Korenaga’s geochemical interpretation is correct, then the accretion (underthrusting) of such depleted oceanic lithosphere beneath Archean oceanic island arcs could have formed the nuclei of the cratonic mantle, and the accretion of these island arcs eventually would have led to the formation of first larger continents and cratons around 3.2 Ga (Windley et al., 2021). The light rare earth element, large ion lithophile element, and Pb-enriched, but Nb-Ta-depleted trace-element systematics

⁴Warren Hamilton, one of the giants of geology in our times, passed away on 26 October 2018, leaving behind an almost complete manuscript of this paper, lacking only the abstract and the conclusions, which Gillian Foulger completed and prepared for publication, earning the eternal gratitude of the geological community. We disagree with Hamilton’s view of the Precambrian tectonics, but his immense knowledge and keen insights have helped us in sharpening our ideas.




Figure 3. Distribution of the cratons and their Archean rocks (shown in black) compared with effective elastic thickness (T_e) of the lithosphere (km) based on the yield strength envelope (Tesrauro et al., 2012). Thick black line indicates borders of cratons as shown in Figure 2. Thin dark gray line indicates regions that are previously thought to be cratons.

of cratonic mantle suggest that accreted oceanic lithosphere beneath Archean island arcs was later overprinted by subduction-derived melts and/or fluids (Polat et al., 2018).

According to McKenzie and Priestley (2008) two possible ways of lithosphere thickening can lead to craton formation. Figure 5A demonstrates homogeneous crustal thickening with constant heat flow at the base of the lithosphere, Figure 5B represents shortening by pure shear, a uniform crustal thickening resulting from bulk shortening, consistent with the model by Sengör (1999; see Fig. 4 herein) and the observations in Priestley et al. (2021). In order to prevent delamination, the mantle lithosphere must be depleted during a previous melting. Figure 5C shows crustal thickening by emplacing a thick crustal layer on top of a craton as a result of thrusting or lower crustal flow. All these model cases are followed by a thermal relaxation phase (Le Pichon et al., 1997). McKenzie and Priestley's (2008) calculation shows that the temperature of the middle crust can rise over 900 °C in 60 Ma because of the crustal radioactivity unless extension occurs. This process ends up creating high pressure and high temperature rocks as already envisioned by Argand in 1922, when he postulated, within the framework of Wegener's theory of continental drift, a large-scale underthrusting of India under Asia (Argand, 1924; see Fig. 6). Note however, that Argand's Tibetan underthrusting does not account for the entire crustal thickening under Tibet; under the Kuen-Lun, he also postulated shortening and thickening of Asian crust.

It has been proposed recently that Tibet is an actualistic analogue of how cratons begin forming (McKenzie and Priestley, 2008, 2016; McKenzie and Tribaldos, 2018). The presence of a thick lithosphere beneath Tibet is inferred using surface wave tomography employing Rayleigh waves, and it indeed resembles that beneath cratons. Although attractive, this model implies that a substantial amount of the thickened crust, nearly 35–40 km, must somehow be removed to bring its surface to near sea level, where most major Archean cratons reside in the present day. Wholesale underthrusting of the Indian plate beneath the entire Tibetan plateau (e.g., Argand, 1924; Powell and Conaghan, 1973; Zuza et al., 2019), would create a persistent buoyancy that would cause erosion, presumably down to sea level, of the crust and when it stabilizes it would reveal only high-grade rocks, namely amphibolite to granulite grade. Based on the possible density distribution of the Indian plate beneath the Tibetan plateau (Hétényi et al., 2007), we have generated a scenario to estimate the metamorphic facies related to a given depth. Figures 7A and 7B show the distribution of metamorphic facies with depth and erosional (tectonic or attritional) surfaces with corresponding rock assemblages that will be exposed. A similar thing would happen if the shortening in Tibet is accommodated purely by bulk shortening of the plateau lithosphere as envisaged by Dewey and Burke (1973), which is also compatible with the observations by Priestley et al. (2021). We now know that a combination of these two processes seems to have happened, the Indian lithosphere having

been claimed to reach as far north as the Banggong Co-Nu Jiang suture, i.e., to the boundary between the Lhasa and the East and West Qiang Tang arc fragments (Zuza et al., 2019). However, the growth of Tibetan topography from its center outwards (Wang et al., 2014a) makes a wholesale Indian underthrusting difficult as does the old observation of the antecedent rivers in the Himalaya (e.g., Medlicott and Blanford, 1879, p. 676–677; Valdiya, 1996), favoring more a manner of bulk crustal thickening as suggested by Dewey and Burke (1973). This is also supported by both widespread folding as clearly visible on Google Earth images of, and some thrusting on, the Lhasa Block (e.g., Sengör, 1981) contrary to some recent claims. The recent observation by Wang et al. (2021) that the entire 80 km thickness of the Tibetan crust in southern Tibet under the Lhasa block has a felsic composition indicates a crustal structure not unlike that present now under the Altiplano in the central Andes, where the entire 65-km-thick crust seems to consist of quartz-rich, felsic rocks (Zandt et al., 1996). In Tibet, a thick magmatic arc crust of the Lhasa Block that includes the Transhimalaya may be underlain by thinned continental margin rocks of India, where shortening results from the convergence that thickens both the crust and the underlying mantle parts of the lithosphere both by thrusting and bulk shortening. However, this mechanism can only operate in length scales less than 1000 km and cannot explain vast cratonic areas generated at the same time interval unless the unlikely scenario of having more than three contiguous Tibet-type plateaux




Figure 4. A cross-sectional model showing how to build a craton by subduction-accretion (Sengör, 1999). A case of homogeneous crustal thickening with constant heat flow at the base of the lithosphere. We added the strain ellipses to indicate the parallelism of Sengör's (1999) interpretation with that of Priestley et al. (2021, fig. 3). In A, the letters a, b, c, and d signify progressively younger subduction accretion prisms together with their now abandoned arcs added to the continent, thereby enlarging it.

that are assumed to form simultaneously with a unified 3000 km underthrusting! By contrast, the largest tectonic detachment underlying an Archean upper crustal assemblage we know of

is that imaged in the Yilgarn Province in Western Australia (Ahmat et al., 1993), the width of which is barely 150 km, and in it such shear zones as that of Bardoc penetrate the sub-de-

tachment basement for more than 20 km across strike (Ahmat et al., 1993). We show below, using surface geological observations, that some Archean cratons had little erosive removal after




Figure 5. Schematic drawings of 2-D scenarios corresponding to (A and B) the uniform thickening of crust and lithosphere and, (C) crustal stacking model based on Le Pichon et al. (1997) and McKenzie and Priestley (2008).

they stabilized, and none ever was a Tibet-type high plateau as a whole.

MYTHS AND FACTS

Many consider that most cratons consist only of polydeformed and metamorphosed crystalline rocks on their surface, as shown by the com-

monly indiscriminate usage of the term "shield" as mentioned above (e.g., Gibbs and Barron, 1993; Hartmann and Delgado, 2001; Furnes et al., 2015). In North America, in the Superior Province and the Slave Province, in India, in the Dharwar, in the Baltic, in Karelia, in the Amazonian, Kaapvaal and Zimbabwe provinces in Africa, and the Pilbara and Yilgarn provinces in

Australia, not only high-temperature–high pressure facies metamorphic rocks of Archean age, but also coeval low-grade or even completely unmetamorphosed and only gently deformed sedimentary rocks crop out at their surface (Table 1). When viewed from this angle, none of the cratons listed above show a uniform Tibet-like structure in their make-up.

The Superior Province, the Canadian Shield

The Superior Province represents the largest Archean exposure on earth, which is surrounded by Proterozoic orogenic belts together forming the Laurentian or the North American Craton (Hoffman, 1988). It consists of a large variety of plutonic, volcano-plutonic, and sedimentary rocks plus their metamorphic equivalents from subgreenschist to granulite facies (Card, 1990; Percival et al., 2012). These rocks are disposed in east-west-trending subprovinces ranging from 3.8 to 2.67 Ga (Polat and Kerrich, 2001; Percival et al., 2012). It is widely accepted that the dominant structural trend was created by subduction-accretion processes during the late Archean (Poulsen et al., 1992; Wyman et al., 2002; Percival et al., 2012) and cratonization took place between 2680 and 2600 Ma (Percival et al., 2012). Among these subprovinces, Wawa-Abitibi is the largest domain of juvenile rocks in the Superior Province and shows the lowest degree of metamorphism nowhere exceeding 4 kb (corresponding to ~12 km depth). The majority of the Superior Province is characterized by greenschist facies rocks. Figures 8 and 9 show examples of sedimentary and volcanic rocks, respectively, metamorphosed in greenschist facies from six different widely separated localities in the Superior Province. Notice that the rocks shown are deformed by medium to zero magnitude strain within the frame of our pictures. None could have possibly come from a depth of 30–40 km as required by the craton formation model of McKenzie and Priestley (2008, 2016) and McKenzie and Tribaldos, (2018).



Figure 6. Cross-section of Tibet (Argand, 1924).




Figure 7. (A) Limits of the metamorphic facies as a function of temperature and pressure. Modified from Bousquet et al. (1997). Orange dashed lines indicate the range of adiabatic thermal gradient, gray band delimits the wet melting zone. Red dashed line indicates the average adiabatic thermal gradient that have been used in our calculations. (B) Metamorphic facies based on the average adiabatic thermal gradient shown with red dashed line in A. In the chart, horizontal red solid lines indicate facies boundaries, vertical black lines indicate adopted crustal thickness and related surface uplift. Blue dashed line corresponds to the erosion surface.

The Barberton Mountain, the Kaapvaal Province

The Kaapvaal province of South Africa is one of the oldest Archean segments in the cratons on the earth. The cratonization started around 3.1 Ga when major granitoid batholiths intruded and essentially has been completed by ca. 2.6 Ga (Nguuri et al., 2001). The Archean rocks of Kaapvaal were subsequently affected by magmatic and accretionary events. It is surrounded

by Proterozoic mobile belts: on the south and east by the Namaqua-Natal belt (1.1–1.9 Ga), on the north by the Limpopo belt (2.3 Ga), and on the west by the Kheis overthrust belt (ca. 2.0 Ga). The north-central Kaapvaal craton was disrupted at ca. 2.05 Ga when the largest known layered mafic intrusion in the world, the Bushveld Complex, emplaced into a stable cratonic setting.

The Barberton Greenstone Belt (3.2–3.53 Ga) of eastern Kaapvaal consists of well-preserved Paleoarchean volcanic and sedimentary rocks

generally at low metamorphic grade and in some places entirely devoid of penetrative deformation (Grosch et al., 2011; de Wit et al., 2011). One of the best examples is the Onverwacht Group which includes the Komati Formation in the lower levels and the Hooggenoeg, Kromberg and Mendon formations in the upper levels (de Wit et al., 2011; Grosch and Slama, 2017). Within the Onverwacht Group the Hooggenoeg Formation comprises a 2.5–3.0-km-thick section of predominantly tholeiitic and komatiitic basalts

TABLE 1. PROVINCES IN CRATONS SHOWING LOW-GRADE METAMORPHISM

Province	Age	Unit name	Lithology	Metamorphic grade	Source
Kaapvaal	3.07 Ga*	Dominion Group	Sandstone along with minor conglomerate and argillaceous horizon followed by thick succession of mafic-intermediate lavas interlayered with felsic lavas and pyroclastic rocks	Lower greenschist facies and minor deformation	Burke et al. (1986); Agangi et al. (2020)
Zimbabwe	2.7 Ga*	Upper Greenstone	Conglomerate, sandstone, siltstone, limestone, cherty limestone, stromatolitic limestone, and minor banded iron formation	Low-grade (sericite and chlorite)	Kusky and Kidd (1992)
Dharwar	2.7 Ga*	The Talya Conglomerate	Conglomerates interbedded with mudstone and sandstone	Greenschist facies	Ojakangas et al. (2014)
Superior Province	2.7 Ga*	Wawa-Abitibi	Calc-alkalic to alkalic metavolcanic and metasedimentary rocks		
Greenland	3.8 Ga†; 2.7 Ga‡	Isua supracrustal belt	Metabasites	Pumpellyite-amphibolite facies	Percival et al. (2012); Lodge et al. (2013)
Yilgarn	2.7–2.6 Ga*	Diemals Formation	Siliciclastic sedimentary rocks	Greenschist to amphibolite facies	Hayashi et al. (2000); Morris et al. (2007)
Pilbara	3426–3350 Ma*	Strelley Pool Formation	Laminated carbonate rock, stromatolitic carbonate, evaporites, sandstone, conglomerate, volcanic ash, volcaniclastic sedimentary rocks	Greenschist-facies	Hickman (2008)
Brazilian Shield	2758 ± 39 Ma†	Grão Pará Group	Basaltic to rhyolitic metavolcanic units at the base, overlain by banded iron formation, and topped by an alluvial sedimentary sequence	Lower-greenschist	Olszewski et al. (1989)
Karelia	2853 ± 11 Ma†	South Vygozersky Greenstone	Plagiogranites	Low to medium grade	Myskova et al. (2015)

*Age of deposition.

†Age of crystallization.

‡Age of metamorphism.

at the base whereas the upper part is dominated by felsic volcaniclastic conglomerates, sandstones, and tuffs. Figure 10 shows various outcrop photographs all taken along its course, and the little to no strain they exhibit is shown particularly well by the well-preserved immiscibility spherules seen in the pillow lavas (Fig. 10A) and the preserved sedimentary structures (Figs. 10B, 10C). Figure 10D shows pillow lavas from the Kromberg Formation which is described as a thick, continuous sequence of komatiite and basaltic volcanic rocks capped by carbonated shallow-water volcaniclastic sediments and cherts. The pillow lavas rest conformably on carbonated tuff. Along the Komati River, both the volcanic rocks and the sedimentary rocks show no metamorphism and very gentle deformation. The low-grade to unmetamorphosed rocks exposed at the surface are inconsistent with Barberton Mountain's having had a full crustal thickness on top of it as required by the model of craton formation by McKenzie and Priestley (2008, 2016) and McKenzie and Tribaldos (2018), and the common misconception that cratons consist typically of metamorphic rocks at their surface.

The Yilgarn Province, Western Australia

The Yilgarn province (Fig. 11) consists mainly of plutonic, volcanic, and sedimentary Archean rocks that formed between 3.73 and 2.60 Ga ago and, with the exception of southwestern Yilgarn high-grade province, was metamorphosed at low grade and in places not at all (Myers, 1993). In the Kalgoorlie region, around the western shores of the great salt pan of Lake Lefroy, there is a highly folded turbidite section sitting on a sequence of tholeiitic basalts and komatiites with interlayered albite cherts (Kalgoorlie Group:

Swager et al., 1992; Nick Archibald, personal commun. in Sengör and Natal'in, 1996). Above these rocks is another turbidite section, considered equivalent to what is called the Black Flags Group (Swager et al., 1992). Turbidites, together with their underlying komatiites are sliced by sled-runner-type thrust faults, which themselves were folded and steepened as a result of an east-west phase of shortening.

Although much of Yilgarn shows the structural and metamorphic pattern described above, the southwestern part of the craton, where extensive granulite facies metamorphism with migmatites and charnockites formed some 2.64 and 2.62 Ga ago, presents a much deeper level of erosion. It is this area that has very old detrital zircons predating any granitic magmatism in this part of the craton. The southwestern high-grade part of the Yilgarn seems to have been attached to the rest of the low-grade craton by 2.6 Ga ago (Cassidy et al., 2006). Within this area Smithies et al. (2018) noted that geochemical signatures indicate northeast-southwest-trending belts that are truncated by the province boundaries and do not correlate with the lower crustal density anomaly. Here geographical variations in proxies for source melting pressure define a broad northeasterly trend that truncates the north-northwesterly trending eastern boundary of the Southwestern Yilgarn Province (Smithies et al., 2018). However, the same northeasterly trends seem to be present also in the rest of the craton, albeit in a much less pronounced manner, although even this, according to Smithies et al. (2018) needs corroboration. If true, however, Smithies et al. (2018) believes that it would invalidate the west-to-east accretion of the various "terrane" in the Yilgarn. However, this suggestion ignores

the very great amounts of strike-slip faulting in the Yilgarn as a whole during the Archean.

We have called the southwestern high-grade region the "McKenzie and Priestley-Land" because it does expose mid to lower crustal rocks as required by their model, in contrast to the "Sengör et alii-Land," where the grade of metamorphism reaches amphibolite grade in only very restricted areas, resembling very much the Altaids (Sengör et al., 2018b) and the Saharides in the Arabian Peninsula and Egypt (Sengör et al., 2020) having undergone all the vicissitudes described in detail by Goscombe et al. (2019) plus much strike-slip faulting in both senses. Yet the entire thing is now one single province. The schematic cross section in our Figure 11 illustrates our suggestion that the McKenzie and Priestley-Land may have been thrust over the Sengör et alii-Land in part and thus became uplifted just like Tibet. How it became denuded is another problem. Whether and how much that thrusting may have also had a strike-slip component is as yet unknown.

The North China Block

The purpose of this section is to provide an example of the inappropriate usage of the term craton for any old terrain exposing Archean and Proterozoic rocks without considering the entire geological history of the region under consideration. Hence the somewhat detailed discussion.

Main Geological Characteristics

What is called the "North China Craton" is one of the most intensively studied Precambrian terrains in the world. Its tectonic history, however, is one of the most controversial subjects in




Figure 8. Contour map (colors) showing general level of erosion across the Superior Province based on igneous and metamorphic pressure estimates. Circled numbers show crustal thickness in kilometers (Percival et al., 2012). Letters represent the localities of photos presented in Figure 9.

Precambrian geology (Kusky et al., 2007a, 2016, 2020; Zhao et al., 2005, 2012; Zhai and Santosh, 2011; Zhai, 2011, 2014; Jiang et al., 2018; Wang et al., 2018; Zhao et al., 2021) in part because of the misleading implications of the term craton as applied to it. We here call it simply the North China Block for the purposes of the following discussion, following the usage of a number of authors before us. We note that the term “North China Block” was also used by Kusky et al. (2007a) on the basis of the assumption that the eastern part of the block lost its cratonic lithosphere in the Mesozoic.

The North China Block is bordered to the north by the Manchurides (Şengör and Natal'in, 1996; Xiao et al., 2003), to the south-southwest by the Qinling-Dabie orogenic belt, to the west by the Qilian part of the Kuen-Lun (Şengör and Okuroğulları, 1991), and to the east by the Su-Lu Belt (Fig. 12), although the Tan-Lu fault (Lu et al., 1983; Wang and Dou, 1997; Huang et al., 2015) and the associated Cainozoic Bo-Hai Basin (Allen et al., 1998; Yang and Xu, 2004) have very largely disrupted its older structure since the Mesozoic. Many models have been proposed to explain the tectonic evolution of what has long

been called the North China Craton and its relationship with the neighboring Proterozoic and Phanerozoic orogenic belts (Kusky and Li, 2003; Zhai et al., 2010; Zhao and Zhai, 2013; Santosh et al., 2016; Kusky et al., 2016, 2020). Our reason for calling it North China Block is because we feel that it does not fulfill the criteria for being a craton for the reasons explained below.

Archean rocks in the North China Block range in age from 3.8 to 2.5 Ga, but most of them formed between 2.9 and 2.5 Ga. The block, however, did not form as a distinct entity until the closure of the ocean between the Eastern




Figure 9. Photographs of outcrops of rocks deformed with middle to no strain from the Superior Province in Canada. (For localities see Figure 8; all photographs are by Ali Polat.)

(A) Bad Vermillion Lake Belt, conglomerates. (B) Cross Lake Sandstone. (C, D) Ca. 2.7 Ga undeformed, greenschist facies pillow breccia and pillow basalts in the Schreiber-Hemlo greenstone belt, Wawa subprovince. (E, F) Autoclastite and komatiite, with spinifex texture, in the Abitibi subprovince, Superior Province. Photograph (C) is from Windley et al. (2021) and photograph (D) is from Polat et al. (2012).

and Western Blocks by 2.43 Ga. (Kusky et al., 2016; Fig. 12). But that date does not signify the end of significant internal deformation of the block and therefore one cannot speak of “cratonization” at 2.43 Ga. The North China Block consists in reality of three main tectonic units including the Eastern Block, Western Block, and the intervening Central Orogenic Belt (Fig. 12). The Eastern Block is predominated by Neoarchean granitoids with subordinate greenstone belts. It has been repeatedly deformed by strike-slip and extensional tectonism since the Mesozoic (Şengör and Natal'in, 1996; Yang and Xu, 2004 and Huang et al., 2015). The geology of the Western Block is poorly known because it is mostly covered by Proterozoic and Phanerozoic sedimentary rocks. The Ordos Basin constitutes

a major part of the Western Block. The Central Orogenic Belt consists mainly of poly-deformed Neoarchean granitoids, mélanges and ophirags (Kusky et al., 2016, 2020).

The North China Block shares the major lithological, structural and geochemical characteristics of many Archean provinces including the Superior Province of Canada, suggesting that it originated through collision of different tectonic units such as island arcs, accretionary complexes, ophirags, oceanic plateaus, and continental blocks (Zhai, 2014; Kusky et al., 2016). The ages of accretion become progressively younger outward from the Central Orogenic Belt to the northwest.

In contrast to real cratons, the North China Block, however, has been affected extensively by Phanerozoic orogenic and taphrogenic

events (Wang et al., 2018; Yang et al., 2019; Sun et al., 2020), which makes it very different from true Archean cratons. The northern margin of the North China Block is marked by the Paleoproterozoic Inner Mongolia–Northern Hebei Orogen, which consists of lithotectonic assemblages of the so-called “Khondalite Belt” to the south-southeast and the Alashan Block (also known as Alxa Block) to the north-northwest (Fig. 12).

According to the tectonic evolution model of Kusky et al. (2016) and more recent studies (Bi et al., 2018; Wang et al., 2018; Yang, et al., 2019; Jia et al., 2020), the major geological events in the North China Block and the neighboring regions started by the assembly of the Eastern Block between 2.7 and 2.6 Ga. This is followed



Figure 10. Photographs from the outcrops of the Barberton Greenstone Belt along the Komati River (all photographs are by A.M.C. Şengör). None of the rocks shown in these photographs show any metamorphism and barely any deformation. (A) Undeformed immiscibility spherules in Archean pillow lavas along the Komati River. (B) Well-preserved soft sediment deformation in turbiditic sandstones. Flame structures of finer-grained sediment (pale) injected into the overlying sandstone (purple-brown). (C) Poorly sorted, matrix-supported conglomerates showing almost «Zero» deformation. (D) Archean pillow lavas from the Kromberg Complex.

by formation of island arcs between the Eastern and Western Blocks between 2.6 and 2.5 Ga which is resulted in the closure of the ocean around 2.43 Ga. Rifting of the newly formed continental block at 2.4–2.35 Ga along its center and northern margin resulted in the development of an Atlantic-type margin to the north. This margin was converted to an Andean-type margin through arc-polarity reversal by 2.3 Ga following a collision with an island arc. The Andean-type margin tectonic, magmatic, sedimentary and metamorphic processes affected much of the North China Block by 1.9 Ga.

The southern margin of the North China Block underwent rift-related magmatic, sedimentary, and metamorphic activities between 2.36 and 2.24 Ga that were followed by an orogenic event between 1.96 and 1.85 Ga and a rifting event between 1.84 and 1.74 Ga (Jia et al., 2020), indicating that the southern margin of the block was tectonically active throughout the early Proterozoic. Sun et al. (2020) proposed the development of an Andean-type margin in the southern part of the block at ca. 1.87 Ga. The eastern margin also underwent multiple rifting, subduction and collision events in the Paleoproterozoic between 2.47 and 1.86 Ga (Li et al., 2011; Yang et al., 2019), resulting in development of an Andean-type margin between 2184 and 2127 Ma (Bi et al., 2018). Continent-continent collision between the northern margin

of the North China Block and the Columbia supercontinent resulted in granulite facies metamorphism and deformation across the block between 1.88 and 1.79 Ga. The North China Block was rifted away from the hypothesized Columbia supercontinent by 1.65 Ga, resulting in the development of a new Atlantic-type margin in the northern part of the block (but see Ding et al., 2020 for palaeomagnetic data suggesting a rifting 1.2 Ga ago⁵).

A Mesozoic orogenic event, nucleated on a Paleozoic extensional structure, and of medial Jurassic to Cainozoic age took place in the Helanshan between the Alashan and Ordos Blocks, which implies that these blocks could not have existed as a single cratonic entity before the Mesozoic (see especially Teilhard de Chardin and Licent, 1924; Yang and Zhang, 1987) or by the late Permian based on the palaeomagnetic results reported by Yuan and Yang (2015).

Deformation of the Luliangshan and Taihangshan in the Mesozoic (Wang et al., 2006) also indicates that an older collisional orogenic belt

(Zhang et al., 2006; Zhao et al., 2006) between the Eastern and Western Blocks was reactivated and overprinted possibly during the medial Triassic (Kobayashi, 1942). Beneath the Mesozoic belt, the entire lithospheric mantle seems refertilized by younger magmatism (Tang et al., 2008). During the Eocene, this belt localized extensional deformation creating such structures as the Fen Wei Rift and the Jizhong Basin to its immediate east, which later became re-shortened (Zhao and Windley, 1990; Sun, 2005).

The entire periphery of the North China Block was affected by Mesozoic magmatism (Xiong'er Shan-WEIFANG Shan granites; Han et al., 2007) resulting from subduction under the North China Block (Wu et al., 2019) and subsequent collision. The Sulu ultrahigh pressure metamorphic belt of Shandong (Zheng et al., 2006) is nothing more than the easterly continuation of the Dabie Shan ultrahigh pressure metamorphic belt related to this northward subduction during the late Paleozoic–earliest Mesozoic (Okay et al., 1989, 1993; Okay and Şengör, 1992; Xu et al., 1992). In the north of the North China Block, the Carboniferous granites extend as far south as 41°N parallel as witnesses to southward subduction below the North China Block (Teilhard de Chardin 1932; Zhang et al., 2007) (Fig. 12).

Early Jurassic (190 Ma) to early Cretaceous (younger than 100 Ma in Liaoning, in the Songliao Basin: Davis, 2003) adakitic and

⁵Ding et al.'s data come from three localities: two within the Yin Shan orogenic belt and one in the so-called “Central Orogenic Belt” separating the Ordos Block from the eastern part of the North China Block (see their fig. 1). Data are not sufficient to tie the Ordos to the entire North China Block 1.2 Ga ago. Thus, we note Ding et al.'s observations, but reserve our judgment about the time of rifting they infer.




Figure 11. A simplified tectonic/metamorphic map of the Yilgarn Province (simplified from Goscombe et al., 2019). The schematic cross section shows the various heights inferred from metamorphic grade while the respective metamorphisms were going on. The areas designated by author's names simply indicate which areas of the Yilgarn are more compatible with which authors' hypotheses and are certainly not attempts to rename the provinces within the Yilgarn nucleus. They refer to discussions in this paper only. The reason we did not use the existing province and/or "terrane" names is that none of them covers what we wish to show. Notice the great difference between the McKenzie and Priestley-Land and the Şengör et alii-Land in terms of inferred elevations. Both now form, however, a single craton with similar crustal thickness. GS—greenschist; UA—upper amphibolite.

coeval non-adakitic magmatism spreads from the northwestern margin of the North China Block (Fig. 12). It seems clear from Davis' discussion of the available observations that these are not products of multiple episodes of lithospheric "delamination" or "decratonicization" but products of the westward subduction, under

thickened continental lithosphere (see also Wu et al., 2019), of a plate in the present space of the western Pacific Ocean much like the similar magmatism in the Andes (Davis, 2003). This episode of shortening in northeast China was followed by one of widespread extension around 130–120 Ma ago in an area exceeding 2 million

km² (Davis, 2003; Davis et al., 2002; Liu et al., 2005), not dissimilar to the early Cainozoic extension that destroyed a North American "Altiplano" during the early Cainozoic and created the Cordilleran extensional metamorphic core complexes similar to those in northeastern China (Davis et al., 2002; Liu et al., 2005). The




Figure 12. Summary map of the North China Block showing the major blocks and sutures. Note that all the tectonic elements shown here were repeatedly redeformed during the Phanerozoic and until about the Jurassic it is not possible with the available observations to be sure of their positions with respect to one another. Abbreviations as follows: HG—Hetao Graben; HS—Helanshan; LL—Luliangshan; TS—Taihangshan; YG—Yinchuan Graben. Redrawn from Deng et al. (2003), Kusky et al. (2016), Bi et al. (2018), Pang et al. (2020).

present-day thin lithosphere along the eastern edge of the North China Block is clearly a result of this late phase stretching (see Zhao and Zheng, 2007). As Şengör (2001) has argued, losing chunks of the lithospheric mantle (the preferred explanation of Zhao and Zheng, 2007) under zones of mantle upwelling is not incompatible with the stretching interpretation (see also Kusky et al., 2007b).

Did the North China Block Have a Thick, Continuous Precambrian Lithospheric Root?

The North China Block is distinct from many Archean cratons with its thinner (80–130 km)

subcontinental lithospheric mantle beneath the eastern part (Kusky et al., 2007b; Wang et al., 2018; Wu et al., 2019; Xia et al., 2020). Based mainly on the presence Ordovician (ca. 480–450 Ma) diamondiferous kimberlite pipes from two locations and on the widespread occurrence of late Mesozoic and Cainozoic spinel lherzolites in the Eastern Block, many studies (Menzies et al., 1993; Zheng et al., 2007; Wu et al., 2019) argued that the cratonic mantle beneath the eastern part of the North China Block was thinned/removed and replaced by the asthenospheric mantle, leading to the decratonicization of the eastern segment of the block. Using geo-

physical data, Xia et al. (2020) showed that most of the cratonic mantle beneath the North China Block was reworked in response to Phanerozoic tectonic and magmatic processes, which reduced the lithospheric thickness to 100–130 km beneath the northern, eastern, and western segments of block. The study by Xia et al. (2020) implies that thinner lithospheric mantle is not restricted to the eastern part but occurs throughout beneath the North China Block.

Although proposed ages for lithospheric thinning/removal range from 2.5 Ga to 120 Ma, most models favor decratonicization and the loss of the lithospheric root in the Mesozoic (Menzies

et al., 1993; Griffin et al., 1998; Zhai et al., 2007; Chen et al., 2008; Gao et al., 2008; Yang et al., 2008; Windley et al., 2010; Li et al., 2011; Zhu et al., 2012; Wang et al., 2018; Wu et al., 2019). A detailed review of these models is beyond the scope of this study. Accordingly, we present a short summary of these models. The majority of these models invoke hydration and weakening of the lithospheric mantle by subduction of oceanic lithosphere to the north and south, and flat subduction of the Paleo-Pacific oceanic crust beneath the North China Block in the Mesozoic (see Davis et al., 2001; Davis, 2003; Kusky et al., 2014; Zheng et al., 2018; Wu et al., 2019 for a review).

Windley et al. (2010) suggested that the Archean lithospheric mantle beneath the supposed craton was hydrated and weakened by the south-dipping slabs of Permian-Triassic Solonker and the Jurassic Mongol-Okhotsk subduction zones to the north, the north-dipping slabs of Permian-Triassic Qinling-Dabishan and Song Ma subductions to the south, and the west-dipping Jurassic-Cretaceous (200–100 Ma) Paleo-Pacific oceanic crust to the east. These processes eventually led to the thinning and delamination of the lithosphere beneath the Eastern Block. Similarly, Zhu et al. (2011, 2012) suggested that the westward subduction of the paleo-Pacific oceanic lithosphere beneath the North China Block played an important role in its lithospheric destruction. Wu et al. (2019) proposed that the lithospheric loss stemmed from Cretaceous orogenic collapse in response to Jurassic crustal thickening resulting from flat subduction of the paleo-Pacific oceanic lithosphere, carrying an oceanic plateau, between 210 and 140 Ma. None of the models above, however, take the timing of structural events into consideration and they are all incompatible with the tectonic calendar as documented by Davis (2003). Phanerozoic tectonics clearly shows that there never was a unified Precambrian entity worthy of the designation craton. This is much like the situation in Europe during the Phanerozoic where repeated tectonism did not allow the formation of a craton (Şengör et al., 2018a).

The question is: did the North China Block have a thick, continuous, rigid cratonic lithosphere before the Mesozoic? In other words, was it cratonized by the Paleoproterozoic (1.7 Ga) and remained intact until the Mesozoic? Although deformation in the early Cretaceous was mainly restricted to the Eastern Block, deformation in the Jurassic extended throughout the block (fig. 4 in Wu et al., 2019). Could a craton with a thick (~200 km) lithospheric mantle root undergo multiple phases of orogenic events and rifting? A record of widespread geological activity in the Proterozoic, Paleozoic, Mesozoic,

and Cainozoic suggest that a thick cratonic lithospheric root beneath the North China Block could not have been fully developed by the end of Paleoproterozoic as commonly assumed. The development of a continuous, rigid lithospheric root might have been impeded by the incidence of the Paleoproterozoic orogenic and rifting events in the northern, southern, and eastern margins of the block, as outlined above, causing only a partial development of a lithospheric root. This partially developed lithospheric root was likely susceptible to destruction by tectonic and thermal erosion by the circum-North China subduction zones of various Paleozoic ages and would have provided the preconditions for its complete removal in the Mesozoic, but the internal taphrogenic and orogenic events along the Helanshan and the Taihangshan clearly show that there never was a united North China craton from under which a lithospheric armor could have been removed.

Although most studies assumed that the North China Block was underlain by a thick (>150 km) lithospheric mantle before the Jurassic (see Wu et al., 2019, for a review), its lateral distribution beneath the block before the Jurassic is not well known. Was such thick lithosphere restricted only to the eastern part of the block where the diamondiferous kimberlites occur, or did it exist beneath the whole block? Was the lithospheric mantle removed from beneath the block in the Mesozoic? If it was removed, how was it removed? These questions are the subjects of intense debate over the past 30 years (Kusky et al., 2007a, 2007b; Wu et al., 2019; Xia et al., 2020). The known diamondiferous kimberlite pipes are restricted to the eastern margin of the Eastern Block. If the whole North China Block was underlain by a thick lithospheric mantle such as that sampled by the Ordovician kimberlite pipes, one would expect to see the occurrence of such kimberlite pipes in other parts of the North China Block. Kimberlite pipes in other parts of the block either have not been discovered or they are not present due to the absence of the thick lithospheric mantle beneath most of the block, which implies that a larger part of the block never developed a lithospheric layer thicker than 150 km to form diamonds.

Collectively, the tectonic evolution of the North China Block indicates that it underwent multiple phases of deformation, magmatism, metamorphism and sedimentation in its northern, southern, and eastern margins plus within it, around the Ordos Basin since the Neoarchean (Teilhard de Chardin and Licent, 1924; Kobayashi, 1941, 1942; He et al., 1983; Yang and Zhang, 1987; Darby et al., 2001; Davis et al., 2002), which raises the question whether the North China Block can be defined as a single, stable

craton. No craton is known to us that exhibits this much tectonism and for such a long time within it. Consequently, we do not recognize the North China Block assembly as a craton: neither now nor any time in the past.

CONCLUSIONS

Ophiolitic remnants, igneous rocks with arc signature, structural thrust belts, juxtaposed rocks of different origin and ages, and the lithospheric structure derived from seismic reflectors in the Archean rock record show a high resemblance to the present-day subduction-accretion systems suggesting that subduction was operating throughout the Archean. This indicates that cratons formed by the amalgamation of magmatic arcs and associated subduction-accretion complexes that were eventually involved in a final continental collision that greatly thickened the underlying mantle lithosphere (Şengör, 1999; Priestley et al., 2021). The amalgamation may involve not only subduction, but also much strike-slip faulting. Both subduction and strike-slip faulting may intercalate episodes of local extension onto different parts of the system thus complicating its geological history. Even the final collision is probably only rarely head-on. This can be supported by the absence of high elevations in these cratonic areas as deduced from the surface geology and denudation rates. This leads us to the conclusion that craton formation does not require total removal of upper crust or exhumation of the entire lower crust. Instead, the granulitization of the roots of arcs may have been responsible for weighing the collided and thickened pieces down. The best candidate of a craton forming event is seen in the Altaids, a Paleozoic superorogenic system which has not been deformed orogenetically since the early Cretaceous (Şengör et al., 2014a, 2014b; 2018b).

We emphasize that when a craton is named, it must be indicated for what time period in the history of the earth that name and the area it implies are meant. Moreover, not every Archean or Proterozoic terrain is a craton. Our Figure 2 shows wide areas of Proterozoic and even Archean regions that are not cratons. Shields are multi-colored showing the ages of the underlying rocks identified in the legend. The platforms are shown in gray, except where sparse basement pieces crop out. The gray-colored platforms have two hues: darker gray indicates true cratonic platforms, about which there is no serious disagreement among geologists. Lighter gray indicates areas considered cratonic by most geologists, but we do not consider them as such because of a protracted deformation history that reaches into our own day, although they display a strength against convergent plate boundary

deformations superior to that in orogenic zones, as defined in Sengör (1984), corresponding with Kober's (1921) orogens. Considering any ancient piece of crust, a craton has led to much confusion in the past as illustrated by the history of research on the Saharides, where an entire orogenic belt has been overlooked because of the abstruse concept of a "metacraton" (Sengör et al., 2020). Another example is illustrated by what is known as "North China Craton." It has undergone multiple episodes of deformation, magmatism, metamorphism, and sedimentation since the Archean, indicating that its geological characteristics are inconsistent with the definition of a craton.

ACKNOWLEDGMENTS

N. Lom acknowledges NWO Vici grant 865.17.001. A. Polat acknowledges a Natural Sciences and Engineering Research Council (Canada) Discovery Grant (RGPIN-2019-04236) and a Chang Jiang Scholarship grant by the Ministry of Education of China. We thank two anonymous reviewers for their incisive, helpful comments, and acknowledge them accordingly.

REFERENCES CITED

- Ackerson, M.R., Trail, D., Buettner, J., 2021. Emergence of peraluminous crustal magmas and implications for the early Earth: *Geochemical Perspectives Letters*, v. 17, p. 50–54, <https://doi.org/10.7185/geochemlet.2114>.
- Agangi, A., Hofmann, A., Hegner, E., Xie, H., Teschner, C., Slabunov, A., and Svetov, S.A., 2020. The Mesoproterozoic Dominion Group and the onset of intracontinental volcanism on the Kaapvaal craton—Geological, geochemical and temporal constraints: *Gondwana Research*, v. 84, p. 131–150, <https://doi.org/10.1016/j.gr.2020.03.005>.
- Agranier, A., Maury, R.C., Geoffroy, L., Chauvet, F., Le Gall, B., and Viana, A.R., 2019. Volcanic record of continental thinning in Baffin Bay margins: Insights from Svarthuk Halvø Peninsula basalts, West Greenland: *Lithos*, v. 334–335, p. 117–140, <https://doi.org/10.1016/j.lithos.2019.03.017>.
- Ahmat, T., Gole, M., Goleby, M., Morris, P., Swager, C., and Wyche, S., 1993. Greenstone terranes and the eastern Goldfields seismic traverse, in Williams, P.R. and Haldane, J.A., eds., An International Conference on Crustal Evolution Metallogeny and Exploration of the Eastern Goldfields Excursion Guidebook Record 1993/53: Canberra, Australian Geological Survey Organisation, p. 1–22.
- Allaby, M., ed., 2008. A Dictionary of Earth Sciences (third ed.): Oxford University Press, 672 p.
- Argand, É., 1924. La tectonique de l'Asie: Congrès Géologiques Internationaux, Comptes Rendus de la XII^e session, Premier Fascicule: Liège: H. Vaillant-Carmanne, p. 171–372.
- Allen, M.B., Macdonald, D.I.M., Zhao, X., Vincent, S.J., and Broeck-Menzies, C., 1998. Transtensional deformation in the evolution of the Bohai Basin, northern China, in Holdsworth, R.E., Strachan, R.A. and Dewey, J.F., eds., Continental Transpressional and Transtensional Tectonics: Geological Society, London, Special Publication 135, p. 215–229, <https://doi.org/10.1144/GSL.SP.1998.135.01.14>.
- Armstrong, R.A., Compston, W., de Wit, M.J., and Williams, I.S., 1990. The stratigraphy of the 3.5–3.2 Ga Barberton Greenstone Belt revisited: a single zircon ion microprobe study: *Earth and Planetary Science Letters*, v. 101, no. 1, p. 90–106, [https://doi.org/10.1016/0012-821X\(90\)90127-J](https://doi.org/10.1016/0012-821X(90)90127-J).
- Arndt, N.T., 2013. The formation and evolution of the continental crust: *Geochemical Perspectives*, v. 2, p. 405–528, <https://doi.org/10.7185/geochempersp.2.3>.
- Artemieva, I.M., 2006. Global 1×1 thermal model TC1 for the continental lithosphere: implications for lithosphere secular evolution: *Tectonophysics*, v. 416, no. 1–4, p. 245–277, <https://doi.org/10.1016/j.tecto.2005.11.022>.
- Artemieva, I.M., 2019. Lithosphere structure in Europe from thermal isostasy: *Earth-Science Reviews*, v. 188, p. 454–468, <https://doi.org/10.1016/j.earscirev.2018.11.004>.
- Artemieva, I.M., and Mooney, W.D., 2001. Thermal thickness and evolution of Precambrian lithosphere: a global study: *Journal of Geophysical Research: Solid Earth*, v. 106, B8, p. 16,387–16,414, <https://doi.org/10.1029/2000JB900439>.
- Artemieva, I.M., and Mooney, W.D., 2002. On the relations between cratonic lithosphere thickness, plate motions, and basal drag: *Tectonophysics*, v. 358, no. 1–4, p. 211–231, [https://doi.org/10.1016/S0040-1951\(02\)00425-0](https://doi.org/10.1016/S0040-1951(02)00425-0).
- Artemieva, I.M., and Vinnik, L.P., 2016. Density structure of the cratonic mantle in southern Africa: 1. Implications for dynamic topography: *Gondwana Research*, v. 39, p. 204–216, <https://doi.org/10.1016/j.gr.2016.03.002>.
- Aulbach, S., 2012. Craton nucleation and formation of thick lithospheric roots: *Lithos*, v. 149, p. 16–30, <https://doi.org/10.1016/j.lithos.2012.02.011>.
- Baltoni, I., and Ballica, C., 2016. Peri-Amazonian provenance of the Euxinic Craton components in Dobrogea and of the North Dobrogean Orogen components (Romania): a detrital zircon study: *Precambrian Research*, v. 278, p. 34–51, <https://doi.org/10.1016/j.precamres.2016.03.008>.
- Baranov, A.A., and Bobrov, A.M., 2018. Crustal structure and properties of Archean cratons of Gondwanaland: similarity and difference: *Russian Geology and Geophysics*, v. 59, no. 5, p. 512–524, <https://doi.org/10.1016/j.rgg.2018.04.005>.
- Bi, J.H., Ge, W.C., Xing, D.H., Yang, H., Dong, Y., Tian, D.X., and Chen, H.J., 2018. Palaeoproterozoic metarhyolite and meta-dacite of the Liaobei Group, Jiao-Liao-Ji Belt, North China Craton: Petrogenesis and implications for tectonic setting: *Precambrian Research*, v. 314, p. 306–324, <https://doi.org/10.1016/j.precamres.2018.06.007>.
- Bousquet, R., Goffé, B., Henry, P., Le Pichon, X., and Chopin, C., 1997. Kinematic, thermal and petrological model of the Central Alps: Lepontine metamorphism in the upper crust and eclogitisation of the lower crust: *Tectonophysics*, v. 273, no. 1–2, p. 105–127, [https://doi.org/10.1016/S0040-1951\(96\)00290-9](https://doi.org/10.1016/S0040-1951(96)00290-9).
- Burchfiel, B.C., and Davis, G.A., 1975. Nature and controls of Cordilleran orogenesis, western United States: extensions of an earlier synthesis: *American Journal of Science (John Rodgers volume)*, v. 275-A, p. 363–396.
- Burke, K., 1996. The African plate: *South African Journal of Geology*, v. 99, p. 339–409.
- Burke, K., and Gunnell, Y., 2008. The African Erosion Surface: A Continental-Scale Synthesis of Geomorphology, Tectonics, and Environmental Change over the past 180 Million Years: *Geological Society of America Memoir* 201, 66 p., <https://doi.org/10.1130/2008.1201>.
- Burke, K., and Kidd, W.S.F., 1978. Were Archean continental geothermal gradients much steeper than those of today?: *Nature*, v. 272, p. 240–241, <https://doi.org/10.1038/272240a0>.
- Burke, K., Kidd, W.S.F., and Kusky, T.M., 1986. Archean foreland basin tectonics in the Witwatersrand, South Africa: *Tectonics*, v. 5, no. 3, p. 439–456, <https://doi.org/10.1029/TC005i003p00439>.
- Byrne, P.K., Klimeczak, C., Sengör, A.M.C., and Solomon, S., 2014. Mercury's global contraction much greater than earlier estimates: *Nature Geoscience*, v. 7, no. 4, <https://doi.org/10.1038/ngeo2097>.
- Calais, E., Dong, L., Wang, M., Shen, Z., and Vergnolle, M., 2006. Continental deformation in Asia from a combined GPS solution: *Geophysical Research Letters*, v. 33, L24319, <https://doi.org/10.1029/2006GL028433>.
- Capitanio, F.A., Nebel, O., and Cawood, P.A., 2020. Thermochemical lithosphere differentiation and the origin of cratonic mantle: *Nature*, v. 588, p. 89–94, <https://doi.org/10.1038/s41586-020-2976-3>.
- Card, K.D., 1990. A review of the Superior Province of the Canadian Shield, a product of Archean accretion: *Precambrian Research*, v. 48, no. 1–2, p. 99–156, [https://doi.org/10.1016/0301-9268\(90\)90059-Y](https://doi.org/10.1016/0301-9268(90)90059-Y).
- Carlson, R.W., Pearson, D.G., and James, D.E., 2005. Physical, chemical, and chronological characteristics of continental mantle: *Reviews of Geophysics*, v. 43, no. 1, 24 p., <https://doi.org/10.1029/2004RG000156>.
- Cassidy, K.F., Champion, D.C., Krapež, B., Barley, M.E., Brown, S.J.A., Blewett, R.S., Groenewald, P.B., and Tyler, I.M., 2006. A revised geological framework for the Yilgarn Craton, Western Australia: *Geological Survey of Western Australia*, Record 2006/8, 8 p.
- Chen, L., Wang, T., Zhao, L., and Zheng, T.Y., 2008. Distinct lateral variation of lithospheric thickness in the north-eastern North China Craton: *Earth and Planetary Science Letters*, v. 267, p. 56–68, <https://doi.org/10.1016/j.epsl.2007.11.024>.
- Clouzet, P., Masson, Y., and Romanowicz, B., 2018. Box Tomography: first application to the imaging of upper-mantle shear velocity and radial anisotropy structure beneath the North American continent: *Geophysical Journal International*, v. 213, no. 3, p. 1849–1875, <https://doi.org/10.1093/gji/ggy078>.
- Collie, M., and Diemer, J., 2004. Murchison's Wanders in Russia—His Geological Exploration of Russia in Europe and the Ural Mountains, 1840 and 1841: *British Geological Survey Occasional Publication No. 2*, xv+474 p., 6 plates.
- Coney, P.J., and Reynolds, S.J., 1977. Cordilleran Benioff zones: *Nature*, v. 270, p. 403–406, <https://doi.org/10.1038/270403a0>.
- Dana, J.D., 1847a. Geological results of the earth's contraction in consequence of cooling: *American Journal of Science and Arts*, 2nd series, v. 3, p. 176–188.
- Dana, J.D., 1847b. Origin of the continents: *American Journal of Science and Arts*, 2nd series, v. 3, p. 94–100.
- Dana, J.D., 1863. Manual of Geology: Treating of the Principles of the Science with Special Reference to American Geological History, for the Use of Colleges, Academies and Schools of Science: Philadelphia, Theodore Bliss, 798 p.
- Darby, B.J., Davis, G.A., and Zheng, Y.D., 2001. Structural evolution of the southwestern Daqing Shan, Yinshan Belt, Inner Mongolia, China, in Hendrix, M.S., and Davis, G.A., eds., Mesozoic Tectonic Evolution of Central Asia. From Continental Assembly to Intracontinental Deformation: *Geological Society of America Memoir* 194, p. 199–214, <https://doi.org/10.1130/0-8137-1194-0.199>.
- Das, R., Saikia, U., and Rai, S.S., 2015. The deep geology of South India inferred from Moho depth and Vp/Vs ratio: *Geophysical Journal International*, v. 203, p. 910–926, <https://doi.org/10.1093/gji/ggv351>.
- Davis, G.A., 2003. The Yanshan Belt of North China: tectonics, adakitic magmatism, and crustal evolution: *Earth Science Frontiers*, v. 10, p. 373–384.
- Davis, G.A., Zheng, Y.D., Wang, C., Darby, B., Zhang, C.H., and Gehrels, G., 2001. Mesozoic tectonic evolution of the Yanshan fold and thrust belt, with emphasis on Hebei and Liaoning provinces, northern China, in Hendrix, M.S. and Davis, G.A., eds., Mesozoic Tectonic Evolution of Central Asia. From Continental Assembly to Intracontinental Deformation: *Geological Society of America Memoir* 194, p. 171–197, <https://doi.org/10.1130/0-8137-1194-0.171>.
- Davis, G.A., Darby, B.J., Zheng, Y.D., and Spell, T.L., 2002. Geometric and temporal evolution of an extensional detachment fault, Hohhot metamorphic core complex, Inner Mongolia, China: *Geology*, v. 30, p. 1003–1006, [https://doi.org/10.1130/0091-7613\(2002\)030<1003:GATEOA>2.0.CO;2](https://doi.org/10.1130/0091-7613(2002)030<1003:GATEOA>2.0.CO;2).
- de Wit, M.J., de Ronde, C.E., Tredoux, M., Roering, C., Hart, R.J., Armstrong, R.A., Green, R.W., Peberdy, E., and Hart, R.A., 1992. Formation of an Archean continent: *Nature*, v. 357, no. 6379, p. 553–562, <https://doi.org/10.1038/35753a0>.
- de Wit, M.J., Furnes, H., and Robins, B., 2011. Geology and tectonostratigraphy of the Onverwacht Suite, Barberton greenstone belt, South Africa: *Precambrian Research*, v. 186, no. 1–4, p. 1–27, <https://doi.org/10.1016/j.precamres.2010.12.007>.
- Deng, Q., Zhang, P., Ran, Y., Yang, X., Min, W., and Chu, Q., 2003. Basic characteristics of active tectonics of China:

- Science in China. Series D, Earth Sciences, v. 46, p. 356–372, 10.1360/03yd9032.
- Dewey, J.F., and Burke, K., 1973, Tibetan, Variscan and Precambrian basement reactivation: products of continental collision: *The Journal of Geology*, v. 81, p. 683–692, <https://doi.org/10.1086/627920>.
- Ding, J.-K., Zhang, S.-H., Zhao, H.-Q., Xian, H.-B., Li, H.Y., Yang, T.-S., Wu, H.-C., and Wang, W., 2020, A combined geochronological and paleomagnetic study on ~1220 Ma mafic dikes in the North China Craton and the implications for the breakup of Nuna and assembly of Rodinia: *American Journal of Science*, v. 320, p. 125–149, <https://doi.org/10.2475/02.2020.02>.
- Dufrénoy, P.A., and Élie de Beaumont, L., 1848, Explication de la Carte Géologique de la France, rédigée sous la direction de M. Brochant de Villiers ...tome deuxième: Imprimerie Nationale, Paris, XII+813 p.
- Élie de Beaumont, L., 1828a, Notice sur un gisement de Végétaux fossiles et de Graphite, situé au col du Chardonnet (département des Hautes Alpes): *Annales des Sciences Naturelles*, v. 14, p. 113–127.
- Élie de Beaumont, L., 1828b, Sur un gisement de Végétaux fossiles et de Graphite, situé au col du Chardonnet (département des Hautes Alpes): *Annales des Sciences Naturelles*, v. 15, p. 353–381.
- Élie de Beaumont, L., 1852, Notice sur les Systèmes de Montagnes, v. III: P. Bertrand, Paris, p. 1069–1543, 4 foldout plates.
- Flowers, R.M., Bowring, S.A., and Reiners, P.W., 2006, Low long-term erosion rates and extreme continental stability documented by ancient (U-Th)/He dates: *Geology*, v. 34, no. 11, p. 925–928, <https://doi.org/10.1130/G22670A.1>.
- Furnes, H., Dilek, Y., and de Wit, M., 2015, Precambrian greenstone sequences represent different ophiolite types: *Gondwana Research*, v. 27, p. 649–685, <https://doi.org/10.1016/j.gr.2013.06.004>.
- Gao, S., Rudnick, R.L., Xu, W.L., Yuan, H.L., Liu, Y.S., Walker, R., Puchtel, I.S., Liu, X., Huang, H., Wang, X.R., and Yang, J., 2008, Recycling deep cratonic lithosphere and generation of intraplate magmatism in the North China Craton: *Earth and Planetary Science Letters*, v. 270, p. 41–53, <https://doi.org/10.1016/j.epsl.2008.03.008>.
- Gibbs, A.K., and Barron, C.N., 1993, The Geology of the Guiana Shield: Oxford Monographs on Geology and Geophysics, no. 22, Clarendon Press, Oxford, 246 p.
- Goscombe, B., Foster, D.A., Blewett, R., Czarnota, K., Wade, B., Groenewald, B., and Gray, D., 2019, Neoarchean metamorphic evolution of the Yilgarn Craton: a record of subduction, accretion, extension and lithospheric delamination: *Precambrian Research*, v. 335, 105441, <https://doi.org/10.1016/j.precamres.2019.105441>.
- Griffin, W.L., Zhang, A.D., O'Reilly, S.Y., and Ryan, C.G., 1998, Phanerozoic evolution of the lithosphere beneath the Sino-Korean craton, in Flower, M.F.J., Chung, S.L., Lo, C.H. and Lee, T.Y., eds., Mantle Dynamics and Plate Interactions in East Asia: Washington, D.C. American Geophysical Union, p. 107–126, <https://doi.org/10.1029/GD027p0107>.
- Grosch, E.G., and Slama, J., 2017, Evidence for 3.3 billion-year-old oceanic crust in the Barberton greenstone, South Africa: *Geology*, v. 45, p. 695–698, <https://doi.org/10.1130/G39035.1>.
- Grosch, E.G., Kosler, J., McLoughlin, N., Drost, K., Slama, J., and Pedersen, R.B., 2011, Paleoproterozoic detrital zircon ages from the earliest tectonic basin in the Barberton Greenstone Belt, Kaapvaal craton, South Africa: *Precambrian Research*, v. 191, p. 85–99, <https://doi.org/10.1016/j.precamres.2011.09.003>.
- Hall, J., 1842, Notes upon the geology of the western states: *American Journal of Science and Arts*, v. 42, p. 51–62.
- Hall, J., 1858, General Geology, in Hall, J. and Whitney, J.D., Report of the Geological Survey of the State of Iowa, v. I, part I: Geology, published by the authority of the Legislature of Iowa, C. van Benthuyzen, Albany, p. 35–44.
- Hall, J., 1859, Introduction: in Hall, J., Geological Survey of New York. Palaeontology: volume III. ...Part I: Text., p. 1–96.
- Hall, A.M., Putkinen, N., Hietala, S., Lindsberg, E., and Holma, M., 2021, Ultra-slow cratonic denudation in Finland since 1.5 Ga indicated by tiered unconformities and impact structures: *Precambrian Research*, v. 352, 106000, <https://doi.org/10.1016/j.precamres.2020.106000>.
- Hamilton, W.B., 2007, Earth's first two billion years—The era of internally mobile crust, in Hatcher, R.D., Jr., Carlson, M.P., McBride, J.H., and Martínez Catalán, J.R., eds., 4-D Framework of Continental Crust: Geological Society of America Memoir 200, p. 233–296, [https://doi.org/10.1130/2007.1200\(13\)](https://doi.org/10.1130/2007.1200(13)).
- Hamilton, W.B., 2019, Toward a myth-free geodynamic history of Earth and its neighbors: *Earth-Science Reviews*, v. 198, 102905, <https://doi.org/10.1016/j.earscirev.2019.102905>.
- Han, Y.G., Zhang, S.H., Piranjo, F., and Zhang, Y.H., 2007, Evolution of the Mesozoic granites in the Xiong'ershan-Waifangshan region, western Henan Province, China, and its tectonic implications: *Acta Geologica Sinica*, v. 81, p. 253–265, <https://doi.org/10.1111/j.1755-6724.2007.tb00949.x>.
- Harrison, T.M., 2009, The Hadean crust: evidence from > 4 Ga zircons: *Annual Review of Earth and Planetary Sciences*, v. 37, p. 479–505, <https://doi.org/10.1146/annurev.earth.031208.100151>.
- Harrison, T.M., 2020, Hadean Earth: Springer xvi+291p. <https://doi.org/10.1007/978-3-030-46687-9>.
- Hartmann, L.A., and Delgado, I., 2001, Cratons and orogenic belts of the Brazilian Shield and their contained gold deposits: *Mineralium Deposita*, v. 36, p. 207–217, <https://doi.org/10.1007/s001260100175>.
- Hawkesworth, C.J., Dhuime, B., Pietranik, A.B., Cawood, P.A., Kemp, A.I., and Storey, C.D., 2010, The generation and evolution of the continental crust: *Journal of the Geological Society*, v. 167, no. 2, p. 229–248, <https://doi.org/10.1144/0016-76492009-072>.
- Hayashi, M., Komiya, T., Nakamura, Y., and Maruyama, S., 2000, Archean regional metamorphism of the Isua supracrustal belt, southern West Greenland: implications for a driving force for Archean plate tectonics: *International Geology Review*, v. 42, no. 12, p. 1055–1115, <https://doi.org/10.1080/00206810009465128>.
- He, G.Q., Wang, S.G., Chi, G.Z., and Shao, J., 1983, On the features of reactivation of the sino-Korean Plate: *Tectonophysics*, v. 100, p. 119–130, [https://doi.org/10.1016/0040-1951\(83\)90181-6](https://doi.org/10.1016/0040-1951(83)90181-6).
- Heaman, L.M., and Pearson, D.G., 2010, Nature and evolution of the Slave Province subcontinental lithospheric mantle: *Canadian Journal of Earth Sciences*, v. 47, p. 369–388, <https://doi.org/10.1139/E09-046>. [This article is one of a series of papers published in this Special Issue on the theme Lithoprobe—Parameters, Processes, and the Evolution of a Continent.]
- Herzberg, C., 2018, From hot oceanic ridges to cool cratons: *Geology*, v. 46, p. 1079–1080, <https://doi.org/10.1130/focus122018.1>.
- Hetényi, G., Cattin, R., Brunet, F., Bollinger, L., Vergne, J., Nábělek, J.L., and Diament, M., 2007, Density distribution of the India plate beneath the Tibetan plateau: Geophysical and petrological constraints on the kinetics of lower-crustal eclogitization: *Earth and Planetary Science Letters*, v. 264, no. 1–2, p. 226–244, <https://doi.org/10.1016/j.epsl.2007.09.036>.
- Hickman, A.H., 2008, Regional review of the 3426–3350 Ma Strelley Pool Formation, Pilbara Craton, Western Australia: *Geological Survey of Western Australia, Record 2008/15*, 27 p.
- Hoffman, P.F., 1988, Early Proterozoic Assembly and Growth of Laurentia: *Annual Review of Earth and Planetary Sciences*, v. 16, p. 543–603, <https://doi.org/10.1146/annurev.ea.16.050188.002551>.
- Hofmann, A., and Kusky, T., 2004, The Belingwe greenstone belt: ensialic or oceanic?: Developments in Precambrian geology, v. 13, p. 487–538.
- Huang, L., Liu, C.Y., and Kusky, T.M., 2015, Cenozoic evolution of the Tan-Lu Fault Zone (East China)—Constraints from seismic data: *Gondwana Research*, v. 28, p. 1079–1095, <https://doi.org/10.1016/j.gr.2014.09.005>.
- Hyung, E., and Jacobsen, S.B., 2020, The $^{142}\text{Nd}/^{144}\text{Nd}$ variations in mantle-derived rocks provide constraints on the stirring rate of the mantle from the Hadean to the present: *Proceedings of the National Academy of Sciences of the United States of America*, v. 117, <https://doi.org/10.1073/pnas.2006950117>.
- Jacobs, J., 2009, A review of two decades (1986–2008) of geochronological work in Heimefrontfjella, and geotectonic interpretation of western Dronning Maud Land, East Antarctica: *Polarforschung*, v. 79, no. 1, p. 47–57.
- James, D.E., and Fouch, M.J., 2002, Formation and evolution of Archaean cratons: insights from southern Africa: *Geological Society, London, Special Publication* 199, no. 1, p. 1–26, <https://doi.org/10.1144/GSL.SP.2002.199.01.01>.
- Jaupart, C., and Mareschal, J.C., 1999, The thermal structure and thickness of continental roots: *Lithos*, v. 48, p. 93–114, [https://doi.org/10.1016/S0024-4937\(99\)00023-7](https://doi.org/10.1016/S0024-4937(99)00023-7).
- Jia, X., Zhai, M., Xiao, W., Li, L., Ratheesh-Kumar, R.T., Wu, J., and Liu, Y., 2020, Mesoarchean to Paleoproterozoic crustal evolution of the Taihua Complex in the southern North China Craton: *Precambrian Research*, v. 337, 105451, <https://doi.org/10.1016/j.precamres.2019.105451>.
- Jiang, S., Cao, W., Li, S., Wang, G., Somerville, I., Zhang, W., Zhao, F., and Chen, H., 2018, Tectonic units of the Early Precambrian basement within the North China Craton: Constraints from gravitational and magnetic anomalies: *Precambrian Research*, v. 318, p. 122–132, <https://doi.org/10.1016/j.precamres.2018.10.002>.
- Jordan, T.H., 1975, Lateral heterogeneity and mantle dynamics: *Nature*, v. 257, p. 745–750, <https://doi.org/10.1038/257745a0>.
- Jordan, T.H., 1988, Structure and formation of the continental tectosphere: *Journal of Petrology*, v. 29, p. 11–37, https://doi.org/10.1093/petrology/Special_Volume.1.11.
- Kley, J., 2013, Saxonische Tektonik im 21. Jahrhundert: *Zeitschrift der Deutschen Gesellschaft für Geowissenschaften*, v. 164, p. 295–311, <https://doi.org/10.1127/1860-1860/2013/0022>.
- Kobayashi, T., 1941, The Sakawa Orogenic Cycle and its bearing on the origin of the Japanese islands: *Journal of the Faculty of Science Imperial University of Tokyo, section II Geology, Mineralogy, Geography, Seismology*, v. 5, part 7, p. 219–578, 4 plates, 10 geological maps in back pocket.
- Kobayashi, T., 1942, The Rakuroan Complex of the Shansi Basin and its surroundings: *Miscellaneous notes on the Cambro-Ordovician geology and palaeontology VIII: Japanese Journal of Geology and Geography*, v. 18, p. 283–306.
- Kober, L., 1921, Der Bau der Erde: Gebrüder Borntraeger, Berlin, [III]+324 p. + 1 foldout map.
- Korenaga, J., 2021, Hadean geodynamics and the nature of early continental crust: *Precambrian Research*, v. 359, 106178, <https://doi.org/10.1016/j.precamres.2021.106178>.
- Küpfer, H., 1965, Quasicraton und Orthogeosynklinal (Ostalpen und Böhmisches Massiv im Kenntnisbild der heutigen Geologie): *Eclogae Geologicae Helvetiae*, v. 58, p. 73–85, 1 foldout map.
- Kusky, T.M., 1989, Accretion of the Archean Slave province: *Geology*, v. 17, p. 63–67, [https://doi.org/10.1130/0091-7613\(1989\)017<0063:AOTASP>2.3.CO;2](https://doi.org/10.1130/0091-7613(1989)017<0063:AOTASP>2.3.CO;2).
- Kusky, T.M., 1993, Collapse of Archean orogens and the generation of late- to postkinematic granitoids: *Geology*, v. 21, p. 925–928, [https://doi.org/10.1130/0091-7613\(1993\)021<0925:COAOAT>2.3.CO;2](https://doi.org/10.1130/0091-7613(1993)021<0925:COAOAT>2.3.CO;2).
- Kusky, T.M., and Kidd, W.S.F., 1992, Remnants of an Archean oceanic plateau, Belingwe greenstone belt, Zimbabwe: *Geology*, v. 20, p. 43–46, [https://doi.org/10.1130/0091-7613\(1992\)020<0043:ROAAP>2.3.CO;2](https://doi.org/10.1130/0091-7613(1992)020<0043:ROAAP>2.3.CO;2).
- Kusky, T.M., and Li, J.H., 2003, Paleoproterozoic tectonic evolution of the North China Craton: *Journal of Asian Earth Sciences*, v. 22, p. 383–397, [https://doi.org/10.1016/S1367-9120\(03\)00071-3](https://doi.org/10.1016/S1367-9120(03)00071-3).
- Kusky, T.M., Windley, B.F., and Zhai, M.G., 2007a, Tectonic evolution of the North China Block: from orogen to craton to orogen, in Zhai, M.G., Windley, B.F., Kusky, T.M., and Meng, Q.R., eds., *Mesozoic Sub-continental Lithospheric Thinning under Eastern Asia: Geological Society, London, Special Publication* 280, p. 1–34, <https://doi.org/10.1144/SP280.1>.

The nature and origin of cratons

- Kusky, T.M., Windley, B.F., and Zhai, M.G., 2007b, Lithospheric thinning in eastern Asia: constraints, evolution and tests of models, in Zhai, M.G., Windley, B.F., Kusky, T.M., and Meng, Q.R., eds., Mesozoic Sub-Continental Lithospheric Thinning Under Eastern Asia: Geological Society, London, Special Publication 280, p. 331–343, <https://doi.org/10.1144/SP280.18>.
- Kusky, T.M., Windley, B.F., Wang, L., Wang, Z., Li, X., and Zhu, P., 2014, Flat slab subduction, trench suction, and craton destruction: Comparison of the North China, Wyoming, and Brazilian cratons: Tectonophysics, v. 630, p. 208–221, <https://doi.org/10.1016/j.tecto.2014.05.028>.
- Kusky, T.M., Polat, A., Windley, B.F., Burke, K.C., Dewey, J.F., Kidd, W.S.F., Maruyama, S., Wang, J.P., Deng, H., Wang, Z.S., Wang, C., Fu, D., Li, X.W., and Peng, H.T., 2016, Insights into the tectonic evolution of the North China Craton through comparative tectonic analysis: A record of outward growth of Precambrian continents: Earth-Science Reviews, v. 162, p. 387–432, <https://doi.org/10.1016/j.earscirev.2016.09.002>.
- Kusky, T., Wang, J., Wang, L., Huang, B., Ning, W., Fu, D., Peng, H., Deng, H., Polat, P., Zhong, Y., and Shi, G., 2020, Mélanges through time: Life cycle of the world's largest Archean mélange compared with Mesozoic and Paleozoic subduction-accretion-collision mélanges: Earth-Science Reviews, v. 209, 103303, <https://doi.org/10.1016/j.earscirev.2020.103303>.
- Lawson, A.C., 1890, Note on the pre-Paleozoic surface of the Archean terranes of Canada: Bulletin of the Geological Society of America, v. 1, p. 163–174, <https://doi.org/10.1130/GSAB-1-163>.
- Le Pichon, X., Henry, P., and Goffé, B., 1997, Uplift of Tibet: from eclogites to granulites—implications for the Andean Plateau and the Variscan belt: Tectonophysics, v. 273, no. 1–2, p. 57–76, [https://doi.org/10.1016/S0040-1951\(96\)00288-0](https://doi.org/10.1016/S0040-1951(96)00288-0).
- Li, S.Z., Zhao, G.C., Santosh, M., Liu, X., and Dai, L.M., 2011, Palaeoproterozoic tectonothermal evolution and deep crustal processes in the Jiao-Liao-Ji Belt, North China Craton: a review: Geological Journal, v. 46, p. 525–543, <https://doi.org/10.1002/gj.1282>.
- Lithgow-Bertelloni, C., and Silver, P.G., 1998, Dynamic topography, plate driving forces and the African superswell: Nature, v. 395, no. 6699, p. 269–272, <https://doi.org/10.1038/26212>.
- Liu, J.L., Davis, G.A., Lin, Z.Y., and Wu, F.Y., 2005, The Liaonan metamorphic core complex, southeastern Liaoning Province, north China: a likely contributor to Cretaceous rotation of eastern Liaoning, Korea and contiguous areas: Tectonophysics, v. 407, p. 65–80, <https://doi.org/10.1016/j.tecto.2005.07.001>.
- Lodge, R.W., Gibson, H.L., Stott, G.M., Hudak, G.J., Jirsa, M.A., and Hamilton, M.A., 2013, New U-Pb geochronology from Timiskaming-type assemblages in the Shebandowan and Vermilion greenstone belts, Wawa subprovince, Superior Craton: Implications for the Neoarchean development of the southwestern Superior Province: Precambrian Research, v. 235, p. 264–277, <https://doi.org/10.1016/j.precamres.2013.06.011>.
- Lu, H.F., Yu, H.N., Ding, Y.W., and Zhang, Q.L., 1983, Changing stress field in the middle segment of the Tan-Lu Fault Zone, Eastern China: Tectonophysics, v. 98, p. 253–270, [https://doi.org/10.1016/0040-1951\(83\)90297-4](https://doi.org/10.1016/0040-1951(83)90297-4).
- Marshak, S., and Paulsen, T., 1996, Midcontinent U.S. fault and fold zones: a legacy of Proterozoic intracontinental extensional tectonism?: Geology, v. 24, p. 151–154, [https://doi.org/10.1130/0091-7613\(1996\)024<0151:MUSFAF>2.3.CO;2](https://doi.org/10.1130/0091-7613(1996)024<0151:MUSFAF>2.3.CO;2).
- Marshak, S., and Paulsen, T., 1997, Structural style, regional distribution, and seismic implications of Midcontinent fault-and-fold zones, United States: Seismological Research Letters, v. 68, p. 511–520, <https://doi.org/10.1785/gssrl.68.4.511>.
- Marshak, S., Hamburger, M., and van der Pluijm, B.A., 1998, Tectonics of continental interiors: GSA Today, v. 4, p. 23–24.
- McKenzie, D., and Priestley, K., 2008, The influence of lithospheric thickness variations on continental evolution: Lithos, v. 102, no. 1–2, p. 1–11, <https://doi.org/10.1016/j.lithos.2007.05.005>.
- McKenzie, D., and Priestley, K., 2016, Speculations on the formation of cratons and cratonic basins: Earth and Planetary Science Letters, v. 435, p. 94–104, <https://doi.org/10.1016/j.epsl.2015.12.010>.
- McKenzie, D., and Tribaldos, V.R., 2018, Lithospheric heating by crustal thickening: a possible origin of the Pará Basin: Geological Society, London, Special Publication 472, no. 1, p. 37–44, <https://doi.org/10.1016/j.spl.2017.09.003>.
- Medlicott, H.B., and Blanford, W.T., 1879, A Manual of the Geology of India—Part II: Extra-Peninsular Area: Office of the Superintendent of Government Printing, Calcutta, viii + p. 445–816, 21 plates.
- Menzies, M.A., Fan, W.M., and Zhang, M., 1993, Palaeozoic and Cenozoic lithoprobe and the loss of >120 km of Archean lithosphere, Sino-Korean craton, China, in Prichard, H.M., Alabaster, T., Harris, N.B.W. and Neary, C.R. eds., Magmatic Processes and Plate Tectonics: Geological Society, London, Special Publication 76, p. 71–81.
- Michaut, C., Jaupart, C., and Mareschal, J.C., 2009, Thermal evolution of cratonic roots: Lithos, v. 109, no. 1–2, p. 47–60, <https://doi.org/10.1016/j.lithos.2008.05.008>.
- Miller, M.S., and Eaton, D.W., 2010, Formation of cratonic mantle keels by arc accretion: Evidence from S receiver functions: Geophysical Research Letters, v. 37, no. 18, <https://doi.org/10.1029/2010GL044366>.
- Mojszis, S.J., Harrison, T.M., and Pidgeon, R.T., 2001, Oxygen-isotope evidence from ancient zircons for liquid water at the Earth's surface 4.300 Myr ago: Nature, v. 409, no. 6817, p. 178–181, <https://doi.org/10.1038/35051557>.
- Morris, P.A., Riganti, A., and Chen, S.F., 2007, Evaluating the provenance of Archean sedimentary rocks of the Diemals Formation (central Yilgarn Craton) using whole-rock chemistry and precise U-Pb zircon chronology: Australian Journal of Earth Sciences, v. 54, no. 8, p. 1123–1136, <https://doi.org/10.1080/08120090701615758>.
- Mukhopadhyay, J., Ghosh, G., Zimmermann, U., Guha, S., and Mukherjee, T., 2012, A 3.51 Ga bimodal volcanics-BIF-ultramafic succession from Singhbhum Craton: implications for Palaeoarchaean geodynamic processes from the oldest greenstone succession of the Indian subcontinent: Geological Journal, v. 47, no. 2–3, p. 284–311, <https://doi.org/10.1002/gj.1314>.
- Myers, J.S., 1993, Precambrian history of the West Australian craton and adjacent orogens: Annual Review of Earth and Planetary Sciences, v. 21, p. 453–485, <https://doi.org/10.1146/annurev.ea.21.050193.002321>.
- Myskova, T.A., Zhitnikova, I.A., and L'vov, P.A., 2015, Late Archean intermediate-felsic magmatism of the South Vygozersky and Kamennozersky greenstone structures of Central Karelia: Stratigraphy and Geological Correlation, v. 23, no. 4, p. 351–374, <https://doi.org/10.1134/S0869593815040061>.
- Neuendorf, K.K.E., Mehl, J.P., Jr., and Jackson, J.A., eds., 2005, Glossary of Geology: American Geological Institute, 799 p.
- Nguiri, T.K., Gore, J., James, D.E., Webb, S.J., Wright, C., Zengeni, T.G., Gwava, O., Snoke, J.A., and Kaapvaal Seismic Group, 2001, Crustal structure beneath southern Africa and its implications for the formation and evolution of the Kaapvaal and Zimbabwe cratons: Geophysical Research Letters, v. 28, no. 13, p. 2501–2504, <https://doi.org/10.1029/2000GL012587>.
- Ojakangas, R.W., Srinivasan, R., Hegde, V.S., Chandrakant, S.M., and Srikantha, S.V., 2014, The Talya conglomerate: An Archean (~2.7 Ga) glaciomarine formation, Western Dharwar Craton, Southern India: Current Science, v. 106, p. 387–396.
- Okay, A.I., and Şengör, A.M.C., 1992, Evidence for intracontinental thrust-related exhumation of the ultra-high-pressure rocks in China: Geology, v. 20, no. 5, p. 411–414, [https://doi.org/10.1130/0091-7613\(1992\)020<0411:EFTIRE>2.3.CO;2](https://doi.org/10.1130/0091-7613(1992)020<0411:EFTIRE>2.3.CO;2).
- Okay, A.I., Xu, S., and Şengör, A.M.C., 1989, Coesite from the Dabie Shan eclogites, central China: European Journal of Mineralogy, v. 1, no. 4, p. 595–598, <https://doi.org/10.1127/ejm/1/4/0595>.
- Okay, A.I., Şengör, A.M.C., and Satir, M., 1993, Tectonics of an ultrahigh-pressure metamorphic terrane: the Dabie Shan/Tongbai Shan orogen, China: Tectonics, v. 12, no. 6, p. 1320–1334, <https://doi.org/10.1029/93TC01544>.
- Olszewski, W.J., Wirth, K.R., Gibbs, A.K., and Gaudette, H.E., 1989, The age, origin, and tectonics of the Grão Pará Group and associated rocks, Serra dos Carajás, Brazil: Archean continental volcanism and rifting: Precambrian Research, v. 42, no. 3–4, p. 229–254, [https://doi.org/10.1016/0301-9268\(89\)90013-2](https://doi.org/10.1016/0301-9268(89)90013-2).
- Pang, L., Zhu, X., Hu, B., Wang, W., Sun, Q., and Zhao, T., 2020, Detrital zircon U-Pb age and Hf isotopic composition and whole-rock geochemical characteristics of the Statherian Huangqikou Formation, western margin of the North China Craton: Implications for provenance and tectonic evolution: Precambrian Research, v. 347, 105840, <https://doi.org/10.1016/j.precamres.2020.105840>.
- Percival, J.A., Skulski, T., Sanborn-Barrie, M., Stott, G.M., Leclair, A.D., Corkery, M.T., and Boily, M., 2012, Geology and tectonic evolution of the Superior Province, Canada: Chapter 6, in Percival, J.A., Cook, F.A. and Clowes, R.M., eds., Tectonic Styles in Canada: The LITHOPROBE Perspective: Geological Association of Canada, Special Paper v. 49, p. 321–378.
- Perry, H.K.C., Eaton, D.W.S., and Forte, A.M., 2002, LITH5.0: a revised crustal model for Canada based on LITHOPROBE results: Geophysical Journal International, v. 150, p. 285–294, <https://doi.org/10.1046/j.1365-246X.2002.01712.x>.
- Pitman, W.C., III, and Golovchenko, X., 1991, The effect of sea level changes on the morphology of mountain belts: Journal of Geophysical Research, v. 96, p. 6879–6891, <https://doi.org/10.1029/91JB00250>.
- Polat, A., and Kerrich, R., 2001, Magnesian andesites, Nb-enriched basalt-andesites, and adakites from late-Archean 2.7 Ga Wawa greenstone belts, Superior Province, Canada: implications for late Archean subduction zone petrogenetic processes: Contributions to Mineralogy and Petrology, v. 141, no. 1, p. 36–52, <https://doi.org/10.1007/s004100000223>.
- Polat, A., Hofmann, A.W., and Rosing, M.T., 2002, Boninite-like volcanic rocks in the 3.7–3.8 Ga Isua greenstone belt, West Greenland: geochemical evidence for intra-oceanic subduction zone processes in the early Earth: Chemical Geology, v. 184, no. 3–4, p. 231–254, [https://doi.org/10.1016/S0009-2541\(01\)00363-1](https://doi.org/10.1016/S0009-2541(01)00363-1).
- Polat, A., Frei, R., Appel, P.W.U., Dilek, Y., Fryer, B., Ordóñez-Calderón, J.C., and Yang, Z., 2008, The origin and compositions of Mesarchaean oceanic crust: Evidence from the 3075 Ma Ivisaartoq greenstone belt, SW Greenland: Lithos, v. 100, p. 293–321, <https://doi.org/10.1016/j.lithos.2007.06.021>.
- Polat, A., Longstaffe, F., Weisener, C., Fryer, B., Frei, R., and Kerrich, R., 2012, Extreme element mobility during transformation of Neoarchean (ca. 2.7 Ga) pillow basalts to a Paleoproterozoic (ca. 1.9 Ga) paleosol, Schreiber Beach, Ontario, Canada: Chemical Geology, v. 326–327, p. 145–173, <https://doi.org/10.1016/j.chemgeo.2012.07.018>.
- Polat, A., Frei, R., Longstaffe, F.J., Thorkelson, D.J., and Friedman, E., 2018, Petrology and geochemistry of the Tasse mantle xenoliths of the Canadian Cordillera: A record of Archean to Quaternary mantle growth, metasomatism, removal, and melting: Tectonophysics, v. 737, p. 1–26, <https://doi.org/10.1016/j.tecto.2018.04.014>.
- Pollack, H.N., 1986, Cratonicization and thermal evolution of the mantle: Earth and Planetary Science Letters, v. 80, p. 175–182, [https://doi.org/10.1016/0012-821X\(86\)90031-2](https://doi.org/10.1016/0012-821X(86)90031-2).
- Poulsen, K.H., Robert, F., and Card, K.D., 1992, Transpressive tectonics and the Archean gold deposits of Superior Province, Canadian Shield, in Bartholomew, M.J., Hyndman, D.W., Mogk, D.W. and Mason, R., eds., Basement Tectonics 8. Proceedings of the International Conferences on Basement Tectonics: Springer, Dordrecht, v. 2, p. 615–623, <https://doi.org/10.1007/978-94-011-1614-5>.
- Powell, C.McA., and Conaghan, P.J., 1973, Plate tectonics and the Himalayas: Earth and Planetary Science Letters, v. 20, p. 1–12, [https://doi.org/10.1016/0012-821X\(73\)90134-9](https://doi.org/10.1016/0012-821X(73)90134-9).
- Powell, J.W., 1876, Report on the Geology of the Eastern Portion of the Uinta Mountains and A Region of Country

- Adjacent Thereto, With Atlas: Department of the Interior. U. S. Geological and Geographical Survey of the Territories. Second Division, J.W. Powell, Geologist in Charge, Government Printing Office, Washington, VII+218 p.
- Praeeda, E., John, B., Srinivasan, C., Singh, Y., Divyalakshmi, K.S., and Samui, P., 2015, Thenmala Fault System, Southern India: implication to neotectonics: Journal of the Geological Society of India, v. 86, p. 391–398, <https://doi.org/10.1007/s12594-015-0327-z>.
- Priestley, K., Ho, T., and McKenzie, D., 2021, The formation of continental roots: *Geology*, v. 49, p. 190–194, <https://doi.org/10.1130/G47696.1>.
- Ranalli, G., and Murphy, D.C., 1987, Rheological stratification of the lithosphere: *Tectonophysics*, v. 132, no. 4, p. 281–295, [https://doi.org/10.1016/0040-1951\(87\)90348-9](https://doi.org/10.1016/0040-1951(87)90348-9).
- Santosh, M., Teng, X.M., He, X.F., Tang, L., and Yang, Q.Y., 2016, Discovery of Neoarchean suprasubduction zone ophiolite suite from Yishui Complex in the North China Craton: *Gondwana Research*, v. 38, p. 1–27, <https://doi.org/10.1016/j.gr.2015.10.017>.
- Sengör, A.M.C., 1981, The geological exploration of Tibet: *Nature*, v. 294, p. 403–404, <https://doi.org/10.1038/294403a0>.
- Sengör, A.M.C., 1984, The Cimmeride Orogenic System and the Tectonics of Eurasia: *Geological Society of America Special Paper* 195, 82 p., <https://doi.org/10.1130/SPE195-p1>.
- Sengör, A.M.C., 1990, Plate tectonics and orogenic research after 25 years: A Tethyan perspective: *Earth-Science Reviews*, v. 27, p. 1–201, [https://doi.org/10.1016/0012-8252\(90\)90002-D](https://doi.org/10.1016/0012-8252(90)90002-D).
- Sengör, A.M.C., 1995, Sedimentation and tectonics of fossil rifts, in Busby, C.J., and Ingersoll, R.V., eds., *Tectonics of Sedimentary Basins*: Blackwell, Oxford, p. 53–117.
- Sengör, A.M.C., 1999, Continental interiors and cratons: any relation?: *Tectonophysics*, v. 305, no. 1–3, p. 1–42, [https://doi.org/10.1016/S0040-1991\(99\)00043-8](https://doi.org/10.1016/S0040-1991(99)00043-8).
- Sengör, A.M.C., 2001, Elevation as indicator of mantle plume activity, in Ernst, R., and Buchan, K., eds., *Mantle Plumes: Their Identification Through Time*: *Geological Society of America Special Paper* 352, p. 183–225, <https://doi.org/10.1130/0-8137-2352-3.183>.
- Sengör, A.M.C., 2003, The Large Wavelength Deformations of the Lithosphere: Materials for a History of the Evolution of Thought from the Earliest Times to Plate Tectonics: *Geological Society of America Memoir* 196, xvii+347 p., 3 folded plates in pocket, <https://doi.org/10.1130/MEM196>.
- Sengör, A.M.C., 2018, What has remained of Hans Stille's tectonics?: *Global Tectonics and Metallogenesis*, v. 10, no. 2–4, p. 77–107, <https://doi.org/10.1127/gtm/2018/0012>.
- Sengör, A.M.C., 2020, Orogenic belts, in Gupta, H.K., ed., *Encyclopedia of Solid Earth Geophysics*: Springer Nature Switzerland, *Encyclopedia of Earth Sciences Series*, https://doi.org/10.1007/978-3-030-10475-7_253-1.
- Sengör, A.M.C., and Natal' in, B.A., 1996, Palaeotectonics of Asia: Fragments of A Synthesis, in Yin, A., and Harrison, M., eds., *The Tectonic Evolution of Asia*: Rubey Colloquium: Cambridge, Cambridge University Press, p. 486–640.
- Sengör, A.M.C., and Okuroğulları, A.H., 1991, The role of accretionary wedges in the growth of continents: Asiatic examples from Argand to plate tectonics: *Eclogae Geologicae Helvetiae*, v. 84, p. 535–597.
- Sengör, A.M.C., Natal' in, B.A., and Burtman, V.S., 1993, Evolution of the Altai tectonic collage and Palaeozoic crustal growth in Eurasia: *Nature*, v. 364, no. 6435, p. 299–307, <https://doi.org/10.1038/364299a0>.
- Sengör, A.M.C., Natal' in, B.A., Sunal, G., and Van der Voo, R., 2014a, A New Look at The Altaiids: A Superorogenic Complex in Northern and Central Asia as a Factory of Continental Crust. Part I: Geological Data Compilation (Exclusive of Palaeomagnetic Observations): Mitteilungen der Österreichischen Geologischen Gesellschaft, v. 107, no. 1, p. 169–232.
- Sengör, A.M.C., Natal' in, B., van der Voo, R., and Sunal, G., 2014b, A new look at the Altaiids: a superorogenic complex in northern and central Asia as a factory of continental crust. Part II: palaeomagnetic data, reconstructions, crustal growth and global sea-level: *Mitteilungen der Österreichischen Geologischen Gesellschaft*, v. 107, no. 2, p. 131–181.
- Sengör, A.M.C., Lom, N., and Sağıdıç, N.G., 2018a, Tectonic inheritance, structure reactivation and lithospheric strength: the relevance of geological history, in Wilson, R.W., Houseman, G.A., McCaffrey, K.J.W., Doré, A.G. and Buitier, S.J.H., eds., *Fifty Years of the Wilson Cycle Concept in Plate Tectonics*: Geological Society, London, Special Publication 470, p. 105–136, <https://doi.org/10.1144/SP470.8>.
- Sengör, A.M.C., Natal' in, B.A., Sunal, G., and van der Voo, R., 2018b, The tectonics of the Altaiids: crustal growth during the construction of the continental lithosphere of Central Asia between ~750 and ~130 Ma Ago: Annual Review of Earth and Planetary Sciences, v. 46, p. 439–494, <https://doi.org/10.1146/annurev-earth-060313-054826>.
- Sengör, A.M.C., Lom, N., Sunal, G., Zabci, C., and Öner, T., 2020, Reconstructing orogens without biostratigraphy: The Saharides and continental growth during the final assembly of Gondwana-Land: *Proceedings of the National Academy of Sciences of the United States of America*, v. 117, no. 51, p. 32,278–32,284, <https://doi.org/10.1073/pnas.2015117117>.
- Sengör, A.M.C., Lom, N., Zabci, C., Sunal, G., and Öner, T., 2021, The Saharides: Turkic-Type Orogeny in Afro-Arabia: *International Journal of Earth Sciences*, <https://doi.org/10.1007/s00531-021-02063-3>.
- Servali, A., and Korenaga, J., 2018, Oceanic origin of continental mantle lithosphere: *Geology*, v. 46, p. 1047–1050, <https://doi.org/10.1130/G45180.1>.
- Shirey, S.B., Richardson, S.H., and Harris, J.W., 2004, Integrated models of diamond formation and craton evolution: *Lithos*, v. 77, no. 1–4, p. 923–944, <https://doi.org/10.1016/j.lithos.2004.04.018>.
- Smithies, R.H., Champion, D.C., Van Kranendonk, M.J., Howard, H.M., and Hickman, A.H., 2005, Modern-style subduction processes in the Mesoarcheans: geochemical evidence from the 3.12 Ga Whundo intra-oceanic arc: *Earth and Planetary Science Letters*, v. 231, no. 3–4, p. 221–237, <https://doi.org/10.1016/j.epsl.2004.12.026>.
- Smithies, R.H., Lu, Y., Gessner, K., Wingate, M.T.D., and Champion, D.C., 2018, Geochemistry of Archean granitic rocks in the South West Terrane of the Yilgarn Craton: *Geological Survey of Western Australia, Record* 2018/10, 13 p.
- Snyder, W.S., Dickinson, W.R., and Silberman, M.L., 1976, Tectonic implications of space–time patterns of Cenozoic magmatism in the western United States: *Earth and Planetary Science Letters*, v. 32, p. 91–106, [https://doi.org/10.1016/0012-821X\(76\)90189-8](https://doi.org/10.1016/0012-821X(76)90189-8).
- Stille, H., 1920, Über Alter und Art variszischer Gebirgsbildung: *Nachrichten von der Gesellschaft für die Wissenschaften zu Göttingen, mathematisch-physikalische Klasse*, year 1920, p. 218–224.
- Stille, H., 1936, Tektonische Beziehungen zwischen Nordamerika und Europa, in *International Geological Congress Rep. XVI Session, U.S.A. 1933*, v. 2: Menasha, Wisconsin, George Banta Publishing Company, p. 829–838.
- Stille, H., 1940, Einführung in den Bau Amerikas: *Gebrüder Borntraeger*, Berlin, XX+717 p.
- Suess, E., 1875, *Die Entstehung der Alpen*: W. Braumüller, Wien, IV+168 p.
- Suess, E., 1883, Das Antlitz der Erde, v. Ia (Erste Abtheilung): F. Tempsky, Prag and G. Freytag, Leipzig, 310 p.
- Suess, E., 1909, The Face of the Earth (Das Antlitz der Erde), translated by H.B.C. Sollas under the direction of W.J. Sollas, v. IV: Clarendon Press, Oxford, viii+678 p.
- Sun, J.M., 2005, Long-term fluvial archives in the Fen Wei Graben, central China, and their bearing on the tectonic history of the India-Asia collision system during the Quaternary: *Quaternary Science Reviews*, v. 24, p. 1279–1286, <https://doi.org/10.1016/j.quascirev.2004.08.018>.
- Sun, Q., Zhou, Y., Zhao, T., Pang, L., and Qiu, Y., 2020, Geochronology and geochemistry of the Paleoproterozoic Fangniushan supracrustal strata in the Xiaoshan area, southern North China Craton: Implications for tectonic evolution: *Precambrian Research*, v. 346, 105789, <https://doi.org/10.1016/j.precamres.2020.105789>.
- Swager, C.P., Witt, W.K., Griffin, T.J., Ahmat, A.L., Hunter, W.M., McGoldrick, P.J., and Wyche, S., 1992, Late Archaean granite-greenstones of the Kalgoorlie terrane, Yilgarn craton, Western Australia, in Glover, J.E. and Ho, S.E., eds., *The Archaean: Terrains, Processes and Metallogeny*, Geology Department (Key Centre) & University Extension: University of Western Australia Publication, v. 22, p. 107–122.
- Tang, Y.J., Zhang, H.F., Ying, J.F., Zhang, J., and Liu, X.M., 2008, Refertilisation of ancient lithospheric mantle beneath central North China Craton: evidence from petrology and geochemistry of peridotite xenoliths: *Lithos*, v. 101, p. 435–452, <https://doi.org/10.1016/j.lithos.2007.09.006>.
- Taylor, S.R., and McLennan, S.M., 1985, *The Continental Crust: Its Composition and Evolution*: Boston, Massachusetts, USA, Blackwell Scientific, 312 p.
- Taylor, S.R., and McLennan, S.M., 1995, The geochemical evolution of the continental crust: *Reviews of Geophysics*, v. 33, no. 2, p. 241–265, <https://doi.org/10.1029/95RG00262>.
- Teilhard de Chardin, P., 1932, *Observations géologiques à travers les déserts d'Asie Centrale de Kalgan à Hami (Mission Citroën-Centre Asie, 1931–1932)*: Revue de Géographie Physique et de Géologie Dynamique, v. 5, p. 365–396+plates 35–38 and one foldout colored geological map.
- Teilhard de Chardin, P., and Licent, E., 1924, *Observations géologiques sur la bordure occidentale et méridionale de l'Ordos: Bulletin de la Société Géologique de France*, sér. 4, v. 24, p. 49–91.
- Tesuaro, M., Kaban, M.K., and Cloetingh, S.A.P.L., 2012, Global strength and elastic thickness of the lithosphere: *Global and Planetary Change*, v. 90–91, p. 51–57, <https://doi.org/10.1016/j.gloplacha.2011.12.003>.
- Tunini, L., Jiménez-Munt, I., Fernandez, M., Vergés, J., and Bird, P., 2017, Neotectonic Deformation in Central Eurasia: A Geodynamic Model Approach: *Journal of Geophysical Research: Solid Earth*, v. 122, p. 9461–9484, <https://doi.org/10.1002/2017JB014487>.
- Turner, S., Rushmer, T., Reagan, M., and Moyen, J.F., 2014, Heading down early? Start of subduction on Earth: *Geology*, v. 42, p. 139–142, <https://doi.org/10.1130/G34886.1>.
- Turner, S., Wilde, S., Wörner, G., Schaefer, B., and Lai, Y.J., 2020, An andesitic source for Jack Hills zircon supports onset of plate tectonics in the Hadean: *Nature Communications*, v. 11, p. 1241, <https://doi.org/10.1038/s41467-020-14857-1>.
- Valdiya, K.S., 1996, Antecedent rivers: *Resonance*, v. 1, no. 8, p. 55–63, <https://doi.org/10.1007/BF02837023>.
- Wang, C.-S., Dai, J.-G., Zhao, X.-X., Li, Y.-L., Graham, S.A., He, D.-F., Ran, B., and Meng, J., 2014a, Outward-growth of the Tibetan Plateau during the Cenozoic: A review: *Tectonophysics*, v. 621, p. 1–43, <https://doi.org/10.1016/j.tecto.2014.01.036>.
- Wang, G.-C., Thybo, H., and Artemieva, I., 2021, No mafic layer in 80 km thick Tibetan crust: *Nature Communications*, v. 12, 1069, <https://doi.org/10.1038/s41467-021-21420-z>.
- Wang, H., van Hunen, J., Pearson, D.G., and Allen, M.B., 2014b, Craton stability and longevity: the roles of composition-dependent rheology and buoyancy: *Earth and Planetary Science Letters*, v. 391, p. 224–233, <https://doi.org/10.1016/j.epsl.2014.01.038>.
- Wang, Q.F., Deng, J., Huang, D.H., Yang, T.Q., Gao, B.F., Xu, H., and Jiang, S.Q., 2006, Tectonic constraints on the transformation of Paleozoic framework of uplift and depression in the Ordos area: *Acta Geologica Sinica*, v. 80, p. 944–953.
- Wang, Y., and Dou, L.R., 1997, Formation time and dynamic characteristics of the northern part of the Tanlu Fault Zone in East China: *Dizhen Dizhi*, v. 19, p. 185–192.
- Wang, Y., Zhou, L., Liu, S., Li, J., and Yang, T., 2018, Post-cratonization deformation processes and tectonic evolution of the North China Craton: *Earth-Science Reviews*, v. 177, p. 320–365, <https://doi.org/10.1016/j.earscirev.2017.11.017>.
- Wilson, A.W.G., 1903, The Laurentian Peneplain: *The Journal of Geology*, v. 11, p. 615–667, <https://doi.org/10.1086/621104>.
- Windley, B.F., Kusky, T., and Polat, A., 2021, Onset of plate tectonics by the Eoarchean: *Precambrian Research*, <https://doi.org/10.1016/j.precamres.2020.105980>.
- Windley, B.F., Maruyama, S., and Xiao, W.J., 2010, Delamination/thinning of sub-continental lithospheric mantle under Eastern China: the role of water and multiple subduction: *American Journal of Science*, v. 310, p. 1250–1293, <https://doi.org/10.2475/10.2010.03>.

The nature and origin of cratons

- Wu, F.Y., Yang, J.H., Xu, Y.G., Wilde, S.A., and Walker, R.J., 2019, Destruction of the North China Craton in the Mesozoic: Annual Review of Earth and Planetary Sciences, v. 47, p. 173–195, <https://doi.org/10.1146/annurev-earth-053018-060342>.
- Wyman, D.A., Kerrich, R., and Polat, A., 2002, Assembly of Archean cratonic mantle lithosphere and crust: plume-arc interaction in the Abitibi–Wawa subduction-accretion complex: Precambrian Research, v. 115, no. 1–4, p. 37–62, [https://doi.org/10.1016/S0301-9268\(02\)00005-0](https://doi.org/10.1016/S0301-9268(02)00005-0).
- Xia, B., Thybo, H., Artemieva, I.M., 2020, Lithosphere mantle density of the North China Craton: Journal of Geophysical Research. Solid Earth, v. 125, <https://doi.org/10.1029/2020JB020296>.
- Xiao, W.J., Windley, B.F., Hao, J., and Zhai, M.G., 2003, Accretion leading to collision and the Permian Solonker suture, Inner Mongolia, China: termination of the Central Asian Orogenic Belt: Tectonics, v. 22, <https://doi.org/10.1029/2002TC001484>.
- Xu, S.T., Okay, A.I., Ji, S., Şengör, A.M.C., Su, W., Liu, Y.C., and Jiang, L.I., 1992, Diamond from the Dabie Shan Metamorphic Rocks and its implication for tectonic setting: Science, v. 256, p. 80–82, <https://doi.org/10.1126/science.256.5053.80>.
- Yang, C., Liu, J., Yang, H., Zhang, C., Feng, J., Lu, T., Sun, Y., and Zhang, J., 2019, Tectonics of the Paleoproterozoic Jiao-Liao-Ji orogenic belt in the Liaodong peninsula, North China Craton: A review: Journal of Asian Earth Sciences, v. 176, p. 141–156, <https://doi.org/10.1016/j.jseaes.2019.01.028>.
- Yang, J.H., Wu, F.Y., Wilde, S.A., Belousova, E., and Griffin, W.L., 2008, Mesozoic decratonization of the North China block: Geology, v. 36, p. 467–470, <https://doi.org/10.1130/G24518A.1>.
- Yang, J.J. and Zhang, B.R., 1987, Overthrust Zone in West Margin of Shan-Gan-Ning Basin and its Hydrocarbon Discovery: Changqing Petroleum Exploration Bureau, Gansu, Qingyang, 14 p., 9 plates.
- Yuan, W., and Yang, Z., 2015, The Alashan Terrane did not amalgamate with North China block by the Late Permian: evidence from Carboniferous and Permian paleomagnetic results: Journal of Asian Earth Sciences, v. 104, p. 145–159, <https://doi.org/10.1016/j.jseaes.2014.02.010>.
- Yang, Y.T., and Xu, T.G., 2004, Hydrocarbon habitat of the offshore Bohai Basin, China: Marine and Petroleum Geology, v. 21, p. 691–708, <https://doi.org/10.1016/j.marpgeo.2004.03.008>.
- Zandt, G., Beck, S.L., Ruppert, S.R., Rock, A.D., Minaya, E., Wallace, T.C., and Silver, P.G., 1996, Anomalous crust of the Bolivian Altiplano, central Andes: Constraints from broadband regional seismic waveforms: Geophysical Research Letters, v. 23, p. 1159–1162, <https://doi.org/10.1029/96GL00967>.
- Zhai, M.G., 2011, Cratonization and the ancient North China continent: a summary and review: Science China. Earth Sciences, v. 54, p. 1110–1120, <https://doi.org/10.1007/s11430-011-4250-x>.
- Zhai, M.G., 2014, Multi-stage crustal growth and cratonization of the North China Craton: Geoscience Frontiers, v. 5, p. 457–469, <https://doi.org/10.1016/j.gsf.2014.01.003>.
- Zhai, M.G., and Santosh, M., 2011, The early Precambrian odyssey of the North China craton: a synoptic overview: Gondwana Research, v. 20, p. 6–25, <https://doi.org/10.1016/j.gr.2011.02.005>.
- Zhai, M.G., Fan, Q.C., Zhang, H.F., and Sui, J.L., 2007, Lower crustal processes leading to Mesozoic lithospheric thinning beneath Eastern North China: underplating, replacement and delamination: Lithos, v. 96, p. 36–54, <https://doi.org/10.1016/j.lithos.2006.09.016>.
- Zhai, M.G., Li, T.S., Peng, P., Hu, B., Liu, F., Zhang, Y.B., and Guo, J.H., 2010, Precambrian key tectonic events and evolution of the North China Craton, in Kusky, T.M., Zhai, M.G. and Xiao, W.J., eds., The Evolving Continents: Geological Society, London, Special Publication 338, p. 235–262, <https://doi.org/10.1144/SP338.12>.
- Zhang, S.H., Zhao, Y., Song, B., Yang, Z.Y., Hu, J.M., and Wu, H., 2007, Carboniferous granitic plutons from the northern margin of the North China block: implications for a late Palaeozoic active continental margin: Journal of the Geological Society, v. 164, p. 451–463, <https://doi.org/10.1144/0016-76492005-190>.
- Zhang, J.A., Zhao, G.C., Li, S.Z., Sun, M., Liu, S.W., Xia, X.P., and He, Y.H., 2006, U-Pb zircon dating of the granitic conglomerates of the Hutuo Group: affinities to the Wutai granitoids and significance to the tectonic evolution of the Trans-North China orogen: Acta Geologica Sinica, v. 80, p. 886–898, <https://doi.org/10.1111/j.1755-6724.2006.tb00310.x>.
- Zhao, G.C., and Zhai, M.G., 2013, Lithotectonic elements of Precambrian basement in the North China Craton: review and tectonic implications: Gondwana Research, v. 23, p. 1207–1240, <https://doi.org/10.1016/j.gr.2012.08.016>.
- Zhao, G.C., Sun, M., Wilde, S.A., and Li, S.Z., 2005, Late Archean to Paleoproterozoic evolution of the North China Craton: key issues revisited: Precambrian Research, v. 136, p. 177–202, <https://doi.org/10.1016/j.precamres.2004.10.002>.
- Zhao, G.C., Liu, S.W., Sun, M., Li, S.Z., Wilde, S., Xia, X.P., Zhang, J., and He, Y.H., 2006, What happened in the Trans-North China orogen in the period 2560–1850?: Acta Geologica Sinica, v. 80, p. 790–806, <https://doi.org/10.1111/j.1755-6724.2006.tb00303.x>.
- Zhao, G.C., Cawood, P.A., Li, S.Z., Wilde, S.A., Sun, M., Zhang, J., He, Y.H., and Yin, C.Q., 2012, Amalgamation of the North China Craton: key issues and discussion: Precambrian Research, v. 222–223, p. 55–76, <https://doi.org/10.1016/j.precamres.2012.09.016>.
- Zhao, L.A., and Zheng, T.Y., 2007, Complex upper-mantle deformation beneath North China Craton: implications for lithospheric thinning: Geophysical Journal International, v. 170, p. 1095–1099, <https://doi.org/10.1111/j.1365-246X.2007.03512.x>.
- Zhao, Y., Li, N., Jiang, Y., and Niu, H., 2021, Petrogenesis and tectonic implications of the ca. 2.5 Ga granitoid assemblage in the eastern Zhongtiao region, southern North China Craton: Lithos, v. 386–387, p. 106007, <https://doi.org/10.1016/j.lithos.2021.106007>.
- Zhao, Z.Y., and Windley, B.F., 1990, Cenozoic tectonic extension and inversion of the Jizhang Basin, northern China: Tectonophysics, v. 185, p. 83–89, [https://doi.org/10.1016/0040-1951\(90\)90406-X](https://doi.org/10.1016/0040-1951(90)90406-X).
- Zheng, J.P., Griffin, W.L., O'Reilly, S.Y., Yang, J.S., and Zhang, R.Y., 2006, A refractory mantle protolith in younger continental crust, east-central China: age and composition of zircon in the Sulu ultrahigh-pressure peridotite: Geology, v. 34, p. 705–708, <https://doi.org/10.1130/G22569.1>.
- Zheng, J.P., Griffin, W.L., O'Reilly, S.Y., Yu, C.M., Zhang, H.F., Pearson, N., and Zhang, M., 2007, Mechanism and timing of lithospheric modification and replacement beneath the eastern North China Craton: Peridotitic xenoliths from the 100 Ma Fuxin basalts and a regional synthesis: Geochimica et Cosmochimica Acta, v. 71, p. 5203–5225, <https://doi.org/10.1016/j.gca.2007.07.028>.
- Zheng, Y., Xu, Z., Zhao, Z., and Dai, L., 2018, Mesozoic mafic magmatism in North China: Implications for thinning and destruction of cratonic lithosphere: Science China. Earth Sciences, v. 61, p. 353–385, <https://doi.org/10.1007/s11430-017-9160-3>.
- Zhu, R.X., Chen, L., Wu, F.Y., and Liu, J.L., 2011, Timing, scale and mechanism of the destruction of the North China Craton: Science China. Earth Sciences, v. 54, p. 789–797, <https://doi.org/10.1007/s11430-011-4203-4>.
- Zhu, R., Xu, Y.G., Zhu, G., Zhang, H.F., Xia, Q.K., and Zheng, T.Y., 2012, Destruction of the North China Craton: Science China. Earth Sciences, v. 55, p. 1565–1587, <https://doi.org/10.1007/s11430-012-4516-y>.
- Zuza, A.V., Wu, C., Wang, Z., Levy, D.A., Li, B., Xiong, X., and Chen, X., 2019, Underthrusting and duplexing beneath the northern Tibetan Plateau and the evolution of the Himalayan-Tibetan orogen: Lithosphere, v. 11, no. 2, p. 209–231, <https://doi.org/10.1130/L1042.1>.

SCIENCE EDITOR: WENJIAO XIAO
ASSOCIATE EDITOR: TIMOTHY KUSKY

MANUSCRIPT RECEIVED 11 FEBRUARY 2021
REVISED MANUSCRIPT RECEIVED 10 JUNE 2021
MANUSCRIPT ACCEPTED 19 JULY 2021

Printed in the USA