

A classification of well-behaved graph clustering schemes

Vilhelm Agdur

28 September 2023

1 Introduction

general stuff about clustering and why it is an interesting problem

some talk about why optimization-based methods have poor theoretical guarantees

some talk about why an axiomatic approach gives algorithms that run in polynomial time, unlike the generally NP-hard optimization problems

soft heuristic statement of what our axioms/properties are and why they might be desirable

statement of our results

2 Definitions of our terms

Definition 1. The category of simple graphs $\mathcal{G}$ has as its objects simple graphs, which we consider to be a finite set V equipped with a reflexive and symmetric binary relation E – that is, our simple graphs are finite, have no double edges, and have loops at every vertex¹. A morphism from $H = (V, E)$ to $G = (V', E')$ is an injective set function $f : V \rightarrow V'$ such that whenever uEv , we also have $f(u)E'f(v)$. That is, a morphism from H to G is a way of realizing H as a subgraph of G .

Definition 2. The category of partitioned sets $\mathcal{P}$ has as its objects sets V with an equivalence relation $\sim$. A morphism in $\mathcal{P}$ from $(V, \sim)$ to $(W, \approx)$ is a set function $f : V \rightarrow W$ such that whenever $v \sim w$, $f(v) \approx f(w)$. So in the case where $V = W$, $f = \text{Id}$ is a morphism if $\approx$ is a coarsening of $\sim$.

Definition 3. A clustering scheme is a function² $\mathfrak{C}$ that sends simple graphs to partitioned sets, with the same underlying set. We say that it is *functorial* if it is a functor from $\mathcal{G}$ to $\mathcal{P}$.

Definition 4. A clustering scheme $\mathfrak{C}$ is *excisive* if, for all graphs $G = (V, E)$, it holds for every equivalence class A of $\mathfrak{C}(G)$ that $\mathfrak{C}(G|_A) = (A, A^2)$. That is, if we take a graph, cluster it, pick out one of the parts, and cluster that part alone, that part will be clustered into a single part.

¹The final property here may seem a bit unusual, but it simplifies our proofs in some places, without really affecting much.

²Technically, of course, neither $\mathcal{G}$ nor $\mathcal{P}$ are small categories, so the word “function” should be read as “class function” here, not in the usual sense of the word. We sweep this issue under the rug here, because it will never affect anything.

3 Representability

Definition 5. Given a not necessarily finite set Ω of simple graphs, we define the endofunctor F_Ω on $\mathcal{G}$ as follows: For any graph $G = (V, E)$, $F_\Omega(G)$ is a graph on the same vertex set, and there is an edge between u and v in $F_\Omega(G)$ whenever there exists some subgraph of G containing both u and v which is isomorphic to some $\omega \in \Omega$.

Equivalently, there is an edge uv in $F_\Omega(G)$ whenever there exists an $\omega \in \Omega$ and a morphism $f : \omega \rightarrow G$ such that $f(\omega) \ni u, v$.

On morphisms, F_Ω is always just the identity.

Lemma 6. *For any representing set Ω of simple graphs, F_Ω is indeed an endofunctor on $\mathcal{G}$.*

Proof. That it always gives a simple graph and that it respects identity morphisms and composition of morphisms is easy to see. The only thing we actually need to check is that if $f : H \rightarrow G$ is a morphism, then $F_\Omega(f) = f : F_\Omega(H) \rightarrow F_\Omega(G)$ is also a morphism.

Unwrapping this statement, what we need is that, for any morphism $f : H \rightarrow G$, whenever uv is an edge in $F_\Omega(H)$, $f(u)f(v)$ is an edge in $F_\Omega(G)$. Now, that uv is an edge in $F_\Omega(H)$ means that there is an $\omega \in \Omega$ and $f_\omega : \omega \rightarrow H$ such that $f_\omega(\omega) \ni u, v$.

But consider what happens if we just compose f_ω and f – this will be a morphism from ω into G , and trivially we must have that $f(f_\omega(\omega)) \ni f(u), f(v)$, and so there must be an edge $f(u)f(v)$ in $F_\Omega(G)$, by definition of F_Ω . $\square$

What is going on in the proof of Lemma 6 is much clearer if we just think about it in terms of things being subgraphs of each other, forgetting the data of *how* exactly they are subgraphs. Then the statement that F_Ω is a functor boils down to the statement that if ω is a subgraph of H and H is a subgraph of G , then certainly ω is a subgraph of G , and so $F_\Omega(G)$ must have all the edges $F_\Omega(H)$ has, since edges in $F_\Omega(G)$ represent the presence of a certain subgraph in G .

Definition 7. The functor $\Pi : \mathcal{G} \rightarrow \mathcal{P}$ which sends a graph to its partition into connected components is called the connected components functor. Equivalently, we can think of it as taking the equivalence relation hull of the binary relation E defining the edges.

Definition 8. A clustering scheme $\mathfrak{C}$ is *representable* if we can write it as $F_\Omega \circ \Pi$ for some representable endofunctor F_Ω . Notice that this means that all representable clustering schemes are functorial, being the composition of two functors.

One somewhat surprising fact is that there are actually representable clustering schemes that are not finitely representable.

Theorem 9. *There exists a representable endofunctor F_Ω which is not finitely representable, that is, it is not equal to F_Γ for any finite set of simple graphs Γ .*

In order to give a proof of this, we first need a structural result about how these endofunctors behave.

Remark 10. For any $\omega \in \Omega$, it is trivial to see that $F_\Omega(\omega)$ must be a clique, since we can just take the identity morphism on ω to show that all edges must exist.

In fact, which graphs are sent to cliques by F_Ω essentially determines what it does as a functor. We will see this even more precisely in the next section, but for now this lemma will suffice for our needs:




Figure 1: The family of graphs $\{\omega_i\}_{i=0}^3$.

Lemma 11. *For any set Ω of simple graphs and any simple graph $G = (V, E)$, it holds that $F_{\Omega \cup \{G\}} = F_\Omega$ if and only if $F_\Omega(G) = (V, V^2)$, that is, if G is sent to a clique on its vertices by F_Ω .*

Proof. Suppose $F_{\Omega \cup \{G\}} = F_\Omega$. That $F_{\Omega \cup \{G\}}(G) = (V, V^2)$ is immediate by Remark 10, and so since $F_{\Omega \cup \{G\}} = F_\Omega$, we also have $F_\Omega(G) = (V, V^2)$, as desired.

Now suppose $F_\Omega(G) = (V, V^2)$, and let H be some arbitrary simple graph. We wish to show that $F_{\Omega \cup \{G\}}(H) = F_\Omega(H)$.

That edges of $F_\Omega(H)$ must also be edges of $F_{\Omega \cup \{G\}}(H)$ is obvious, so all we need to show is that edges of $F_{\Omega \cup \{G\}}(H)$ are also edges of $F_\Omega(H)$. So suppose uv is an edge of $F_{\Omega \cup \{G\}}(H)$ – that this is so must be because there is some $\omega \in \Omega \cup \{G\}$ and an $f : \omega \rightarrow H$ such that $f(\omega) \ni u, v$.

If this ω is not G , then we are done. If it is G , we recall that since $F_\Omega(G)$ is a clique, there must be an edge $f^{-1}(u)f^{-1}(v)$ in it. Therefore there must exist some $\omega \in \Omega$ and some $g : \omega \rightarrow G$ such that $f(\omega) \ni f^{-1}(u), f^{-1}(v)$. But then the composition $g \circ f : \omega \rightarrow H$ will be a morphism from an $\omega \in \Omega$ into H whose image contains u and v , proving that uv is an edge of $F_\Omega(H)$. $\square$

With this result in hand, we can now give a proof of the existence of a representable but not finitely representable endofunctor.

Proof of Theorem 9. Let, for each i , ω_i consist of a triangle with a tail of i vertices, as shown in Figure 3, and let $\Omega = \{\omega_i \mid i \in \mathbb{N}\}$. We claim that F_Ω is not equal to F_Γ for any finite set of simple graphs Γ .

To prove this, let us first define

$$\bar{\Omega} = \{G \in \mathcal{G} \mid F_\Omega(G) = (V_G, V_G^2)\}$$

as the set of all things which F_Ω sends to cliques. By Lemma 11, $F_\Omega = F_{\bar{\Omega}}$, and $\bar{\Omega}$ is the greatest set with this property.

A little bit of reflection will make it clear that $\bar{\Omega}$ is precisely the set of connected graphs which contain a triangle. Particularly, for any edge in such a graph, we can map the tail end of some ω_i onto that edge, the rest of the tail onto a path to the triangle in the graph, and the triangle of the ω_i onto the triangle of the graph. That F_Ω will send any triangle-free graph to a graph with no edges is clear, since there is no morphism from any triangle-containing graph such as ω_i into a triangle-free graph. Likewise, that it cannot send a disconnected graph to a clique is obvious.

Now suppose for contradiction that Γ is a finite set of graphs such that $F_\Gamma = F_{\bar{\Omega}}$. Lemma 11 implies that we must have $\Gamma \subset \bar{\Omega}$, since if there were a $\gamma \in \Gamma \setminus \bar{\Omega}$ it'd be sent to a clique by F_Γ , and thus also by $F_{\bar{\Omega}}$, since we've assumed they are equal – but we chose $\bar{\Omega}$ so that it already contains everything sent to a clique by $F_{\bar{\Omega}}$.

So, let

$$\delta = \max_{\gamma \in \Gamma} \max \{d(u, v) \mid u, v \in V_\gamma, v \text{ is in a triangle}\}.$$

That is, δ is the furthest any vertex in any graph in Γ is from a triangle. We claim that F_Γ will not send $\omega_{\delta+1}$ to a clique, proving it is not equal to $F_{\bar{\Omega}}$, giving our contradiction.

That this is so is actually easy to see – if $\omega_{\delta+1}$ were sent to a clique, that must in particular mean there is an edge in $F_\Gamma(\omega_{\delta+1})$ between the tail vertex and its neighbour. So there is some morphism from some $\gamma \in \Gamma$ that hits both vertices. However, such a morphism must also map the triangle in γ onto the triangle in $\omega_{\delta+1}$, since that is the only place the triangle can go. Then, however, it cannot also reach to the tail – the tail is further away from the triangle than any vertex in any γ is from the triangle. So no such morphism can exist, and we are done. $\square$

Remark 12. The construction in our proof of course works equally as well replacing the triangle by any connected graph that is not a line. So we have found a whole lot of examples of classes of graphs that give representable but not finitely representable endofunctors.

The question of which natural classes of graphs are representable or finitely representable may be interesting. The connected graphs, for example, are of course represented by just a K_2 .

4 Proof of equivalence of representability and excisiveness

We are able to exactly characterize the relationship between the three concepts of functoriality, representability, and excisiveness for clustering schemes.

Theorem 13. *A representable clustering scheme is excisive. An excisive and functorial clustering scheme is representable. No other implications hold between these concepts.*

We divide the proof of Theorem 13 into two lemmas and a collection of examples.

Lemma 14. *A representable clustering scheme is excisive.*

Proof. Let Ω be some set of simple graphs, and G be some simple graph. Assume the equivalence classes of $(F_\Omega \circ \Pi)(G)$ are $\{A_1, A_2, \dots, A_k\}$, and let $H = G|_{A_1}$ be the induced subgraph of G on A_1 . What we need to show is that $(F_\Omega \circ \Pi)(H) = (A_1, A_1^2)$, that is, that every vertex is equivalent in the clustering of H .

It is obvious that this is equivalent to showing that $F_\Omega(H)$ is a connected graph. We will in fact show something stronger – that $F_\Omega(H) = F_\Omega(G_{A_1}) = (F_\Omega(G))|_{A_1}$. Since A_1 is by assumption a connected component of $F_\Omega(G)$, this will show what we want.

First, let $f : H \hookrightarrow G$ be the subgraph inclusion morphism of H into G . By functoriality of F_Ω , f is also a morphism from $F_\Omega(H)$ into $F_\Omega(G)$, showing that $F_\Omega(H)$ is a subgraph of $F_\Omega(G)$, so every edge of $F_\Omega(H)$ is an edge of $F_\Omega(G)$.

Now, let uv be an arbitrary edge of $F_\Omega(G)|_{A_1}$. We wish to show this is already an edge of $F_\Omega(H)$. That it is an edge of $F_\Omega(G)$ means there exists some $\omega \in \Omega$ and a morphism $f_\omega : \omega \rightarrow G$ such that $\text{im}(f_\omega) \ni u, v$.

We claim that in fact $\text{im}(f_\omega) \subset A_1$, so that f_ω restricts to a morphism into H . If we can show this, this will show that there is an edge uv also in $F_\Omega(H)$, and we will be done.

So let w be an arbitrary vertex in $\text{im}(f_\omega)$. Now f_ω is also a morphism from something in Ω that hits both w and v , so there is an edge vw in $F_\Omega(G)$. But this of course means that w and v are in the same connected component – and this connected component is precisely A_1 . So $w \in A_1$, and we are done. $\square$

Lemma 15. *An excisive and functorial clustering scheme is representable.*

Proof. Let $\mathfrak{C}$ be some excisive and functorial clustering scheme, and let

$$\Omega = \{G \in \mathcal{G} \mid \mathfrak{C}(G) = (V_G, V_G^2)\},$$

that is, Ω is the set of all graphs clustered by $\mathfrak{C}$ into a single part. We will show that in fact $\mathfrak{C} = F_\Omega$.

So, let G be some graph. We need to show two things:

1. Whenever u and v are clustered in the same part by $\mathfrak{C}$, they are in the same connected component of $F_\Omega(G)$.
2. Whenever uv is an edge of $F_\Omega(G)$, u and v are clustered in the same part by $\mathfrak{C}$. This of course implies that whenever u and v are in the same connected component, they are also clustered in the same part by $\mathfrak{C}$, which is what we really need.

We start with the first direction, letting u and v be two vertices that are sent to the same part by $\mathfrak{C}$ – call this part A , and let $H = G|_A$. By excisiveness of $\mathfrak{C}$, we have that $\mathfrak{C}$ sends H to a single part, so $H \in \Omega$. Therefore, the inclusion morphism $f : H \hookrightarrow G$ is a morphism from something in Ω that hits both u and v , showing there is an edge uv in $F_\Omega(G)$, as desired.

In the second direction, assume that uv is an edge of $F_\Omega(G)$. Thus, there is some $\omega \in \Omega$ and $f_\omega : \omega \rightarrow G$ such that $\text{im}(f_\omega) \ni u, v$. By functoriality of $\mathfrak{C}$, f will also be a morphism from $\mathfrak{C}(\omega)$ into $\mathfrak{C}(G)$ – which by how we defined morphisms in the category of partitioned sets means that any two vertices equivalent in $\mathfrak{C}(\omega)$ must also be equivalent in $\mathfrak{C}(G)$. However, we defined Ω as precisely the set of things such that $\mathfrak{C}(\omega)$ has only a single part – so everything is equivalent in $\mathfrak{C}(\omega)$ and in particular $u \sim_{\mathfrak{C}(\omega)} v$, and so they are also equivalent in $\mathfrak{C}(G)$, as desired. $\square$

Finally, let us think about what we need to show to demonstrate that there are no other implications between these concepts. An implication between these concepts is logically the same thing as a proof that some combination of them is impossible – so what we will do is just to draw a two-by-two-by-two table of all combinations, and fill in which cells our theorems already rule out, and give examples of clustering schemes for every other cell.

Functorial		
Not functorial	Excisive	Not Excisive
Representable	Yes, Ex. 16	No, by Lm. 14
	No, by Def. 8	No, by Def. 8 (or by Lm. 14)
Not Representable	No, by Lm. 15	Yes, Ex. 18
	Yes, Ex. 17	Yes, Ex. 19

Table 1: Table of all the possible combinations of properties, with examples or proofs of impossibility.

Example 16. The connected components functor Π is representable and excisive – we can take its representation to be $\Omega = \{K_2\}$.

Example 17. For a scheme that is excisive but not functorial (and thus not representable), consider the scheme that partitions all graphs into a single part except K_2 , which it partitions into two parts.

Example 18. For a scheme which is functorial but neither excisive nor representable, cluster all connected components on more than two vertices as well as all isolated vertices as their own parts.

If there is exactly one component on two vertices, i.e. an isolated edge, cluster it as two parts, otherwise cluster all isolated edges as single parts.

Example 19. Finally, for a scheme which has none of the properties, cluster all graphs as a single part except for K_2 , which we cluster as two parts, and the disjoint union of two copies of K_2 , which we cluster as two parts – that is, each of the copies is its own part.

Maybe add a remark outlining the results for the non-injective morphisms case.

5 Representability for hierarchical clustering

Definition 20. A functorial hierarchical clustering scheme is a functor from the product category $\mathcal{G} \times \mathbb{R}$ into $\mathcal{P}$, where we make $\mathbb{R}^+ = [0, \infty)$ into a category by saying there is a morphism from a to b whenever $a \geq b$.

There is actually a natural extension of our definition of a representable clustering scheme also to this setting, but it requires that we set up some more machinery.

Definition 21. The category $\mathcal{G}^w$ of weighted simple graphs has as its objects simple graphs together with a function assigning a real-valued positive weight to each edge of the graph. A morphism from (V, E, w) to (V', E', w') is an injective set function $f : V \rightarrow V'$ such that whenever uEv , $f(u)E'f(v)$ and $w(uv) \leq w'(f(u)f(v))$.

Definition 22. The scissors functor Σ from $\mathcal{G}^w \times \mathbb{R}^+$ into $\mathcal{G}$ takes in a weighted graph $G = (V, E, w)$ and a non-negative real number τ , removes all edges whose weight is below τ , and then forgets all the remaining weights, getting just a simple graph.

Lemma 23. *The scissors functor is actually a functor.*

Proof. Similarly to how we proved that the representable endofunctors are in fact functors, most of the required properties are obvious. There are only two things we need to check:

1. If f is a morphism of weighted graphs sending $G = (V, E, w)$ to $G' = (V', E', w')$, then for every $\tau \in \mathbb{R}^+$, f is also a morphism in $\mathcal{G}$ from $\Sigma(G, \tau)$ to $\Sigma(G', \tau)$,
2. and if $\alpha \geq \beta \geq 0$, then for any weighted graph $G = (V, E, w)$, the identity function $\text{Id} : V \rightarrow V$ is a morphism from $\Sigma(G, \alpha)$ to $\Sigma(G, \beta)$.

Finish proof. □

Definition 24. A representable functor F_Ω from $\mathcal{G}$ to $\mathcal{G}^w$ is determined by a set Ω of pairs (ω, w) of simple graphs ω and positive weights w . For any graph G , there is an edge uv of $F_\Omega(G)$ if there is some subgraph of G containing u and v which is isomorphic to some graph in Ω . The weight of this edge is then given by

$$w(uv) = \sum_{(\omega, w) \in \Omega} \sum_{f \in \text{Hom}(\omega, G)} \frac{w}{|\text{Aut}(\omega)|} \mathbb{1}_{u, v \in \text{im}(f)}.$$

Lemma 25. *The representable functors from $\mathcal{G}$ to $\mathcal{G}^w$ are actually functors.*

Proof. Fill in proof. □

Definition 26. A representable hierarchical clustering scheme is a functor from $\mathcal{G} \times \mathbb{R}^+$ into $\mathcal{P}$ that can be written as $(F_\Omega \times \text{Id}) \circ \Sigma \circ \Pi$ for some representable functor F_Ω from $\mathcal{G}$ to $\mathcal{G}^w$.

Remark 27. We can of course turn such a representable hierarchical clustering scheme into a non-hierarchical clustering functor by just fixing a threshold. However, unfortunately, it turns out that this will *not* in general result in an excisive, and thus also not a representable, functor!

Fill in example

Figure out the right notion of excisiveness for hierarchical clustering schemes – I have some vague ideas for this.