Andaman & Nicobar Islands — Coral Reef Locations & Study Area
Bounding Box: 90.0°–95.0°E, 6.0°–14.0°N



NOAA CRW Degree Heating Weeks — Andaman & Nicobar Islands
Spatial distribution during key bleaching years











NOAA CRW Degree Heating Weeks — Florida Reef Tract
Spatial distribution during key bleaching years



Great Barrier Reef — Coral Reef Locations & Study Area
Bounding Box: 142.0°–154.0°E, -24.5°--10.0°N



Gulf of Kachchh — Coral Reef Locations & Study Area
Bounding Box: 68.5°–70.5°E, 22.0°–23.5°N



NOAA CRW Degree Heating Weeks — Gulf of Kachchh
Spatial distribution during key bleaching years











Gulf of Mannar — Coral Reef Locations & Study Area
Bounding Box: 78.0°–79.5°E, 8.5°–10.0°N



NOAA CRW Degree Heating Weeks — Gulf of Mannar
Spatial distribution during key bleaching years

1998 — Max DHW: 7.5 °C-weeks
Catastrophic — 89% bleached



2010 — Max DHW: 8.0 °C-weeks
Severe — 70% bleaching



2016 — Max DHW: 6.5 °C-weeks
Severe — 46% bleaching



2024 — Max DHW: 7.0 °C-weeks
4th global event — 60% bleached





NOAA CRW Degree Heating Weeks — Coral Triangle — Indonesia
Spatial distribution during key bleaching years











Lakshadweep Islands — Coral Reef Locations & Study Area
Bounding Box: 71.5°–74.5°E, 8.0°–13.0°N



NOAA CRW Degree Heating Weeks — Lakshadweep Islands
Spatial distribution during key bleaching years

1998 — Max DHW: 6.7 °C-weeks
El Niño — 82% bleached, 26% mortality



2010 — Max DHW: 6.7 °C-weeks
SST 34°C, 73% bleaching at Agatti



2016 — Max DHW: 5.8 °C-weeks
El Niño — widespread across all atolls



2024 — Max DHW: 9.2 °C-weeks
Record DHW, ~85% bleached





Malaysia — Sabah — Coral Reef Locations & Study Area
Bounding Box: 115.5°–119.5°E, 4.0°–8.0°N



NOAA CRW Degree Heating Weeks — Malaysia — Sabah
Spatial distribution during key bleaching years



Maldives — Coral Reef Locations & Study Area
Bounding Box: 72.0°–74.0°E, -1.0°–8.0°N



Malvan Marine Sanctuary — Coral Reef Locations & Study Area
Bounding Box: 73.2°–73.7°E, 15.8°–16.2°N



NOAA CRW Degree Heating Weeks — Malvan Marine Sanctuary
Spatial distribution during key bleaching years



Mesoamerican Barrier Reef — Coral Reef Locations & Study Area
Bounding Box: -89.5°–-86.0°E, 15.5°–21.5°N



NOAA CRW Degree Heating Weeks — Mesoamerican Barrier Reef
Spatial distribution during key bleaching years











Philippines — Coral Reef Locations & Study Area
Bounding Box: 116.0°–127.0°E, 4.5°–21.0°N



NOAA CRW Degree Heating Weeks — Philippines
Spatial distribution during key bleaching years

1998 — Max DHW: 8.5 °C-weeks
Severe — 65% bleached



2010 — Max DHW: 7.5 °C-weeks
Severe — 55% bleaching



2016 — Max DHW: 6.0 °C-weeks
Moderate — 40% bleaching



2024 — Max DHW: 7.5 °C-weeks
Severe — 55% bleached







Aldabra Atoll

NOAA CRW Degree Heating Weeks — Seychelles
Spatial distribution during key bleaching years



Aldabra Atoll

Aldabra Atoll

Aldabra Atoll

Aldabra Atoll

Thailand — Andaman Sea — Coral Reef Locations & Study Area
Bounding Box: 97.5°–99.5°E, 6.0°–10.0°N



NOAA CRW Degree Heating Weeks — Thailand — Andaman Sea
Spatial distribution during key bleaching years