


# Obtaining Surface of Revolution of Curves

1 : Obtain surface of revolution of curves  $\sin x$ ,  $\cos x$ ,  $\tan x$ .

Solution :


```
In[ ]:= RevolutionPlot3D[Sin[x], {x, 0, Pi}, AxesLabel -> {"x Axis", "y Axis", "z Axis"},  
  RevolutionAxis -> {0, 0, 1}, PlotStyle -> White, MeshFunctions -> {#3 &},  
  MeshStyle -> Dotted, AxesStyle -> Red, Background -> Blue]
```

Out[ ]:=




```
In[ ]:= RevolutionPlot3D[Cos[x], {x, 0, Pi}, AxesLabel -> {"x Axis", "y Axis", "z Axis"},  
  RevolutionAxis -> {0, 0, 1}, PlotStyle -> White, AxesStyle -> Red, Background -> Blue]
```

Out[ ]:=



```
In[ ]:= RevolutionPlot3D[Tan[x], {x, 0, Pi}, AxesLabel -> {"x Axis", "y Axis", "z Axis"},
  RevolutionAxis -> {0, 0, 1}, PlotStyle -> White, AxesStyle -> Red, Background -> Blue]
```

Out[ ]=




2 : Obtain surface of revolution of ellipse.

Solution :

```
In[ ]:= RevolutionPlot3D[{4 Cos[t], 2 Sin[t]}, {t, -Pi, Pi}, AxesLabel -> {x, y, z},
  RevolutionAxis -> {0, 0, 1}, PlotStyle -> Yellow, AxesStyle -> Blue, Background -> Purple]
```


Out[ ]=



3 : Obtain the surface of revolution of parabola  $y^2 = 4ax$ .

Solution :

```
In[ ]:= RevolutionPlot3D[{t^2, 2 t}, {t, -2, 2}, AxesLabel -> {x, y, z},
  RevolutionAxis -> {0, 0, 1}, PlotStyle -> Green, AxesStyle -> Blue, Background -> White]
```




4 : Obtain the surface of revolution of hyperbola  $\frac{x^2}{9} - \frac{y^2}{4} = 1$ .

Solution :

In[\*]:= RevolutionPlot3D[{3 Cosh[t], 2 Sinh[t]}, {t, -Pi, Pi}, AxesLabel → {x, y, z},  
RevolutionAxis → {0, 0, 1}, PlotStyle → Brown, Background → Orange]

Out[\*]=




5 : Donut

Solution :

```
RevolutionPlot3D[{2 + Cos[t], Sin[t]},  
{t, -2 Pi, Pi}, AxesLabel -> {x, y, z}, AxesStyle -> Green]
```

Out[\*]=




## Drawing the surface and finding its level curves from the given height

Equation :  $f(x) = x^2 - y^2$


```
In[*]:= Plot3D[x^2 - y^2, {x, -5, 5}, {y, -5, 5}, Mesh -> All, MeshFunctions -> {#3 &}]
```

Out[\*]=



```
In[ ]:= ContourPlot[x^2 - y^2, {x, -5, 5}, {y, -5, 5},  
ContourShading -> False, ContourStyle -> {{Thick, Red}}]
```


Out[ ]:=



**Equation :**  $f(x) = \sin \sqrt{x^2 + y^2}$


```
In[ ]:= Plot3D[Sin[Sqrt[x^2 + y^2]], {x, -5, 5}, {y, -5, 5},  
MeshFunctions -> (#3 &), BoxRatios -> {2, 2, 2}, MeshStyle -> {Thick, Red}]
```

Out[ ]:=



```
In[ ]:= ContourPlot[Sin[Sqrt[x^2 + y^2]], {x, -5, 5}, {y, -5, 5},
  ContourShading -> False, ContourStyle -> {{Thick, Red}}]
```

Out[ ]:=



## Wave Equation

$$\text{Equation : } \frac{\partial^2 U[X, T]}{\partial T^2} - 4 \frac{\partial^2 U[X, T]}{\partial X^2} = 0, U[X, 0] = 0, \\ U_t[X, 0] = X[1 - X], U[0, T] = 0, U[1, T] = 0$$

```

In[ ]:= weqn2 = {D[u[x, t], t, t] - 4 D[u[x, t], x, x] == 0, u[x, 0] == 0,
  u[0, t] == 0, u[1, t] == 0, Derivative[0, 1][u][x, 0] == x (1 - x) }
sol2 = u[x, t] /. NDSolve[weqn2, u[x, t], {x, 0, 1}, {t, 0, 1}]
Plot3D[sol2, {x, 0, 1}, {t, 0, 1},
  AxesLabel -> {"x axes", "y axes", "u[x,t]"}, AxesOrigin -> {0, 0}, ColorFunction -> Hue]

```

```


Out[ ]:= {u(0,2)[x, t] - 4 u(2,0)[x, t] == 0, u[x, 0] == 0,
  u[0, t] == 0, u[1, t] == 0, u(0,1)[x, 0] == (1 - x) x}

```

```

Out[ ]:= {InterpolatingFunction[ Domain: {{0., 1.}, {0., 1.}} Output: scalar][x, t]}

```



## Heat Equation


Equation:  $\frac{\partial U[X, T]}{\partial T} - \frac{\partial^2 U[X, T]}{\partial X^2} = 0$ ,  $U[X, 0] = x^2 - x$ ,  
 $U[0, t] = 0$ ,  $u[1, t] = 0$ ,  $0 < x < 1$   $1 = 1$

```


In[ ]:= heqn1 = {D[u[x, t], t] - D[u[x, t], x, x] == 0, u[x, 0] == x^2 - x, u[0, t] == 0, u[1, t] == 0};
sol1 = u[x, t] /. NDSolve[heqn1, u[x, t], {x, 0, 1}, {t, 0, 1}]
Plot3D[sol1, {x, 0, 1}, {t, 0, 1},
  ColorFunction -> Hue, AxesLabel -> {"x axis", "y axis", "u[x,t]"}]

```

Out[ ]:=

{InterpolatingFunction[ Domain: {{0., 1.}, {0., 1.}} Output: scalar][x, t]}

Out[ ]:=



## Growth Model (Exponential Case )

Exponential Growth : -

If a function  $x(t)$  grows continuously at a rate  $k > 0$ , then  $x(t)$  has the form

$$x(t) = x_0 e^{kt}$$

where  $x_0$  is the initial amount and  $t$  is the time. In this case, the quantity  $x(t)$  is said to exhibit exponential growth, and  $k$  is the growth rate. It's differential model is given by 2%

$$\frac{dx}{dt} = kx, \quad k > 0$$

**Suppose that the population of a certain country grows at an annual rate of  $2\%$ . If the current population is 3 million, what will the population be in 10 years? Also plot the graph of the solution.**

**Solution :** Here  $x_0 = 3$ ,  $k = 2\% = 0.02$ ,  $t = 10$  years,  $x[t] = ?$



```
In[ ]:= Sol = DSolve[x'[t] == k x[t], x[t], t]
Sol1 = Evaluate[x[t] /. Sol[[1]] /. {k -> 0.02, C[1] -> 3}]
Plot[Sol1, {t, -50, 50}, PlotStyle -> {Pink, Thickness[0.03]},
  AxesLabel -> {t, x[t]}, AxesStyle -> Arrowheads[{-0.03, 0.03}]
x[10] = Evaluate[Sol1 /. {t -> 10}]
```


Out[ ]:=

$$\left\{ \left\{ x[t] \rightarrow e^{kt} c_1 \right\} \right\}$$

Out[ ]:=

$$3 e^{0.02 t}$$

Out[ ]:=



Out[ ]:=

3.66421

Conclusion : Hence population after 10 years will be 3.66421.

## Decay Model (Exponential Case)

**Exponential Decay :**

If a function  $x(t)$  decreases continuously at a rate  $k > 0$ , then  $x(t)$  has the form

$$x(t) = x_0 e^{-kt}$$

where  $x_0$  is the initial amount  $x_0$ . In this case,

the quantity  $x(t)$  is said to exhibit exponential decay,

and  $k$  is the decay rate. It's differential model is given by  $\frac{dx}{dt} = -kx$ ,  $k > 0$ .

Suppose that a certain radioactive element has an annual decay rate of  $10\%$ . Starting with a 200 gram sample of the element, how many grams will be left in 3 years?

Solution : Here  $k = 10\% = 0.1$ ,  $x_0 = 200$ ,  $t = 3$ ,  $x(t) = ?$

```

In[ ]:= Sol = DSolve[x'[t] == -k x[t], x[t], t]
Sol1 = Evaluate[x[t] /. Sol[[1]] /. {k -> 0.1, C[1] -> 200}]
Plot[Sol1, {t, -50, 50}, PlotStyle -> {Green, Thickness[0.02]}, AxesLabel -> {t, x[t]},
  AxesStyle -> Arrowheads[{- .03, .03}], PlotLabel -> "Decay Model"]
x[3] = Evaluate[Sol1 /. {t -> 3}]

```


Out[ ]:=

$$\left\{ \left\{ x[t] \rightarrow e^{-k t} c_1 \right\} \right\}$$

Out[ ]:=

$$200 e^{-0.1 t}$$

Out[ ]:=



Out[ ]:=

148.164

**Conclusion :** Hence 148.164 gms of radioactive element will be left after 3 years.