Catalogues: How Astronomers Keep Track of Billions of Objects

What you will be able to do

1. Explain why astronomers need catalogues, and trace their history from Hipparchus to Gaia.
2. Recognise the major catalogue prefixes: **HD, HIP, Gaia, M, NGC, IC, C** (Caldwell).
3. Tell the story of the **Messier catalogue**.
4. Name the types of **deep-sky objects** with an example of each.
5. Plan an observing list of objects visible from India, with the right instrument and season.

1. Why catalogues?

Names like "Sirius" cover only a few hundred stars. About 9,000 stars are visible to the naked eye, *billions* are within reach of modern telescopes, and there are countless galaxies and nebulae. A catalogue gives each object a **unique label** plus its **position (RA/Dec), brightness, type and distance**, so anyone, anywhere, can find it and compare results.

2. Star catalogues through history

Catalogue	Date	Stars	Notes
Hipparchus	~130 BCE	~850	first known star catalogue; brightness classes
Ptolemy, *Almagest*	~150 CE	1,022	positions and magnitudes in 48 constellations
Al-Sufi, *Book of Fixed Stars*	964 CE	~1,000	revised magnitudes; Arabic star names
Ulugh Beg (Samarkand)	1437	~1,000	best pre-telescope catalogue after Tycho
Tycho Brahe	1598	~1,000	naked-eye positions accurate to ~ 1′
Bayer, *Uranometria*	1603	~1,500	Greek-letter names
Flamsteed	1725	~2,900	Flamsteed numbers
Madras Catalogue (T. G. Taylor, Madras Observatory)	1844	~11,000	an important early **Indian** contribution
Henry Draper (HD)	1918–24	225,300	**spectral types (O B A F G K M)**, classified mainly by **Annie Jump Cannon**
Bright Star Catalogue (HR)	1908 on	~9,100	every star to magnitude 6.5
Hipparcos (HIP) satellite	1997	118,218	precise parallaxes ⇒ true distances
Gaia satellite (DR3)	2022	~1.8 billion	positions, distances, motions, colours — a 3-D map of the Milky Way

One star, many names: Betelgeuse = α Orionis = 58 Orionis = HR 2061 = HD 39801 = HIP 27989.

© Vaibhav Yenare

P.T.O.

3. Deep-sky objects: what's out there?

Anything beyond the solar system that is not a single star is a **deep-sky object** (**DSO**).




Open cluster — Pleiades (M45)
NASA, ESA, AURA/Caltech, Palomar; PD




Globular cluster — M13
NASA, ESA, Hubble Heritage Team; PD




Emission nebula — Orion Nebula (M42)
NASA, ESA, M. Robberto; PD



Planetary nebula — Ring Nebula (M57)
Hubble Heritage Team (AURA/STScI/NASA); PD



Supernova remnant — Crab Nebula (M1)
NASA, ESA, J. Hester & A. Loll; PD



Galaxy — Andromeda (M31)
Benjamin Inouye, CC BY 4.0

The main types of deep-sky object, as photographed by large telescopes (via Wikimedia Commons). Through a small telescope they look much fainter and mostly grey — the eye cannot see colour in dim light.

Type	What it is	Example	In a small telescope
Open cluster	young group of tens to thousands of stars born together, in the Milky Way's disc	Pleiades, Beehive (M44)	sparkling loose group
Globular cluster	ancient ball of 100,000+ stars in the galaxy's halo	M13, Omega Centauri	fuzzy ball
Emission nebula	gas cloud glowing from hot young stars inside	Orion Nebula, Lagoon (M8)	grey-green glow
Reflection nebula	dust reflecting starlight	around the Pleiades	very faint
Planetary nebula	gas shell puffed off by a dying Sun-like star — nothing to do with planets	Ring (M57), Dumbbell (M27)	small ring or disc
Supernova remnant	debris of an exploded star	Crab Nebula (supernova of 1054 CE)	faint smudge
Galaxy	a separate "island universe" of billions of stars	Andromeda, 2.5 million ly away	oval glow
Double star	two stars close together in the sky	Mizar–Alcor, Albireo	colourful pairs

4. The Messier catalogue: a comet-hunter's "do not confuse" list

Charles Messier (1730–1817), a French astronomer obsessed with discovering **comets** (he found 13), was repeatedly fooled by fuzzy patches. So he listed things that **look like comets but are not**: 45 objects in 1774, 103 by 1781; later astronomers added M104–M110 from his and Pierre Méchain's notes, making **110**. Messier's small telescope picked up only the *brightest* objects — so his list of nuisances became the most popular list of beautiful objects in the sky. **M1** is the Crab Nebula, **M31** the Andromeda Galaxy, **M42** the Orion Nebula, **M45** the Pleiades.

India is perfect for the "Messier Marathon"

Around the **New Moon of late March or early April**, all 110 Messier objects can be seen in **one night**, from dusk to dawn. This works best between about 20° and 25° N — much of India, including Mumbai, Nagpur, Bhopal and Kolkata.

5. NGC, IC and Caldwell

Catalogue	Compiled by	Year	Objects	Notes
NGC (New General Catalogue)	J. L. E. Dreyer	1888	7,840	built on the work of William, Caroline and John Herschel
IC (Index Catalogue)	J. L. E. Dreyer	1895, 1908	5,386	many found by photography
Caldwell (C)	Patrick Moore	1995	109	bright objects Messier missed, including the southern sky

Most Messier objects also have NGC numbers: M42 = NGC 1976, M31 = NGC 224, M13 = NGC 6205. Non-Messier gems for Indian observers: **Omega Centauri** (NGC 5139 = C80), the **Double Cluster** in Perseus (NGC 869/884 = C14), the **Carina Nebula** (NGC 3372 = C92).

6. Reading a catalogue entry — M42, the Orion Nebula

Field	Value	Meaning
Designation	M42 = NGC 1976	Messier no. 42; NGC no. 1976
Type	emission nebula (H II region)	glowing hydrogen gas
Constellation	Orion	
RA / Dec (J2000)	$05^{h}35.3^{m}$, $-05°23'$	position
Magnitude	4.0	total brightness, spread over a large area
Size	$85' \times 60'$	larger than the Full Moon ($30'$)
Distance	~1,350 ly	

Magnitude caution

For extended objects the magnitude is the *total* light spread over the object. A galaxy of magnitude 8 can be *harder* to see than a star of magnitude 11, because its light is spread out — it has a low **surface brightness**.

7. Observing list: 20 deep-sky treasures visible from India

#	Object	Type	Con.	RA	Dec	Mag	Best	Instrument
1	**M45 Pleiades (Krittika)**	open cl.	Tau	3h47m	+24°07′	1.6	Nov–Feb	eye / binoculars
2	**M42 Orion Nebula**	emission	Ori	5h35m	−05°23′	4.0	Dec–Feb	eye / any
3	**M31 Andromeda Galaxy**	galaxy	And	0h43m	+41°16′	3.4	Oct–Dec	eye (dark) / binoculars
4	**M44 Beehive (Pushya)**	open cl.	Cnc	8h40m	+19°59′	3.7	Feb–Apr	binoculars
5	Hyades	open cl.	Tau	4h27m	+15°52′	0.5	Nov–Feb	eye / binoculars
6	**M41**	open cl.	CMa	6h46m	−20°46′	4.5	Jan–Feb	binoculars (4° below Sirius)
7	M35	open cl.	Gem	6h09m	+24°20′	5.3	Dec–Feb	binoculars
8	**M7 Ptolemy's Cluster**	open cl.	Sco	17h54m	−34°48′	3.3	Jun–Aug	eye / binoculars
9	M6 Butterfly Cluster	open cl.	Sco	17h40m	−32°15′	4.2	Jun–Aug	binoculars
10	**M8 Lagoon Nebula**	emission	Sgr	18h04m	−24°23′	6.0	Jun–Sep	binoculars
11	M22	globular	Sgr	18h36m	−23°54′	5.1	Jun–Sep	binoculars / telescope
12	**M13 Hercules Cluster**	globular	Her	16h42m	+36°28′	5.8	May–Aug	binoculars / telescope
13	M11 Wild Duck Cluster	open cl.	Sct	18h51m	−06°16′	5.8	Jul–Sep	telescope
14	**M57 Ring Nebula**	planetary	Lyr	18h54m	+33°02′	8.8	Jul–Sep	telescope, 100×+
15	M27 Dumbbell Nebula	planetary	Vul	20h00m	+22°43′	7.5	Jul–Oct	telescope
16	**Omega Centauri** (NGC 5139)	globular	Cen	13h27m	−47°29′	3.9	Apr–Jun	eye / binoculars (low S)
17	**Double Cluster** (NGC 869/884)	open cl.	Per	2h20m	+57°08′	3.7	Oct–Jan	binoculars
18	Carina Nebula (NGC 3372)	emission	Car	10h45m	−59°52′	∼1	Feb–Apr	binoculars (south India)
19	M81 & M82	galaxies	UMa	9h56m	+69°04′	6.9/8.4	Feb–May	telescope (dark site)
20	M104 Sombrero Galaxy	galaxy	Vir	12h40m	−11°37′	8.0	Apr–Jun	telescope (dark site)

Check visibility from your latitude with the transit-altitude formula $h = 90^\circ - \varphi + \delta$ (Lecture 2.2): objects that never rise more than $\sim 15^\circ$ are usually lost in haze.

8. Using catalogue numbers in practice

- **Stellarium / SkySafari:** type "M42" or "NGC 5139" in the search box to see where it is tonight.
- **GoTo mounts:** choose "Messier" or "NGC" in the hand controller and enter the number.
- **Star-hop to it** (better for learning!): M41 is 4° straight south of Sirius; M13 is a third of the way from η to ζ Herculis on the Keystone's western side.
- **Log it:** catalogue number, date, instrument, magnification, sky conditions and a sketch. Seeing all 110 Messier objects is a classic amateur goal — many clubs award a certificate!

Key points to remember

- Catalogues give each object a unique ID + position + data; one object can have many IDs.
- Star catalogues: Hipparchus → Ptolemy → al-Sufi → Tycho → HD (spectral types, Annie Jump Cannon) → Hipparcos → Gaia (∼1.8 billion).
- Messier (110): a comet-hunter's list of "not comets", now the favourite list of bright deep-sky objects; NGC (Dreyer 1888, 7,840), IC, Caldwell (109).
- DSO types: open and globular clusters; emission, reflection and planetary nebulae; supernova remnants; galaxies.
- India ($\sim 20^\circ$–25° N) is ideal for a Messier Marathon around the March–April New Moon.

Check yourself

1. Why did Messier make his catalogue?
2. What kind of object is M57, and what is it made of?
3. Which catalogue first classified stars by spectral type, and who did most of the classifying?
4. Give the Messier and NGC numbers of the Andromeda Galaxy.

Answers: 1. To avoid mistaking fuzzy objects for comets. 2. A planetary nebula: a gas shell from a dying Sun-like star. 3. The Henry Draper Catalogue; Annie Jump Cannon. 4. M31 = NGC 224.

Further reading (free / public)

- NASA, *Hubble's Messier Catalog* — https://science.nasa.gov/mission/hubble/science/explore-the-night-sky/hubble-messier-catalog/
- SEDS Messier Database — http://www.messier.seds.org
- ESA Gaia — https://www.cosmos.esa.int/web/gaia
- S. J. O'Meara, *Deep-Sky Companions: The Messier Objects* (Cambridge University Press).