

ASSIGNMENT 3

Vaibhav Chhabra
AI20BTECH11022

Download all python codes from

<https://github.com/vaibhavchhabra25/EE3900-course/blob/main/Assignment-3/codes>

and latex-tikz codes from

<https://github.com/vaibhavchhabra25/EE3900-course/blob/main/Assignment-3/main.tex>

1 PROBLEM

(Ramsey-4.2-Tangent and Normal-Q.21)

Verify that the perpendicular bisector of the chord joining two points $\mathbf{x}_1, \mathbf{x}_2$ on the circle

$$\mathbf{x}^\top \mathbf{x} + 2(g \ f) \mathbf{x} + c = 0 \quad (1.0.1)$$

passes through the centre.

2 SOLUTION

Since \mathbf{x}_1 and \mathbf{x}_2 lie on the circle,

$$\mathbf{x}_1^\top \mathbf{x}_1 + 2(g \ f) \mathbf{x}_1 + c = 0 \quad (2.0.1)$$

$$\mathbf{x}_2^\top \mathbf{x}_2 + 2(g \ f) \mathbf{x}_2 + c = 0 \quad (2.0.2)$$

Subtracting the two, we get,

$$\mathbf{x}_2^\top \mathbf{x}_2 - \mathbf{x}_1^\top \mathbf{x}_1 = -2(g \ f)(\mathbf{x}_2 - \mathbf{x}_1) \quad (2.0.3)$$

The midpoint of the chord $\mathbf{x}_1 \mathbf{x}_2$ is

$$\mathbf{M} = \frac{\mathbf{x}_1 + \mathbf{x}_2}{2} \quad (2.0.4)$$

Since perpendicular bisector of $\mathbf{x}_1 \mathbf{x}_2$ is perpendicular to $\mathbf{x}_1 \mathbf{x}_2$ and passes through \mathbf{M} , any general point \mathbf{P} on the perpendicular bisector can be given using the relation

$$(\mathbf{x}_2 - \mathbf{x}_1)^\top (\mathbf{P} - \mathbf{M}) = 0 \quad (2.0.5)$$

$$\Rightarrow (\mathbf{x}_2 - \mathbf{x}_1)^\top \mathbf{P} - (\mathbf{x}_2^\top - \mathbf{x}_1^\top) \left(\frac{\mathbf{x}_1 + \mathbf{x}_2}{2} \right) = 0 \quad (2.0.6)$$

$$\Rightarrow (\mathbf{x}_2 - \mathbf{x}_1)^\top \mathbf{P} = \frac{\mathbf{x}_2^\top \mathbf{x}_2 - \mathbf{x}_1^\top \mathbf{x}_1 + \mathbf{x}_2^\top \mathbf{x}_1 - \mathbf{x}_1^\top \mathbf{x}_2}{2} \quad (2.0.7)$$

Since $\mathbf{x}_1^\top \mathbf{x}_2 = \mathbf{x}_2^\top \mathbf{x}_1$, the equation reduces to

$$2(\mathbf{x}_2 - \mathbf{x}_1)^\top \mathbf{P} = \mathbf{x}_2^\top \mathbf{x}_2 - \mathbf{x}_1^\top \mathbf{x}_1 \quad (2.0.8)$$

Using (2.0.3), we get

$$2(\mathbf{x}_2 - \mathbf{x}_1)^\top \mathbf{P} = -2(g \ f)(\mathbf{x}_2 - \mathbf{x}_1) \quad (2.0.9)$$

$$\Rightarrow (\mathbf{x}_2 - \mathbf{x}_1)^\top \mathbf{P} = (\mathbf{x}_2 - \mathbf{x}_1)^\top \begin{pmatrix} -g \\ -f \end{pmatrix} \quad (2.0.10)$$

Since the centre of the circle is $\mathbf{C} = \begin{pmatrix} -g \\ -f \end{pmatrix}$, we can clearly see that $\mathbf{P} = \mathbf{C}$ satisfies the equation of perpendicular bisector.

Hence, the perpendicular bisector of any chord of a circle passes through the centre of the circle.




Fig. 0: Example figure