

Out of Syllabus Problems (2005–2026 Archive)

Category: Legacy & Discontinued Topics • Legacy Problem Collection

TOTAL QUESTIONS

87

TOTAL MARKS

277 M

TEST DURATION

261 Min

PAPER PATTERN

MCQ • MSQ • NAT
(2005–20

Section / Type	Questions	Marking Scheme	Negative Marking	Format / Details
Multiple Choice (MCQ)	63	+1 / +2 / +6 Marks	-1/3 / -2/3 / -2 Marks	4 Options (1 Correct)
Multiple Select (MSQ)	12	+2 Marks	Nil (0)	4 Options (1+ Correct)
Numerical Answer (NAT)	12	+1 / +2 Marks	Nil (0)	Real Decimal Value

IMPORTANT INSTRUCTIONS FOR CANDIDATES

- Historical & Discontinued Topics:** This collection contains authentic official IIT JAM Mathematics questions from historical topics (such as Vector Calculus including Green's, Stokes', and Gauss Divergence Theorems, Mechanics, Statics, and Dynamics) that were tested in earlier years (2005–2021) but have been phased out of the modern JAM MA syllabus.
- Problem-Solving & Entrance Value:** These problems remain exceptionally valuable for competitive mathematical examinations (TIFR, GATE MA, CSIR-NET, NBHM, and CMI/ISI admissions) and deepening advanced analytical intuition.
- Section A (MCQ):** Four options (A), (B), (C), (D) where **only one** is correct. Incorrect attempts incur standard negative marking penalty.
- Section B (MSQ):** Four options (A), (B), (C), (D) where **one or more** options may be correct. Full marks awarded only if all correct choices and zero incorrect choices are chosen. No partial or negative marking.
- Section C (NAT):** Numerical answers are real numbers within the official accepted interval. No negative marking.
- Master Answer Key:** Complete official answers and acceptance bounds are provided on the final page(s) of this document.
- Online Interactive Simulator:** Practice full papers and topic test series at vaibhavgeometer.github.io.

IIT JAM Mathematics Mock Test Series • Curated by Vaibhav Geometer

Question 1 (JAM 2021 • Q58 • ID: JAM_2021_Q58)

NAT

+2 M

Neg: Nil (0)

Q. 58 Let S be the surface defined by

$$\{(x, y, z) \in \mathbb{R}^3 : z = 1 - x^2 - y^2, z \geq 0\}.$$

Let $\vec{F} = -y\hat{i} + (x-1)\hat{j} + z^2\hat{k}$ and $\hat{n}$ be the continuous unit normal field to the surface S with positive z -component. Then the value of

$$\frac{1}{\pi} \iint_S (\nabla \times \vec{F}) \cdot \hat{n} dS$$

is ____.

/ 0 1 2 \

Question 2 (JAM 2021 • Q49 • ID: JAM_2021_Q49)

NAT

+1 M

Neg: Nil (0)

Q. 49 Let $\vec{F} = (y+1)e^y \cos(x)\hat{i} + (y+2)e^y \sin(x)\hat{j}$ be a vector field in $\mathbb{R}^2$ and C be a continuously differentiable path with the starting point $(0, 1)$ and the end point $(\frac{\pi}{2}, 0)$. Then

$$\int_C \vec{F} \cdot d\vec{r}$$

equals ____.

Question 3 (JAM 2021 • Q44 • ID: JAM_2021_Q44)

NAT

+1 M

Neg: Nil (0)

Q. 44 Consider the subset $S = \{(x, y) : x^2 + y^2 > 0\}$ of $\mathbb{R}^2$. Let

$$P(x, y) = \frac{y}{x^2 + y^2} \text{ and } Q(x, y) = -\frac{x}{x^2 + y^2}$$

for $(x, y) \in S$. If C denotes the unit circle traversed in the counter-clockwise direction, then the value of

$$\frac{1}{\pi} \int_C (Pdx + Qdy)$$


is ____.

Question 4 (JAM 2021 • Q37 • ID: JAM_2021_Q37)

MSQ

+2 M

Neg: Nil (0)

Q. 37 Which of the following subsets of $\mathbb{R}$ is/are connected?(A) The set $\{x \in \mathbb{R} : x \text{ is irrational}\}$.(B) The set $\{x \in \mathbb{R} : x^3 - 1 \geq 0\}$.(C) The set $\{x \in \mathbb{R} : x^3 + x + 1 \geq 0\}$.(D) The set $\{x \in \mathbb{R} : x^3 - 2x + 1 \geq 0\}$.

MA

12 / 17

Question 5 (JAM 2021 • Q13 • ID: JAM_2021_Q13)

MCQ

+2 M

Neg: -0.67

Q. 13 Consider the surface $S = \{(x, y, xy) \in \mathbb{R}^3 : x^2 + y^2 \leq 1\}$. Let $\vec{F} = y\hat{i} + x\hat{j} + \hat{k}$. If $\hat{n}$ is the continuous unit normal field to the surface S with positive z -component, then

$$\iint_S \vec{F} \cdot \hat{n} dS$$

equals

(A) $\frac{\pi}{4}$.(B) $\frac{\pi}{2}$.(C) π .(D) 2π .

Question 6 (JAM 2021 • Q8 • ID: JAM_2021_Q8)

MCQ

+1 M

Neg: -0.33

Q. 8 Which one of the following subsets of $\mathbb{R}$ has a non-empty interior?(A) The set of all irrational numbers in $\mathbb{R}$.(B) The set $\{a \in \mathbb{R} : \sin(a) = 1\}$.(C) The set $\{b \in \mathbb{R} : x^2 + bx + 1 = 0 \text{ has distinct roots}\}$.(D) The set of all rational numbers in $\mathbb{R}$.

Question 7 (JAM 2020 • Q55 • ID: JAM_2020_Q55)

NAT

+2 M

Neg: Nil (0)

Q. 55 Let C be the boundary of the square with vertices $(0, 0)$, $(1, 0)$, $(1, 1)$ and $(0, 1)$ oriented in the counter clockwise sense. Then, the value of the line integral

$$\oint_C x^2 y^2 dx + (x^2 - y^2) dy$$

is _____. (rounded off to two decimal places)

Question 8 (JAM 2020 • Q42 • ID: JAM_2020_Q42)

NAT

+1 M

Neg: Nil (0)

Q. 42 Let $\vec{F} = x\hat{i} + y\hat{j} + z\hat{k}$ and S be the sphere given by $(x - 2)^2 + (y - 2)^2 + (z - 2)^2 = 4$. If $\hat{n}$ is the unit outward normal to S , then

$$\frac{1}{\pi} \iint_S \vec{F} \cdot \hat{n} \, dS$$

is _____.

Question 9 (JAM 2020 • Q27 • ID: JAM_2020_Q27)

MCQ

+2 M

Neg: -0.67

Q. 27 Let $D = \{(x, y) \in \mathbb{R}^2 : |x| + |y| \leq 1\}$ and $f : D \rightarrow \mathbb{R}$ be a non-constant continuous function. Which of the following is TRUE?

- (A) The range of f is unbounded
- (B) The range of f is a union of open intervals
- (C) The range of f is a closed interval
- (D) The range of f is a union of at least two disjoint closed intervals

Question 10 (JAM 2020 • Q26 • ID: JAM_2020_Q26)

MCQ

+2 M

Neg: -0.67

JAM 2020

MATHEMATICS - MA

Q. 26 Let $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ and $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$, $x, y, z \in \mathbb{R}$. Which of the following is FALSE?

(A) $\nabla(\vec{a} \cdot \vec{r}) = \vec{a}$

(B) $\nabla \cdot (\vec{a} \times \vec{r}) = 0$

(C) $\nabla \times (\vec{a} \times \vec{r}) = \vec{a}$

(D) $\nabla \cdot ((\vec{a} \cdot \vec{r})\vec{r}) = 4(\vec{a} \cdot \vec{r})$

Question 11 (JAM 2019 • Q44 • ID: JAM_2019_Q44)

NAT

+1 M

Neg: Nil (0)

Q.44 Let $\vec{F}(x, y) = -y\hat{i} + x\hat{j}$ and let C be the ellipse

$$\frac{x^2}{16} + \frac{y^2}{9} = 1$$

oriented counter clockwise. Then the value of $\oint_C \vec{F} \cdot d\vec{r}$ (round off to 2 decimal places) is _____

Question 12 (JAM 2019 • Q35 • ID: JAM_2019_Q35)

MSQ

+2 M

Neg: Nil (0)

Q.35 Consider the intervals $S = (0, 2]$ and $T = [1, 3)$. Let S° and T° be the sets of interior points of S and T , respectively. Then the set of interior points of $S \setminus T$ is equal to

(A) $S \setminus T^\circ$

(B) $S \setminus T$

(C) $S^\circ \setminus T^\circ$

(D) $S^\circ \setminus T$

Question 13 (JAM 2019 • Q34 • ID: JAM_2019_Q34)

MSQ

+2 M

Neg: Nil (0)

Q.34 Let $\vec{F}$ and $\vec{G}$ be differentiable vector fields and let g be a differentiable scalar function. Then

(A) $\nabla \cdot (\vec{F} \times \vec{G}) = \vec{G} \cdot \nabla \times \vec{F} - \vec{F} \cdot \nabla \times \vec{G}$

(B) $\nabla \cdot (\vec{F} \times \vec{G}) = \vec{G} \cdot \nabla \times \vec{F} + \vec{F} \cdot \nabla \times \vec{G}$

(C) $\nabla \cdot (g\vec{F}) = g\nabla \cdot \vec{F} - \nabla g \cdot \vec{F}$

(D) $\nabla \cdot (g\vec{F}) = g\nabla \cdot \vec{F} + \nabla g \cdot \vec{F}$

Question 14 (JAM 2019 • Q25 • ID: JAM_2019_Q25)

MCQ

+2 M

Neg: -0.67

Q.25 Let $\vec{F}(x, y, z) = 2y\hat{i} + x^2\hat{j} + xy\hat{k}$ and let $\mathcal{C}$ be the curve of intersection of the plane $x + y + z = 1$ and the cylinder $x^2 + y^2 = 1$. Then the value of

$$\left| \oint_{\mathcal{C}} \vec{F} \cdot d\vec{r} \right|$$

is

(A) π (B) $\frac{3\pi}{2}$ (C) 2π (D) 3π

MA

6/13

Question 15 (JAM 2019 • Q24 • ID: JAM_2019_Q24)

MCQ

+2 M

Neg: -0.67

Q.24 Let $\mathcal{C}$ be the circle $(x - 1)^2 + y^2 = 1$, oriented counter clockwise. Then the value of the line integral

$$\oint_{\mathcal{C}} -\frac{4}{3}xy^3 dx + x^4 dy$$

is

(A) 6π (B) 8π (C) 12π (D) 14π

Question 16 (JAM 2019 • Q14 • ID: JAM_2019_Q14)

MCQ

+2 M

Neg: -0.67

Q.14 The set $\left\{ \frac{1}{m} + \frac{1}{n} : m, n \in \mathbb{N} \right\} \cup \{0\}$, as a subset of $\mathbb{R}$, is

(A) compact and open

(B) compact but not open

(C) not compact but open

(D) neither compact nor open

Question 17 (JAM 2019 • Q13 • ID: JAM_2019_Q13)

MCQ

+2 M

Neg: -0.67

Q.13 The set $\left\{\frac{x}{1+x} : -1 < x < 1\right\}$, as a subset of $\mathbb{R}$, is

- (A) connected and compact
- (B) connected but not compact
- (C) not connected but compact
- (D) neither connected nor compact

Question 18 (JAM 2019 • Q5 • ID: JAM_2019_Q5)

MCQ

+1 M

Neg: -0.33

Q.5 Let S be the set of all limit points of the set $\left\{\frac{n}{\sqrt{2}} + \frac{\sqrt{2}}{n} : n \in \mathbb{N}\right\}$. Let $\mathbb{Q}_+$ be the set of all positive rational numbers. Then

- (A) $\mathbb{Q}_+ \subseteq S$
- (B) $S \subseteq \mathbb{Q}_+$
- (C) $S \cap (\mathbb{R} \setminus \mathbb{Q}_+) \neq \emptyset$
- (D) $S \cap \mathbb{Q}_+ \neq \emptyset$

MA

3/13

Question 19 (JAM 2018 • Q40 • ID: JAM_2018_Q40)

MSQ

+2 M

Neg: Nil (0)

Q.40 Let S be a subset of $\mathbb{R}$ such that 2018 is an interior point of S . Which of the following is (are) TRUE?

- (A) S contains an interval
- (B) There is a sequence in S which does not converge to 2018
- (C) There is an element $y \in S$, $y \neq 2018$ such that y is also an interior point of S
- (D) There is a point $z \in S$, such that $|z - 2018| = 0.002018$

SECTION – C

NUMERICAL ANSWER TYPE (NAT)

Q 41 – Q 50 carry one mark each

Question 20 (JAM 2018 • Q39 • ID: JAM_2018_Q39)

MSQ

+2 M

Neg: Nil (0)

Q.39 Which of the following subsets of $\mathbb{R}$ is (are) connected?

- (A) $\{x \in \mathbb{R} \mid x^2 + x > 4\}$ (B) $\{x \in \mathbb{R} \mid x^2 + x < 4\}$
 (C) $\{x \in \mathbb{R} \mid |x| < |x - 4|\}$ (D) $\{x \in \mathbb{R} \mid |x| > |x - 4|\}$

MA

9/12

Question 21 (JAM 2018 • Q38 • ID: JAM_2018_Q38)

MSQ

+2 M

Neg: Nil (0)

Q.38 If $\vec{F}(x, y, z) = (2x + 3yz)\hat{i} + (3xz + 2y)\hat{j} + (3xy + 2z)\hat{k}$ for $(x, y, z) \in \mathbb{R}^3$, then which among the following is (are) TRUE?

- (A) $\nabla \times \vec{F} = \vec{0}$
 (B) $\oint_C \vec{F} \cdot d\vec{r} = 0$ along any simple closed curve C
 (C) There exists a scalar function $\phi: \mathbb{R}^3 \rightarrow \mathbb{R}$ such that $\nabla \cdot \vec{F} = \phi_{xx} + \phi_{yy} + \phi_{zz}$
 (D) $\nabla \cdot \vec{F} = 0$

Question 22 (JAM 2018 • Q34 • ID: JAM_2018_Q34)

MSQ

+2 M

Neg: Nil (0)

Q.34 Let P and Q be two non-empty disjoint subsets of $\mathbb{R}$. Which of the following is (are) FALSE?

- (A) If P and Q are compact, then $P \cup Q$ is also compact
 (B) If P and Q are not connected, then $P \cup Q$ is also not connected
 (C) If $P \cup Q$ and P are closed, then Q is closed
 (D) If $P \cup Q$ and P are open, then Q is open

MA

8/12

Question 23 (JAM 2018 • Q20 • ID: JAM_2018_Q20)

MCQ

+2 M

Neg: -0.67

Q.20 If $\vec{F}(x, y) = (3x - 8y)\hat{i} + (4y - 6xy)\hat{j}$ for $(x, y) \in \mathbb{R}^2$, then $\oint_C \vec{F} \cdot d\vec{r}$, where C is the boundary of the triangular region bounded by the lines $x = 0, y = 0$ and $x + y = 1$ oriented in the anti-clockwise direction, is

(A) $\frac{5}{2}$

(B) 3

(C) 4

(D) 5

Question 24 (JAM 2018 • Q7 • ID: JAM_2018_Q7)

MCQ

+1 M

Neg: -0.33

Q.7 Let $f: \mathbb{R}^3 \rightarrow \mathbb{R}$ be a scalar field, $\vec{v}: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a vector field and let $\vec{a} \in \mathbb{R}^3$ be a constant vector. If $\vec{r}$ represents the position vector $x\hat{i} + y\hat{j} + z\hat{k}$, then which one of the following is FALSE?

(A) $\text{curl}(f \vec{v}) = \text{grad}(f) \times \vec{v} + f \text{curl}(\vec{v})$

(B) $\text{div}(\text{grad}(f)) = \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) f$

(C) $\text{curl}(\vec{a} \times \vec{r}) = 2 |\vec{a}| \vec{r}$

(D) $\text{div}\left(\frac{\vec{r}}{|\vec{r}|^3}\right) = 0$, for $\vec{r} \neq \vec{0}$

Question 25 (JAM 2017 • Q52 • ID: JAM_2017_Q52)

NAT

+2 M

Neg: Nil (0)

Q.52 Let T be the smallest positive real number such that the tangent to the helix

$$\cos t \hat{i} + \sin t \hat{j} + \frac{t}{\sqrt{2}} \hat{k}$$

at $t = T$ is orthogonal to the tangent at $t = 0$. Then the line integral of $\vec{F} = x\hat{j} - y\hat{i}$ along the section of the helix from $t = 0$ to $t = T$ is _____

Question 26 (JAM 2017 • Q37 • ID: JAM_2017_Q37)

MSQ

+2 M

Neg: Nil (0)

Q.37 Let S be the set of all rational numbers in $(0,1)$. Then which of the following statements is / are TRUE?

- (A) S is a closed subset of $\mathbb{R}$
- (B) S is not a closed subset of $\mathbb{R}$
- (C) S is an open subset of $\mathbb{R}$
- (D) Every $x \in (0,1) \setminus S$ is a limit point of S

Question 27 (JAM 2017 • Q24 • ID: JAM_2017_Q24)

MCQ

+2 M

Neg: -0.67

Q.24 Let S be an infinite subset of $\mathbb{R}$ such that $S \setminus \{\alpha\}$ is compact for some $\alpha \in S$. Then which one of the following is TRUE?

- (A) S is a connected set
- (B) S contains no limit points
- (C) S is a union of open intervals
- (D) Every sequence in S has a subsequence converging to an element in S

Question 28 (JAM 2017 • Q18 • ID: JAM_2017_Q18)

MCQ

+2 M

Neg: -0.67

Q.18 The flux of $\vec{F} = y\hat{i} - x\hat{j} + z^2\hat{k}$ along the outward normal, across the surface of the solid

$$\left\{ (x, y, z) \in \mathbb{R}^3 \mid 0 \leq x \leq 1, 0 \leq y \leq 1, 0 \leq z \leq \sqrt{2 - x^2 - y^2} \right\}$$

is equal to

- (A) $\frac{2}{3}$
- (B) $\frac{5}{3}$
- (C) $\frac{8}{3}$
- (D) $\frac{4}{3}$

Question 29 (JAM 2017 • Q17 • ID: JAM_2017_Q17)

MCQ

+2 M

Neg: -0.67

Q.17 For $a > 0, b > 0$, let $\vec{F} = \frac{x\hat{j} - y\hat{i}}{b^2x^2 + a^2y^2}$ be a planar vector field. Let

$$C = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = a^2 + b^2\}$$

be the circle oriented anti-clockwise. Then

$$\oint_C \vec{F} \cdot d\vec{r} =$$

(A) $\frac{2\pi}{ab}$

(B) 2π

(C) $2\pi ab$

(D) 0

Question 30 (JAM 2017 • Q15 • ID: JAM_2017_Q15)

MCQ

+2 M

Neg: -0.67

Q.15 The line integral of the vector field

$$\vec{F} = zx\hat{i} + xy\hat{j} + yz\hat{k}$$

along the boundary of the triangle with vertices $(1,0,0)$, $(0,1,0)$ and $(0,0,1)$, oriented anti-clockwise, when viewed from the point $(2,2,2)$, is

(A) $\frac{-1}{2}$

(B) -2

(C) $\frac{1}{2}$

(D) 2

Question 31 (JAM 2017 • Q14 • ID: JAM_2017_Q14)

MCQ

+2 M

Neg: -0.67

Q.14 Let $\vec{F} = (3 + 2xy)\hat{i} + (x^2 - 3y^2)\hat{j}$ and let L be the curve

$$\vec{r}(t) = e^t \sin t \hat{i} + e^t \cos t \hat{j}, \quad 0 \leq t \leq \pi.$$

Then

$$\int_L \vec{F} \cdot d\vec{r} =$$

(A) $e^{-3\pi} + 1$

(B) $e^{-6\pi} + 2$

(C) $e^{6\pi} + 2$

(D) $e^{3\pi} + 1$

Question 32 (JAM 2017 • Q11 • ID: JAM_2017_Q11)

MCQ

+2 M

Neg: -0.67

Q.11 The flux of the vector field

$$\vec{F} = \left(2\pi x + \frac{2x^2 y^2}{\pi} \right) \hat{i} + \left(2\pi xy - \frac{4y}{\pi} \right) \hat{j}$$

along the outward normal, across the ellipse $x^2 + 16y^2 = 4$ is equal to

- (A) $4\pi^2 - 2$ (B) $2\pi^2 - 4$ (C) $\pi^2 - 2$ (D) 2π

Question 33 (JAM 2016 • Q55 • ID: JAM_2016_Q55)

NAT

+2 M

Neg: Nil (0)

Q.55 Let S be the closed surface forming the boundary of the region V bounded by $x^2 + y^2 = 3$, $z = 0$, $z = 6$. A vector field $\vec{F}$ is defined over V with $\nabla \cdot \vec{F} = 2y + z + 1$. Then the value of

$$\frac{1}{\pi} \iint_S \vec{F} \cdot \hat{n} \, ds,$$

where $\hat{n}$ is the unit outward drawn normal to the surface S , is _____,

Question 34 (JAM 2016 • Q54 • ID: JAM_2016_Q54)

NAT

+2 M

Neg: Nil (0)

Q.54 Let C be the boundary of the region enclosed by $y = x^2$, $y = x + 2$, and $x = 0$. Then the value of the line integral

$$\oint_C (xy - y^2)dx - x^3 dy,$$

where C is traversed in the counter clockwise direction, is _____

Question 35 (JAM 2016 • Q45 • ID: JAM_2016_Q45)

NAT

+1 M

Neg: Nil (0)

Q.45 Let $\vec{F} = \sqrt{x} \hat{i} + (x + y^3) \hat{j}$ be a vector field for all (x, y) with $x \geq 0$ and $\vec{r} = x\hat{i} + y\hat{j}$. Then the value of the line integral $\int_C \vec{F} \cdot d\vec{r}$ from $(0, 0)$ to $(1, 1)$ along the path $C: x = t^2, y = t^3, 0 \leq t \leq 1$ is _____

Question 36 (JAM 2016 • Q40 • ID: JAM_2016_Q40)

MSQ

+2 M

Neg: Nil (0)

Q.40 Let $S = \left\{ \frac{1}{3^n} + \frac{1}{7^m} \mid n, m \in \mathbb{N} \right\}$. Then which of the following statement(s) is(are) TRUE?

- (A) S is closed
- (B) S is not open
- (C) S is connected
- (D) 0 is a limit point of S

SECTION – C

NUMERICAL ANSWER TYPE (NAT)

Q. 41 – Q. 50 carry one mark each.

Question 37 (JAM 2016 • Q36 • ID: JAM_2016_Q36)

MSQ

+2 M

Neg: Nil (0)

Q.36 Which of the following statement(s) is(are) TRUE?

- (A) There exists a connected set in $\mathbb{R}$ which is not compact
- (B) Arbitrary union of closed intervals in $\mathbb{R}$ need not be compact
- (C) Arbitrary union of closed intervals in $\mathbb{R}$ is always closed
- (D) Every bounded infinite subset V of $\mathbb{R}$ has a limit point in V itself

Question 38 (JAM 2016 • Q34 • ID: JAM_2016_Q34)

MSQ

+2 M

Neg: Nil (0)

Q.34 Consider the vector field $\vec{F} = x\hat{i} + y\hat{j}$ on an open connected set $S \subset \mathbb{R}^2$. Then which of the following statement(s) is(are) TRUE ?

- (A) Divergence of $\vec{F}$ is zero on S
- (B) The line integral of $\vec{F}$ is independent of path in S
- (C) $\vec{F}$ can be expressed as a gradient of a scalar function on S
- (D) The line integral of $\vec{F}$ is zero around any piecewise smooth closed path in S

Question 39 (JAM 2016 • Q23 • ID: JAM_2016_Q23)

MCQ

+2 M

Neg: -0.67

Q.23 Let $S \subset \mathbb{R}$ and ∂S denote the set of points x in $\mathbb{R}$ such that every neighbourhood of x contains some points of S as well as some points of complement of S . Further, let $\bar{S}$ denote the closure of S . Then which one of the following is FALSE?

- (A) $\partial \mathbb{Q} = \mathbb{R}$
- (B) $\partial(\mathbb{R} \setminus T) = \partial T$, $T \subset \mathbb{R}$
- (C) $\partial(T \cup V) = \partial T \cup \partial V$, $T, V \subset \mathbb{R}$, $T \cap V \neq \emptyset$
- (D) $\partial T = \bar{T} \cap (\mathbb{R} \setminus T)$, $T \subset \mathbb{R}$

Question 40 (JAM 2016 • Q20 • ID: JAM_2016_Q20)

MCQ

+2 M

Neg: -0.67

Q.20 Let S be the surface of the cone $z = \sqrt{x^2 + y^2}$ bounded by the planes $z = 0$ and $z = 3$. Further, let C be the closed curve forming the boundary of the surface S . A vector field $\vec{F}$ is such that $\nabla \times \vec{F} = -x\hat{i} - y\hat{j}$. The absolute value of the line integral $\oint_C \vec{F} \cdot d\vec{r}$, where $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ and $r = |\vec{r}|$, is

- (A) 0
- (B) 9π
- (C) 15π
- (D) 18π

Question 41 (JAM 2016 • Q19 • ID: JAM_2016_Q19)

MCQ

+2 M

Neg: -0.67

Q.19 Consider the vector field $\vec{F} = r^\beta(y\hat{i} - x\hat{j})$, where $\beta \in \mathbb{R}$, $\vec{r} = x\hat{i} + y\hat{j}$ and $r = |\vec{r}|$. If the absolute value of the line integral $\oint_C \vec{F} \cdot d\vec{r}$ along the closed curve $C: x^2 + y^2 = a^2$ (oriented counter clockwise) is 2π , then β is

- (A) -2
- (B) -1
- (C) 1
- (D) 2

Question 42 (JAM 2016 • Q10 • ID: JAM_2016_Q10)

MCQ

+1 M

Neg: -0.33

Q.10 Let S be a closed subset of $\mathbb{R}$, T a compact subset of $\mathbb{R}$ such that $S \cap T \neq \emptyset$. Then $S \cap T$ is

- (A) closed but not compact
- (B) not closed
- (C) compact
- (D) neither closed nor compact

Q. 11 – Q. 30 carry two marks each.

Question 43 (JAM 2016 • Q7 • ID: JAM_2016_Q7)

MCQ

+1 M

Neg: -0.33

Q.7 Let $\vec{r} = (x\hat{i} + y\hat{j} + z\hat{k})$ and $r = |\vec{r}|$. If $f(r) = \ln r$ and $g(r) = \frac{1}{r}$, $r \neq 0$, satisfy $2\nabla f + h(r)\nabla g = \vec{0}$, then $h(r)$ is

(A) r (B) $\frac{1}{r}$ (C) $2r$ (D) $\frac{2}{r}$

Question 44 (JAM 2015 • Q51 • ID: JAM_2015_Q51)

NAT

+2 M

Neg: Nil (0)

Q. 11 – Q. 20 carry two marks each.

Q.11 Let R be the planar region bounded by the lines $x = 0$, $y = 0$ and the curve $x^2 + y^2 = 4$, in the first quadrant. Let C be the boundary of R , oriented counter-clockwise. Then the value of

$$\oint_C x(1-y) dx + (x^2 - y^2) dy$$

is _____

Question 45 (JAM 2015 • Q41 • ID: JAM_2015_Q41)

NAT

+1 M

Neg: Nil (0)

Q. 1 – Q. 10 carry one mark each.

Q.1

Let C be the straight line segment from $P(0, \pi)$ to $Q\left(4, \frac{\pi}{2}\right)$, in the xy -plane. Then the value of $\int_C e^x (\cos y dx - \sin y dy)$ is _____

Question 46 (JAM 2015 • Q34 • ID: JAM_2015_Q34)

MSQ

+2 M

Neg: Nil (0)

Q.4

Let $\vec{F}$ be a vector field given by $\vec{F}(x, y, z) = -y\hat{i} + 2xy\hat{j} + z^3\hat{k}$, for $(x, y, z) \in \mathbb{R}^3$. If C is the curve of intersection of the surfaces $x^2 + y^2 = 1$ and $y + z = 2$, then which of the following is (are) equal to $\left| \int_C \vec{F} \cdot d\vec{r} \right|$?

(A) $\int_0^{2\pi} \int_0^1 (1 + 2r \sin \theta) r dr d\theta$

(B) $\int_0^{2\pi} \left(\frac{1}{2} + \frac{2}{3} \sin \theta \right) d\theta$

(C) $\int_0^{2\pi} \int_0^1 (1 + 2r \sin \theta) dr d\theta$

(D) $\int_0^{2\pi} (1 + \sin \theta) d\theta$

Question 47 (JAM 2015 • Q29 • ID: JAM_2015_Q29)

MCQ

+2 M

Neg: -0.67

Q.29

Let G and H be nonempty subsets of $\mathbb{R}$, where G is connected and $G \cup H$ is not connected. Which one of the following statements is true for all such G and H ?

- (A) If $G \cap H = \emptyset$, then H is connected
- (B) If $G \cap H = \emptyset$, then H is not connected
- (C) If $G \cap H \neq \emptyset$, then H is connected
- (D) If $G \cap H \neq \emptyset$, then H is not connected

MA

10/16

Question 48 (JAM 2015 • Q20 • ID: JAM_2015_Q20)

MCQ

+2 M

Neg: -0.67

Q.20

Let $S = \bigcap_{n=1}^{\infty} \left(\left[0, \frac{1}{2n+1} \right] \cup \left[\frac{1}{2n}, 1 \right] \right)$. Which one of the following statements is **FALSE**?

- (A) There exist sequences $\{a_n\}$ and $\{b_n\}$ in $[0,1]$ such that $S = [0,1] \setminus \bigcup_{n=1}^{\infty} (a_n, b_n)$
- (B) $[0,1] \setminus S$ is an open set
- (C) If A is an infinite subset of S , then A has a limit point
- (D) There exists an infinite subset of S having no limit points

Question 49 (JAM 2015 • Q10 • ID: JAM_2015_Q10)

MCQ

+1 M

Neg: -0.33

Q.10 Let A be a nonempty subset of $\mathbb{R}$. Let $I(A)$ denote the set of interior points of A . Then $I(A)$ can be

- (A) empty
- (B) singleton
- (C) a finite set containing more than one element
- (D) countable but not finite

Q. 11 – Q. 30 carry two marks each.

Question 50 (JAM 2015 • Q4 • ID: JAM_2015_Q4)

MCQ

+1 M

Neg: -0.33

Q.4 Let S be a nonempty subset of $\mathbb{R}$. If S is a finite union of disjoint bounded intervals, then which one of the following is true?

- (A) If S is not compact, then $\sup S \notin S$ and $\inf S \notin S$
- (B) Even if $\sup S \in S$ and $\inf S \in S$, S need not be compact
- (C) If $\sup S \in S$ and $\inf S \in S$, then S is compact
- (D) Even if S is compact, it is not necessary that $\sup S \in S$ and $\inf S \in S$

Question 51 (JAM 2014 • Q35 • ID: JAM_2014_Q35)

MCQ

+2 M

Neg: -0.67

Q.35 Let $S = \{x \in \mathbb{R} : x^6 - x^5 \leq 100\}$ and $T = \{x^2 - 2x : x \in (0, \infty)\}$. The set $S \cap T$ is

- (A) closed and bounded in $\mathbb{R}$
- (B) closed but NOT bounded in $\mathbb{R}$
- (C) bounded but NOT closed in $\mathbb{R}$
- (D) neither closed nor bounded in $\mathbb{R}$

Question 52 (JAM 2014 • Q32 • ID: JAM_2014_Q32)

MCQ

+2 M

Neg: -0.67

Q.32 Let S be the oriented surface $x^2 + y^2 + z^2 = 1$ with the unit normal $\mathbf{n}$ pointing outward. For the vector field $\mathbf{F}(x, y, z) = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$, the value of $\iint_S \mathbf{F} \cdot \mathbf{n} \, dS$ is

- (A) $\frac{\pi}{3}$
- (B) 2π
- (C) $\frac{4\pi}{3}$
- (D) 4π

Question 53 (JAM 2014 • Q29 • ID: JAM_2014_Q29)

MCQ

+2 M

Neg: -0.67

Q.29 Let C be the boundary of the region $R = \{(x, y) \in \mathbb{R}^2 : -1 \leq y \leq 1, 0 \leq x \leq 1 - y^2\}$ oriented in the counterclockwise direction. Then the value of $\oint_C y \, dx + 2x \, dy$ is

- (A) $-\frac{4}{3}$
- (B) $-\frac{2}{3}$
- (C) $\frac{2}{3}$
- (D) $\frac{4}{3}$

Question 54 (JAM 2014 • Q28 • ID: JAM_2014_Q28)

MCQ

+2 M

Neg: -0.67

Q.28 If C is a smooth curve in $\mathbb{R}^3$ from $(-1, 0, 1)$ to $(1, 1, -1)$, then the value of $\int_C (2xy + z^2)dx + (x^2 + z)dy + (y + 2xz)dz$ is

- (A) 0 (B) 1 (C) 2 (D) 3

Question 55 (JAM 2014 • Q21 • ID: JAM_2014_Q21)

MCQ

+2 M

Neg: -0.67

Q.21 The set of all limit points of the set $\left\{ \frac{2}{x+1} : x \in (-1, 1) \right\}$ in $\mathbb{R}$ is

- (A) $[1, \infty)$ (B) $(1, \infty)$ (C) $[-1, 1]$ (D) $[-1, \infty)$

Question 56 (JAM 2014 • Q20 • ID: JAM_2014_Q20)

MCQ

+2 M

Neg: -0.67

Q.20 The set $\left\{ \frac{x^2}{1+x^2} : x \in \mathbb{R} \right\}$ is

- (A) connected but NOT compact in $\mathbb{R}$ (B) compact but NOT connected in $\mathbb{R}$
 (C) compact and connected in $\mathbb{R}$ (D) neither compact nor connected in $\mathbb{R}$

Question 57 (JAM 2013 • Q3 • ID: JAM_2013_Q3)

MCQ

+2 M

Neg: -0.67

Q.3 Let A and B be subsets of $\mathbb{R}$. Which of the following is **NOT** necessarily true?

- (A) $(A \cap B)^o \subseteq A^o \cap B^o$ (B) $A^o \cup B^o \subseteq (A \cup B)^o$
 (C) $\overline{A \cup B} \subseteq \overline{A} \cup \overline{B}$ (D) $\overline{A \cap B} \subseteq \overline{A} \cap \overline{B}$

Question 58 (JAM 2013 • Q2 • ID: JAM_2013_Q2)

MCQ

+2 M

Neg: -0.67

Q.2 The value of n for which the divergence of the function $\vec{F} = \frac{\vec{r}}{|\vec{r}|^n}$, $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$, $|\vec{r}| \neq 0$, vanishes is

- (A) 1 (B) -1 (C) 3 (D) -3

Question 59 (JAM 2012 • Q15 • ID: JAM_2012_Q15)

MCQ

+6 M

Neg: -2.0

Q.15 If C is a smooth curve in $\mathbb{R}^3$ from $(0,0,0)$ to $(2,1,-1)$, then the value of

$$\int_C (2xy + z)dx + (z + x^2)dy + (x + y)dz$$

is

(A) -1

(B) 0

(C) 1

(D) 2

MA-3 / 36

Question 60 (JAM 2012 • Q14 • ID: JAM_2012_Q14)

MCQ

+6 M

Neg: -2.0

Q.14 If $Y = \left\{ \frac{x}{1+|x|} \mid x \in \mathbb{R} \right\}$, then the set of all limit points of Y is

(A) $(-1,1)$

(B) $(-1,1]$

(C) $[0,1]$

(D) $[-1,1]$

Question 61 (JAM 2012 • Q9 • ID: JAM_2012_Q9)

MCQ

+6 M

Neg: -2.0

Q.9 Which one of the following is TRUE ?

(A) The characteristic of the ring $6\mathbb{Z}$ is 6

(B) The ring $6\mathbb{Z}$ has a zero divisor

(C) The characteristic of the ring $(\mathbb{Z}/6\mathbb{Z}) \times 6\mathbb{Z}$ is zero

(D) The ring $6\mathbb{Z} \times 6\mathbb{Z}$ is an integral domain

Question 62 (JAM 2011 • Q15 • ID: JAM_2011_Q15)

MCQ

+6 M

Neg: -2.0

Q.15 Consider the following subsets of $\mathbb{R}$:

$$E = \left\{ \frac{n}{n+1} : n \in \mathbb{N} \right\}, \quad F = \left\{ \frac{1}{1-x} : 0 \leq x < 1 \right\}.$$

Then

- (A) Both E and F are closed
 (B) E is closed and F is NOT closed
 (C) E is NOT closed and F is closed
 (D) Neither E nor F is closed

MA-3 / 40

Question 63 (JAM 2011 • Q8 • ID: JAM_2011_Q8)

MCQ

+6 M

Neg: -2.0

Q.8 Consider the vector field $\vec{F} = (ax + y + a)\hat{i} + \hat{j} - (x + y)\hat{k}$, where a is a constant. If $\vec{F} \cdot \text{curl } \vec{F} = 0$, then the value of a is

- (A) -1 (B) 0 (C) 1 (D) $\frac{3}{2}$

Question 64 (JAM 2011 • Q7 • ID: JAM_2011_Q7)

MCQ

+6 M

Neg: -2.0

Q.7 Let $\vec{F} = ay\hat{i} + z\hat{j} + x\hat{k}$ and C be the positively oriented closed curve given by $x^2 + y^2 = 1, z = 0$. If $\oint_C \vec{F} \cdot d\vec{r} = \pi$, then the value of a is

- (A) -1 (B) 0 (C) $\frac{1}{2}$ (D) 1

Question 65 (JAM 2010 • Q8 • ID: JAM_2010_Q8)

MCQ

+6 M

Neg: -2.0

Q.8 Let $\mathbf{R}[X]$ be the ring of real polynomials in the variable X . The number of ideals in the quotient ring $\mathbf{R}[X]/(X^2 - 3X + 2)$ is

- (A) 2 (B) 3 (C) 4 (D) 6

Question 66 (JAM 2010 • Q7 • ID: JAM_2010_Q7)

MCQ

+6 M

Neg: -2.0

Q.7 Let $\vec{F} = xy\hat{i} + y\hat{j} - yz\hat{k}$ denote the force field on a particle traversing the path L from $(0, 0, 0)$ to $(1, 1, 1)$ along the curve of intersection of the cylinder $y = x^2$ and the plane $z = x$. The work done by $\vec{F}$ is

- (A) 0 (B) $\frac{1}{4}$ (C) $\frac{1}{2}$ (D) 1

Question 67 (JAM 2010 • Q6 • ID: JAM_2010_Q6)

MCQ

+6 M

Neg: -2.0

Q.6 Let $\vec{F} = 2xyz e^{x^2}\hat{i} + z e^{x^2}\hat{j} + y e^{x^2}\hat{k}$ be the gradient of a scalar function. The value of $\int_L \vec{F} \cdot d\vec{r}$ along the oriented path L from $(0, 0, 0)$ to $(1, 0, 2)$ and then to $(1, 1, 2)$ is

- (A) 0 (B) $2e$ (C) e (D) e^2

Question 68 (JAM 2010 • Q4 • ID: JAM_2010_Q4)

MCQ

+6 M

Neg: -2.0

Q.4 Let S be an infinite subset of $\mathbf{R}$ such that $S \cap \mathbf{Q} = \emptyset$. Which of the following statements is true?

- (A) S must have a limit point which belongs to $\mathbf{Q}$
 (B) S must have a limit point which belongs to $\mathbf{R} \setminus \mathbf{Q}$
 (C) S cannot be a closed set in $\mathbf{R}$
 (D) $\mathbf{R} \setminus S$ must have a limit point which belongs to S

Question 69 (JAM 2009 • Q15 • ID: JAM_2009_Q15)

MCQ

+6 M

Neg: -2.0

Q.15 The value of c for which there exists a twice differentiable vector field $\vec{F}$ with $\text{curl } \vec{F} = 2x\hat{i} - 7y\hat{j} + cz\hat{k}$ is

- (A) 0 (B) 2 (C) 5 (D) 7

Question 70 (JAM 2009 • Q13 • ID: JAM_2009_Q13)

MCQ

+6 M

Neg: -2.0

Q.13 Suppose $\vec{V} = p(x, y)\hat{i} + q(x, y)\hat{j}$ is a continuously differentiable vector field defined in a domain D in $\mathbb{R}^2$. Which one of the following statements is NOT equivalent to the remaining ones?

- (A) There exists a function $\phi(x, y)$ such that $\frac{\partial \phi}{\partial x} = p(x, y)$ and $\frac{\partial \phi}{\partial y} = q(x, y)$ for all $(x, y) \in D$
- (B) $\frac{\partial q}{\partial x} = \frac{\partial p}{\partial y}$ holds at all points of D
- (C) $\oint_C \vec{V} \cdot d\vec{r} = 0$ for every piecewise smooth closed curve C in D
- (D) The differential $pdx + qdy$ is exact in D

Question 71 (JAM 2009 • Q2 • ID: JAM_2009_Q2)

MCQ

+6 M

Neg: -2.0

Q.2 Let R be the ring of all functions from $\mathbb{R}$ to $\mathbb{R}$ under point-wise addition and multiplication. Let $I = \{f: \mathbb{R} \rightarrow \mathbb{R} \mid f \text{ is a bounded function}\}$, $J = \{f: \mathbb{R} \rightarrow \mathbb{R} \mid f(3) = 0\}$. Then

- (A) J is an ideal of R but I is not an ideal of R
- (B) I is an ideal of R but J is not an ideal of R
- (C) both I and J are ideals of R
- (D) neither I nor J is an ideal of R

Question 72 (JAM 2008 • Q10 • ID: JAM_2008_Q10)

MCQ

+6 M

Neg: -2.0

Q.10 The set $R = \{f \mid f \text{ is a function from } \mathbb{Z} \text{ to } \mathbb{R}\}$ under the binary operations $+$ and $\cdot$ defined as $(f+g)(n) = f(n) + g(n)$ and $(f \cdot g)(n) = f(n)g(n)$ for all $n \in \mathbb{Z}$ forms a ring. Let $S_1 = \{f \in R \mid f(-n) = f(n) \text{ for all } n \in \mathbb{Z}\}$ and $S_2 = \{f \in R \mid f(0) = 0\}$. Then

- (A) S_1 and S_2 are both ideals in R
 (C) S_2 is an ideal in R while S_1 is not

- (B) S_1 is an ideal in R while S_2 is not
 (D) neither S_1 nor S_2 is an ideal in R

Question 73 (JAM 2008 • Q6 • ID: JAM_2008_Q6)

MCQ

+6 M

Neg: -2.0

Q.6 Let $V = \{(x, y, z) \in \mathbb{R}^3 : \frac{1}{4} \leq x^2 + y^2 + z^2 \leq 1\}$ and $\vec{F} = \frac{x\hat{i} + y\hat{j} + z\hat{k}}{(x^2 + y^2 + z^2)^2}$ for $(x, y, z) \in V$. Let $\hat{n}$ denote the outward unit normal vector to the boundary of V and S denote the part $\{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 = \frac{1}{4}\}$ of the boundary of V . Then $\iint_S \vec{F} \cdot \hat{n} \, dS$ is equal to

- (A) -8π (B) -4π (C) 4π (D) 8π

Question 74 (JAM 2008 • Q5 • ID: JAM_2008_Q5)

MCQ

+6 M

Neg: -2.0

Q.5 The set of all boundary points of $\mathbb{Q}$ in $\mathbb{R}$ is

- (A) $\mathbb{R}$ (B) $\mathbb{R} \setminus \mathbb{Q}$ (C) $\mathbb{Q}$ (D) $\emptyset$

Question 75 (JAM 2008 • Q4 • ID: JAM_2008_Q4)

MCQ

+6 M

Neg: -2.0

Q.4 The work done by the force $\vec{F} = 4y\hat{i} - 3xy\hat{j} + z^2\hat{k}$ in moving a particle over the circular path $x^2 + y^2 = 1, z = 0$ from $(1, 0, 0)$ to $(0, 1, 0)$ is

- (A) $\pi + 1$ (B) $\pi - 1$ (C) $-\pi + 1$ (D) $-\pi - 1$

Question 76 (JAM 2007 • Q15 • ID: JAM_2007_Q15)

MCQ

+6 M

Neg: -2.0

15. Let C denote the boundary of the semi-circular disk $D = \{(x, y) \in \mathbf{R}^2 : x^2 + y^2 \leq 1, y \geq 0\}$ and let $\varphi(x, y) = x^2 + y$ for $(x, y) \in D$. If $\hat{n}$ is the outward unit normal to C , then the integral $\oint_C (\vec{\nabla} \varphi) \cdot \hat{n} ds$, evaluated counter-clockwise over C , is equal to

- (A) 0
- (B) $\pi - 2$
- (C) π
- (D) $\pi + 2$

Question 77 (JAM 2007 • Q13 • ID: JAM_2007_Q13)

MCQ

+6 M

Neg: -2.0

13. Let $\vec{u} = (ae^x \sin y - 4x)\hat{i} + (2y + e^x \cos y)\hat{j} + az\hat{k}$, where a is a constant. If the line integral $\oint_C \vec{u} \cdot d\vec{r}$ over every closed curve C is zero, then a is equal to

- (A) -2
- (B) -1
- (C) 0
- (D) 1

Question 78 (JAM 2007 • Q4 • ID: JAM_2007_Q4)

MCQ

+6 M

Neg: -2.0

4. Let R be the ring of polynomials over $\mathbf{Z}_2$ and let I be the ideal of R generated by the polynomial $x^3 + x + 1$. Then the number of elements in the quotient ring R/I is
- (A) 2
 - (B) 4
 - (C) 8
 - (D) 16

Question 79 (JAM 2006 • Q14 • ID: JAM_2006_Q14)

MCQ

+6 M

Neg: -2.0

14. Let R be the ring of all 2×2 matrices with integer entries. Which of the following subsets of R is an integral domain?

- (A) $\left\{ \begin{pmatrix} 0 & x \\ y & 0 \end{pmatrix} : x, y \in \mathbf{Z} \right\}$
- (B) $\left\{ \begin{pmatrix} x & 0 \\ 0 & y \end{pmatrix} : x, y \in \mathbf{Z} \right\}$
- (C) $\left\{ \begin{pmatrix} x & 0 \\ 0 & x \end{pmatrix} : x \in \mathbf{Z} \right\}$
- (D) $\left\{ \begin{pmatrix} x & y \\ y & z \end{pmatrix} : x, y, z \in \mathbf{Z} \right\}$

Question 80 (JAM 2006 • Q9 • ID: JAM_2006_Q9)

MCQ

+6 M

Neg: -2.0

9. Let S be a closed surface for which $\iint_S \vec{r} \cdot \hat{n} d\sigma = 1$. Then the volume enclosed by the surface is

- (A) 1
- (B) $1/3$
- (C) $2/3$
- (D) 3

Question 81 (JAM 2006 • Q7 • ID: JAM_2006_Q7)

MCQ

+6 M

Neg: -2.0

7. Let r be the distance of a point $P(x, y, z)$ from the origin O . Then ∇r is a vector

- (A) orthogonal to $\overrightarrow{OP}$
- (B) normal to the level surface of r at P
- (C) normal to the surface of revolution generated by OP about x-axis
- (D) normal to the surface of revolution generated by OP about y-axis

Question 82 (JAM 2006 • Q6 • ID: JAM_2006_Q6)

MCQ

+6 M

Neg: -2.0

6. Let C be the circle $x^2 + y^2 = 1$ taken in the anti-clockwise sense. Then the value of the integral

$$\int_C \left[(2xy^3 + y)dx + (3x^2y^2 + 2x)dy \right]$$

equals

- (A) 1
- (B) $\pi/2$
- (C) π
- (D) 0

Question 83 (JAM 2006 • Q4 • ID: JAM_2006_Q4)

MCQ

+6 M

Neg: -2.0

4. Let G be the set of all irrational numbers. The interior and the closure of G are denoted by G^0 and $\overline{G}$, respectively. Then

- (A) $G^0 = \emptyset$, $\overline{G} = G$
- (B) $G^0 = \mathbb{I}$, $\overline{G} = \mathbb{I}$
- (C) $G^0 = \emptyset$, $\overline{G} = \mathbb{I}$
- (D) $G^0 = G$, $\overline{G} = \mathbb{I}$

Question 84 (JAM 2005 • Q13 • ID: JAM_2005_Q13)

MCQ

+6 M

Neg: -2.0

13. Let $f: \mathbf{R} \longrightarrow \mathbf{R}$ be defined by $f(t) = t^2$ and let U be any non-empty open subset of $\mathbf{R}$. Then

- (A) $f(U)$ is open
- (B) $f^{-1}(U)$ is open
- (C) $f(U)$ is closed
- (D) $f^{-1}(U)$ is closed

Question 85 (JAM 2005 • Q9 • ID: JAM_2005_Q9)

MCQ

+6 M

Neg: -2.0

9. A ring R has maximal ideals

- (A) if R is infinite
- (B) if R is finite
- (C) if R is finite with at least 2 elements
- (D) only if R is finite

Question 86 (JAM 2005 • Q5 • ID: JAM_2005_Q5)

MCQ

+6 M

Neg: -2.0

5. Let $\vec{F} = x\hat{i} + 2y\hat{j} + 3z\hat{k}$, S be the surface of the sphere $x^2 + y^2 + z^2 = 1$ and $\hat{n}$ be the inward unit normal vector to S . Then $\iint_S \vec{F} \cdot \hat{n} dS$ is equal to

- (A) 4π
- (B) -4π
- (C) 8π
- (D) -8π

Question 87 (JAM 2005 • Q4 • ID: JAM_2005_Q4)

MCQ

+6 M

Neg: -2.0

4. Let $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$. If $\phi(x, y, z)$ is a solution of the Laplace equation then the vector field $\left(\vec{\nabla} \phi + \vec{r} \right)$ is

- (A) neither solenoidal nor irrotational
- (B) solenoidal but not irrotational
- (C) both solenoidal and irrotational
- (D) irrotational but not solenoidal

OFFICIAL MASTER ANSWER KEY & EVALUATION RUBRIC

Out of Syllabus Problems (2005–2026 Archive) • All Official Answers & Acceptance Ranges

Q. No.	Type	Marks	Official Key	Q. No.	Type	Marks	Official Key	Q. No.	Type	Marks	Official Key
1	NAT	+2	2	30	MCQ	+2	C	59	MCQ	+6	C
2	NAT	+1	1	31	MCQ	+2	D	60	MCQ	+6	D
3	NAT	+1	−2	32	MCQ	+2	B	61	MCQ	+6	C
4	MSQ	+2	B, C	33	NAT	+2	72.0 to 72.0	62	MCQ	+6	C
5	MCQ	+2	B	34	NAT	+2	0.8 to 1.9	63	MCQ	+6	D
6	MCQ	+1	C	35	NAT	+1	1.49 to 1.55	64	MCQ	+6	A
7	NAT	+2	0.65 to 0.67	36	MSQ	+2	B;D	65	MCQ	+6	C
8	NAT	+1	32 to 32	37	MSQ	+2	A;B	66	MCQ	+6	C
9	MCQ	+2	C	38	MSQ	+2	B;C;D	67	MCQ	+6	B
10	MCQ	+2	C	39	MCQ	+2	C	68	MCQ	+6	D
11	NAT	+1	75.35 to 75.45	40	MCQ	+2	MTA	69	MCQ	+6	C
12	MSQ	+2	B, D	41	MCQ	+2	A	70	MCQ	+6	B
13	MSQ	+2	A, D	42	MCQ	+1	C	71	MCQ	+6	A
14	MCQ	+2	C	43	MCQ	+1	C	72	MCQ	+6	C
15	MCQ	+2	B	44	NAT	+2	8	73	MCQ	+6	A
16	MCQ	+2	B	45	NAT	+1	1	74	MCQ	+6	A
17	MCQ	+2	B	46	MSQ	+2	A;B	75	MCQ	+6	D
18	MCQ	+1	B	47	MCQ	+2	D	76	MCQ	+6	C
19	MSQ	+2	A;B;C	48	MCQ	+2	D	77	MCQ	+6	D
20	MSQ	+2	B;C;D	49	MCQ	+1	A	78	MCQ	+6	C
21	MSQ	+2	A;B;C	50	MCQ	+1	B	79	MCQ	+6	C
22	MSQ	+2	B;C;D	51	MCQ	+2	A	80	MCQ	+6	B
23	MCQ	+2	B	52	MCQ	+2	D	81	MCQ	+6	B
24	MCQ	+1	C	53	MCQ	+2	D	82	MCQ	+6	C
25	NAT	+2	2.0 to 2.2	54	MCQ	+2	C	83	MCQ	+6	C
26	MSQ	+2	B, D	55	MCQ	+2	A	84	MCQ	+6	B
27	MCQ	+2	D	56	MCQ	+2	A	85	MCQ	+6	C
28	MCQ	+2	D	57	MCQ	+2	D	86	MCQ	+6	D
29	MCQ	+2	A	58	MCQ	+2	C	87	MCQ	+6	D

SCORING METHODOLOGY & EVALUATION GUIDELINES

- **Multiple Choice Questions (MCQ):** Full marks for correct single choice. Negative marks deducted for incorrect attempts. Unattempted questions award zero.
- **Multiple Select Questions (MSQ):** Full marks if and only if all correct choices are marked and no incorrect choice is selected. Zero marks for partial selection or any incorrect choice. No negative marking.
- **Numerical Answer Type (NAT):** Any numerical value falling strictly within the specified official range receives full credit. No negative marking.

- **Marks to All (MTA):** Questions officially designated as MTA award full marks to all candidates regardless of attempt.