

Matrices and Systems of Linear Equations (2005–2026 Archive)

Category: Linear Algebra and Algebra • Official Mock Test Series

TOTAL QUESTIONS

91

TOTAL MARKS

183 M

TEST DURATION

273 Min

PAPER PATTERN

MCQ • MSQ • NAT
(2005–20)

Section / Type	Questions	Marking Scheme	Negative Marking	Format / Details
Multiple Choice (MCQ)	52	+1 / +2 Marks	$-1/3$ / $-2/3$ Marks	4 Options (1 Correct)
Multiple Select (MSQ)	12	+2 Marks	Nil (0)	4 Options (1+ Correct)
Numerical Answer (NAT)	27	+1 / +2 Marks	Nil (0)	Real Decimal Value

IMPORTANT INSTRUCTIONS FOR CANDIDATES

- Authentic Examination PYQs:** This mock paper contains verified official IIT JAM Mathematics questions and high-resolution problem statements.
- Section A (MCQ):** Each question has four choices (A), (B), (C), (D), out of which **only one** is correct. Wrong answers carry negative penalty ($1/3$ of positive marks).
- Section B (MSQ):** Each question has four choices (A), (B), (C), (D), out of which **one or more than one** choice(s) may be correct. Full marks are awarded only if all correct choices and zero incorrect choices are chosen. No partial marks and no negative marks.
- Section C (NAT):** Numerical answers are real numbers entered via virtual keyboard. Full credit is awarded if the numerical answer falls within the official accepted interval. No negative marking.
- Master Answer Key:** Complete official answers and acceptance bounds are provided at the end of this document.
- Online Interactive Simulator:** Practice this test with virtual calculator, timers, and instant scoring analytics at vaibhavgeometer.github.io.

IIT JAM Mathematics Mock Test Series • Curated by Vaibhav Geometer

Question 1 (JAM 2026 • Q59 • ID: JAM_2026_Q59)

NAT

+2 M

Neg: Nil (0)

Q.59

Let A be a 3×3 real matrix such that given any column vector $x \in \mathbb{R}^3$, the column vector Ax is the reflection of x about the plane

$$\{(a, b, -a - b) : a, b \in \mathbb{R}\}.$$

Then the sum of the diagonal elements of A is _____ (rounded off to one decimal place).

Question 2 (JAM 2026 • Q57 • ID: JAM_2026_Q57)

NAT

+2 M

Neg: Nil (0)

Q.57

Let $a, b \in \mathbb{R}$. If the system of linear equations

$$x + 2y + 2z = 1$$

$$2x + 3y + z = 2$$

$$ax + 5y + bz = b$$

has infinitely many solutions, then $a + b =$ _____ (rounded off to one decimal place).

Question 3 (JAM 2026 • Q56 • ID: JAM_2026_Q56)

NAT

+2 M

Neg: Nil (0)

Q.56

Let $M \in M_3(\mathbb{R})$. If

$$M \begin{pmatrix} \sqrt{15} \\ \sqrt{15} \\ 0 \end{pmatrix} = \begin{pmatrix} \sqrt{3} \\ \sqrt{2} \\ -5 \end{pmatrix}, \quad M \begin{pmatrix} 0 \\ 0 \\ \sqrt{30} \end{pmatrix} = \begin{pmatrix} \sqrt{15} \\ \sqrt{10} \\ \sqrt{5} \end{pmatrix}, \quad M \begin{pmatrix} -\sqrt{15} \\ \sqrt{15} \\ 0 \end{pmatrix} = \begin{pmatrix} \sqrt{12} \\ -\sqrt{18} \\ 0 \end{pmatrix}$$

then $\det(M) =$ _____ (rounded off to one decimal place).

Question 4 (JAM 2026 • Q50 • ID: JAM_2026_Q50)

NAT

+1 M

Neg: Nil (0)

Q.50

Let P be a 5×5 real matrix with $\det(P) = 2$. Let Q be the matrix of cofactors of P . Then $\det(Q) = \underline{\hspace{2cm}}$ (rounded off to one decimal place).

Question 5 (JAM 2026 • Q49 • ID: JAM_2026_Q49)

NAT

+1 M

Neg: Nil (0)

Q.49

Let $A \in M_3(\mathbb{C})$. Suppose the column vector $v = \begin{pmatrix} \sqrt{5}i \\ 2i \\ x \end{pmatrix}$ in $\mathbb{C}^3$ belongs to the intersection of $\text{nullspace}(A)$ and $\text{rangespace}(A^T)$.
Then $|x| = \underline{\hspace{2cm}}$ (rounded off to one decimal place).

Question 6 (JAM 2026 • Q47 • ID: JAM_2026_Q47)

NAT

+1 M

Neg: Nil (0)

Q.47

Let C_n denote a cyclic group having n elements. If there is a surjective group homomorphism from C_n to C_{30} , then the total number of such distinct surjective homomorphisms is $\underline{\hspace{2cm}}$.

Question 7 (JAM 2026 • Q39 • ID: JAM_2026_Q39)

MSQ

+2 M

Neg: Nil (0)

Q.39	Consider matrices P of order 4×6 and Q of order 6×4 with real entries such that $PQ = 0$ and $QP = 0$. Which of the following statements is/are TRUE?
(A)	$\text{rangespace}(P) \subseteq \text{nullspace}(Q)$ and $\text{rangespace}(Q) \subseteq \text{nullspace}(P)$.
(B)	$\text{rank}(P) + \text{rank}(Q) \leq 4$.
(C)	If $\text{rangespace}(P) = \text{nullspace}(Q)$, then $\text{rank}(P) + \text{rank}(Q) = 4$.
(D)	$\text{rangespace}(Q) = \text{nullspace}(P)$.

Question 8 (JAM 2026 • Q37 • ID: JAM_2026_Q37)

MSQ

+2 M

Neg: Nil (0)

Q.37

$$\text{Let } P = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 \end{bmatrix} \in M_5(\mathbb{C}).$$

Which of the following statements is/are TRUE?

- (A) $\text{nullity}(P - I) \geq 2$, where I is the 5×5 identity matrix.
- (B) P has 4 distinct eigenvalues in $\mathbb{C}$.
- (C) P has 3 distinct eigenvalues in $\mathbb{R}$.
- (D) If λ is an eigenvalue of P , then there exists a positive integer n such that $\lambda^n = 1$.

Question 9 (JAM 2026 • Q30 • ID: JAM_2026_Q30)

MCQ

+2 M

Neg: -0.67

Q.30	Let G be the power set of $\{1,2,3,4\}$. Let the binary operation Δ on G be $A \Delta B = (A \setminus B) \cup (B \setminus A).$ Which ONE of the following statements is TRUE?
(A)	$\{1\}$ is an identity element of (G, Δ) .
(B)	(G, Δ) is an abelian group but not cyclic.
(C)	(G, Δ) is a group and has an element of order 4.
(D)	(G, Δ) is a group and has an element of order 8.

Question 10 (JAM 2026 • Q25 • ID: JAM_2026_Q25)

MCQ

+2 M

Neg: -0.67

Q.25	Let P be a nonzero complex $n \times n$ matrix such that $v^* P v \geq 0$ for all column vectors v in $\mathbb{C}^n$. Which ONE of the following statements is necessarily TRUE?
(A)	All eigenvalues of P are negative real numbers.
(B)	If v_1, v_2 are column vectors in $\mathbb{C}^n$ such that $P v_1 = v_2$ and $P v_2 = v_1$, then $v_1 = v_2$.
(C)	$v^* P v \neq 0$ for all nonzero column vectors v in $\mathbb{C}^n$.
(D)	All eigenvalues of P^* are negative real numbers.

Question 11 (JAM 2026 • Q14 • ID: JAM_2026_Q14)

MCQ

+2 M

Neg: -0.67

Q.14	Let A be a 5×3 real matrix of rank 2 and b be a non-zero 5×1 real column vector. Suppose $\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ and $\begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix}$ are two solutions of the system of linear equations $Ax = b$. Which ONE of the following is also a solution of $Ax = b$?
(A)	$\begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix}$
(B)	$\begin{pmatrix} 5 \\ 7 \\ 9 \end{pmatrix}$
(C)	$\begin{pmatrix} 5 \\ 6 \\ 9 \end{pmatrix}$
(D)	$\begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix}$

Question 12 (JAM 2026 • Q8 • ID: JAM_2026_Q8)

MCQ

+1 M

Neg: -0.33

Q.8	Let A be an invertible 5×5 real matrix. Let B be the matrix obtained by interchanging the second and third columns of A and then adding 3 times the third column to the fifth column. Which ONE of the following statements is TRUE?
(A)	B^{-1} is obtained by interchanging the second and third rows of A^{-1} , and then adding 3 times the third row to the fifth row.
(B)	B^{-1} is obtained by interchanging the second and third rows of A^{-1} , and then adding -3 times the fifth row to the third row.
(C)	B^{-1} is obtained by adding -3 times the third row to the fifth row, and then interchanging the second and third rows of A^{-1} .
(D)	B^{-1} is obtained by adding 3 times the fifth row to the third row, and then interchanging the second and third rows of A^{-1} .

Question 13 (JAM 2025 • Q60 • ID: JAM_2025_Q60)

NAT

+2 M

Neg: Nil (0)

Q.60 The number of surjective (onto) group homomorphisms from S_4 to $\mathbb{Z}_6$ is equal to _____.

Question 14 (JAM 2025 • Q59 • ID: JAM_2025_Q59)

NAT

+2 M

Neg: Nil (0)

Q.59 Let G be an abelian group of order 35. Let m denote the number of elements of order 5 in G , and let n denote the number of elements of order 7 in G .
Then, the value of $m + n$ is equal to _____.

Question 15 (JAM 2025 • Q52 • ID: JAM_2025_Q52)

NAT

+2 M

Neg: Nil (0)

Q.52 Let $\sigma \in S_4$ be the permutation defined by $\sigma(1) = 2$, $\sigma(2) = 3$, $\sigma(3) = 1$ and $\sigma(4) = 4$. The number of elements in the set $\{\tau \in S_4 : \tau\sigma\tau^{-1} = \sigma\}$ is equal to _____.

Question 16 (JAM 2025 • Q39 • ID: JAM_2025_Q39)

MSQ

+2 M

Neg: Nil (0)

Q.39 For $n \in \mathbb{N}$, consider the set $U(n) = \{\bar{x} \in \mathbb{Z}_n : \gcd(x, n) = 1\}$ as a group under multiplication modulo n .

Then, which of the following is/are TRUE?

- (A) $U(8)$ is a cyclic group
- (B) $U(5)$ is a cyclic group
- (C) $U(12)$ is a cyclic group
- (D) $U(9)$ is a cyclic group

Question 17 (JAM 2025 • Q23 • ID: JAM_2025_Q23)

MCQ

+2 M

Neg: -0.67

Q.23 Let $\mathbb{R}/\mathbb{Z}$ denote the quotient group, where $\mathbb{Z}$ is considered as a subgroup of the additive group of real numbers $\mathbb{R}$.

Let m denote the number of injective (one-one) group homomorphisms from $\mathbb{Z}_3$ to $\mathbb{R}/\mathbb{Z}$ and n denote the number of group homomorphisms from $\mathbb{R}/\mathbb{Z}$ to $\mathbb{Z}_3$.

Then, which one of the following is TRUE?

(A) $m = 2$ and $n = 1$

(B) $m = 3$ and $n = 3$

(C) $m = 2$ and $n = 3$

(D) $m = 1$ and $n = 1$

Question 18 (JAM 2025 • Q22 • ID: JAM_2025_Q22)

MCQ

+2 M

Neg: -0.67

Q.22

Let $M = (m_{ij})$ be a 3×3 real, invertible matrix and $\sigma \in S_3$ be the permutation defined by $\sigma(1) = 2$, $\sigma(2) = 3$ and $\sigma(3) = 1$. The matrix $M_\sigma = (n_{ij})$ is defined by $n_{ij} = m_{i\sigma(j)}$ for all $i, j \in \{1, 2, 3\}$.

Then, which one of the following is TRUE?

(A) $\det(M) = \det(M_\sigma)$, and nullity of the matrix $M - M_\sigma$ is 0

(B) $\det(M) = -\det(M_\sigma)$, and nullity of the matrix $M - M_\sigma$ is 1

(C) $\det(M) = \det(M_\sigma)$, and nullity of the matrix $M - M_\sigma$ is 1

(D) $\det(M) = -\det(M_\sigma)$, and nullity of the matrix $M - M_\sigma$ is 0

Question 19 (JAM 2025 • Q16 • ID: JAM_2025_Q16)

MCQ

+2 M

Neg: -0.67

Q.16

Let $M = \begin{pmatrix} -2 & 0 & 0 \\ 3 & 2 & 3 \\ 4 & -1 & x \end{pmatrix}$ for some real number x . Suppose that -2 and 3 are eigenvalues of M . If $M^3 \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 125 \\ 125 \end{pmatrix}$, then which one of the following is TRUE?

- (A) $x = 5$, and the matrix $M^2 + M$ is invertible
- (B) $x \neq 5$, and the matrix $M^2 + M$ is invertible
- (C) $x = 5$, and the matrix $M^2 + M$ is NOT invertible
- (D) $x \neq 5$, and the matrix $M^2 + M$ is NOT invertible

Question 20 (JAM 2025 • Q15 • ID: JAM_2025_Q15)

MCQ

+2 M

Neg: -0.67

Q.15

Let $M = \begin{pmatrix} 6 & 2 & -6 & 8 \\ 5 & 3 & -9 & 8 \\ 3 & 1 & -2 & 4 \end{pmatrix}$. Consider the system S of linear equations given by

$$\begin{aligned} 6x_1 + 2x_2 - 6x_3 + 8x_4 &= 8 \\ 5x_1 + 3x_2 - 9x_3 + 8x_4 &= 16 \\ 3x_1 + x_2 - 2x_3 + 4x_4 &= 32 \end{aligned}$$

where x_1, x_2, x_3, x_4 are unknowns.

Then, which one of the following is TRUE?

- (A) The rank of M is 3, and the system S has a solution
- (B) The rank of M is 3, and the system S does NOT have a solution
- (C) The rank of M is 2, and the system S has a solution
- (D) The rank of M is 2, and the system S does NOT have a solution

Question 21 (JAM 2025 • Q13 • ID: JAM_2025_Q13)

MCQ

+2 M

Neg: -0.67

Q.13 Let $X = \{x \in S_4 : x^3 = id\}$ and $Y = \{x \in S_4 : x^2 \neq id\}$.

If m and n denote the number of elements in X and Y , respectively, then which one of the following is TRUE?

- (A) m is even and n is even
- (B) m is odd and n is even
- (C) m is even and n is odd
- (D) m is odd and n is odd

Question 22 (JAM 2025 • Q9 • ID: JAM_2025_Q9)

MCQ

+1 M

Neg: -0.33

Q.9 Let G be a finite abelian group of order 10. Let x_0 be an element of order 2 in G .

If $X = \{x \in G : x^3 = x_0\}$, then which one of the following is TRUE?

- (A) X has exactly one element
- (B) X has exactly two elements
- (C) X has exactly three elements
- (D) X is an empty set

Question 23 (JAM 2025 • Q7 • ID: JAM_2025_Q7)

MCQ

+1 M

Neg: -0.33

Q.7

Let $M = \begin{pmatrix} 2 & 0 & -1 \\ 4 & 1 & -4 \\ 2 & 0 & x \end{pmatrix}$ for some real number x .

If 0 is an eigenvalue of M , then $(M^4 + M) \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$ is equal to

(A) $\begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$

(B) $\begin{pmatrix} 2 \\ 0 \\ 2 \end{pmatrix}$

(C) $\begin{pmatrix} 5 \\ 0 \\ 5 \end{pmatrix}$

(D) $\begin{pmatrix} 17 \\ 0 \\ 17 \end{pmatrix}$

Question 24 (JAM 2024 • Q50 • ID: JAM_2024_Q50)

NAT

+1 M

Neg: Nil (0)

Q.50 Let

$$M = \begin{pmatrix} 0 & 0 & 0 & 0 & -1 \\ 2 & 0 & 0 & 0 & -4 \\ 0 & 2 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 3 \\ 0 & 0 & 0 & 2 & 2 \end{pmatrix}.$$

If $p(x)$ is the characteristic polynomial of M , then $p(2) - 1$ equals _____

Question 25 (JAM 2024 • Q44 • ID: JAM_2024_Q44)

NAT

+1 M

Neg: Nil (0)

Q.44 Consider the 4×4 matrix

$$M = \begin{pmatrix} 0 & 1 & 2 & 3 \\ 1 & 0 & 1 & 2 \\ 2 & 1 & 0 & 1 \\ 3 & 2 & 1 & 0 \end{pmatrix}.$$

If $a_{i,j}$ denotes the $(i,j)^{\text{th}}$ entry of M^{-1} , then $a_{4,1}$ equals _____ (rounded off to two decimal places).

Question 26 (JAM 2024 • Q37 • ID: JAM_2024_Q37)

MSQ

+2 M

Neg: Nil (0)

Q.37 For a matrix M , let $\text{Rowspace}(M)$ denote the linear span of the rows of M and $\text{Colspace}(M)$ denote the linear span of the columns of M . Which of the following hold(s) for all $A, B, C \in M_{10}(\mathbb{R})$ satisfying $A = BC$?

- (A) $\text{Rowspace}(A) \subseteq \text{Rowspace}(B)$
- (B) $\text{Rowspace}(A) \subseteq \text{Rowspace}(C)$
- (C) $\text{Colspace}(A) \subseteq \text{Colspace}(B)$
- (D) $\text{Colspace}(A) \subseteq \text{Colspace}(C)$

Question 27 (JAM 2024 • Q16 • ID: JAM_2024_Q16)

MCQ

+2 M

Neg: -0.67

Q.16 Let A be a 6×5 matrix with entries in $\mathbb{R}$ and B be a 5×4 matrix with entries in $\mathbb{R}$. Consider the following two statements.

P: For all such nonzero matrices A and B , there is a nonzero matrix Z such that AZB is the 6×4 zero matrix.

Q: For all such nonzero matrices A and B , there is a nonzero matrix Y such that BYA is the 5×5 zero matrix.

Which one of the following holds?

- (A) Both P and Q are true
- (B) P is true but Q is false
- (C) P is false but Q is true
- (D) Both P and Q are false

Question 28 (JAM 2024 • Q15 • ID: JAM_2024_Q15)

MCQ

+2 M

Neg: -0.67

Q.15

Let $a = \begin{bmatrix} \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{6}} \\ 0 \end{bmatrix}$. Consider the following two statements.

P: The matrix $I_4 - aa^T$ is invertible.

Q: The matrix $I_4 - 2aa^T$ is invertible.

Then, which one of the following holds?

- (A) P is false but Q is true
- (B) P is true but Q is false
- (C) Both P and Q are true
- (D) Both P and Q are false

Question 29 (JAM 2024 • Q9 • ID: JAM_2024_Q9)

MCQ

+1 M

Neg: -0.33

Q.9 Consider the following statements.

P: If a system of linear equations $Ax = b$ has a unique solution, where A is an $m \times n$ matrix and b is an $m \times 1$ matrix, then $m = n$.

Q: For a subspace W of a nonzero vector space V , whenever $u \in V \setminus W$ and $v \in V \setminus W$, then $u + v \in V \setminus W$.

Which one of the following holds?

- (A) Both P and Q are true
- (B) P is true but Q is false
- (C) P is false but Q is true
- (D) Both P and Q are false

Question 30 (JAM 2024 • Q6 • ID: JAM_2024_Q6)

MCQ

+1 M

Neg: -0.33

Q.6 For a positive integer n , let $U(n) = \{\bar{r} \in \mathbb{Z}_n : \gcd(r, n) = 1\}$ be the group under multiplication modulo n . Then, which one of the following statements is TRUE?

- (A) $U(5)$ is isomorphic to $U(8)$
- (B) $U(10)$ is isomorphic to $U(12)$
- (C) $U(8)$ is isomorphic to $U(10)$
- (D) $U(8)$ is isomorphic to $U(12)$

Question 31 (JAM 2024 • Q5 • ID: JAM_2024_Q5)

MCQ

+1 M

Neg: -0.33

Q.5 Let G be a group of order 39 such that it has exactly one subgroup of order 3 and exactly one subgroup of order 13. Then, which one of the following statements is TRUE?

- (A) G is necessarily cyclic
- (B) G is abelian but need not be cyclic
- (C) G need not be abelian
- (D) G has 13 elements of order 13

Question 32 (JAM 2023 • Q58 • ID: JAM_2023_Q58)

NAT

+2 M

Neg: Nil (0)

Q. 58 The maximum number of linearly independent eigenvectors of the matrix

$$\begin{pmatrix} 1 & 1 & 0 & 0 \\ 2 & 2 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 1 & 3 \end{pmatrix}$$

is equal to _____.

Question 33 (JAM 2023 • Q56 • ID: JAM_2023_Q56)

NAT

+2 M

Neg: Nil (0)

Q. 56 Let

$$A = \begin{pmatrix} 1 & 1 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 3 \\ 1 & 1 & 4 & 4 & 4 \end{pmatrix}$$

and B be a 5×5 real matrix such that AB is the zero matrix. Then the maximum possible rank of B is equal to _____.

Question 34 (JAM 2023 • Q14 • ID: JAM_2023_Q14)

MCQ

+2 M

Neg: -0.67

Q. 14 The system of linear equations in x_1, x_2, x_3

$$\begin{pmatrix} 1 & 1 & 1 \\ 0 & -1 & 1 \\ 2 & 3 & \alpha \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3 \\ 1 \\ \beta \end{pmatrix}$$

where $\alpha, \beta \in \mathbb{R}$, has

- (A) at least one solution for any α and β
- (B) a unique solution for any β when $\alpha \neq 1$
- (C) NO solution for any α when $\beta \neq 5$
- (D) infinitely many solutions for any α when $\beta = 5$

Question 35 (JAM 2023 • Q13 • ID: JAM_2023_Q13)

MCQ

+2 M

Neg: -0.67

Q. 13 Let

$$A = \begin{pmatrix} 1 & -1 & 0 \\ 0 & 0 & 0 \\ -2 & 2 & 2 \end{pmatrix}$$

and $B = A^5 + A^4 + I_3$. Which of the following is NOT an eigenvalue of B ?

- (A) 1
- (B) 2
- (C) 49
- (D) 3

Question 36 (JAM 2023 • Q2 • ID: JAM_2023_Q2)

MCQ

+1 M

Neg: -0.33

Q. 2 Let $\mathbf{v}_1, \dots, \mathbf{v}_9$ be the column vectors of a non-zero 9×9 real matrix A . Let $a_1, \dots, a_9 \in \mathbb{R}$, not all zero, be such that $\sum_{i=1}^9 a_i \mathbf{v}_i = \mathbf{0}$. Then the system $A\mathbf{x} = \sum_{i=1}^9 \mathbf{v}_i$ has

- (A) no solution
- (B) a unique solution
- (C) more than one but only finitely many solutions
- (D) infinitely many solutions

Question 37 (JAM 2022 • Q57 • ID: JAM_2022_Q57)

NAT

+2 M

Neg: Nil (0)

Q.57

Consider the 4×4 matrix $M = \begin{pmatrix} 11 & 10 & 10 & 10 \\ 10 & 11 & 10 & 10 \\ 10 & 10 & 11 & 10 \\ 10 & 10 & 10 & 11 \end{pmatrix}$. Then the value of the determinant of M is equal to _____.

Question 38 (JAM 2022 • Q53 • ID: JAM_2022_Q53)

NAT

+2 M

Neg: Nil (0)

Q.53

Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ -1 & 1 \end{pmatrix}$ and let A^T denote the transpose of A . Let $u = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix}$ and $v = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}$ be column vectors with entries in $\mathbb{R}$ such that $u_1^2 + u_2^2 = 1$ and $v_1^2 + v_2^2 + v_3^2 = 1$. Suppose

$$Au = \sqrt{2} v \quad \text{and} \quad A^T v = \sqrt{2} u.$$

Then $|u_1 + 2\sqrt{2} v_1|$ is equal to _____. (Rounded off to two decimal places)

Question 39 (JAM 2022 • Q40 • ID: JAM_2022_Q40)

MSQ

+2 M

Neg: Nil (0)

Q.40	Let P be a fixed 3×3 matrix with entries in $\mathbb{R}$. Which of the following maps from $M_3(\mathbb{R})$ to $M_3(\mathbb{R})$ is/are linear?
(A)	$T_1: M_3(\mathbb{R}) \rightarrow M_3(\mathbb{R})$ given by $T_1(M) = MP - PM$ for $M \in M_3(\mathbb{R})$.
(B)	$T_2: M_3(\mathbb{R}) \rightarrow M_3(\mathbb{R})$ given by $T_2(M) = M^2P - P^2M$ for $M \in M_3(\mathbb{R})$.
(C)	$T_3: M_3(\mathbb{R}) \rightarrow M_3(\mathbb{R})$ given by $T_3(M) = MP^2 + P^2M$ for $M \in M_3(\mathbb{R})$.
(D)	$T_4: M_3(\mathbb{R}) \rightarrow M_3(\mathbb{R})$ given by $T_4(M) = MP^2 - PM^2$ for $M \in M_3(\mathbb{R})$.
Section C: Q.41 – Q.50 Carry ONE mark each.	

Question 40 (JAM 2022 • Q39 • ID: JAM_2022_Q39)

MSQ

+2 M

Neg: Nil (0)

Q.39	Let V be the real vector space consisting of all polynomials in one variable with real coefficients and having degree at most 5, together with the zero polynomial. Let $T: V \rightarrow \mathbb{R}$ be the linear map defined by $T(1) = 1$ and $T(x(x-1)\cdots(x-k+1)) = 1 \quad \text{for } 1 \leq k \leq 5.$ Then which of the following is/are true?
(A)	$T(x^4) = 15.$
(B)	$T(x^3) = 5.$
(C)	$T(x^4) = 14.$
(D)	$T(x^3) = 3.$

Question 41 (JAM 2022 • Q30 • ID: JAM_2022_Q30)

MCQ

+2 M

Neg: -0.67

Q.30	Let P be a 3×3 real matrix having eigenvalues $\lambda_1 = 0, \lambda_2 = 1$ and $\lambda_3 = -1$. Further, $v_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$, $v_2 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$ and $v_3 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$ are eigenvectors of the matrix P corresponding to the eigenvalues λ_1, λ_2 and λ_3 , respectively. Then the entry in the first row and the third column of P is
(A)	0
(B)	1
(C)	-1
(D)	2

Question 42 (JAM 2022 • Q26 • ID: JAM_2022_Q26)

MCQ

+2 M

Neg: -0.67


Q.26

For a 4×4 matrix $M \in M_4(\mathbb{C})$, let $\bar{M}$ denote the matrix obtained from M by replacing each entry of M by its complex conjugate. Consider the real vector space

$$H = \{M \in M_4(\mathbb{C}) : M^T = \bar{M}\}$$

where M^T denotes the transpose of M . The dimension of H as a vector space over $\mathbb{R}$ is equal to

- (A) 6
- (B) 16
- (C) 15
- (D) 12



Question 43 (JAM 2022 • Q20 • ID: JAM_2022_Q20)

MCQ

+2 M

Neg: -0.67

Q.20	For $P \in M_5(\mathbb{R})$ and $i, j \in \{1, 2, \dots, 5\}$, let p_{ij} denote the $(i, j)^{\text{th}}$ entry of P. Let $S = \{P \in M_5(\mathbb{R}) : p_{ij} = p_{rs} \text{ for } i, j, r, s \in \{1, 2, \dots, 5\} \text{ with } i + r = j + s\}.$ Then which one of the following is FALSE?
(A)	S is a subspace of the vector space over $\mathbb{R}$ of all 5×5 symmetric matrices.
(B)	The dimension of S over $\mathbb{R}$ is 5.
(C)	The dimension of S over $\mathbb{R}$ is 11.
(D)	If $P \in S$ and all the entries of P are integers, then 5 divides the sum of all the diagonal entries of P .

Question 44 (JAM 2022 • Q18 • ID: JAM_2022_Q18)

MCQ

+2 M

Neg: -0.67

Q.18

Consider the system of linear equations

$$\begin{aligned}x + y + t &= 4, \\2x - 4t &= 7, \\x + y + z &= 5, \\x - 3y - z - 10t &= \lambda,\end{aligned}$$

where x, y, z, t are variables and λ is a constant. Then which one of the following is true?

- (A) If $\lambda = 1$, then the system has a unique solution.
- (B) If $\lambda = 2$, then the system has infinitely many solutions.
- (C) If $\lambda = 1$, then the system has infinitely many solutions.
- (D) If $\lambda = 2$, then the system has a unique solution.

Question 45 (JAM 2022 • Q17 • ID: JAM_2022_Q17)

MCQ

+2 M

Neg: -0.67

Q.17	For $X, Y \in M_2(\mathbb{R})$, define $(X, Y) = XY - YX$. Let $\mathbf{0} \in M_2(\mathbb{R})$ denote the zero matrix. Consider the two statements: $P : (X, (Y, Z)) + (Y, (Z, X)) + (Z, (X, Y)) = \mathbf{0}$ for all $X, Y, Z \in M_2(\mathbb{R})$. $Q : (X, (Y, Z)) = ((X, Y), Z)$ for all $X, Y, Z \in M_2(\mathbb{R})$. Then which one of the following is correct?
(A)	Both P and Q are true.
(B)	P is true, but Q is false.
(C)	P is false, but Q is true.
(D)	Both P and Q are false.

Question 46 (JAM 2022 • Q16 • ID: JAM_2022_Q16)

MCQ

+2 M

Neg: -0.67

Q.16	Let $P \in M_4(\mathbb{R})$ be such that P^4 is the zero matrix, but P^3 is a nonzero matrix. Then which one of the following is FALSE?
(A)	For every nonzero vector $v \in \mathbb{R}^4$, the subset $\{v, Pv, P^2v, P^3v\}$ of the real vector space $\mathbb{R}^4$ is linearly independent.
(B)	The rank of P^k is $4 - k$ for every $k \in \{1, 2, 3, 4\}$.
(C)	0 is an eigenvalue of P .
(D)	If $Q \in M_4(\mathbb{R})$ is such that Q^4 is the zero matrix, but Q^3 is a nonzero matrix, then there exists a nonsingular matrix $S \in M_4(\mathbb{R})$ such that $S^{-1}QS = P$.

Question 47 (JAM 2022 • Q3 • ID: JAM_2022_Q3)

MCQ

+1 M

Neg: -0.33

Q.3	Let V be the real vector space consisting of all polynomials in one variable with real coefficients and having degree at most 6, together with the zero polynomial. Then which one of the following is true?
(A)	$\{f \in V : f(1/2) \notin \mathbb{Q}\}$ is a subspace of V .
(B)	$\{f \in V : f(1/2) = 1\}$ is a subspace of V .
(C)	$\{f \in V : f(1/2) = f(1)\}$ is a subspace of V .
(D)	$\{f \in V : f'(1/2) = 1\}$ is a subspace of V .

Question 48 (JAM 2022 • Q2 • ID: JAM_2022_Q2)

MCQ

+1 M

Neg: -0.33

Q.2	The rank of the 4×6 matrix $\begin{pmatrix} 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 \end{pmatrix}$ with entries in $\mathbb{R}$, is
(A)	1
(B)	2
(C)	3
(D)	4

Question 49 (JAM 2022 • Q1 • ID: JAM_2022_Q1)

MCQ

+1 M

Neg: -0.33

Q.1	Consider the 2×2 matrix $M = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \in M_2(\mathbb{R})$. If the eighth power of M satisfies $M^8 \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix}$, then the value of x is
(A)	21
(B)	22
(C)	34
(D)	35

Question 50 (JAM 2021 • Q59 • ID: JAM_2021_Q59)

NAT

+2 M

Neg: Nil (0)

Q. 59 Let $A = \begin{pmatrix} 2 & -1 & 3 \\ 2 & -1 & 3 \\ 3 & 2 & -1 \end{pmatrix}$. Then the largest eigenvalue of A is ____.

Question 51 (JAM 2021 • Q56 • ID: JAM_2021_Q56)

NAT

+2 M

Neg: Nil (0)

JAM 2021

MATHEMATICS - MA

Q. 56 The determinant of the matrix

$$\begin{pmatrix} 2021 & 2020 & 2020 & 2020 \\ 2021 & 2021 & 2020 & 2020 \\ 2021 & 2021 & 2021 & 2020 \\ 2021 & 2021 & 2021 & 2021 \end{pmatrix}$$

is ____.

Question 52 (JAM 2021 • Q40 • ID: JAM_2021_Q40)

MSQ

+2 M

Neg: Nil (0)

Q. 40 Let $m > 1$ and $n > 1$ be integers. Let A be an $m \times n$ matrix such that for some $m \times 1$ matrix b_1 , the equation $Ax = b_1$ has infinitely many solutions. Let b_2 denote an $m \times 1$ matrix different from b_1 . Then $Ax = b_2$ has

- (A) infinitely many solutions for some b_2 . (B) a unique solution for some b_2 .
 (C) no solution for some b_2 . (D) finitely many solutions for some b_2 .



MA

13 / 17

Question 53 (JAM 2021 • Q22 • ID: JAM_2021_Q22)

MCQ

+2 M

Neg: -0.67

Q. 22 Let A be an $n \times n$ invertible matrix and C be an $n \times n$ nilpotent matrix. If $X = \begin{pmatrix} X_{11} & X_{12} \\ X_{21} & X_{22} \end{pmatrix}$ is a $2n \times 2n$ matrix (each X_{ij} being $n \times n$) that commutes with the $2n \times 2n$ matrix $B = \begin{pmatrix} A & 0 \\ 0 & C \end{pmatrix}$, then

- (A) X_{11} and X_{22} are necessarily zero matrices.
- (B) X_{12} and X_{21} are necessarily zero matrices.
- (C) X_{11} and X_{21} are necessarily zero matrices.
- (D) X_{12} and X_{22} are necessarily zero matrices.

Admission test for Masters 2021
Institute:
of Science Bangalore

Question 54 (JAM 2021 • Q10 • ID: JAM_2021_Q10)

MCQ

+1 M

Neg: -0.33

JAM 2021

MATHEMATICS - MA

Q. 10 Let $n > 1$ be an integer. Consider the following two statements for an arbitrary $n \times n$ matrix A with complex entries.

- I. If $A^k = I_n$ for some integer $k \geq 1$, then all the eigenvalues of A are k^{th} roots of unity.
II. If, for some integer $k \geq 1$, all the eigenvalues of A are k^{th} roots of unity, then $A^k = I_n$.

Then

- (A) both I and II are TRUE. (B) I is TRUE but II is FALSE.
(C) I is FALSE but II is TRUE. (D) neither I nor II is TRUE.



MA

4 / 17

Question 55 (JAM 2020 • Q58 • ID: JAM_2020_Q58)

NAT

+2 M

Neg: Nil (0)

Q. 58 Let $M = \begin{bmatrix} 9 & 2 & 7 & 1 \\ 0 & 7 & 2 & 1 \\ 0 & 0 & 11 & 6 \\ 0 & 0 & -5 & 0 \end{bmatrix}$. Then, the value of $\det((8I - M)^3)$ is _____.

Question 56 (JAM 2020 • Q36 • ID: JAM_2020_Q36)

MSQ

+2 M

Neg: Nil (0)

Q. 36 Consider the following system of linear equations

$$x + y + 5z = 3, \quad x + 2y + mz = 5 \quad \text{and} \quad x + 2y + 4z = k.$$

The system is consistent if

(A) $m \neq 4$ (B) $k \neq 5$ (C) $m = 4$ (D) $k = 5$

MA

12 / 17

Question 57 (JAM 2020 • Q20 • ID: JAM_2020_Q20)

MCQ

+2 M

Neg: -0.67

Q. 20 Let M be an $n \times n$ ($n \geq 2$) non-zero real matrix with $M^2 = 0$ and let $\alpha \in \mathbb{R} \setminus \{0\}$. Then

(A) α is the only eigenvalue of $(M + \alpha I)$ and $(M - \alpha I)$ (B) α is the only eigenvalue of $(M + \alpha I)$ and $(\alpha I - M)$ (C) $-\alpha$ is the only eigenvalue of $(M + \alpha I)$ and $(M - \alpha I)$ (D) $-\alpha$ is the only eigenvalue of $(M + \alpha I)$ and $(\alpha I - M)$

MA

7 / 17

Question 58 (JAM 2020 • Q16 • ID: JAM_2020_Q16)

MCQ

+2 M

Neg: -0.67

JAM 2020

MATHEMATICS - MA

Q. 16 Let S be the surface of the portion of the sphere with centre at the origin and radius 4, above the xy -plane. Let $\vec{F} = y\hat{i} - x\hat{j} + yx^3\hat{k}$. If $\hat{n}$ is the unit outward normal to S , then

$$\iint_S (\nabla \times \vec{F}) \cdot \hat{n} \, dS$$

equals

(A) -32π (B) -16π (C) 16π (D) 32π

Question 59 (JAM 2020 • Q5 • ID: JAM_2020_Q5)

MCQ

+1 M

Neg: -0.33

JAM 2020

MATHEMATICS - MA

Q. 5 If the equation of the tangent plane to the surface $z = 16 - x^2 - y^2$ at the point $P(1, 3, 6)$ is $ax + by + cz + d = 0$, then the value of $|d|$ is

(A) 16

(B) 26

(C) 36

(D) 46

Question 60 (JAM 2019 • Q58 • ID: JAM_2019_Q58)

NAT

+2 M

Neg: Nil (0)

Q.58 Let M be a 3×3 matrix with real entries such that $M^2 = M + 2I$, where I denotes the 3×3 identity matrix. If α, β and γ are eigenvalues of M such that $\alpha\beta\gamma = -4$, then $\alpha + \beta + \gamma$ is equal to _____

Question 61 (JAM 2019 • Q57 • ID: JAM_2019_Q57)

NAT

+2 M

Neg: Nil (0)

Q.57 Let M and N be any two 4×4 matrices with integer entries satisfying

$$MN = 2 \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Then the maximum value of $\det(M) + \det(N)$ is _____

Question 62 (JAM 2019 • Q56 • ID: JAM_2019_Q56)

NAT

+2 M

Neg: Nil (0)

Q.56 If $\begin{pmatrix} 2 \\ y \\ z \end{pmatrix}$, $y, z \in \mathbb{R}$, is an eigenvector corresponding to a real eigenvalue of the matrix $\begin{pmatrix} 0 & 0 & 2 \\ 1 & 0 & -4 \\ 0 & 1 & 3 \end{pmatrix}$ then $z - y$ is equal to _____

Question 63 (JAM 2019 • Q43 • ID: JAM_2019_Q43)

NAT

+1 M

Neg: Nil (0)

Q.43 Consider the following system of three linear equations in four unknowns x_1, x_2, x_3 and x_4

$$\begin{aligned} x_1 + x_2 + x_3 + x_4 &= 4, \\ x_1 + 2x_2 + 3x_3 + 4x_4 &= 5, \\ x_1 + 3x_2 + 5x_3 + kx_4 &= 5. \end{aligned}$$

If the system has no solutions, then $k =$ _____

Question 64 (JAM 2019 • Q27 • ID: JAM_2019_Q27)

MCQ

+2 M

Neg: -0.67

Q.27 The set of eigenvalues of which one of the following matrices is NOT equal to the set of eigenvalues of $\begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix}$?

(A) $\begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$

(B) $\begin{pmatrix} 3 & 2 \\ 4 & 1 \end{pmatrix}$

(C) $\begin{pmatrix} 3 & 4 \\ 2 & 1 \end{pmatrix}$

(D) $\begin{pmatrix} 2 & 3 \\ 1 & 4 \end{pmatrix}$

Question 65 (JAM 2019 • Q16 • ID: JAM_2019_Q16)

MCQ

+2 M

Neg: -0.67

Q.16 Let $f(x) = (\ln x)^2$, $x > 0$. Then

- (A) $\lim_{x \rightarrow \infty} \frac{f(x)}{x}$ does not exist
 (B) $\lim_{x \rightarrow \infty} f'(x) = 2$
 (C) $\lim_{x \rightarrow \infty} (f(x+1) - f(x)) = 0$
 (D) $\lim_{x \rightarrow \infty} (f(x+1) - f(x))$ does not exist

Question 66 (JAM 2019 • Q5 • ID: JAM_2019_Q5)

MCQ

+1 M

Neg: -0.33

Q.5 Let S be the set of all limit points of the set $\left\{\frac{n}{\sqrt{2}} + \frac{\sqrt{2}}{n} : n \in \mathbb{N}\right\}$. Let $\mathbb{Q}_+$ be the set of all positive rational numbers. Then

- (A) $\mathbb{Q}_+ \subseteq S$ (B) $S \subseteq \mathbb{Q}_+$
 (C) $S \cap (\mathbb{R} \setminus \mathbb{Q}_+) \neq \emptyset$ (D) $S \cap \mathbb{Q}_+ \neq \emptyset$

MA

3/13

Question 67 (JAM 2018 • Q37 • ID: JAM_2018_Q37)

MSQ

+2 M

Neg: Nil (0)

Q.37 Let $m, n \in \mathbb{N}$, $m < n$, $P \in M_{n \times m}(\mathbb{R})$, $Q \in M_{m \times n}(\mathbb{R})$. Then which of the following is (are) NOT possible?

- (A) $\text{rank}(PQ) = n$
 (B) $\text{rank}(QP) = m$
 (C) $\text{rank}(PQ) = m$
 (D) $\text{rank}(QP) = \left\lceil \frac{m+n}{2} \right\rceil$, the smallest integer larger than or equal to $\frac{m+n}{2}$

Question 68 (JAM 2018 • Q36 • ID: JAM_2018_Q36)

MSQ

+2 M

Neg: Nil (0)

Q.36 Suppose $\alpha, \beta, \gamma \in \mathbb{R}$. Consider the following system of linear equations.

$x + y + z = \alpha$, $x + \beta y + z = \gamma$, $x + y + \alpha z = \beta$. If this system has at least one solution, then which of the following statements is (are) TRUE?

- (A) If $\alpha = 1$ then $\gamma = 1$ (B) If $\beta = 1$ then $\gamma = \alpha$
(C) If $\beta \neq 1$ then $\alpha = 1$ (D) If $\gamma = 1$ then $\alpha = 1$

Question 69 (JAM 2018 • Q27 • ID: JAM_2018_Q27)

MCQ

+2 M

Neg: -0.67

Q.27 Let I denote the 4×4 identity matrix. If the roots of the characteristic polynomial of a 4×4 matrix

M are $\pm \sqrt{\frac{1 \pm \sqrt{5}}{2}}$, then $M^8 =$

- (A) $I + M^2$ (B) $2I + M^2$ (C) $2I + 3M^2$ (D) $3I + 2M^2$

Question 70 (JAM 2018 • Q17 • ID: JAM_2018_Q17)

MCQ

+2 M

Neg: -0.67

Q.17 Let $f(x, y) = \begin{cases} \frac{xy}{(x^2+y^2)^\alpha}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$

Then which one of the following is TRUE for f at the point $(0, 0)$?

- (A) For $\alpha = 1$, f is continuous but not differentiable
(B) For $\alpha = \frac{1}{2}$, f is continuous and differentiable
(C) For $\alpha = \frac{1}{4}$, f is continuous and differentiable
(D) For $\alpha = \frac{3}{4}$, f is neither continuous nor differentiable

Question 71 (JAM 2018 • Q7 • ID: JAM_2018_Q7)

MCQ

+1 M

Neg: -0.33

Q.7 Let $f: \mathbb{R}^3 \rightarrow \mathbb{R}$ be a scalar field, $\vec{v}: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a vector field and let $\vec{a} \in \mathbb{R}^3$ be a constant vector.

If $\vec{r}$ represents the position vector $x\hat{i} + y\hat{j} + z\hat{k}$, then which one of the following is FALSE?

(A) $\text{curl}(f \vec{v}) = \text{grad}(f) \times \vec{v} + f \text{curl}(\vec{v})$

(B) $\text{div}(\text{grad}(f)) = \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) f$

(C) $\text{curl}(\vec{a} \times \vec{r}) = 2 |\vec{a}| \vec{r}$

(D) $\text{div}\left(\frac{\vec{r}}{|\vec{r}|^3}\right) = 0$, for $\vec{r} \neq \vec{0}$

Question 72 (JAM 2017 • Q58 • ID: JAM_2017_Q58)

NAT

+2 M

Neg: Nil (0)

Q.58 Let $\alpha, \beta, \gamma, \delta$ be the eigenvalues of the matrix

$$\begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 2 \end{bmatrix}$$

Then $\alpha^2 + \beta^2 + \gamma^2 + \delta^2 = \underline{\hspace{2cm}}$

Question 73 (JAM 2017 • Q48 • ID: JAM_2017_Q48)

NAT

+1 M

Neg: Nil (0)

Q.48 Let P be a 7×7 matrix of rank 4 with real entries. Let $\mathbf{a} \in \mathbb{R}^7$ be a column vector. Then the rank of $P + \mathbf{a}\mathbf{a}^T$ is at least $\underline{\hspace{2cm}}$

Question 74 (JAM 2017 • Q38 • ID: JAM_2017_Q38)

MSQ

+2 M

Neg: Nil (0)

Q.38 Let M be an $n \times n$ matrix with real entries such that $M^3 = I$. Suppose that $Mv \neq v$ for any non-zero vector v . Then which of the following statements is / are TRUE?

- (A) M has real eigenvalues
- (B) $M + M^{-1}$ has real eigenvalues
- (C) n is divisible by 2
- (D) n is divisible by 3

Question 75 (JAM 2017 • Q34 • ID: JAM_2017_Q34)

MSQ

+2 M

Neg: Nil (0)

Q.34 If X and Y are $n \times n$ matrices with real entries, then which of the following is/are TRUE?

- (A) If $P^{-1}XP$ is diagonal for some real invertible matrix P , then there exists a basis for $\mathbb{R}^n$ consisting of eigenvectors of X
- (B) If X is diagonal with distinct diagonal entries and $XY = YX$, then Y is also diagonal
- (C) If X^2 is diagonal, then X is diagonal
- (D) If X is diagonal and $XY = YX$ for all Y , then $X = \lambda I$ for some $\lambda \in \mathbb{R}$

Question 76 (JAM 2017 • Q12 • ID: JAM_2017_Q12)

MCQ

+2 M

Neg: -0.67

Q.12 Let $\mathcal{M}$ be the set of all invertible 5×5 matrices with entries 0 and 1. For each $M \in \mathcal{M}$, let $n_1(M)$ and $n_0(M)$ denote the number of 1's and 0's in M , respectively. Then

$$\min_{M \in \mathcal{M}} |n_1(M) - n_0(M)| =$$

- (A) 1
- (B) 3
- (C) 5
- (D) 15

Question 77 (JAM 2016 • Q53 • ID: JAM_2016_Q53)

NAT

+2 M

Neg: Nil (0)

Q.53 Let

$$M = \begin{bmatrix} \alpha & 1 & 1 \\ 1 & \beta & 1 \\ 1 & 1 & \gamma \end{bmatrix}, \quad \alpha\beta\gamma = 1, \quad \alpha, \beta, \gamma \in \mathbb{R} \quad \text{and} \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \in \mathbb{R}^3.$$

Then $M\mathbf{x} = \mathbf{0}$ has infinitely many solutions if $\text{trace}(M)$ is _____

Question 78 (JAM 2016 • Q43 • ID: JAM_2016_Q43)

NAT

+1 M

Neg: Nil (0)

Q.43 Let $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \in \mathbb{R}^3$ be a non-zero vector and $A = \frac{\mathbf{x}\mathbf{x}^T}{\mathbf{x}^T\mathbf{x}}$. Then the dimension of the vector space $\left\{ \mathbf{y} \in \mathbb{R}^3 \mid A\mathbf{y} = \mathbf{0} \right\}$ over $\mathbb{R}$ is _____

Question 79 (JAM 2016 • Q26 • ID: JAM_2016_Q26)

MCQ

+2 M

Neg: -0.67

Q.26 The matrix $M = \begin{bmatrix} \cos \alpha & \sin \alpha \\ i \sin \alpha & i \cos \alpha \end{bmatrix}$ is a unitary matrix when α is

(A) $(2n+1)\frac{\pi}{2}, n \in \mathbb{Z}$

(B) $(3n+1)\frac{\pi}{3}, n \in \mathbb{Z}$

(C) $(4n+1)\frac{\pi}{4}, n \in \mathbb{Z}$

(D) $(5n+1)\frac{\pi}{5}, n \in \mathbb{Z}$

Question 80 (JAM 2016 • Q16 • ID: JAM_2016_Q16)

MCQ

+2 M

Neg: -0.67

Q.16 The surface area of the portion of the plane $y + 2z = 2$ within the cylinder $x^2 + y^2 = 3$ is

(A) $\frac{3\sqrt{5}}{2}\pi$

(B) $\frac{5\sqrt{5}}{2}\pi$

(C) $\frac{7\sqrt{5}}{2}\pi$

(D) $\frac{9\sqrt{5}}{2}\pi$

Question 81 (JAM 2015 • Q60 • ID: JAM_2015_Q60)

NAT

+2 M

Neg: Nil (0)

Q.20 Let P and Q be two real matrices of size 4×6 and 5×4 , respectively. If $\text{rank}(Q) = 4$ and $\text{rank}(QP) = 2$, then $\text{rank}(P)$ is equal to _____

END OF THE QUESTION PAPER

MA

16/16

Question 82 (JAM 2015 • Q16 • ID: JAM_2015_Q16)

MCQ

+2 M

Neg: -0.67

Q.16

Let $A = \begin{bmatrix} 0 & 1-i \\ -1-i & i \end{bmatrix}$ and $B = A^T \bar{A}$. Then

- (A) an eigenvalue of B is purely imaginary
- (B) an eigenvalue of A is zero
- (C) all eigenvalues of B are real
- (D) A has a non-zero real eigenvalue

Question 83 (JAM 2015 • Q5 • ID: JAM_2015_Q5)

MCQ

+1 M

Neg: -0.33

Q.5 Let $\{x_n\}$ be a convergent sequence of real numbers. If $x_1 > \pi + \sqrt{2}$ and $x_{n+1} = \pi + \sqrt{x_n - \pi}$ for $n \geq 1$, then which one of the following is the limit of this sequence?

- (A) $\pi + 1$
- (B) $\pi + \sqrt{2}$
- (C) π
- (D) $\pi + \sqrt{\pi}$

Question 84 (JAM 2014 • Q12 • ID: JAM_2014_Q12)

MCQ

+2 M

Neg: -0.67

Q.12 Let $A = \begin{bmatrix} a & -1 & 4 \\ 0 & b & 7 \\ 0 & 0 & 3 \end{bmatrix}$ be a matrix with real entries. If the sum and the product of all the eigenvalues of A are 10 and 30 respectively, then $a^2 + b^2$ equals

- (A) 29
- (B) 40
- (C) 58
- (D) 65

Question 85 (JAM 2014 • Q11 • ID: JAM_2014_Q11)

MCQ

+2 M

Neg: -0.67

Q.11 The system of linear equations

$$x - y + 2z = b_1$$

$$x + 2y - z = b_2$$

$$2y - 2z = b_3$$

is inconsistent when (b_1, b_2, b_3) equals

- (A) $(2, 2, 0)$ (B) $(0, 3, 2)$ (C) $(2, 2, 1)$ (D) $(2, -1, -2)$

Question 86 (JAM 2012 • Q12 • ID: JAM_2012_Q12)

MCQ

+6 M

Neg: -2.0

Q.12 Let P be a 4×4 matrix whose determinant is 10. The determinant of the matrix $-3P$ is

- (A) -810 (B) -30 (C) 30 (D) 810

Question 87 (JAM 2011 • Q11 • ID: JAM_2011_Q11)

MCQ

+6 M

Neg: -2.0

Q.11 Let A be a 3×3 matrix with $\text{trace}(A) = 3$ and $\det(A) = 2$. If 1 is an eigenvalue of A , then the eigenvalues of the matrix $A^2 - 2I$ are

- (A) $1, 2(i-1), -2(i+1)$ (B) $-1, 2(i-1), 2(i+1)$
 (C) $1, 2(i+1), -2(i+1)$ (D) $-1, 2(i-1), -2(i+1)$

Question 88 (JAM 2008 • Q2 • ID: JAM_2008_Q2)

MCQ

+6 M

Neg: -2.0

Q.2 Let $S = \{T : \mathbb{R}^3 \rightarrow \mathbb{R}^3 \mid T \text{ is a linear transformation with } T(1,0,1) = (1,2,3), T(1,2,3) = (1,0,1)\}$. Then S is

- (A) a singleton set (B) a finite set containing more than one element
 (C) a countably infinite set (D) an uncountable set

Question 89 (JAM 2007 • Q11 • ID: JAM_2007_Q11)

MCQ

+6 M

Neg: -2.0

11. The rank of the matrix $\begin{bmatrix} 1 & 4 & 8 \\ 2 & 10 & 22 \\ 0 & 4 & 12 \end{bmatrix}$ is

- (A) 3
- (B) 2
- (C) 1
- (D) 0

Question 90 (JAM 2006 • Q11 • ID: JAM_2006_Q11)

MCQ

+6 M

Neg: -2.0

11. If A and B are 3×3 real matrices such that $\text{rank}(AB)=1$, then $\text{rank}(BA)$ **cannot** be

- (A) 0
- (B) 1
- (C) 2
- (D) 3

Question 91 (JAM 2005 • Q6 • ID: JAM_2005_Q6)

MCQ

+6 M

Neg: -2.0

6. Let A be a 3×3 matrix with eigenvalues 1, -1 and 3. Then

- (A) $A^2 + A$ is non-singular
- (B) $A^2 - A$ is non-singular
- (C) $A^2 + 3A$ is non-singular
- (D) $A^2 - 3A$ is non-singular

OFFICIAL MASTER ANSWER KEY & EVALUATION RUBRIC

Matrices and Systems of Linear Equations (2005–2026 Archive) • All Official Answers & Acceptance Ranges

Q. No.	Type	Marks	Official Key	Q. No.	Type	Marks	Official Key	Q. No.	Type	Marks	Official Key
1	NAT	+2	1.0 to 1.0	32	NAT	+2	3 to 3	63	NAT	+1	7 to 7
2	NAT	+2	6.0 to 6.0	33	NAT	+2	2 to 2	64	MCQ	+2	D
3	NAT	+2	−1.0 to −1.0	34	MCQ	+2	B	65	MCQ	+2	C
4	NAT	+1	16.0 to 16.0	35	MCQ	+2	B	66	MCQ	+1	B
5	NAT	+1	3.0 to 3.0	36	MCQ	+1	D	67	MSQ	+2	A;D
6	NAT	+1	8 to 8	37	NAT	+2	41 to 41	68	MSQ	+2	A;B
7	MSQ	+2	A;B;C	38	NAT	+2	3.00 to 3.00	69	MCQ	+2	C
8	MSQ	+2	A;B;D	39	MSQ	+2	A,C	70	MCQ	+2	C
9	MCQ	+2	B	40	MSQ	+2	A,B	71	MCQ	+1	C
10	MCQ	+2	B	41	MCQ	+2	C	72	NAT	+2	5.9 to 6.1
11	MCQ	+2	D	42	MCQ	+2	B	73	NAT	+1	2.9 to 3.1
12	MCQ	+1	B	43	MCQ	+2	C	74	MSQ	+2	B, C
13	NAT	+2	0 to 0	44	MCQ	+2	C	75	MSQ	+2	A, B, D
14	NAT	+2	10 to 10	45	MCQ	+2	B	76	MCQ	+2	A
15	NAT	+2	3 to 3	46	MCQ	+2	A	77	NAT	+2	3.0 to 3.0
16	MSQ	+2	B;D	47	MCQ	+1	C	78	NAT	+1	2.0 to 2.0
17	MCQ	+2	A	48	MCQ	+1	D	79	MCQ	+2	A
18	MCQ	+2	C	49	MCQ	+1	C	80	MCQ	+2	A
19	MCQ	+2	B	50	NAT	+2	4	81	NAT	+2	2
20	MCQ	+2	A	51	NAT	+2	2021	82	MCQ	+2	C
21	MCQ	+2	B	52	MSQ	+2	MTA	83	MCQ	+1	A
22	MCQ	+1	A	53	MCQ	+2	B	84	MCQ	+2	A
23	MCQ	+1	B	54	MCQ	+1	B	85	MCQ	+2	C
24	NAT	+1	31 to 31	55	NAT	+2	−216 to −216	86	MCQ	+6	D
25	NAT	+1	0.15 to 0.18	56	MSQ	+2	A;D	87	MCQ	+6	D
26	MSQ	+2	B;C	57	MCQ	+2	B	88	MCQ	+6	D
27	MCQ	+2	A	58	MCQ	+2	A	89	MCQ	+6	B
28	MCQ	+2	A	59	MCQ	+1	Marks to All	90	MCQ	+6	D
29	MCQ	+1	D	60	NAT	+2	3 to 3	91	MCQ	+6	C
30	MCQ	+1	D	61	NAT	+2	17 to 17				
31	MCQ	+1	A	62	NAT	+2	3 to 3				

SCORING METHODOLOGY & EVALUATION GUIDELINES

- **Multiple Choice Questions (MCQ):** Full marks for correct single choice. Negative marks deducted for each incorrect attempt (−1/3 for 1-mark questions, −2/3 for 2-mark questions). Unattempted questions award zero.

- **Multiple Select Questions (MSQ):** Full marks if and only if all correct choices are marked and no incorrect choice is selected. Zero marks for partial selection or any incorrect choice. No negative marking.
- **Numerical Answer Type (NAT):** Any numerical value falling strictly within the specified official range (e.g., $[a, b]$) receives full credit. No negative marking.
- **Marks to All (MTA):** Questions officially designated as MTA award full marks to all candidates regardless of attempt.