

IIT JAM 2025 Official Paper

Category: CBT Era (2015–2026) • Official Mock Test Series

TOTAL QUESTIONS

60

TOTAL MARKS

100 M

TEST DURATION

180 Min

PAPER PATTERN

MCQ • MSQ • NAT
(60 Ques)

Section / Type	Questions	Marking Scheme	Negative Marking	Format / Details
Multiple Choice (MCQ)	30	+1 / +2 Marks	−1/3 / −2/3 Marks	4 Options (1 Correct)
Multiple Select (MSQ)	10	+2 Marks	Nil (0)	4 Options (1+ Correct)
Numerical Answer (NAT)	20	+1 / +2 Marks	Nil (0)	Real Decimal Value

IMPORTANT INSTRUCTIONS FOR CANDIDATES

- Authentic Examination PYQs:** This mock paper contains verified official IIT JAM Mathematics questions and high-resolution problem statements.
- Section A (MCQ):** Each question has four choices (A), (B), (C), (D), out of which **only one** is correct. Wrong answers carry negative penalty (1/3 of positive marks).
- Section B (MSQ):** Each question has four choices (A), (B), (C), (D), out of which **one or more than one** choice(s) may be correct. Full marks are awarded only if all correct choices and zero incorrect choices are chosen. No partial marks and no negative marks.
- Section C (NAT):** Numerical answers are real numbers entered via virtual keyboard. Full credit is awarded if the numerical answer falls within the official accepted interval. No negative marking.
- Master Answer Key:** Complete official answers and acceptance bounds are provided at the end of this document.
- Online Interactive Simulator:** Practice this test with virtual calculator, timers, and instant scoring analytics at vaibhavgeometer.github.io.

IIT JAM Mathematics Mock Test Series • Curated by Vaibhav Geometer

Question 1 (JAM 2025 • Q1 • ID: JAM_2025_Q1)

MCQ

+1 M

Neg: -0.33

Q.1 The sum of the infinite series

$$\sum_{n=1}^{\infty} (-1)^{n+1} \frac{\pi^{2n+1}}{2^{2n+1}(2n)!}$$

is equal to

(A) $-\pi$

(B) $\frac{\pi}{4}$

(C) $\frac{\pi}{2}$

(D) $-\frac{\pi}{4}$

Question 2 (JAM 2025 • Q2 • ID: JAM_2025_Q2)

MCQ

+1 M

Neg: -0.33

Q.2 For which one of the following choices of $N(x, y)$, is the equation

$$(e^x \sin y - 2y \sin x) dx + N(x, y) dy = 0$$

an exact differential equation?

(A) $N(x, y) = e^x \sin y + 2 \cos x$

(B) $N(x, y) = e^x \cos y + 2 \cos x$

(C) $N(x, y) = e^x \cos y + 2 \sin x$

(D) $N(x, y) = e^x \sin y + 2 \sin x$

Question 3 (JAM 2025 • Q3 • ID: JAM_2025_Q3)

MCQ

+1 M

Neg: -0.33

Q.3 Let $f, g: \mathbb{R} \rightarrow \mathbb{R}$ be two functions defined by

$$f(x) = \begin{cases} x|x| \left| \sin \frac{1}{x} \right| & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

and

$$g(x) = \begin{cases} x^2 \sin \frac{1}{x} + x \cos \frac{1}{x} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0. \end{cases}$$

Then, which one of the following is TRUE?

- (A) f is differentiable at $x = 0$, and g is NOT differentiable at $x = 0$
- (B) f is NOT differentiable at $x = 0$, and g is differentiable at $x = 0$
- (C) f is differentiable at $x = 0$, and g is differentiable at $x = 0$
- (D) f is NOT differentiable at $x = 0$, and g is NOT differentiable at $x = 0$

Question 4 (JAM 2025 • Q4 • ID: JAM_2025_Q4)

MCQ

+1 M

Neg: -0.33

Q.4 Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ be two functions defined by

$$f(x) = \begin{cases} |x|^{1/8} \left| \sin \frac{1}{x} \right| \cos x & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

and

$$g(x) = \begin{cases} e^x \cos \frac{1}{x} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}.$$

Then, which one of the following is TRUE?

- (A) f is continuous at $x = 0$, and g is NOT continuous at $x = 0$
- (B) f is NOT continuous at $x = 0$, and g is continuous at $x = 0$
- (C) f is continuous at $x = 0$, and g is continuous at $x = 0$
- (D) f is NOT continuous at $x = 0$, and g is NOT continuous at $x = 0$

Question 5 (JAM 2025 • Q5 • ID: JAM_2025_Q5)

MCQ

+1 M

Neg: -0.33

Q.5 Which one of the following is the general solution of the differential equation

$$\frac{d^2y}{dx^2} - 8\frac{dy}{dx} + 16y = 2e^{4x} ?$$

- (A) $\alpha_1 e^{4x} + \alpha_2 x e^{4x} + x^2 e^{4x}$, where $\alpha_1, \alpha_2 \in \mathbb{R}$
- (B) $\alpha_1 e^{4x} + \alpha_2 x e^{4x} + 2x^2 e^{4x}$, where $\alpha_1, \alpha_2 \in \mathbb{R}$
- (C) $\alpha_1 e^{-4x} + \alpha_2 e^{4x} + 2x^2 e^{4x}$, where $\alpha_1, \alpha_2 \in \mathbb{R}$
- (D) $\alpha_1 x e^{-4x} + \alpha_2 x^2 e^{-4x} + x^2 e^{4x}$, where $\alpha_1, \alpha_2 \in \mathbb{R}$

Question 6 (JAM 2025 • Q6 • ID: JAM_2025_Q6)

MCQ

+1 M

Neg: -0.33

Q.6 Define $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ by

$$T(x, y, z) = (x + z, 2x + 3y + 5z, 2y + 2z), \text{ for all } (x, y, z) \in \mathbb{R}^3.$$

Then, which one of the following is TRUE?

- (A) T is one-one and T is NOT onto
- (B) T is NOT one-one and T is onto
- (C) T is one-one and T is onto
- (D) T is NOT one-one and T is NOT onto

Question 7 (JAM 2025 • Q7 • ID: JAM_2025_Q7)

MCQ

+1 M

Neg: -0.33

Q.7

Let $M = \begin{pmatrix} 2 & 0 & -1 \\ 4 & 1 & -4 \\ 2 & 0 & x \end{pmatrix}$ for some real number x .

If 0 is an eigenvalue of M , then $(M^4 + M) \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$ is equal to

(A) $\begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$

(B) $\begin{pmatrix} 2 \\ 0 \\ 2 \end{pmatrix}$

(C) $\begin{pmatrix} 5 \\ 0 \\ 5 \end{pmatrix}$

(D) $\begin{pmatrix} 17 \\ 0 \\ 17 \end{pmatrix}$

Question 8 (JAM 2025 • Q8 • ID: JAM_2025_Q8)

MCQ

+1 M

Neg: -0.33

Q.8 Let $T : P_2(\mathbb{R}) \rightarrow P_2(\mathbb{R})$ be the linear transformation defined by $T(p(x)) = p(x+1)$, for all $p(x) \in P_2(\mathbb{R})$. If M is the matrix representation of T with respect to the ordered basis $\{1, x, x^2\}$ of $P_2(\mathbb{R})$, then which one of the following is TRUE?

- (A) The determinant of M is 2
- (B) The rank of M is 2
- (C) 1 is the only eigenvalue of M
- (D) The nullity of M is 2

Question 9 (JAM 2025 • Q9 • ID: JAM_2025_Q9)

MCQ

+1 M

Neg: -0.33

Q.9 Let G be a finite abelian group of order 10. Let x_0 be an element of order 2 in G . If $X = \{x \in G : x^3 = x_0\}$, then which one of the following is TRUE?

- (A) X has exactly one element
- (B) X has exactly two elements
- (C) X has exactly three elements
- (D) X is an empty set

Question 10 (JAM 2025 • Q10 • ID: JAM_2025_Q10)

MCQ

+1 M

Neg: -0.33

Q.10 The value of

$$\int_0^1 \left(\int_{\sqrt{y}}^1 3e^{x^3} dx \right) dy$$

is equal to

(A) $e - 1$

(B) $\frac{e-1}{2}$

(C) $\sqrt{e} - 1$

(D) $\frac{\sqrt{e}-1}{2}$

Question 11 (JAM 2025 • Q11 • ID: JAM_2025_Q11)

MCQ

+2 M

Neg: -0.67

Q.11 Let $\mathcal{C}$ denote the family of curves described by $yx^2 = \lambda$, for $\lambda \in (0, \infty)$ and lying in the first quadrant of the xy plane. Let $\mathcal{O}$ denote the family of orthogonal trajectories of $\mathcal{C}$.

Which one of the following curves is a member of $\mathcal{O}$, and passes through the point $(2, 1)$?

- (A) $y = \frac{x^2}{4}, x > 0, y > 0$
- (B) $x^2 - 2y^2 = 2, x > 0, y > 0$
- (C) $x - y = 1, x > 0, y > 0$
- (D) $2x - y^2 = 3, x > 0, y > 0$

Question 12 (JAM 2025 • Q12 • ID: JAM_2025_Q12)

MCQ

+2 M

Neg: -0.67

Q.12 Let $\varphi : (0, \infty) \rightarrow \mathbb{R}$ be the solution of the differential equation

$$x \frac{dy}{dx} = (\ln y - \ln x)y,$$

satisfying $\varphi(1) = e^2$. Then, the value of $\varphi(2)$ is equal to

- (A) e^2
- (B) $2e^3$
- (C) $3e^2$
- (D) $6e^3$

Question 13 (JAM 2025 • Q13 • ID: JAM_2025_Q13)

MCQ

+2 M

Neg: -0.67

Q.13 Let $X = \{x \in S_4 : x^3 = id\}$ and $Y = \{x \in S_4 : x^2 \neq id\}$.

If m and n denote the number of elements in X and Y , respectively, then which one of the following is TRUE?

- (A) m is even and n is even
- (B) m is odd and n is even
- (C) m is even and n is odd
- (D) m is odd and n is odd

Question 14 (JAM 2025 • Q14 • ID: JAM_2025_Q14)

MCQ

+2 M

Neg: -0.67

Q.14 Let $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ be the solution of the differential equation

$$\frac{dy}{dx} = (y - 1)(y - 3),$$

satisfying $\varphi(0) = 2$. Then, which one of the following is TRUE?

- (A) $\lim_{x \rightarrow \infty} \varphi(x) = 0$
- (B) $\lim_{x \rightarrow \ln \sqrt{2}} \varphi(x) = 1$
- (C) $\lim_{x \rightarrow -\infty} \varphi(x) = 3$
- (D) $\lim_{x \rightarrow \ln \frac{1}{\sqrt{2}}} \varphi(x) = 6$

Question 15 (JAM 2025 • Q15 • ID: JAM_2025_Q15)

MCQ

+2 M

Neg: -0.67

Q.15

Let $M = \begin{pmatrix} 6 & 2 & -6 & 8 \\ 5 & 3 & -9 & 8 \\ 3 & 1 & -2 & 4 \end{pmatrix}$. Consider the system S of linear equations given by

$$\begin{aligned} 6x_1 + 2x_2 - 6x_3 + 8x_4 &= 8 \\ 5x_1 + 3x_2 - 9x_3 + 8x_4 &= 16 \\ 3x_1 + x_2 - 2x_3 + 4x_4 &= 32 \end{aligned}$$

where x_1, x_2, x_3, x_4 are unknowns.

Then, which one of the following is TRUE?

- (A) The rank of M is 3, and the system S has a solution
- (B) The rank of M is 3, and the system S does NOT have a solution
- (C) The rank of M is 2, and the system S has a solution
- (D) The rank of M is 2, and the system S does NOT have a solution

Question 16 (JAM 2025 • Q16 • ID: JAM_2025_Q16)

MCQ

+2 M

Neg: -0.67

Q.16

Let $M = \begin{pmatrix} -2 & 0 & 0 \\ 3 & 2 & 3 \\ 4 & -1 & x \end{pmatrix}$ for some real number x . Suppose that -2 and 3 are eigenvalues of M . If $M^3 \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 125 \\ 125 \end{pmatrix}$, then which one of the following is TRUE?

- (A) $x = 5$, and the matrix $M^2 + M$ is invertible
- (B) $x \neq 5$, and the matrix $M^2 + M$ is invertible
- (C) $x = 5$, and the matrix $M^2 + M$ is NOT invertible
- (D) $x \neq 5$, and the matrix $M^2 + M$ is NOT invertible

Question 17 (JAM 2025 • Q17 • ID: JAM_2025_Q17)

MCQ

+2 M

Neg: -0.67

Q.17

Let $f(x) = 10x^2 + e^x - \sin(2x) - \cos x$, $x \in \mathbb{R}$. The number of points at which the function f has a local minimum is

- (A) 0
- (B) 1
- (C) 2
- (D) greater than or equal to 3

Question 18 (JAM 2025 • Q18 • ID: JAM_2025_Q18)

MCQ

+2 M

Neg: -0.67

Q.18 For $n \in \mathbb{N}$, define x_n and y_n by

$$x_n = (-1)^n \cos \frac{1}{n} \quad \text{and} \quad y_n = \sum_{k=1}^n \frac{1}{n+k}.$$

Then, which one of the following is TRUE?

- (A) $\sum_{n=1}^{\infty} x_n$ converges, and $\sum_{n=1}^{\infty} y_n$ does NOT converge
- (B) $\sum_{n=1}^{\infty} x_n$ does NOT converge, and $\sum_{n=1}^{\infty} y_n$ converges
- (C) $\sum_{n=1}^{\infty} x_n$ converges, and $\sum_{n=1}^{\infty} y_n$ converges
- (D) $\sum_{n=1}^{\infty} x_n$ does NOT converge, and $\sum_{n=1}^{\infty} y_n$ does NOT converge

Question 19 (JAM 2025 • Q19 • ID: JAM_2025_Q19)

MCQ

+2 M

Neg: -0.67

Q.19 Let $x_1 = \frac{5}{2}$. For $n \in \mathbb{N}$, define

$$x_{n+1} = \frac{1}{5}(x_n^2 + 6).$$

Then, which one of the following is TRUE?

- (A) (x_n) is an increasing sequence, and (x_n) is NOT a bounded sequence
- (B) (x_n) is NOT an increasing sequence, and (x_n) is NOT a bounded sequence
- (C) (x_n) is NOT a decreasing sequence, and (x_n) is a bounded sequence
- (D) (x_n) is a decreasing sequence, and (x_n) is a bounded sequence

Question 20 (JAM 2025 • Q20 • ID: JAM_2025_Q20)

MCQ

+2 M

Neg: -0.67

Q.20

Let $x_1 = 2$ and $x_{n+1} = 2 + \frac{1}{2x_n}$ for all $n \in \mathbb{N}$.

Then, which one of the following is TRUE?

- (A) $x_{n+1} \geq \frac{4}{x_n}$ for all $n \in \mathbb{N}$, and (x_n) is a Cauchy sequence
- (B) $x_{n+1} < \frac{4}{x_n}$ for some $n \in \mathbb{N}$, and (x_n) is a Cauchy sequence
- (C) $x_{n+1} \geq \frac{4}{x_n}$ for all $n \in \mathbb{N}$, and (x_n) is NOT a Cauchy sequence
- (D) $x_{n+1} < \frac{4}{x_n}$ for some $n \in \mathbb{N}$, and (x_n) is NOT a Cauchy sequence

Question 21 (JAM 2025 • Q21 • ID: JAM_2025_Q21)

MCQ

+2 M

Neg: -0.67

Q.21 For $n \in \mathbb{N}$, define x_n and y_n by

$$x_n = (-1)^n \frac{3^n}{n^3} \text{ and } y_n = (4^n + (-1)^n 3^n)^{1/n}.$$

Then, which one of the following is TRUE?

- (A) (x_n) has a convergent subsequence, and NO subsequence of (y_n) is convergent
- (B) NO subsequence of (x_n) is convergent, and (y_n) has a convergent subsequence
- (C) (x_n) has a convergent subsequence, and (y_n) has a convergent subsequence
- (D) NO subsequence of (x_n) is convergent, and NO subsequence of (y_n) is convergent

Question 22 (JAM 2025 • Q22 • ID: JAM_2025_Q22)

MCQ

+2 M

Neg: -0.67

Q.22 Let $M = (m_{ij})$ be a 3×3 real, invertible matrix and $\sigma \in S_3$ be the permutation defined by $\sigma(1) = 2$, $\sigma(2) = 3$ and $\sigma(3) = 1$. The matrix $M_\sigma = (n_{ij})$ is defined by $n_{ij} = m_{i\sigma(j)}$ for all $i, j \in \{1, 2, 3\}$.

Then, which one of the following is TRUE?

- (A) $\det(M) = \det(M_\sigma)$, and nullity of the matrix $M - M_\sigma$ is 0
- (B) $\det(M) = -\det(M_\sigma)$, and nullity of the matrix $M - M_\sigma$ is 1
- (C) $\det(M) = \det(M_\sigma)$, and nullity of the matrix $M - M_\sigma$ is 1
- (D) $\det(M) = -\det(M_\sigma)$, and nullity of the matrix $M - M_\sigma$ is 0

Question 23 (JAM 2025 • Q23 • ID: JAM_2025_Q23)

MCQ

+2 M

Neg: -0.67

Q.23 Let $\mathbb{R}/\mathbb{Z}$ denote the quotient group, where $\mathbb{Z}$ is considered as a subgroup of the additive group of real numbers $\mathbb{R}$.

Let m denote the number of injective (one-one) group homomorphisms from $\mathbb{Z}_3$ to $\mathbb{R}/\mathbb{Z}$ and n denote the number of group homomorphisms from $\mathbb{R}/\mathbb{Z}$ to $\mathbb{Z}_3$.

Then, which one of the following is TRUE?

(A) $m = 2$ and $n = 1$

(B) $m = 3$ and $n = 3$

(C) $m = 2$ and $n = 3$

(D) $m = 1$ and $n = 1$

Question 24 (JAM 2025 • Q24 • ID: JAM_2025_Q24)

MCQ

+2 M

Neg: -0.67

Q.24 Let f_1, f_2, f_3 be nonzero linear transformations from $\mathbb{R}^4$ to $\mathbb{R}$ and

$$\ker(f_1) \subseteq \ker(f_2) \cap \ker(f_3).$$

Let $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be the linear transformation defined by

$$T(v) = (f_1(v), f_2(v), f_3(v)), \quad \text{for all } v \in \mathbb{R}^4.$$

Then, the nullity of T is equal to

(A) 1

(B) 2

(C) 3

(D) 4

Question 25 (JAM 2025 • Q25 • ID: JAM_2025_Q25)

MCQ

+2 M

Neg: -0.67

Q.25 Let $x_1 = 1$. For $n \in \mathbb{N}$, define

$$x_{n+1} = \left(\frac{1}{2} + \frac{\sin^2 n}{n} \right) x_n.$$

Then, which one of the following is TRUE?

- (A) $\sum_{n=1}^{\infty} x_n$ converges
- (B) $\sum_{n=1}^{\infty} x_n$ does NOT converge
- (C) $\sum_{n=1}^{\infty} x_n^2$ does NOT converge
- (D) $\sum_{n=1}^{\infty} x_n x_{n+1}$ does NOT converge

Question 26 (JAM 2025 • Q26 • ID: JAM_2025_Q26)

MCQ

+2 M

Neg: -0.67

Q.26 Let $x_1 > 0$. For $n \in \mathbb{N}$, define $x_{n+1} = x_n + 4$. If

$$\lim_{n \rightarrow \infty} \left(\frac{1}{x_2 x_3} + \frac{1}{x_3 x_4} + \cdots + \frac{1}{x_{n+1} x_{n+2}} \right) = \frac{1}{24},$$

then the value of x_1 is equal to

- (A) 1
- (B) 2
- (C) 3
- (D) 8

Question 27 (JAM 2025 • Q27 • ID: JAM_2025_Q27)

MCQ

+2 M

Neg: -0.67

Q.27 Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by

$$f(x, y) = e^y(x^2 + y^2) \quad \text{for all } (x, y) \in \mathbb{R}^2.$$

Then, which one of the following is TRUE?

- (A) The number of points at which f has a local minimum is 2
- (B) The number of points at which f has a local maximum is 2
- (C) The number of points at which f has a local minimum is 1
- (D) The number of points at which f has a local maximum is 1

Question 28 (JAM 2025 • Q28 • ID: JAM_2025_Q28)

MCQ

+2 M

Neg: -0.67

Q.28 Let Ω be the bounded region in $\mathbb{R}^3$ lying in the first octant ($x \geq 0$, $y \geq 0$, $z \geq 0$), and bounded by the surfaces $z = x^2 + y^2$, $z = 4$, $x = 0$ and $y = 0$.

Then, the volume of Ω is equal to

- (A) π
- (B) 2π
- (C) 3π
- (D) 4π

Question 29 (JAM 2025 • Q29 • ID: JAM_2025_Q29)

MCQ

+2 M

Neg: -0.67

Q.29 Let $\varphi : [0, \infty) \rightarrow \mathbb{R}$ be the continuous function satisfying

$$\varphi(x) = \left(\int_0^x \varphi(t) dt \right) + \sin x, \text{ for all } x \in [0, \infty).$$

Then, the value of $\lim_{x \rightarrow \pi/2} (2\varphi(x) - e^x)$ is equal to

(A) 1

(B) 2

(C) 3

(D) 4

Question 30 (JAM 2025 • Q30 • ID: JAM_2025_Q30)

MCQ

+2 M

Neg: -0.67

Q.30 The number of elements in the set

$$\{x \in \mathbb{R} : 8x^2 + x^4 + x^8 = \cos x\}$$

is equal to

(A) 0

(B) 1

(C) 2

(D) greater than or equal to 3

Question 31 (JAM 2025 • Q31 • ID: JAM_2025_Q31)

MSQ

+2 M

Neg: Nil (0)

Q.31 Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by

$$f(x, y) = \begin{cases} \frac{xy^2 + y^5}{x^2 + y^4} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$

Then, which of the following is/are TRUE?

- (A) The iterated limits $\lim_{x \rightarrow 0} \left(\lim_{y \rightarrow 0} f(x, y) \right)$ and $\lim_{y \rightarrow 0} \left(\lim_{x \rightarrow 0} f(x, y) \right)$ exist
- (B) Exactly one of the partial derivatives $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ exists at $(0, 0)$
- (C) Both the partial derivatives $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ exist at $(0, 0)$
- (D) f is NOT differentiable at $(0, 0)$

Question 32 (JAM 2025 • Q32 • ID: JAM_2025_Q32)

MSQ

+2 M

Neg: Nil (0)

Q.32 If $M, N, \mu, w: \mathbb{R}^2 \rightarrow \mathbb{R}$ are differentiable functions with continuous partial derivatives, satisfying

$$\mu(x, y)M(x, y)dx + \mu(x, y)N(x, y)dy = dw$$

then which of the following is/are TRUE?

- (A) μw is an integrating factor for $M(x, y)dx + N(x, y)dy = 0$
- (B) μw^2 is an integrating factor for $M(x, y)dx + N(x, y)dy = 0$
- (C) $w(x, y) = w(0, 0) + \int_0^x (\mu M)(s, 0)ds + \int_0^y (\mu N)(x, t)dt$, for all $(x, y) \in \mathbb{R}^2$
- (D) $w(x, y) = w(0, 0) + \int_0^x (\mu M)(s, y)ds + \int_0^y (\mu N)(0, t)dt$, for all $(x, y) \in \mathbb{R}^2$

Question 33 (JAM 2025 • Q33 • ID: JAM_2025_Q33)

MSQ

+2 M

Neg: Nil (0)

Q.33 Let $\varphi : (-1, \infty) \rightarrow (0, \infty)$ be the solution of the differential equation

$$\frac{dy}{dx} - 2ye^x = 2e^x\sqrt{y},$$

satisfying $\varphi(0) = 1$.

Then, which of the following is/are TRUE?

(A) φ is an unbounded function

(B) $\lim_{x \rightarrow \ln 2} \varphi(x) = (2e - 1)^2$

(C) $\lim_{x \rightarrow \ln 2} \varphi(x) = \sqrt{2e - 1}$

(D) φ is a strictly increasing function on the interval $(0, \infty)$

Question 34 (JAM 2025 • Q34 • ID: JAM_2025_Q34)

MSQ

+2 M

Neg: Nil (0)

Q.34 Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by

$$f(x, y) = \begin{cases} \frac{(x^2 + \sin x)y^2}{x^2 + y^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$

Then, which of the following is/are TRUE?

- (A) $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$ exists and $\lim_{(x,y) \rightarrow (0,0)} f(x, y) = 1$
- (B) $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$ exists and $\lim_{(x,y) \rightarrow (0,0)} f(x, y) = 0$
- (C) f is differentiable at $(0, 0)$
- (D) f is NOT differentiable at $(0, 0)$

Question 35 (JAM 2025 • Q35 • ID: JAM_2025_Q35)

MSQ

+2 M

Neg: Nil (0)

Q.35 Let

$$u_1 = (1, 0, 0, -1), u_2 = (0, 2, 0, -1), u_3 = (0, 0, 1, -1) \text{ and } u_4 = (0, 0, 0, 1)$$

be elements in the real vector space $\mathbb{R}^4$.

Then, which of the following is/are TRUE?

- (A) $\{u_1, u_2, u_3, u_4\}$ is a linearly independent set in $\mathbb{R}^4$
- (B) $\{u_1 - u_2, u_2 - u_3, u_3 - u_4, u_4 - u_1\}$ is NOT a linearly independent set in $\mathbb{R}^4$
- (C) $\{u_1, -u_2, u_3, -u_4\}$ is NOT a linearly independent set in $\mathbb{R}^4$
- (D) $\{u_1 + u_2, u_2 + u_3, u_3 + u_4, u_4 + u_1\}$ is a linearly independent set in $\mathbb{R}^4$

Question 36 (JAM 2025 • Q36 • ID: JAM_2025_Q36)

MSQ

+2 M

Neg: Nil (0)

Q.36 For $n \in \mathbb{N}$, let

$$x_n = \sum_{k=1}^n \frac{k}{n^2 + k}.$$

Then, which of the following is/are TRUE?

- (A) The sequence (x_n) converges
- (B) The series $\sum_{n=1}^{\infty} x_n$ converges
- (C) The series $\sum_{n=1}^{\infty} x_n$ does NOT converge
- (D) The series $\sum_{n=1}^{\infty} x_n^n$ converges

Question 37 (JAM 2025 • Q37 • ID: JAM_2025_Q37)

MSQ

+2 M

Neg: Nil (0)

Q.37 Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a twice differentiable function such that

$$f(0) = 0, f'(0) = 2 \text{ and } f(1) = -3.$$

Then, which of the following is/are TRUE?

- (A) $|f'(x)| \leq 2$ for all $x \in [0, 1]$
- (B) $|f'(x_1)| > 2$ for some $x_1 \in [0, 1]$
- (C) $|f''(x)| < 10$ for all $x \in [0, 1]$
- (D) $|f''(x_2)| \geq 10$ for some $x_2 \in [0, 1]$

Question 38 (JAM 2025 • Q38 • ID: JAM_2025_Q38)

MSQ

+2 M

Neg: Nil (0)

Q.38 Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a twice differentiable function such that

$$f(0) = 4, f(1) = -2, f(2) = 8 \text{ and } f(3) = 2.$$

Then, which of the following is/are TRUE?

- (A) $|f'(x)| < 5$ for all $x \in [0, 1]$
- (B) $|f'(x_1)| \geq 5$ for some $x_1 \in [0, 1]$
- (C) $f'(x_2) = 0$ for some $x_2 \in [0, 3]$
- (D) $f''(x_3) = 0$ for some $x_3 \in [0, 3]$

Question 39 (JAM 2025 • Q39 • ID: JAM_2025_Q39)

MSQ

+2 M

Neg: Nil (0)

Q.39 For $n \in \mathbb{N}$, consider the set $U(n) = \{\bar{x} \in \mathbb{Z}_n : \gcd(x, n) = 1\}$ as a group under multiplication modulo n .

Then, which of the following is/are TRUE?

- (A) $U(8)$ is a cyclic group
- (B) $U(5)$ is a cyclic group
- (C) $U(12)$ is a cyclic group
- (D) $U(9)$ is a cyclic group

Question 40 (JAM 2025 • Q40 • ID: JAM_2025_Q40)

MSQ

+2 M

Neg: Nil (0)

Q.40 Consider the following subspaces of the real vector space $\mathbb{R}^3$:

$$V_1 = \text{span} \{(1, 2, 3), (1, 1, 0)\},$$

$$V_2 = \text{span} \{(1, -1, 0)\},$$

$$V_3 = \text{span} \{(1, 1, 1)\},$$

$$V_4 = \text{span} \{(1, 3, 6)\} \text{ and}$$

$$V_5 = \text{span} \{(1, 0, -3)\}.$$

Then, which of the following is/are TRUE?

(A) $V_1 \cup V_2$ is a subspace of $\mathbb{R}^3$

(B) $V_1 \cup V_3$ is a subspace of $\mathbb{R}^3$

(C) $V_1 \cup V_4$ is a subspace of $\mathbb{R}^3$

(D) $V_1 \cup V_5$ is a subspace of $\mathbb{R}^3$

Question 41 (JAM 2025 • Q41 • ID: JAM_2025_Q41)

NAT

+1 M

Neg: Nil (0)

Q.41 The radius of convergence of the power series

$$\sum_{n=1}^{\infty} \frac{\left(x + \frac{1}{4}\right)^n}{(-2)^n n^2}$$

about $x = -\frac{1}{4}$, is equal to _____ (rounded off to two decimal places).

Question 42 (JAM 2025 • Q42 • ID: JAM_2025_Q42)

NAT

+1 M

Neg: Nil (0)

Q.42 The value of

$$\lim_{n \rightarrow \infty} 8n \left(e^{\left(\frac{1}{2n}\right)} - 1 \right) \left(\sin \frac{1}{2n} + \left| \cos \frac{1}{2n} \right| \right)$$

is equal to _____ (rounded off to two decimal places).

Question 43 (JAM 2025 • Q43 • ID: JAM_2025_Q43)

NAT

+1 M

Neg: Nil (0)

Q.43 Let α be the real number such that

$$\lim_{x \rightarrow 0} \frac{(1 - \cos x)(2^{2+x} - 4)}{x^3} = \alpha \ln 2.$$

Then, the value of α is equal to _____ (rounded off to two decimal places).

Question 44 (JAM 2025 • Q44 • ID: JAM_2025_Q44)

NAT

+1 M

Neg: Nil (0)

Q.44 Let $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ be the solution of the differential equation

$$4 \frac{d^2 y}{dx^2} + 16 \frac{dy}{dx} + 25y = 0$$

satisfying $\varphi(0) = 1$ and $\varphi'(0) = -\frac{1}{2}$.Then, the value of $\lim_{x \rightarrow \pi/6} e^{2x} \varphi(x)$ is equal to _____ (rounded off to two decimal places).

Question 45 (JAM 2025 • Q45 • ID: JAM_2025_Q45)

NAT

+1 M

Neg: Nil (0)

Q.45 Let S be the surface area of the portion of the plane $z = x + y + 3$, which lies inside the cylinder $x^2 + y^2 = 1$.

Then, the value of $\left(\frac{S}{\pi}\right)^2$ is equal to _____ (rounded off to two decimal places).

Question 46 (JAM 2025 • Q46 • ID: JAM_2025_Q46)

NAT

+1 M

Neg: Nil (0)

Q.46 Consider the following subspaces of $\mathbb{R}^4$:

$$V_1 = \{(x, y, z, w) \in \mathbb{R}^4 : x + y + 2w = 0\},$$

$$V_2 = \{(x, y, z, w) \in \mathbb{R}^4 : 2y + z + w = 0\},$$

$$V_3 = \{(x, y, z, w) \in \mathbb{R}^4 : x + 3y + z + 3w = 0\}.$$

Then, the dimension of the subspace $V_1 \cap V_2 \cap V_3$ is equal to _____.

Question 47 (JAM 2025 • Q47 • ID: JAM_2025_Q47)

NAT

+1 M

Neg: Nil (0)

Q.47 Consider the real vector space $\mathbb{R}^3$. Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a linear transformation such that

$$T(1, 1, 1) = 0, \quad T(1, -1, 1) = 0 \quad \text{and} \quad T(0, 0, 1) = 16.$$

Then, the value of $T\left(\frac{1}{2}, \frac{2}{3}, \frac{3}{4}\right)$ is equal to _____ (rounded off to two decimal places).

Question 48 (JAM 2025 • Q48 • ID: JAM_2025_Q48)

NAT

+1 M

Neg: Nil (0)

Q.48 Let T denote the triangle in the xy plane bounded by the x axis and the lines $y = x$ and $x = 1$. The value of the double integral (over T)

$$\iint_T (5 - y) dx dy$$

is equal to _____ (rounded off to two decimal places).

Question 49 (JAM 2025 • Q49 • ID: JAM_2025_Q49)

NAT

+1 M

Neg: Nil (0)

Q.49 Let $T, S : P_4(\mathbb{R}) \rightarrow P_4(\mathbb{R})$ be the linear transformations defined by

$$T(p(x)) = xp'(x) \text{ and } S(p(x)) = (x+1)p'(x)$$

for all $p(x) \in P_4(\mathbb{R})$.

Then, the nullity of the composition $S \circ T$ is _____.

Question 50 (JAM 2025 • Q50 • ID: JAM_2025_Q50)

NAT

+1 M

Neg: Nil (0)

Q.50 Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by

$$f(x, y) = \begin{cases} \frac{(x^2 - y^2)xy}{x^2 + y^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$

Then, the value of $\frac{\partial f}{\partial y}(1, 0) - \frac{\partial f}{\partial x}(0, 2)$ is equal to _____ (rounded off to two decimal places).

Question 51 (JAM 2025 • Q51 • ID: JAM_2025_Q51)

NAT

+2 M

Neg: Nil (0)

Q.51 Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function satisfying

$$\int_0^{\pi/4} \left(\sin(x) f(x) + \cos(x) \int_0^x f(t) dt \right) dx = \sqrt{2}.$$

Then, the value of $\int_0^{\pi/4} f(x) dx$ is equal to _____ (rounded off to two decimal places).

Question 52 (JAM 2025 • Q52 • ID: JAM_2025_Q52)

NAT

+2 M

Neg: Nil (0)

Q.52 Let $\sigma \in S_4$ be the permutation defined by $\sigma(1) = 2$, $\sigma(2) = 3$, $\sigma(3) = 1$ and $\sigma(4) = 4$. The number of elements in the set

$$\{\tau \in S_4 : \tau\sigma\tau^{-1} = \sigma\}$$

is equal to _____.

Question 53 (JAM 2025 • Q53 • ID: JAM_2025_Q53)

NAT

+2 M

Neg: Nil (0)

Q.53 Let $f(x) = 2x - \sin x$, for all $x \in \mathbb{R}$. Let $k \in \mathbb{N}$ be such that

$$\lim_{x \rightarrow 0} \left(\frac{1}{x} \sum_{i=1}^k i^2 f\left(\frac{x}{i}\right) \right) = 45.$$

Then, the value of k is equal to _____.

Question 54 (JAM 2025 • Q54 • ID: JAM_2025_Q54)

NAT

+2 M

Neg: Nil (0)

Q.54 The value of the infinite series

$$\sum_{n=1}^{\infty} n \left(\frac{3}{4}\right)^{2(n-1)}$$

is equal to _____ (rounded off to two decimal places)

Question 55 (JAM 2025 • Q55 • ID: JAM_2025_Q55)

NAT

+2 M

Neg: Nil (0)

Q.55 Let $\varphi: (0, \infty) \rightarrow \mathbb{R}$ be the solution of the differential equation

$$x^2 \frac{d^2 y}{dx^2} - x \frac{dy}{dx} + y = 6x \ln x$$

satisfying $\varphi(1) = -3$ and $\varphi(e) = 0$.Then, the value of $|\varphi'(1)|$ is equal to _____ (rounded off to two decimal places).

Question 56 (JAM 2025 • Q56 • ID: JAM_2025_Q56)

NAT

+2 M

Neg: Nil (0)

Q.56 Let $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ be the solution of the differential equation

$$\frac{dy}{dx} + 2xy = 2 + 4x^2$$

satisfying $\varphi(0) = 0$.Then, the value of $\varphi(2)$ is equal to _____ (rounded off to two decimal places).

Question 57 (JAM 2025 • Q57 • ID: JAM_2025_Q57)

NAT

+2 M

Neg: Nil (0)

Q.57 Let Ω be the solid bounded by the planes $z = 0$, $y = 0$, $x = \frac{1}{2}$, $2y = x$ and $2x + y + z = 4$.

If V is the volume of Ω , then the value of $64V$ is equal to _____
(rounded off to two decimal places).

Question 58 (JAM 2025 • Q58 • ID: JAM_2025_Q58)

NAT

+2 M

Neg: Nil (0)

Q.58 Let the subspace H of $P_3(\mathbb{R})$ be defined as

$$H = \{p(x) \in P_3(\mathbb{R}) : xp'(x) = 3p(x)\}.$$

Then, the dimension of H is equal to _____.

Question 59 (JAM 2025 • Q59 • ID: JAM_2025_Q59)

NAT

+2 M

Neg: Nil (0)

Q.59 Let G be an abelian group of order 35. Let m denote the number of elements of order 5 in G , and let n denote the number of elements of order 7 in G .

Then, the value of $m + n$ is equal to _____.

Question 60 (JAM 2025 • Q60 • ID: JAM_2025_Q60)

NAT

+2 M

Neg: Nil (0)

Q.60 The number of surjective (onto) group homomorphisms from S_4 to $\mathbb{Z}_6$ is equal to _____.

OFFICIAL MASTER ANSWER KEY & EVALUATION RUBRIC

IIT JAM 2025 Official Paper • All Official Answers & Acceptance Ranges

Q. No.	Type	Marks	Official Key	Q. No.	Type	Marks	Official Key	Q. No.	Type	Marks	Official Key
1	MCQ	+1	C	21	MCQ	+2	B	41	NAT	+1	1.98 to 2.02
2	MCQ	+1	B	22	MCQ	+2	C	42	NAT	+1	3.98 to 4.02
3	MCQ	+1	A	23	MCQ	+2	A	43	NAT	+1	1.98 to 2.02
4	MCQ	+1	A	24	MCQ	+2	C	44	NAT	+1	1.39 to 1.43
5	MCQ	+1	A	25	MCQ	+2	A	45	NAT	+1	2.98 to 3.02
6	MCQ	+1	D	26	MCQ	+2	B	46	NAT	+1	2 to 2
7	MCQ	+1	B	27	MCQ	+2	C	47	NAT	+1	3.98 to 4.02
8	MCQ	+1	C	28	MCQ	+2	B	48	NAT	+1	2.31 to 2.35
9	MCQ	+1	A	29	MCQ	+2	A	49	NAT	+1	1 to 1
10	MCQ	+1	A	30	MCQ	+2	C	50	NAT	+1	2.98 to 3.02
11	MCQ	+2	B	31	MSQ	+2	A;C;D	51	NAT	+2	1.98 to 2.02
12	MCQ	+2	B	32	MSQ	+2	A;B;C	52	NAT	+2	3 to 3
13	MCQ	+2	B	33	MSQ	+2	A;B;D	53	NAT	+2	9 to 9
14	MCQ	+2	C	34	MSQ	+2	B;D	54	NAT	+2	5.20 to 5.24
15	MCQ	+2	A	35	MSQ	+2	A;B	55	NAT	+2	0.98 to 1.02
16	MCQ	+2	B	36	MSQ	+2	A;C;D	56	NAT	+2	3.98 to 4.02
17	MCQ	+2	B	37	MSQ	+2	B;D	57	NAT	+2	MTA
18	MCQ	+2	D	38	MSQ	+2	B;C;D	58	NAT	+2	1 to 1
19	MCQ	+2	D	39	MSQ	+2	B;D	59	NAT	+2	10 to 10
20	MCQ	+2	A	40	MSQ	+2	C;D	60	NAT	+2	0 to 0

SCORING METHODOLOGY & EVALUATION GUIDELINES

- **Multiple Choice Questions (MCQ):** Full marks for correct single choice. Negative marks deducted for each incorrect attempt ($-1/3$ for 1-mark questions, $-2/3$ for 2-mark questions). Unattempted questions award zero.
- **Multiple Select Questions (MSQ):** Full marks if and only if all correct choices are marked and no incorrect choice is selected. Zero marks for partial selection or any incorrect choice. No negative marking.
- **Numerical Answer Type (NAT):** Any numerical value falling strictly within the specified official range (e.g., $[a, b]$) receives full credit. No negative marking.
- **Marks to All (MTA):** Questions officially designated as MTA award full marks to all candidates regardless of attempt.