


PAST PAPER QUESTION FROM REVISION LECTURE

Q1 Consider the following three-joint manipulator geometry –



The waist and shoulder joints are revolute and the radial joint is prismatic. Joint 2 is located a distance d_1 above the robot origin, i.e. the TCP is at a height of d_1 in the above diagram. The TCP is located a distance d_2 from Joint 2 when the radial joint is fully retracted, i.e. when the arm extension is 0.

- (a) Derive the inverse kinematic equations for the three joints of this manipulator.
[Hint: A geometric solution can be used here] (9)
- (b) Hence derive the inverse Jacobian for the manipulator. (11)

SOLUTION

(a) Derive the inverse kinematic equations for the three joints of this manipulator.
[Hint: A geometric solution can be used here]

Let the joint angles be θ_1 for the waist rotation, θ_2 for the shoulder rotation and r for the radial extension

Viewing the arm from above we obtain the waist rotation –

$$\tan(\theta_1) = \frac{y}{x}$$

$$\therefore \theta_1 = \text{ATAN2}(y, x)$$

3 MARKS

Viewing the side elevation we can determine the shoulder rotation –

$$\tan(\theta_2) = \frac{z - d_1}{\sqrt{x^2 + y^2}}$$

$$\therefore \theta_2 = \text{ATAN2}\left(z - d_1, \sqrt{x^2 + y^2}\right)$$

3 MARKS

Also from the side elevation we can determine the radial extension –

$$r + d_2 = \sqrt{(z - d_1)^2 + x^2 + y^2}$$

$$\therefore r = \sqrt{(z - d_1)^2 + x^2 + y^2} - d_2$$

3 MARKS

b). Hence derive the inverse Jacobian for the manipulator.

We differentiate each of the inverse kinematics equations with respect to time –

$$\tan(\theta_1) = \frac{y}{x}$$

$$\therefore \sec^2(\theta_1) \frac{\partial \theta_1}{\partial t} = \frac{-y}{x^2} \frac{\partial x}{\partial t} + \frac{1}{x} \frac{\partial y}{\partial t}$$

Noting that $\sec^2(\theta_1) = \frac{(r + d_2)^2}{x^2}$ because $\sec(x) = 1/\cos(x)$

$$\therefore \dot{\theta}_1 = \frac{-y}{x^2 \left(\frac{(r + d_2)^2}{x^2} \right)} \dot{x} + \frac{1}{x \left(\frac{(r + d_2)^2}{x^2} \right)} \dot{y}$$

$$\therefore \dot{\theta}_1 = \frac{-y}{(r + d_2)^2} \dot{x} + \frac{x}{(r + d_2)^2} \dot{y}$$

3 MARKS

$$\tan(\theta_2) = \frac{z - d_1}{\sqrt{x^2 + y^2}}$$

$$\therefore \sec^2(\theta_2) \frac{\partial \theta_2}{\partial t} = (z - d_1) \frac{\partial \left(\frac{1}{\sqrt{x^2 + y^2}} \right)}{\partial t} + \frac{1}{\sqrt{x^2 + y^2}} \frac{\partial z}{\partial t}$$

$$\therefore \dot{\theta}_2 \sec^2(\theta_2) = \frac{1}{\sqrt{x^2 + y^2}} \dot{z} + (z - d_1) \left[\frac{-x\dot{x} - y\dot{y}}{(x^2 + y^2)^{3/2}} \right]$$

Noting that $\sec(\theta_2) = \frac{1}{\cos(\theta_2)} = \frac{1}{\frac{\sqrt{x^2 + y^2}}{r + d_2}} = \frac{r + d_2}{\sqrt{x^2 + y^2}} \quad \therefore \sec^2(\theta_2) = \frac{(r + d_2)^2}{x^2 + y^2}$

$$\therefore \dot{\theta}_2 = \frac{\sqrt{x^2 + y^2}}{(r + d_2)^2} \dot{z} + \frac{z - d_1}{(r + d_2)^2 \sqrt{x^2 + y^2}} [-x\dot{x} - y\dot{y}]$$

3 MARKS

$$(r + d_2)^2 = (z - d_1)^2 + x^2 + y^2$$

$$(2r + 2d_2) \frac{\partial r}{\partial t} = (2z - 2d_1) \frac{\partial z}{\partial t} + 2x \frac{\partial x}{\partial t} + 2y \frac{\partial y}{\partial t}$$

$$\therefore \dot{r} = \frac{z - d_1}{r + d_2} \dot{z} + \frac{x}{r + d_2} \dot{x} + \frac{y}{r + d_2} \dot{y}$$

3 MARKS

So the inverse Jacobian is therefore –

$$\begin{pmatrix} \dot{\theta}_1 \\ \dot{\theta}_2 \\ \dot{r} \end{pmatrix} = \begin{pmatrix} \frac{-y}{(r + d_2)^2} & \frac{x}{(r + d_2)^2} & 0 \\ \frac{-(z - d_1)x}{(r + d_2)^2 \sqrt{x^2 + y^2}} & \frac{-(z - d_1)y}{(r + d_2)^2 \sqrt{x^2 + y^2}} & \frac{\sqrt{x^2 + y^2}}{(r + d_2)^2} \\ \frac{x}{r + d_2} & \frac{y}{r + d_2} & \frac{z - d_1}{r + d_2} \end{pmatrix} \begin{pmatrix} \dot{x} \\ \dot{y} \\ \dot{z} \end{pmatrix}$$

2 MARKS