

Lecture 18 - Natural Deduction

Vaishnavi Sundararajan

COL703/COL7203 - Logic for Computer Science

Proof rules for propositional fragment

Introduction rule	Elimination rule
$\frac{\Gamma \vdash \varphi_0 \quad \Gamma \vdash \varphi_1}{\Gamma \vdash \varphi_0 \wedge \varphi_1} \wedge i$	$\frac{\Gamma \vdash \varphi_0 \wedge \varphi_1}{\Gamma \vdash \varphi_j} \wedge e_j$
$\frac{\Gamma \vdash \varphi_j}{\Gamma \vdash \varphi_0 \vee \varphi_1} \vee i_j$	$\frac{\Gamma \vdash \varphi_0 \vee \varphi_1 \quad \Gamma, \varphi_0 \vdash \psi \quad \Gamma, \varphi_1 \vdash \psi}{\Gamma \vdash \psi} \vee e$
$\frac{\Gamma, \varphi \vdash \psi}{\Gamma \vdash \varphi \supset \psi} \supset i$	$\frac{\Gamma \vdash \varphi \supset \psi \quad \Gamma \vdash \varphi}{\Gamma \vdash \psi} \supset e$
$\frac{\Gamma, \varphi \vdash \neg \psi \quad \Gamma, \varphi \vdash \psi}{\Gamma \vdash \neg \varphi} \neg i$	$\frac{\Gamma \vdash \neg \neg \varphi}{\Gamma \vdash \varphi} \neg e$

Proof rules for \exists and \forall

Introduction rule	Elimination rule
$\frac{\Gamma \vdash \varphi\{y/x\}}{\Gamma \vdash \forall x. [\varphi]} \forall i \text{ (} y \text{ fresh)}$	$\frac{\Gamma \vdash \forall x. [\varphi]}{\Gamma \vdash \varphi\{t/x\}} \forall e$
$\frac{\Gamma \vdash \varphi\{t/x\}}{\Gamma \vdash \exists x. [\varphi]} \exists i$	$\frac{\Gamma \vdash \exists x. [\varphi] \quad \Gamma, \varphi\{y/x\} \vdash \psi}{\Gamma \vdash \psi} \exists e \text{ (} y \text{ fresh)}$

where t is a term in the language, and $y \in \mathcal{V}$ is fresh if $y \notin \text{vars}(\Gamma \cup \{\varphi, \psi\})$.

$$\frac{}{\Gamma \vdash \varphi} Ax \quad (\varphi \in \Gamma)$$

We say that $\Gamma \vdash_{\mathcal{E}} \varphi$ if there is a proof of φ from assumptions Γ using Ax and the rules in both the above tables.

Example o

Show that $\forall x. [P(x)] \vdash_{\mathcal{G}} \exists x. [P(x)]$

$$\frac{\frac{\frac{}{\forall x. [P(x)] \vdash \forall x. [P(x)]} \text{Ax}}{\forall x. [P(x)] \vdash P(t)} \text{Ve}}{\forall x. [P(x)] \vdash \exists x. [P(x)]} \text{Ei}$$

Example 1

Let $\Gamma = \{\forall x. [L(x) \supset \neg U(x)], \forall x. [P(x) \supset \neg T(x)], L(a) \vee T(a)\}$. Show that $\Gamma \vdash_{\mathcal{G}} P(a) \supset \neg U(a)$.

Example 1

Let $\Gamma = \{\forall x. [L(x) \supset \neg U(x)], \forall x. [P(x) \supset \neg T(x)], L(a) \vee T(a)\}$. Show that $\Gamma \vdash_{\mathcal{L}} P(a) \supset \neg U(a)$. Let $\Gamma' = \Gamma \cup \{P(a)\}$.

$$\begin{array}{c}
 \frac{}{\Gamma', T(a), \neg L(a) \vdash \forall x. [P(x) \supset \neg T(x)]} \text{Ax} \\
 \frac{}{\Gamma', T(a), \neg L(a) \vdash P(a) \supset \neg T(a)} \text{Ve} \quad \frac{}{\Gamma', T(a), \neg L(a) \vdash P(a)} \text{Ax} \\
 \frac{}{\Gamma', T(a), \neg L(a) \vdash \neg T(a)} \supset e \quad \frac{}{\Gamma', T(a), \neg L(a) \vdash T(a)} \text{Ax} \\
 \frac{}{\Gamma', T(a) \vdash \neg \neg L(a)} \neg i \\
 \frac{}{\Gamma', T(a) \vdash L(a)} \neg e \\
 \vdots \\
 \frac{}{\Gamma' \vdash \forall x. [L(x) \supset \neg U(x)]} \text{Ax} \quad \frac{}{\Gamma' \vdash L(a) \vee T(a)} \text{Ax} \quad \frac{}{\Gamma', L(a) \vdash L(a)} \text{Ax} \quad \vdots \\
 \frac{}{\Gamma' \vdash L(a) \supset \neg U(a)} \text{Ve} \quad \frac{}{\Gamma' \vdash L(a)} \text{Ve} \\
 \frac{}{\Gamma' \vdash \neg U(a)} \supset e \\
 \frac{}{\Gamma \vdash P(a) \supset \neg U(a)} \supset i
 \end{array}$$

Example 2

Show that $\vdash_{\mathcal{G}} (\neg\phi \vee \neg\psi) \supset \neg(\phi \wedge \psi)$.

Example 2

Show that $\vdash_{\mathcal{L}} (\neg\phi \vee \neg\psi) \supset \neg(\phi \wedge \psi)$.

Let $\Gamma = \{\neg\phi \vee \neg\psi\}$ and $\Delta = \Gamma \cup \{\neg\phi\}$.

Example 3

Show that $\vdash_{\mathcal{G}} \neg(\varphi \wedge \psi) \supset (\neg\varphi \vee \neg\psi)$.

Example 3

Show that $\vdash_{\mathcal{L}} \neg(\varphi \wedge \psi) \supset (\neg\varphi \vee \neg\psi)$. Let $\alpha := \neg(\varphi \wedge \psi)$.

$$\frac{\begin{array}{c} ??? \\ \vdots \\ \alpha \vdash \neg\varphi \vee \neg\psi \end{array}}{\vdash \alpha \supset (\neg\varphi \vee \neg\psi)} \supset i$$

- **Tip:** If you have to prove $\Gamma \vdash_{\mathcal{L}} \alpha \vee \beta$, where $\Gamma \not\vdash_{\mathcal{L}} \alpha$ and $\Gamma \not\vdash_{\mathcal{L}} \beta$, use $\neg e$!
- Same for if you have to prove $\Gamma \vdash_{\mathcal{L}} \exists x. [\alpha]$, but $\Gamma \not\vdash_{\mathcal{L}} \alpha(t)$ for any t .
- If all else fails, look to $\neg e$ for help!

Example 3

Show that $\vdash_{\mathcal{L}} \neg(\varphi \wedge \psi) \supset (\neg\varphi \vee \neg\psi)$. Let $\alpha := \neg(\varphi \wedge \psi)$.

$$\frac{\begin{array}{c} ??? \\ \vdots \\ \alpha \vdash \neg\neg(\neg\varphi \vee \neg\psi) \end{array}}{\alpha \vdash \neg\varphi \vee \neg\psi} \neg e$$

The only way we know to get a “brand new” expression headed by \neg is $\neg i$! Suppose we had a formula β such that the following held, then done.

$$\frac{\begin{array}{c} \vdots \\ \alpha, \neg(\neg\varphi \vee \neg\psi) \vdash \neg\beta \end{array} \quad \begin{array}{c} \vdots \\ \alpha, \neg(\neg\varphi \vee \neg\psi) \vdash \beta \end{array}}{\alpha \vdash \neg\neg(\neg\varphi \vee \neg\psi)} \neg i$$

$$\frac{\alpha \vdash \neg\neg(\neg\varphi \vee \neg\psi)}{\alpha \vdash \neg\varphi \vee \neg\psi} \neg e$$

But what is this β supposed to be?

Example 3

Show that $\vdash_{\mathcal{L}} \neg(\varphi \wedge \psi) \supset (\neg\varphi \vee \neg\psi)$. Let $\alpha := \neg(\varphi \wedge \psi)$.

$$\frac{\begin{array}{c} ??? \\ \vdots \\ \alpha \vdash \neg\neg(\neg\varphi \vee \neg\psi) \end{array}}{\alpha \vdash \neg\varphi \vee \neg\psi} \neg e$$

The only way we know to get a “brand new” expression headed by \neg is $\neg i$! Suppose we had a formula β such that the following held, then done.

$$\frac{\begin{array}{c} \vdots \\ \alpha, \neg(\neg\varphi \vee \neg\psi) \vdash \neg\beta \end{array} \quad \begin{array}{c} \vdots \\ \alpha, \neg(\neg\varphi \vee \neg\psi) \vdash \beta \end{array}}{\alpha \vdash \neg\neg(\neg\varphi \vee \neg\psi)} \neg i$$

$$\frac{\alpha \vdash \neg\neg(\neg\varphi \vee \neg\psi)}{\alpha \vdash \neg\varphi \vee \neg\psi} \neg e$$

But what is this β supposed to be? What can we prove from this context?

Example 3

Show that $\vdash_{\mathcal{G}} \neg(\varphi \wedge \psi) \supset (\neg\varphi \vee \neg\psi)$. Let $\alpha := \neg(\varphi \wedge \psi)$.

$$\begin{array}{c}
 \frac{}{\neg(\neg\varphi \vee \neg\psi), \neg\varphi \vdash \neg\varphi} \text{Ax} \\
 \frac{}{\neg(\neg\varphi \vee \neg\psi), \neg\psi \vdash \neg\psi} \text{Ax} \\
 \frac{}{\neg(\neg\varphi \vee \neg\psi), \neg\varphi \vdash \neg\varphi \vee \neg\psi} \vee i_0 \quad \frac{}{\neg(\neg\varphi \vee \neg\psi), \neg\psi \vdash \neg\varphi \vee \neg\psi} \vee i_1 \\
 \frac{}{\neg(\neg\varphi \vee \neg\psi) \vdash \neg(\neg\varphi \vee \neg\psi)} \neg i \\
 \frac{}{\neg(\neg\varphi \vee \neg\psi) \vdash \neg\neg\varphi} \neg e \\
 \frac{}{\neg(\neg\varphi \vee \neg\psi) \vdash \varphi} \neg e
 \end{array}$$

Similarly, $\neg(\neg\varphi \vee \neg\psi) \vdash \psi$. **Exercise:** Draw this proof tree.

Can use Monotonicity and $\wedge i$ to get a proof π of

$$\alpha, \neg(\neg\varphi \vee \neg\psi) \vdash_{\mathcal{G}} \varphi \wedge \psi$$

Example 3

Show that $\vdash_{\mathcal{G}} \neg(\varphi \wedge \psi) \supset (\neg\varphi \vee \neg\psi)$. Let $\alpha := \neg(\varphi \wedge \psi)$.

$$\frac{\frac{\alpha, \neg(\neg\varphi \vee \neg\psi) \vdash \alpha \quad \text{Ax} \quad \alpha, \neg(\neg\varphi \vee \neg\psi) \vdash (\varphi \wedge \psi)}{\alpha \vdash \neg\neg(\neg\varphi \vee \neg\psi)} \neg i}{\alpha \vdash \neg\varphi \vee \neg\psi} \neg e$$

Exercise: Prove that $\forall x. [P(x) \vee \neg P(x)]$.

Example 4

Show that $\vdash_{\mathcal{L}} \neg \forall x. [P(x)] \supset \exists x. [\neg P(x)]$.

Example 4

Show that $\vdash_{\mathcal{L}} \neg \forall x. [P(x)] \supset \exists x. [\neg P(x)]$.

$$\frac{\begin{array}{c} ??? \\ \vdots \\ \neg \forall x. [P(x)] \vdash \exists x. [\neg P(x)] \end{array}}{\vdash \neg \forall x. [P(x)] \supset \exists x. [\neg P(x)]} \supset i$$

Example 4

Show that $\vdash_{\mathcal{L}} \neg \forall x. [P(x)] \supset \exists x. [\neg P(x)]$.

$$\begin{array}{c}
 \begin{array}{c} ??? \\ \vdots \\ \neg \forall x. [P(x)], \neg \exists x. [\neg P(x)] \vdash \neg \gamma \end{array} \quad \begin{array}{c} ??? \\ \vdots \\ \neg \forall x. [P(x)], \neg \exists x. [\neg P(x)] \vdash \gamma \end{array} \\
 \hline
 \neg \forall x. [P(x)] \vdash \neg \neg \exists x. [\neg P(x)] \quad \neg i \\
 \hline
 \neg \forall x. [P(x)] \vdash \exists x. [\neg P(x)] \quad \neg e \\
 \hline
 \vdash \neg \forall x. [P(x)] \supset \exists x. [\neg P(x)] \quad \supset i
 \end{array}$$

Why move to $\vdash_{\mathcal{G}}$?

- One main reason for moving to $\vdash_{\mathcal{G}}$ was intuitiveness
- Easier proofs, as we just saw
- Another reason is convenience for automation
- Proof search in $\vdash_{\mathcal{H}}$ is not syntactically decidable (even for PL)
 - Have to search through (infinitely many possible) instantiations of axiom schema which might appear in a proof
- Is $\vdash_{\mathcal{G}}$ better?
- We will see that $\vdash_{\mathcal{G}}$ enjoys some nice properties.
- Monotonicity and cut hold as usual.
- Is there anything that helps with proof search?

Unnecessary detours in proofs

Consider a proof of the following sort.

$$\frac{\frac{\text{Ax}}{\varphi, \psi \vdash \varphi} \quad \frac{\text{Ax}}{\varphi, \psi \vdash \psi}}{\varphi, \psi \vdash \varphi \wedge \psi} \wedge i$$
$$\frac{\varphi, \psi \vdash \varphi \wedge \psi}{\varphi, \psi \vdash \varphi} \wedge e_0$$

Unnecessary detours in proofs

Consider a proof of the following sort.

$$\frac{\frac{\frac{}{\varphi, \psi \vdash \varphi} \text{Ax}}{\varphi, \psi \vdash \varphi \wedge \psi} \wedge i}{\varphi, \psi \vdash \varphi} \wedge e_0$$

We first introduce an \wedge , and then immediately eliminate it.

Could have replaced this entire proof by the following, smaller proof without any such wasteful detours involving large expressions.

$$\frac{}{\varphi, \psi \vdash \varphi} \text{Ax}$$

Unnecessary detours in proofs

- **Exercise:** What does a proof involving a detour on an \vee or a \supset look like?
- Detours on these operators involve the introduction rule for that operator, immediately followed by the elimination rule.
- What about \neg ?

Unnecessary detours in proofs

- **Exercise:** What does a proof involving a detour on an \vee or a \supset look like?
- Detours on these operators involve the introduction rule for that operator, immediately followed by the elimination rule.
- What about \neg ?
- Clearly \neg_i followed by \neg_e is **not** an unnecessary detour.
- We could not have done the earlier proofs without using this combo!
- However, the expressions we used for \neg_i were informed by the context and the expected conclusion.
- Can we eliminate all unnecessary detours?
- Given Γ, α , is there some finite set to which every expression occurring in any proof of $\Gamma \vdash_{\mathcal{F}} \alpha$ belongs?