

clear history

```
clearvars all;
clc;
```

part a

```
% given
```

```
I = diag([10; 20; 30])
```

```
I = 3x3
    10     0     0
     0    20     0
     0     0    30
```

```
omega0_deg = [10; 0; 30];
omega0 = deg2rad(omega0_deg);
```

```
% solution
```

```
H0 = I*omega0;
H0_mag = norm(H0);
T0 = 1/2*transpose(omega0)*I*omega0;
```

```
% display
```

```
H0_mag, T0
```

```
H0_mag =
15.8046
T0 =
4.2646
```

part b

```
% solution
```

```
dwdt = @(t, omega) inv(I)*(-cross(omega, I*omega));
tspan = [0, 100];
oopts = odeset('RelTol',1e-9,'AbsTol',1e-9);
```

```
[t1, w1] = ode45(dwdt, tspan, omega0, oopts);
w1_deg = rad2deg(w1);
```

```
% plot
```


```
fig = figure;
plot(t1, w1_deg(:,1), 'r', t1, w1_deg(:,2), 'g', t1, w1_deg(:,3), 'b', 'LineWidth',
1.5);
title('Angular Velocity vs Time: \omega_0 = (10;0;30) [deg/s]');
legend('\omega_x', '\omega_y', '\omega_z');
xlabel('Time [s]');
```

```

ylabel('Angular Velocity [deg/s]');
ylim([-15 45]);
grid on;

% save
saveas(fig, "./images/s02b.png");

```



part c

```


% solution
H1 = transpose(I * transpose(w1));
H1_mag = sqrt(sum(H1.^2,2));
T1 = 1/2*sum(w1.*H1, 2);

dH = H1_mag-H0_mag;
dT = T1-T0;

% plot
fig = figure;
plot(t1, dH, 'k', t1, dT, 'm', 'LineWidth', 1.5);
title('Deviation of |H| and T');
legend('|H|-|H_0|', 'T-T_0', 'Location', 'southeast');
xlabel('Time [s]');
ylabel('Deviation');
grid on;

```

```
% save
saveas(fig, "./images/s02c.png");
```



I believe my code is working, as the both the angular momentum deviation and kinetic energy deviation are extremely minimal, being on the order of 10^{-8} .

part d

```
% solution
[Xs, Ys, Zs] = sphere(20);
Xs = H0_mag*Xs;
Ys = H0_mag*Ys;
Zs = H0_mag*Zs;

radii = sqrt(2*T0.*diag(I));
[Xe, Ye, Ze] = ellipsoid(0,0,0, radii(1), radii(2), radii(3), 60);

% plot
fig = figure;
hold on;
% sphere
surf(Xs, Ys, Zs, 'FaceAlpha',0.3, 'EdgeColor','b', 'FaceColor', [0.8 0.8 1]);
% ellipsoid
surf(Xe, Ye, Ze, 'FaceAlpha',0.3, 'EdgeColor','g', 'FaceColor', [0.2 0.8 0.2]);
% polhode points (body-fixed H)
plot3(H1(:,1), H1(:,2), H1(:,3), '.r', 'MarkerSize', 5);
```


```

title('Polhode Plot: \omega_0 = (10;0;30) [deg/s]');
legend('Angular Momentum [kg*m^2/s]', 'Kinetic Energy [kg*m^2/s]', 'Body Fixed  
Angular Momentum [kg*m^2/s]', 'Location', 'southoutside');
xlabel('H_x [kg*m^2/s]');
ylabel('H_y [kg*m^2/s]');
zlabel('H_z [kg*m^2/s]');
axis equal;
grid on;
view([-43.3 33.8])
hold off;

% save
saveas(fig, './images/s02d.png');

```

Polhode Plot: $\omega_0 = (10;0;30)$ [deg/s]



part e

```

% solution
omega2_deg = [1; 15; 0];
omega2 = deg2rad(omega2_deg);
[t2, w2] = ode45(dwdt, tspan, omega2, oopts);
w2_deg = rad2deg(w2);

% plot
fig = figure;
plot(t2, w2_deg(:,1), 'r', t2, w2_deg(:,2), 'g', t2, w2_deg(:,3), 'b', 'LineWidth',
1.5);


```

```

title('Angular Velocity vs Time: \omega_0 = (1;15;0) [deg/s]');
legend('\omega_x', '\omega_y', '\omega_z');
xlabel('Time [s]');
ylabel('Angular Velocity [deg/s]');
ylim([-20 30]);
grid on;

% save
saveas(fig, "./images/s02e.png");

```



part f

```

% solution
H2_0 = I*omega2;
H2_0_mag = norm(H2_0);
T2_0 = 1/2*transpose(omega2)*I*omega2;

H2 = transpose(I*transpose(w2));
H2_mag = sqrt(sum(H2.^2,2));
T2 = 1/2*sum(w2.*H2, 2);

[Xs, Ys, Zs] = sphere(20);
Xs = H2_0_mag*Xs;
Ys = H2_0_mag*Ys;
Zs = H2_0_mag*Zs;

```


```

radii = sqrt(2*T2_0.*diag(I));
[Xe, Ye, Ze] = ellipsoid(0,0,0, radii(1), radii(2), radii(3), 60);

% plot
fig = figure;
hold on;
% sphere
surf(Xs, Ys, Zs, 'FaceAlpha',0.3, 'EdgeColor','b', 'FaceColor', [0.8 0.8 1]);
% ellipsoid
surf(Xe, Ye, Ze, 'FaceAlpha',0.3, 'EdgeColor','g', 'FaceColor', [0.2 0.8 0.2]);
% polhode points (body-fixed H)
plot3(H2(:,1), H2(:,2), H2(:,3), '.r', 'MarkerSize', 5);
title('Polhode Plot: \omega_0 = (1;15;0) [deg/s]');
legend('Angular Momentum [kg*m^2/s]', 'Kinetic Energy [kg*m^2/s]', 'Body Fixed
Angular Momentum [kg*m^2/s]', 'Location', 'southoutside');
xlabel('H_x [kg*m^2/s]');
ylabel('H_y [kg*m^2/s]');
zlabel('H_z [kg*m^2/s]');
axis equal;
grid on;
view([40.6 14.7]);
hold off;

% save
saveas(fig, "./images/s02f.png");

```



The spacecraft in part b and d has a much higher kinetic energy in the Z axis than in the X and Y axes, and as well varies each axis in only small oscillations. This is far different from the spacecraft in e and f, who has far more energy-variance within each axis, as well as having its overall spread be much closer together. These two differences result in the Polhode plot of the first spacecraft being confined to a small area high in the positive Z axis, while the second spacecraft's plot ventures all over the place.