

A Hard Satisfiable Problem with 160 Variables

Valentin Mayer-Eichberger
NICTA and
University of New South Wales
Valentin.Mayer-Eichberger@nicta.com.au

Abstract—In trying to solve a hard graph colouring problem we ran into an interesting SAT formula. The encoding uses just 160 variables and defines a special case of a rectangle-free coloring of a 18×18 grid using four colors. Rectangle-free means that the corners of every rectangle in the grid cannot have all the same colour. Such structured satisfiable problems pose a real challenge to SAT solvers.

I. INTRODUCTION

In 2011 a blog post of

`blog.computationalcomplexity.org`

announced a reward of 289 \$ for a solution to the problem of 4-colouring a 17×17 grid such that for each rectangle in the grid all its corners consist of at least two different colours. A solution to 16×16 was known to exist and all grids 19×19 and larger were proven to not contain such a colouring. In 2012 Steinbach and Posthoff presented a solution to 17×17 and 18×18 [1] (every solution of larger grids generates solutions to smaller). We provide the SAT competition with an interesting encoding for this problem which is similar to the approach they used. It will be valuable for the community to see if any SAT solver is able to solve this hard problem within the time-out. Our own experiments show that CDCL solvers tend to spend several hours to find a solution. By such a benchmark we might identify advantages of non-standard SAT solver techniques.

II. ENCODING

Naive encodings for this problem can solve grids up to 14×14 almost instantly and do not put a challenge to a SAT solver. With some advancements and symmetry breaking one can also solve 15×15 and 16×16 . However, no direct approach seems to tackle the hard cases of 17×17 and 18×18 . In this section we explain the tricks that made it possible.

We identify a special case that can be extended to a full solution. If such a solution would exist then the problem is solved, but a negative result would not give much insight. Luckily, it turns out that the simplification does indeed lead to solutions.

We simplify the problem to find a two coloring. We denote the two colours as primary and secondary, and the secondary colour represents the three other colours of the original problem. A solution to this problem can be extended to a solution if

- only the primary colour needs to be rectangle-free,
- $1/4$ of all positions are filled with the primary colour,

- rotating the solution by 90, 180, and 270 degrees will not map a position containing a primary colour onto another.

We can then take a solution of this problem and fill for each rotation the mapped positions of the primary colour with one of the remaining one. Since there are no collisions and rectangle-free is preserved under rotation, we generate a full solution.

A natural choice would be to define for each position in the board a Boolean variable that is true if that position contains the primary colour. We reduce the number of variables by using the restriction that each orbit wrt. to the 90 degree rotations should have exactly one primary colour. This can be encoded in a logarithmic fashion such that two Boolean variables identify for each orbit in which of the four half section of the grid it exists. Furthermore, we break symmetries by forcing the upper left position to contain a primary colour. By these reductions we get a formula that only uses 160 variables.

III. BENCHMARK

The set contains 4 encodings of the same problems. They have been generated by shuffling the variables, literals and order of clauses of the encoding described above.

ACKNOWLEDGEMENT

NICTA is funded by the Australian Government as represented by the Department of Broadband, Communications and the Digital Economy and the Australian Research Council through the ICT Centre of Excellence program.

REFERENCES

- [1] B. Steinbach and C. Posthoff, "Extremely Complex 4-Colored Rectangle-Free Grids: Solution of Open Multiple-Valued Problems," in *ISMVL*, 2012, pp. 37–44.