
Candidate 000000: DM = 25.50 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
75.004
75.002
75.000
74.998
74.996
74.994
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000001: DM = 23.50 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
100.010
100.005
100.000
99.995
99.990
99.985
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000002: DM = 23.50 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
100.035
100.030
100.025
100.020
100.015
100.010
100.005
100.000
99.995
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000003: DM = 24.50 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
186.675
186.650
186.625
186.600
186.575
186.550
186.525
186.500
186.475
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000004: DM = 24.80 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
186.68
186.66
186.64
186.62
186.60
186.58
186.56
186.54
186.52
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000005: DM = 25.10 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
186.68
186.66
186.64
186.62
186.60
186.58
186.56
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000006: DM = 24.80 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
186.66
186.65
186.64
186.63
186.62
186.61
Barycentric frequency (Hz)
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000007: DM = 24.80 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
186.6600
186.6575
186.6550
186.6525
186.6500
186.6475
186.6450
186.6425
186.6400
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000008: DM = 24.80 pc cm^{-3}, $N_T = 16$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
186.656
186.654
186.652
186.650
186.648
186.646
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000009: DM = 24.80 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
186.67
186.66
186.65
186.64
186.63
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000010: DM = 24.50 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
186.71
186.70
186.69
186.68
186.67
186.66
186.65
186.64
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000011: DM = 24.80 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
186.76
186.74
186.72
186.70
186.68
186.66
186.64
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000012: DM = 24.70 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
186.800
186.775
186.750
186.725
186.700
186.675
186.650
186.625
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000013: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
200.02
200.00
199.98
199.96
199.94
199.92
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000014: DM = 24.80 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
200.01
200.00
199.99
199.98
199.97
199.96
199.95
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000015: DM = 24.60 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
200.000
199.998
199.996
199.994
199.992
199.990
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000016: DM = 24.60 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
200.010
200.005
200.000
199.995
199.990
199.985
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000017: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
200.02
200.01
200.00
199.99
199.98
199.97
199.96
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000018: DM = 25.00 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
200.04
200.03
200.02
200.01
200.00
199.99
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000019: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
200.08
200.06
200.04
200.02
200.00
199.98
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000020: DM = 23.70 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
208.174
208.172
208.170
208.168
208.166
208.164
208.162
208.160
208.158
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000021: DM = 24.50 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
216.65
216.60
216.55
216.50
216.45
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000022: DM = 23.90 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
216.675
216.650
216.625
216.600
216.575
216.550
216.525
216.500
216.475
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000023: DM = 23.90 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
216.68
216.66
216.64
216.62
216.60
216.58
216.56
216.54
216.52
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000024: DM = 23.90 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
216.68
216.66
216.64
216.62
216.60
216.58
216.56
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000025: DM = 23.90 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
216.68
216.66
216.64
216.62
216.60
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000026: DM = 23.90 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
216.67
216.66
216.65
216.64
216.63
216.62
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000027: DM = 23.90 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
216.670
216.665
216.660
216.655
216.650
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000028: DM = 23.90 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
216.666
216.664
216.662
216.660
216.658
216.656
216.654
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000029: DM = 23.90 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
216.69
216.68
216.67
216.66
216.65
216.64
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000030: DM = 25.20 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
216.71
216.70
216.69
216.68
216.67
216.66
216.65
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000031: DM = 23.90 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
216.72
216.70
216.68
216.66
216.64
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000032: DM = 23.90 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
216.76
216.74
216.72
216.70
216.68
216.66
216.64
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000033: DM = 23.90 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
216.800
216.775
216.750
216.725
216.700
216.675
216.650
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000034: DM = 25.20 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
216.85
216.80
216.75
216.70
216.65
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000035: DM = 23.90 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
216.90
216.85
216.80
216.75
216.70
216.65
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000036: DM = 24.40 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
250.000
249.998
249.996
249.994
249.992
249.990
249.988
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000037: DM = 24.40 pc cm^{-3}, $N_T = 16$
Barycentric; J0024-7204I; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
286.952
286.950
286.948
286.946
286.944
286.942
286.940
286.938
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000038: DM = 24.70 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
299.998
299.996
299.994
299.992
299.990
299.988
299.986
299.984
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000039: DM = 24.80 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204as; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
373.30
373.25
373.20
373.15
373.10
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000040: DM = 24.50 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
373.32
373.30
373.28
373.26
373.24
373.22
373.20
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000041: DM = 24.50 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
373.32
373.30
373.28
373.26
373.24
373.22
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000042: DM = 24.70 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
373.315
373.310
373.305
373.300
373.295
Barycentric frequency (Hz)
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000043: DM = 24.70 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
373.320
373.315
373.310
373.305
373.300
373.295
373.290
373.285
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000044: DM = 24.70 pc cm^{-3}, $N_T = 16$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
373.308
373.306
373.304
373.302
373.300
373.298
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000045: DM = 24.30 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
373.34
373.33
373.32
373.31
373.30
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000046: DM = 24.90 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
373.40
373.38
373.36
373.34
373.32
373.30
373.28
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000047: DM = 24.80 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
373.44
373.42
373.40
373.38
373.36
373.34
373.32
373.30
373.28
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000048: DM = 24.70 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
373.50
373.45
373.40
373.35
373.30
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000049: DM = 24.40 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
378.31
378.30
378.29
378.28
378.27
378.26
378.25
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000050: DM = 24.30 pc cm^{-3}, $N_T = 16$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
378.308
378.306
378.304
378.302
378.300
378.298
378.296
378.294
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000051: DM = 24.40 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
378.34
378.33
378.32
378.31
378.30
378.29
378.28
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000052: DM = 24.40 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
378.325
378.320
378.315
378.310
378.305
378.300
378.295
378.290
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000053: DM = 24.30 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
378.35
378.34
378.33
378.32
378.31
378.30
378.29
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000054: DM = 24.50 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204F; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
381.170
381.165
381.160
381.155
381.150
381.145
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000055: DM = 24.40 pc cm^{-3}, $N_T = 16$
Barycentric; J0024-7204F; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
381.164
381.162
381.160
381.158
381.156
381.154
381.152
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000056: DM = 24.90 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
399.994
399.992
399.990
399.988
399.986
399.984
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000057: DM = 24.80 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
400.000
399.995
399.990
399.985
399.980
Barycentric frequency (Hz)
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000058: DM = 25.50 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
433.30
433.25
433.20
433.15
433.10
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000059: DM = 25.50 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
433.34
433.32
433.30
433.28
433.26
433.24
433.22
433.20
433.18
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000060: DM = 25.50 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
433.34
433.32
433.30
433.28
433.26
433.24
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000061: DM = 24.40 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
433.33
433.32
433.31
433.30
433.29
433.28
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000062: DM = 25.50 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
433.35
433.34
433.33
433.32
433.31
433.30
433.29
433.28
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000063: DM = 25.50 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
433.330
433.328
433.326
433.324
433.322
433.320
433.318
433.316
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000064: DM = 25.50 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
433.335
433.330
433.325
433.320
433.315
433.310
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000065: DM = 25.50 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
433.36
433.35
433.34
433.33
433.32
433.31
433.30
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000066: DM = 25.50 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
433.37
433.36
433.35
433.34
433.33
433.32
433.31
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000067: DM = 25.50 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
433.40
433.38
433.36
433.34
433.32
433.30
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000068: DM = 25.50 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
433.46
433.44
433.42
433.40
433.38
433.36
433.34
433.32
433.30
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000069: DM = 24.90 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
433.475
433.450
433.425
433.400
433.375
433.350
433.325
433.300
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000070: DM = 25.00 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
433.55
433.50
433.45
433.40
433.35
433.30
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000071: DM = 25.30 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
437.576
437.574
437.572
437.570
437.568
437.566
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000072: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
476.075
476.050
476.025
476.000
475.975
475.950
475.925
475.900
475.875
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000073: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
476.08
476.06
476.04
476.02
476.00
475.98
475.96
475.94
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000074: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
476.08
476.06
476.04
476.02
476.00
475.98
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000075: DM = 24.60 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
476.07
476.06
476.05
476.04
476.03
476.02
476.01
476.00
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000076: DM = 24.60 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
476.07
476.06
476.05
476.04
476.03
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000077: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
476.08
476.07
476.06
476.05
476.04
476.03
476.02
476.01
476.00
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000078: DM = 24.60 pc cm^{-3}, $N_T = 16$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
476.064
476.062
476.060
476.058
476.056
476.054
476.052
476.050
476.048
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000079: DM = 24.60 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
476.09
476.08
476.07
476.06
476.05
476.04
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000080: DM = 24.80 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
476.10
476.08
476.06
476.04
476.02
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000081: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
476.14
476.12
476.10
476.08
476.06
476.04
476.02
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000082: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
476.18
476.16
476.14
476.12
476.10
476.08
476.06
476.04
476.02
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000083: DM = 25.50 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
499.994
499.992
499.990
499.988
499.986
499.984
499.982
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000084: DM = 25.50 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
500.000
499.995
499.990
499.985
499.980
499.975
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000085: DM = 24.70 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
559.98
559.96
559.94
559.92
559.90
559.88
559.86
559.84
559.82
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000086: DM = 24.70 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
559.98
559.96
559.94
559.92
559.90
559.88
559.86
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000087: DM = 24.90 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
559.98
559.96
559.94
559.92
559.90
Barycentric frequency (Hz)
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000088: DM = 24.70 pc cm^{-3}, $N_T = 16$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
559.960
559.958
559.956
559.954
559.952
559.950
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000089: DM = 24.70 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
559.9650
559.9625
559.9600
559.9575
559.9550
559.9525
559.9500
559.9475
559.9450
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000090: DM = 24.70 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
559.98
559.97
559.96
559.95
559.94
Barycentric frequency (Hz)
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000091: DM = 25.00 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
560.00
559.99
559.98
559.97
559.96
559.95
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000092: DM = 24.70 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
560.04
560.02
560.00
559.98
559.96
559.94
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000093: DM = 24.70 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
560.08
560.06
560.04
560.02
560.00
559.98
559.96
559.94
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000094: DM = 24.20 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
599.995
599.990
599.985
599.980
599.975
599.970
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000095: DM = 25.50 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
599.990
599.988
599.986
599.984
599.982
599.980
599.978
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000096: DM = 24.10 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
650.00
649.95
649.90
649.85
649.80
Barycentric frequency (Hz)
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000097: DM = 25.20 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
650.000
649.975
649.950
649.925
649.900
649.875
649.850
649.825
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000098: DM = 25.20 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
650.00
649.98
649.96
649.94
649.92
649.90
649.88
649.86
649.84
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000099: DM = 25.20 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
650.00
649.98
649.96
649.94
649.92
Barycentric frequency (Hz)
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000100: DM = 25.20 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
649.99
649.98
649.97
649.96
649.95
649.94
649.93
Barycentric frequency (Hz)
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000101: DM = 25.20 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
650.01
650.00
649.99
649.98
649.97
649.96
Barycentric frequency (Hz)
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000102: DM = 25.20 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
649.988
649.986
649.984
649.982
649.980
649.978
649.976
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000103: DM = 25.20 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
649.990
649.985
649.980
649.975
649.970
Barycentric frequency (Hz)
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000104: DM = 25.20 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
650.03
650.02
650.01
650.00
649.99
649.98
649.97
649.96
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000105: DM = 23.60 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
650.03
650.02
650.01
650.00
649.99
649.98
649.97
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000106: DM = 25.20 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
650.06
650.04
650.02
650.00
649.98
649.96
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000107: DM = 25.20 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
650.10
650.08
650.06
650.04
650.02
650.00
649.98
649.96
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000108: DM = 25.20 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
650.150
650.125
650.100
650.075
650.050
650.025
650.000
649.975
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000109: DM = 25.20 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
650.15
650.10
650.05
650.00
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000110: DM = 23.60 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
699.986
699.984
699.982
699.980
699.978
699.976
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000111: DM = 24.90 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
746.61
746.60
746.59
746.58
746.57
746.56
746.55
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000112: DM = 24.80 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
746.6175
746.6150
746.6125
746.6100
746.6075
746.6050
746.6025
746.6000
746.5975
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000113: DM = 24.80 pc cm^{-3}, $N_T = 16$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
746.612
746.610
746.608
746.606
746.604
746.602
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000114: DM = 23.50 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
749.986
749.984
749.982
749.980
749.978
749.976
749.974
749.972
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000115: DM = 24.40 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
756.650
756.625
756.600
756.575
756.550
756.525
756.500
756.475
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000116: DM = 24.40 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
756.64
756.62
756.60
756.58
756.56
756.54
756.52
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000117: DM = 24.40 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
756.61
756.60
756.59
756.58
756.57
756.56
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000118: DM = 24.40 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
756.64
756.62
756.60
756.58
756.56
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000119: DM = 24.40 pc cm^{-3}, $N_T = 16$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
756.608
756.606
756.604
756.602
756.600
756.598
756.596
756.594
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000120: DM = 24.40 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
756.65
756.64
756.63
756.62
756.61
756.60
756.59
756.58
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000121: DM = 24.40 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
756.620
756.615
756.610
756.605
756.600
756.595
756.590
756.585
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000122: DM = 24.50 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
756.65
756.64
756.63
756.62
756.61
756.60
756.59
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000123: DM = 24.40 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
756.66
756.64
756.62
756.60
756.58
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000124: DM = 24.40 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
756.70
756.68
756.66
756.64
756.62
756.60
756.58
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000125: DM = 24.40 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204O; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
756.750
756.725
756.700
756.675
756.650
756.625
756.600
756.575
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000126: DM = 24.40 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204F; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
762.330
762.325
762.320
762.315
762.310
762.305
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000127: DM = 24.40 pc cm^{-3}, $N_T = 16$
Barycentric; J0024-7204F; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
762.322
762.320
762.318
762.316
762.314
762.312
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000128: DM = 24.50 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
799.990
799.985
799.980
799.975
799.970
799.965
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000129: DM = 24.50 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
799.984
799.982
799.980
799.978
799.976
799.974
799.972
799.970
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000130: DM = 25.10 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
866.650
866.625
866.600
866.575
866.550
866.525
866.500
866.475
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000131: DM = 25.10 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
866.66
866.64
866.62
866.60
866.58
866.56
866.54
866.52
866.50
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000132: DM = 25.10 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
866.66
866.64
866.62
866.60
866.58
866.56
866.54
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000133: DM = 25.10 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
866.65
866.64
866.63
866.62
866.61
866.60
866.59
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000134: DM = 25.10 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
866.655
866.650
866.645
866.640
866.635
Barycentric frequency (Hz)
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000135: DM = 25.10 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
866.650
866.648
866.646
866.644
866.642
866.640
866.638
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000136: DM = 25.10 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
866.68
866.67
866.66
866.65
866.64
866.63
866.62
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000137: DM = 24.60 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
866.70
866.69
866.68
866.67
866.66
866.65
866.64
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000138: DM = 25.10 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
866.74
866.72
866.70
866.68
866.66
866.64
866.62
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000139: DM = 25.10 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
866.800
866.775
866.750
866.725
866.700
866.675
866.650
866.625
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000140: DM = 25.10 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_family (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
866.85
866.80
866.75
866.70
866.65
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000141: DM = 23.90 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_comb (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
899.984
899.982
899.980
899.978
899.976
899.974
899.972
899.970
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000142: DM = 24.70 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
933.27
933.26
933.25
933.24
933.23
933.22
933.21
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000143: DM = 24.70 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
933.270
933.265
933.260
933.255
933.250
933.245
Barycentric frequency (Hz)
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000144: DM = 24.70 pc cm^{-3}, $N_T = 16$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
933.264
933.262
933.260
933.258
933.256
933.254
Barycentric frequency (Hz)
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000145: DM = 24.80 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204D; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
933.30
933.29
933.28
933.27
933.26
933.25
Barycentric frequency (Hz)
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000146: DM = 23.60 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
938.460
938.458
938.456
938.454
938.452
938.450
938.448
938.446
938.444
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000147: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
952.10
952.05
952.00
951.95
951.90
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000148: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
952.125
952.100
952.075
952.050
952.025
952.000
951.975
951.950
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000149: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
952.12
952.10
952.08
952.06
952.04
952.02
952.00
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000150: DM = 24.50 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
952.10
952.09
952.08
952.07
952.06
952.05
952.04
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000151: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
952.14
952.12
952.10
952.08
952.06
952.04
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000152: DM = 24.60 pc cm^{-3}, $N_T = 16$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
952.0775
952.0750
952.0725
952.0700
952.0675
952.0650
952.0625
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000153: DM = 24.60 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
952.12
952.11
952.10
952.09
952.08
952.07
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000154: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
952.14
952.12
952.10
952.08
952.06
952.04
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000155: DM = 24.60 pc cm^{-3}, $N_T = 16$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
952.126
952.124
952.122
952.120
952.118
952.116
952.114
952.112
952.110
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000156: DM = 24.60 pc cm^{-3}, $N_T = 32$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
952.14
952.13
952.12
952.11
952.10
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000157: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
952.16
952.14
952.12
952.10
952.08
952.06
Barycentric frequency (Hz)
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000158: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
952.20
952.18
952.16
952.14
952.12
952.10
952.08
952.06
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000159: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; J0024-7204J; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
952.225
952.200
952.175
952.150
952.125
952.100
952.075
952.050
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000160: DM = 24.60 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: none (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
952.25
952.20
952.15
952.10
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000161: DM = 25.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
996
995
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000162: DM = 23.60 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
996
995
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000163: DM = 25.30 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
996
995
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000164: DM = 24.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
996
995
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000165: DM = 23.80 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
996
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000166: DM = 25.20 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
996
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000167: DM = 25.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
996
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000168: DM = 25.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
996
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000169: DM = 25.30 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
996
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000170: DM = 24.30 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
996
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000171: DM = 24.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
996
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000172: DM = 25.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000173: DM = 25.30 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000174: DM = 24.10 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
996
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000175: DM = 24.20 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000176: DM = 24.20 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000177: DM = 24.10 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
999
998
997
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000178: DM = 24.10 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
998.0
997.5
997.0
996.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000179: DM = 23.80 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
998.0
997.5
997.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000180: DM = 24.10 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
997.5
997.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000181: DM = 24.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
998.0
997.5
997.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000182: DM = 24.30 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
997.5
997.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000183: DM = 23.60 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
998.0
997.5
997.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000184: DM = 24.20 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
997.5
997.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000185: DM = 25.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
998.0
997.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000186: DM = 23.70 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
997.5
997.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000187: DM = 23.60 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
997.5
997.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000188: DM = 24.00 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
997.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000189: DM = 23.60 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
997.5
997.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000190: DM = 25.30 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
998.0
997.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000191: DM = 24.00 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
997.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000192: DM = 23.80 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
997.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000193: DM = 23.90 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
998.0
997.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000194: DM = 24.00 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
997.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000195: DM = 23.60 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
997.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000196: DM = 24.00 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
997.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000197: DM = 24.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
998.0
997.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000198: DM = 23.80 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
998.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000199: DM = 25.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
998.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000200: DM = 23.80 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000201: DM = 24.30 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000202: DM = 24.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
998.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000203: DM = 24.30 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000204: DM = 23.90 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
998.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000205: DM = 23.90 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.5
999.0
998.5
998.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000206: DM = 25.20 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
998.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000207: DM = 23.80 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.25
1000.00
999.75
999.50
999.25
999.00
998.75
998.50
998.25
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000208: DM = 24.30 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.25
1000.00
999.75
999.50
999.25
999.00
998.75
998.50
998.25
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000209: DM = 23.60 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.25
1000.00
999.75
999.50
999.25
999.00
998.75
998.50
998.25
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000210: DM = 24.70 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
998.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000211: DM = 24.00 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.25
1000.00
999.75
999.50
999.25
999.00
998.75
998.50
998.25
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000212: DM = 23.80 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.25
1000.00
999.75
999.50
999.25
999.00
998.75
998.50
998.25
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000213: DM = 24.70 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.25
1000.00
999.75
999.50
999.25
999.00
998.75
998.50
998.25
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000214: DM = 24.70 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000215: DM = 23.70 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.00
999.75
999.50
999.25
999.00
998.75
998.50
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000216: DM = 23.90 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000217: DM = 23.60 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.0
999.5
999.0
998.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000218: DM = 23.90 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.0
999.8
999.6
999.4
999.2
999.0
998.8
998.6
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000219: DM = 24.80 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.50
1000.25
1000.00
999.75
999.50
999.25
999.00
998.75
998.50
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000220: DM = 24.70 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.0
999.8
999.6
999.4
999.2
999.0
998.8
998.6
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000221: DM = 24.00 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.25
1000.00
999.75
999.50
999.25
999.00
998.75
998.50
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000222: DM = 25.30 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.50
1000.25
1000.00
999.75
999.50
999.25
999.00
998.75
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000223: DM = 24.70 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.8
999.6
999.4
999.2
999.0
998.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000224: DM = 24.20 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.8
999.6
999.4
999.2
999.0
998.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000225: DM = 23.80 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.0
999.8
999.6
999.4
999.2
999.0
998.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000226: DM = 23.70 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.8
999.6
999.4
999.2
999.0
998.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000227: DM = 24.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.50
1000.25
1000.00
999.75
999.50
999.25
999.00
998.75
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000228: DM = 24.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.50
1000.25
1000.00
999.75
999.50
999.25
999.00
998.75
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000229: DM = 24.60 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.8
999.6
999.4
999.2
999.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000230: DM = 25.50 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.0
999.8
999.6
999.4
999.2
999.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000231: DM = 25.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.4
1000.2
1000.0
999.8
999.6
999.4
999.2
999.0
998.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000232: DM = 24.00 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.8
999.6
999.4
999.2
999.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000233: DM = 23.80 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.0
999.8
999.6
999.4
999.2
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000234: DM = 24.10 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.8
999.6
999.4
999.2
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000235: DM = 25.50 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.8
999.6
999.4
999.2
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000236: DM = 24.70 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.75
1000.50
1000.25
1000.00
999.75
999.50
999.25
999.00
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000237: DM = 24.50 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.9
999.8
999.7
999.6
999.5
999.4
999.3
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000238: DM = 24.00 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.0
999.8
999.6
999.4
999.2
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000239: DM = 23.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.6
1000.4
1000.2
1000.0
999.8
999.6
999.4
999.2
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000240: DM = 24.30 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.1
1000.0
999.9
999.8
999.7
999.6
999.5
999.4
999.3
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000241: DM = 24.30 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000242: DM = 23.80 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.9
999.8
999.7
999.6
999.5
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000243: DM = 24.20 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.1
1000.0
999.9
999.8
999.7
999.6
999.5
999.4
999.3
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000244: DM = 25.40 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000245: DM = 25.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.6
1000.4
1000.2
1000.0
999.8
999.6
999.4
999.2
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000246: DM = 25.40 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.9
999.8
999.7
999.6
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000247: DM = 24.70 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.9
999.8
999.7
999.6
999.5
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000248: DM = 23.80 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.4
1000.2
1000.0
999.8
999.6
999.4
999.2
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000249: DM = 23.60 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.9
999.8
999.7
999.6
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000250: DM = 24.70 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.1
1000.0
999.9
999.8
999.7
999.6
999.5
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000251: DM = 24.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.4
1000.2
1000.0
999.8
999.6
999.4
999.2
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000252: DM = 23.50 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.9
999.8
999.7
999.6
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000253: DM = 25.50 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.1
1000.0
999.9
999.8
999.7
999.6
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000254: DM = 23.80 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.9
999.8
999.7
999.6
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000255: DM = 23.90 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.4
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000256: DM = 23.60 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.05
1000.00
999.95
999.90
999.85
999.80
999.75
999.70
999.65
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000257: DM = 25.50 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0
999.9
999.8
999.7
999.6
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000258: DM = 25.00 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.00
999.95
999.90
999.85
999.80
999.75
999.70
999.65
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000259: DM = 23.50 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.1
1000.0
999.9
999.8
999.7
999.6
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000260: DM = 24.10 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.6
1000.4
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000261: DM = 24.70 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.1
1000.0
999.9
999.8
999.7
999.6
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000262: DM = 24.40 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.00
999.95
999.90
999.85
999.80
999.75
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000263: DM = 23.80 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.1
1000.0
999.9
999.8
999.7
999.6
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000264: DM = 24.70 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.6
1000.4
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000265: DM = 23.90 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.05
1000.00
999.95
999.90
999.85
999.80
999.75
999.70
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000266: DM = 24.20 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.000
999.975
999.950
999.925
999.900
999.875
999.850
999.825
999.800
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000267: DM = 24.10 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.00
999.95
999.90
999.85
999.80
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000268: DM = 23.90 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.05
1000.00
999.95
999.90
999.85
999.80
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000269: DM = 24.70 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.1
1000.0
999.9
999.8
999.7
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000270: DM = 23.80 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.4
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000271: DM = 25.10 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.025
1000.000
999.975
999.950
999.925
999.900
999.875
999.850
999.825
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000272: DM = 24.70 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.05
1000.00
999.95
999.90
999.85
999.80
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000273: DM = 24.40 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
999.8925
999.8900
999.8875
999.8850
999.8825
999.8800
999.8775
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000274: DM = 25.20 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.00
999.98
999.96
999.94
999.92
999.90
999.88
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000275: DM = 23.70 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.1
1000.0
999.9
999.8
999.7
999.6
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000276: DM = 23.70 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.10
1000.05
1000.00
999.95
999.90
999.85
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000277: DM = 23.60 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.1
1000.0
999.9
999.8
999.7
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000278: DM = 24.00 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.4
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000279: DM = 24.30 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.00
999.98
999.96
999.94
999.92
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000280: DM = 25.30 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
999.944
999.942
999.940
999.938
999.936
999.934
999.932
999.930
999.928
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000281: DM = 23.90 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
999.98
999.97
999.96
999.95
999.94
999.93
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000282: DM = 23.90 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.1
1000.0
999.9
999.8
999.7
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000283: DM = 24.00 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.05
1000.00
999.95
999.90
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000284: DM = 24.00 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.02
1000.00
999.98
999.96
999.94
999.92
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000285: DM = 24.00 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.00
999.99
999.98
999.97
999.96
999.95
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000286: DM = 23.90 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
999.980
999.975
999.970
999.965
999.960
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000287: DM = 24.00 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
999.980
999.978
999.976
999.974
999.972
999.970
999.968
999.966
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000288: DM = 24.00 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.05
1000.00
999.95
999.90
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000289: DM = 24.50 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.3
1000.2
1000.1
1000.0
999.9
999.8
999.7
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000290: DM = 23.90 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.4
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000291: DM = 24.50 pc cm^{-3}, $N_T = 32$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 449.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.02
1000.01
1000.00
999.99
999.98
999.97
999.96
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000292: DM = 24.30 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.04
1000.02
1000.00
999.98
999.96
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000293: DM = 23.90 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.06
1000.04
1000.02
1000.00
999.98
999.96
999.94
999.92
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000294: DM = 24.80 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.0375
1000.0350
1000.0325
1000.0300
1000.0275
1000.0250
1000.0225
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000295: DM = 25.40 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.10
1000.05
1000.00
999.95
999.90
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000296: DM = 25.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.4
1000.2
1000.0
999.8
999.6
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000297: DM = 24.00 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.125
1000.100
1000.075
1000.050
1000.025
1000.000
999.975
999.950
999.925
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000298: DM = 25.30 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.125
1000.100
1000.075
1000.050
1000.025
1000.000
999.975
999.950
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000299: DM = 23.90 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.15
1000.10
1000.05
1000.00
999.95
999.90
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000300: DM = 25.50 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.2
1000.1
1000.0
999.9
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000301: DM = 24.50 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.114
1000.112
1000.110
1000.108
1000.106
1000.104
1000.102
1000.100
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000302: DM = 24.30 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.20
1000.15
1000.10
1000.05
1000.00
999.95
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000303: DM = 23.90 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.20
1000.15
1000.10
1000.05
1000.00
999.95
999.90
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000304: DM = 25.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.6
1000.4
1000.2
1000.0
999.8
999.6
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000305: DM = 25.40 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.25
1000.20
1000.15
1000.10
1000.05
1000.00
999.95
999.90
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000306: DM = 23.50 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.25
1000.20
1000.15
1000.10
1000.05
1000.00
999.95
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000307: DM = 24.70 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.3
1000.2
1000.1
1000.0
999.9
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000308: DM = 25.50 pc cm^{-3}, $N_T = 64$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 225.00$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.20
1000.15
1000.10
1000.05
1000.00
999.95
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000309: DM = 23.80 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.4
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000310: DM = 24.40 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.3
1000.2
1000.1
1000.0
999.9
999.8
999.7
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000311: DM = 23.70 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.25
1000.20
1000.15
1000.10
1000.05
1000.00
999.95
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000312: DM = 24.40 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.30
1000.25
1000.20
1000.15
1000.10
1000.05
1000.00
999.95
999.90
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000313: DM = 25.40 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.3
1000.2
1000.1
1000.0
999.9
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000314: DM = 24.10 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.6
1000.4
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000315: DM = 23.70 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.4
1000.3
1000.2
1000.1
1000.0
999.9
999.8
999.7
999.6
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000316: DM = 25.40 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.3
1000.2
1000.1
1000.0
999.9
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000317: DM = 25.20 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.3
1000.2
1000.1
1000.0
999.9
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000318: DM = 25.30 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.4
1000.3
1000.2
1000.1
1000.0
999.9
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000319: DM = 24.00 pc cm^{-3}, $N_T = 16$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 899.99$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.336
1000.334
1000.332
1000.330
1000.328
1000.326
1000.324
1000.322
1000.320
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000320: DM = 23.50 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.4
1000.3
1000.2
1000.1
1000.0
999.9
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000321: DM = 23.80 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.6
1000.4
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000322: DM = 25.40 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.4
1000.3
1000.2
1000.1
1000.0
999.9
999.8
999.7
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000323: DM = 23.70 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.4
1000.2
1000.0
999.8
999.6
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000324: DM = 25.40 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.4
1000.3
1000.2
1000.1
1000.0
999.9
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000325: DM = 24.00 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.4
1000.3
1000.2
1000.1
1000.0
999.9
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000326: DM = 25.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.6
1000.4
1000.2
1000.0
999.8
999.6
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000327: DM = 25.50 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.6
1000.5
1000.4
1000.3
1000.2
1000.1
1000.0
999.9
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000328: DM = 24.20 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.6
1000.5
1000.4
1000.3
1000.2
1000.1
1000.0
999.9
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000329: DM = 25.40 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.4
1000.3
1000.2
1000.1
1000.0
999.9
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000330: DM = 24.00 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000331: DM = 25.40 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.6
1000.4
1000.2
1000.0
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000332: DM = 24.50 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.6
1000.5
1000.4
1000.3
1000.2
1000.1
1000.0
999.9
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000333: DM = 23.70 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.5
1000.4
1000.3
1000.2
1000.1
1000.0
999.9
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000334: DM = 24.40 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.6
1000.4
1000.2
1000.0
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000335: DM = 24.80 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000336: DM = 23.90 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.6
1000.4
1000.2
1000.0
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000337: DM = 24.00 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.6
1000.4
1000.2
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000338: DM = 23.70 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000339: DM = 24.00 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000340: DM = 23.60 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.7
1000.6
1000.5
1000.4
1000.3
1000.2
1000.1
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000341: DM = 24.20 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.8
1000.6
1000.4
1000.2
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000342: DM = 23.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.0
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000343: DM = 23.90 pc cm^{-3}, $N_T = 128$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 112.50$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.8
1000.6
1000.4
1000.2
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000344: DM = 24.70 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.8
1000.6
1000.4
1000.2
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000345: DM = 24.20 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000.8
1000.6
1000.4
1000.2
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000346: DM = 23.90 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.0
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
999.6
999.4
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000347: DM = 24.30 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.0
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000348: DM = 25.20 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.0
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000349: DM = 25.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.25
1001.00
1000.75
1000.50
1000.25
1000.00
999.75
999.50
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000350: DM = 23.60 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.0
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000351: DM = 23.90 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.2
1001.0
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000352: DM = 25.40 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.2
1001.0
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000353: DM = 24.80 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.25
1001.00
1000.75
1000.50
1000.25
1000.00
999.75
999.50
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000354: DM = 24.30 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.2
1001.0
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000355: DM = 24.00 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.50
1001.25
1001.00
1000.75
1000.50
1000.25
1000.00
999.75
999.50
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000356: DM = 23.60 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.2
1001.0
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000357: DM = 24.70 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.4
1001.2
1001.0
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000358: DM = 24.10 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.4
1001.2
1001.0
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000359: DM = 25.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.50
1001.25
1001.00
1000.75
1000.50
1000.25
1000.00
999.75
999.50
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000360: DM = 23.60 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.4
1001.2
1001.0
1000.8
1000.6
1000.4
1000.2
1000.0
999.8
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000361: DM = 23.80 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.25
1001.00
1000.75
1000.50
1000.25
1000.00
999.75
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000362: DM = 23.80 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000363: DM = 24.70 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.50
1001.25
1001.00
1000.75
1000.50
1000.25
1000.00
999.75
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000364: DM = 24.00 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.50
1001.25
1001.00
1000.75
1000.50
1000.25
1000.00
999.75
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000365: DM = 25.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000366: DM = 23.90 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.50
1001.25
1001.00
1000.75
1000.50
1000.25
1000.00
999.75
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000367: DM = 24.30 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.75
1001.50
1001.25
1001.00
1000.75
1000.50
1000.25
1000.00
999.75
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000368: DM = 23.90 pc cm^{-3}, $N_T = 256$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 56.25$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1001.50
1001.25
1001.00
1000.75
1000.50
1000.25
1000.00
999.75
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000369: DM = 23.70 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000370: DM = 24.20 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000371: DM = 23.60 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.0
1001.5
1001.0
1000.5
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000372: DM = 25.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000373: DM = 23.80 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.0
1001.5
1001.0
1000.5
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000374: DM = 25.20 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000375: DM = 24.00 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.0
1001.5
1001.0
1000.5
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000376: DM = 23.80 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.0
1001.5
1001.0
1000.5
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000377: DM = 24.70 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000378: DM = 25.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000379: DM = 24.00 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000380: DM = 23.60 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000381: DM = 24.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000382: DM = 24.00 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000383: DM = 23.90 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000384: DM = 23.90 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000385: DM = 24.30 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000386: DM = 24.20 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000387: DM = 24.10 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000388: DM = 24.20 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000389: DM = 25.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000390: DM = 23.90 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000391: DM = 23.80 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000392: DM = 23.60 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000393: DM = 24.30 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.5
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000394: DM = 25.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000395: DM = 24.20 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.5
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000396: DM = 24.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000397: DM = 24.10 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.5
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000398: DM = 25.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000399: DM = 24.20 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.5
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000400: DM = 24.70 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.5
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000401: DM = 24.10 pc cm^{-3}, $N_T = 512$
Barycentric; J0024-7204af; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
1001
1002
1003
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000402: DM = 23.90 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.5
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000403: DM = 24.10 pc cm^{-3}, $N_T = 512$
Barycentric; J0024-7204af; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
1001
1002
1003
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000404: DM = 25.20 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1003.5
1003.0
1002.5
1002.0
1001.5
1001.0
1000.5
1000.0
999.5
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000405: DM = 25.40 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1004
1003
1002
1001
1000
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000406: DM = 25.30 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1004
1003
1002
1001
1000
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000407: DM = 25.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1004
1003
1002
1001
1000
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000408: DM = 23.90 pc cm^{-3}, $N_T = 512$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 28.12$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
1001
1002
1003
1004
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000409: DM = 24.20 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1004
1003
1002
1001
1000
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000410: DM = 24.70 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
1001
1002
1003
1004
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000411: DM = 25.50 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1004
1003
1002
1001
1000
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000412: DM = 25.20 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1004
1003
1002
1001
1000
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000413: DM = 25.20 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1000
1001
1002
1003
1004
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)


Candidate 000414: DM = 24.10 pc cm^{-3}, $N_T = 1024$
Barycentric; unclassified; RFI flag: rfi_swarm (retained)
Viterbi path ($T_{coh} = 14.06$ s)
Polynomial fit (order 5)
Circular Kepler fit
Barycentric frequency (Hz)
1004
1003
1002
1001
1000
0
2000
4000
6000
8000
10000
12000
14000
Time from barycentric input epoch (s)