

# Parameter Estimation in Diffusion Processes on the Space of Shapes

Valentina Staneva

joint work with Laurent Younes

Applied Mathematics & Statistics Department  
Center for Imaging Science  
Johns Hopkins University<sup>1</sup>


September 4, 2014

---

<sup>1</sup>This research is partially supported by the ONR

# Introduction

Our goal is to study temporal changes of shapes observed in videos.



We need to:

- construct stochastic processes for modeling the evolution of 2D shapes.
- provide methods for parameter estimation from a sequence of observations
- ensure the solutions are applicable to situations when observations are sparse

- “The diffusion of Shape” (Kendall’77) introduces Brownian motion on the space of landmarks[1]
- extensions to an Ornstein - Uhlenbeck process (Ball’08)[2]
- modeling biological growth by diffeomorphisms (Grenander’06)[3]
- stochastic diffeomorphic flows for shapes (Vialard’12)[4]

# The Geometry of Shape Space

- The *shape manifold*  $\mathcal{M}$  consists of simple closed plane curves  $\gamma$ 
  - here  $\gamma = \{x_1, \dots, x_m \in \mathbb{R}^2\}$ .

# The Geometry of Shape Space

- The *shape manifold*  $\mathcal{M}$  consists of simple closed plane curves  $\gamma$ 
  - here  $\gamma = \{x_1, \dots, x_m \in \mathbb{R}^2\}$ .
- The *tangent space*  $\mathcal{TM}$  contains collections of 2D vectors
  - $\mathbf{v} = \{v_1, \dots, v_m \in \mathbb{R}^2\}$

# The Geometry of Shape Space

- The *shape manifold*  $\mathcal{M}$  consists of simple closed plane curves  $\gamma$ 
  - here  $\gamma = \{x_1, \dots, x_m \in \mathbb{R}^2\}$ .
- The *tangent space*  $\mathcal{TM}$  contains collections of 2D vectors
  - $\mathbf{v} = \{v_1, \dots, v_m \in \mathbb{R}^2\}$
- *Riemannian metric*:  $\|\mathbf{v}\|_\gamma = \mathbf{v} \mathbf{K}(\gamma)^{-1} \mathbf{v}$ 
  - $K(x_i, x_j) = \exp(\|x_i - x_j\|^2 / 2\sigma^2)$

# The Geometry of Shape Space

- The *shape manifold*  $\mathcal{M}$  consists of simple closed plane curves  $\gamma$ 
  - here  $\gamma = \{x_1, \dots, x_m \in \mathbb{R}^2\}$ .
- The *tangent space*  $\mathcal{TM}$  contains collections of 2D vectors
  - $\mathbf{v} = \{v_1, \dots, v_m \in \mathbb{R}^2\}$
- *Riemannian metric*:  $\|\mathbf{v}\|_\gamma = \mathbf{v} \mathbf{K}(\gamma)^{-1} \mathbf{v}$ 
  - $K(x_i, x_j) = \exp(\|x_i - x_j\|^2 / 2\sigma^2)$
- *Exponential map*:  $\gamma_1 = \exp_{\gamma_0}(\mathbf{v})$ 
  - can be computed by solving a system of ODE's

# The Geometry of Shape Space

- The *shape manifold*  $\mathcal{M}$  consists of simple closed plane curves  $\gamma$ 
  - here  $\gamma = \{x_1, \dots, x_m \in \mathbb{R}^2\}$ .
- The *tangent space*  $\mathcal{TM}$  contains collections of 2D vectors
  - $\mathbf{v} = \{v_1, \dots, v_m \in \mathbb{R}^2\}$
- *Riemannian metric*:  $\|\mathbf{v}\|_\gamma = \mathbf{v} \mathbf{K}(\gamma)^{-1} \mathbf{v}$ 
  - $K(x_i, x_j) = \exp(\|x_i - x_j\|^2 / 2\sigma^2)$
- *Exponential map*:  $\gamma_1 = \exp_{\gamma_0}(\mathbf{v})$ 
  - can be computed by solving a system of ODE's
- *Logarithm map*:  $\mathbf{v} = \log_{\gamma_0}(\gamma_1)$ 
  - we assume it exists locally



# The Geometry of Shape Space

- The *shape manifold*  $\mathcal{M}$  consists of simple closed plane curves  $\gamma$ 
  - here  $\gamma = \{x_1, \dots, x_m \in \mathbb{R}^2\}$ .
- The *tangent space*  $\mathcal{TM}$  contains collections of 2D vectors
  - $\mathbf{v} = \{v_1, \dots, v_m \in \mathbb{R}^2\}$
- *Riemannian metric*:  $\|\mathbf{v}\|_\gamma = \mathbf{v} \mathbf{K}(\gamma)^{-1} \mathbf{v}$ 
  - $K(x_i, x_j) = \exp(\|x_i - x_j\|^2 / 2\sigma^2)$
- *Exponential map*:  $\gamma_1 = \exp_{\gamma_0}(\mathbf{v})$ 
  - can be computed by solving a system of ODE's
- *Logarithm map*:  $\mathbf{v} = \log_{\gamma_0}(\gamma_1)$ 
  - we assume it exists locally

**Note:** in our setting we assume  $\mathbf{v}(\cdot) = \sum_{i=1}^n K(x_i, \cdot) \alpha_i$ , where  $\alpha_i \in \mathbb{R}^2$  and  $\{x_i\}_{i=1}^n$  is a subset of the points on  $\gamma$  ( $n < m$ ). This turns  $\mathcal{M}$  into a *sub-Riemannian manifold*.

# Diffusions on Manifolds

Let  $\mathcal{M}$  be a  $m$ -dimensional manifold. Need to define

$$dX_t = A(X_t, \theta)dt + B(X_t)dW_t, \quad X_t \in \mathcal{M}.$$

- Itô and Stratonovich equations can be defined locally in a chart.
- Global formulation needs to take care of transformations between charts.

# Diffusions on Manifolds

Let  $\mathcal{M}$  be a  $m$ -dimensional manifold. Need to define

$$dX_t = A(X_t, \theta)dt + B(X_t)dW_t, \quad X_t \in \mathcal{M}.$$

- Itô and Stratonovich equations can be defined locally in a chart.
- Global formulation needs to take care of transformations between charts.

Let  $\{A, B_1, \dots, B_n\}$  be vector fields on  $\mathcal{M}$ .

## Stratonovich Equations on $\mathcal{M}$

The solution is a process  $X_t$  on  $\mathcal{M}$ , which satisfies for any smooth compactly supported  $f$

$$f(X_T) - f(X_0) = \int_0^T Af(X_t)dt + \int_0^T \sum_{k=1}^n B_k f(X_t) \circ dw_k(t)$$

**Note:** Stratonovich equations transform as tangent vectors!

# Itô Equations on a Manifold

Itô equations transform according to Itô's Rule, not as tangent vectors.

# Itô Equations on a Manifold

Itô equations transform according to Itô's Rule, not as tangent vectors.

$$f(X_T) - f(X_0) = \int_0^T Af(X_t)dt + \int_0^T \sum_{k=1}^n B_k f(X_t) \cdot dw_k(t)$$

# Itô Equations on a Manifold

Itô equations transform according to Itô's Rule, not as tangent vectors.

$$f(X_T) - f(X_0) = \int_0^T Af(X_t)dt + \int_0^T \sum_{k=1}^n B_k f(X_t) \cdot dw_k(t)$$

Instead we can define them through the exponential map.

Itô Equations in Belopolskaya-Daletsky form:

$$dX_t = \exp_{X_t}(A(X_t, \theta)dt + \sum_{k=1}^n B_k \cdot dw_k(t)),$$

# Generating Diffusions

Consider the following vector fields on the shape manifold  $K(\cdot, x_1)\mathbf{e}, \dots, K(\cdot, x_n)\mathbf{e}$ . We can obtain from them orthonormal vector fields:  $E_1, \dots, E_n$ .

Generating Brownian motion on  $\mathcal{M}$

$$X_{t+dt} = \exp_{X_t} \left( \sqrt{dt} \sum_{k=1}^n E_k(X_t) \varepsilon_k \right),$$

where  $\{\varepsilon_k\}_{k=1}^n \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_n)$ .

Constant with respect to the  $\mathbf{K}$  basis

$$dX_t = \sum_{k=1}^n \theta_k K(x_k, X_t) dt + dW_t$$

**Note:** the drift will not be constant wrt to other bases.



# Mean-reverting Drift

Ornstein-Uhlenbeck process on  $\mathbb{R}^n$

$$dX_t = -\theta \nabla_{X_t} [(X_t - \mu)^2] + dW_t$$

Let  $\mu$  be a template shape.

Mean-reverting process on  $\mathcal{M}$

$$dX_t = -\theta \nabla_{X_t} \text{dist}(X_t, \mu) + dW_t$$

Instead of Riemannian distance we use:

$$\text{dist}(X_t, \mu) = \text{area of mismatch of the two shapes}$$



Suppose we don't have  $\mu$  but only estimates for its length  $L_\mu$  and area  $A_\mu$ .

Shape process



$$dX_t = -\frac{1}{2}\theta_1 \nabla_{X_t} |L_{X_t} - L_\mu|^2 - \frac{1}{2}\theta_2 \nabla_{X_t} |A_{X_t} - A_\mu|^2 + dW_t$$

The above gradients can be explicitly computed using the notion of a shape gradient (Zolesio'81)[5].



# Driftless Diffusion





# Constant Drift Diffusion



# Mean-reverting Drift Diffusion



# Length-Area Drift Diffusion



# Parameter Estimation in Diffusions on $\mathbb{R}^n$

Let  $X_t$  be a diffusion process on  $\mathbb{R}^n$ :

$$\begin{aligned}\mu_X : \quad dX_t &= A(X_t, \theta)dt + dW_t \\ \mu_Y : \quad dY_t &= dW_t\end{aligned}$$

## Girsanov Theorem:

Under some boundedness conditions on  $A(X_t, \theta)$  we have  $\mu_X \sim \mu_Y$  and

$$\frac{d\mu_X}{d\mu_Y}(X_t) = \exp \left( \int_0^T A(X_t, \theta)^T \cdot dX_t - \frac{1}{2} \int_0^T A(X_t, \theta)^T A(X_t, \theta) dt \right).$$

Parameter estimate of  $\theta$  can be obtained by maximizing  $\frac{d\mu_X}{d\mu_Y}(X_t)$  wrt  $\theta$ .

# Parameter Estimation in Diffusions on $\mathbb{R}^n$

Let  $X_t$  be a diffusion process on  $\mathbb{R}^n$ :

$$\begin{aligned}\mu_X : \quad dX_t &= A(X_t, \theta)dt + dW_t \\ \mu_Y : \quad dY_t &= dW_t\end{aligned}$$

## Girsanov Theorem:

Under some boundedness conditions on  $A(X_t, \theta)$  we have  $\mu_X \sim \mu_Y$  and

$$\frac{d\mu_X}{d\mu_Y}(X_t) = \exp \left( \int_0^T A(X_t, \theta)^T \cdot dX_t - \frac{1}{2} \int_0^T A(X_t, \theta)^T A(X_t, \theta) dt \right).$$

Parameter estimate of  $\theta$  can be obtained by maximizing  $\frac{d\mu_X}{d\mu_Y}(X_t)$  wrt  $\theta$ .

**Note:** when  $A$  is linear wrt  $\theta$ , the exponent above becomes a quadratic function and there is a closed form solution for  $\theta$ .



# Girsanov Theorem on Manifolds (Elworthy'82)[6]

$$\frac{d\mu_X}{d\mu_Y}(X_t) = \exp \left( \int_0^T \langle A(X_t, \theta), \cdot dX_t \rangle_{X_t} - \frac{1}{2} \int_0^T \langle A(X_t, \theta), A(X_t, \theta) \rangle_{X_t} dt \right),$$

where  $\cdot dX_t$  should be interpreted as  $\log_{X_t}(X_{t+dt})$

# Parameter Estimates

Given a sequence of observations  $X_1, \dots, X_N$  we can approximate the stochastic integrals.

- constant drift:

$$\hat{\theta} = \frac{1}{T} \sum_{i=1}^N (K_{X_i}^{-1/2})^T \log(X_i, X_{i+1})$$





- mean-reverting drift:

$$\hat{\theta} = \frac{\sum_{i=1}^{N-1} \langle \nabla \text{dist}(X_i, \mu), \log(X_i, X_{i+1}) \rangle}{\sum_{i=1}^N \|\nabla \text{dist}(X_i, \mu)\|^2 dt}.$$

- length-area drift

$$\begin{bmatrix} \hat{\theta}_1 \\ \hat{\theta}_2 \end{bmatrix} = \left( \sum_{i=1}^{N-1} M_i dt \right)^{-1} \begin{bmatrix} \sum_{i=1}^{N-1} \langle \nabla |L(X_i) - L_\mu|^2, \log(X_i, X_{i+1}) \rangle \\ \sum_{i=1}^{N-1} \langle \nabla |A(X_i) - A_\mu|^2, \log(X_i, X_{i+1}) \rangle \end{bmatrix},$$

where  $M_i$  as the Grammian matrix of  $\nabla |L(X_i) - L|^2$  and  $\nabla |A(X_i) - A|^2$ .



**Figure:** Top left: constant drift coefficients; top right: distance drift coefficient; bottom left: area drift coefficient; bottom right: length drift coefficient.

# Current and Future Work

- parameter estimation from sparse observations
  - EM formulation to treat the missing observations
  - importance sampling approximation of the expectation step
- including higher order shape terms: e.g. curvature.
- considering the distance to several template shapes
- classification of time series

# References



D. Kendall, "The diffusion of shape," Advances in Applied Probability, vol. 9, no. 3, pp. 428–430, 1977.



F. Ball, I. Dryden, and M. Golalizadeh, "Brownian motion and Ornstein-Uhlenbeck processes in planar shape space," Methodology and Computing in Applied Probability, vol. 10, no. 1, pp. 1–22, 2008.



U. Grenander, A. Srivastava, and S. Saini, "Characterization of biological growth using iterated diffeomorphisms," IEEE International Symposium on Biological Imaging, pp. 1136–1139, 2006.



F.-X. Vialard, "Extension to infinite dimensions of a stochastic second-order model associated with shape splines," Stochastic Processes and their Applications, vol. 123, no. 6, pp. 2110 – 2157, 2013.



J. Zolesio, "The material derivative (or speed) method for shape optimization," Optimization of distributed parameter structures II, N.A.T.O. Asi Series, Series E: Applied Sciences, no. 50, pp. 1089–1151, 1981.



K. Elworthy, Stochastic Differential Equations on Manifolds.