

Curve-constrained Gaussian Random Fields

Valentina Staneva

Applied Mathematics & Statistics Department
Center for Imaging Science
Johns Hopkins University

May 11, 2012

Introduction


We would like to build models for random deformations of curves, which

- have natural geometric properties
- allow for easy approximation

Introduction

We would like to build models for random deformations of curves, which

- 1 select a vector field defined over \mathbb{R}^2
- 2 deform an initial curve along the flow of this vector field




- have natural geometric properties
- allow for easy approximation

Introduction

We would like to build models for random deformations of curves, which

- 1 select a vector field defined over \mathbb{R}^2
- 2 deform an initial curve along the flow of this vector field




- have natural geometric properties
 - select random vector fields which drive neighboring points to move together
- allow for easy approximation

Introduction

We would like to build models for random deformations of curves, which

- 1 select a vector field defined over \mathbb{R}^2
- 2 deform an initial curve along the flow of this vector field



- have natural geometric properties
 - select random vector fields which drive neighboring points to move together
- allow for easy approximation
 - use the structure of reproducing kernel Hilbert spaces (RKHS)

Motivating example: GRF on \mathbb{R}^2

Let $\xi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a centered Gaussian random vector field with covariance

$$C(x, y) = \frac{1}{2\pi\sigma_1^2} e^{-\frac{\|x-y\|_2^2}{2\sigma_1^2}} \mathbb{I}_2, \quad (1)$$

and let $V(C)$ be the associated RKHS with reproducing kernel C .

Motivating example: GRF on \mathbb{R}^2

Let $\xi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a centered Gaussian random vector field with covariance

$$C(x, y) = \frac{1}{2\pi\sigma_1^2} e^{-\frac{\|x-y\|_2^2}{2\sigma_1^2}} \mathbb{I}_2, \quad (1)$$

and let $V(C)$ be the associated RKHS with reproducing kernel C .

Q: Do the realizations of ξ belong to $V(C)$?

Motivating example: GRF on \mathbb{R}^2

Let $\xi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a centered Gaussian random vector field with covariance

$$C(x, y) = \frac{1}{2\pi\sigma_1^2} e^{-\frac{\|x-y\|_2^2}{2\sigma_1^2}} \mathbb{I}_2, \quad (1)$$

and let $V(C)$ be the associated RKHS with reproducing kernel C .

Q: Do the realizations of ξ belong to $V(C)$?

A: No.

Motivating example: GRF on \mathbb{R}^2

Let $\xi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a centered Gaussian random vector field with covariance

$$C(x, y) = \frac{1}{2\pi\sigma_1^2} e^{-\frac{\|x-y\|_2^2}{2\sigma_1^2}} \mathbb{I}_2, \quad (1)$$

and let $V(C)$ be the associated RKHS with reproducing kernel C .

Q: Do the realizations of ξ belong to $V(C)$?

A: No.

Let $V(K)$ be the RKHS with reproducing kernel K

$$K(x, y) = \frac{1}{2\pi\sigma_0^2} e^{-\frac{\|x-y\|_2^2}{2\sigma_0^2}} \mathbb{I}_2. \quad (2)$$

Motivating example: GRF on \mathbb{R}^2

Let $\xi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a centered Gaussian random vector field with covariance

$$C(x, y) = \frac{1}{2\pi\sigma_1^2} e^{-\frac{\|x-y\|_2^2}{2\sigma_1^2}} \mathbb{I}_2, \quad (1)$$

and let $V(C)$ be the associated RKHS with reproducing kernel C .

Q: Do the realizations of ξ belong to $V(C)$?

A: No.

Let $V(K)$ be the RKHS with reproducing kernel K

$$K(x, y) = \frac{1}{2\pi\sigma_0^2} e^{-\frac{\|x-y\|_2^2}{2\sigma_0^2}} \mathbb{I}_2. \quad (2)$$

Q: Do the realizations of ξ belong to $V(K)$?

Motivating example: GRF on \mathbb{R}^2

Let $\xi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a centered Gaussian random vector field with covariance

$$C(x, y) = \frac{1}{2\pi\sigma_1^2} e^{-\frac{\|x-y\|_2^2}{2\sigma_1^2}} \mathbb{I}_2, \quad (1)$$

and let $V(C)$ be the associated RKHS with reproducing kernel C .

Q: Do the realizations of ξ belong to $V(C)$?

A: No.

Let $V(K)$ be the RKHS with reproducing kernel K

$$K(x, y) = \frac{1}{2\pi\sigma_0^2} e^{-\frac{\|x-y\|_2^2}{2\sigma_0^2}} \mathbb{I}_2. \quad (2)$$

Q: Do the realizations of ξ belong to $V(K)$?

A: No.

Conditions¹

Conditions for the realizations to belong to an RKHS (I)

$$P(\xi \in V(K)) = 1 \quad \text{if} \quad \sup_{\chi_n} \text{tr}(C(\chi_n)K(\chi_n)^{-1}) < \infty \quad (3)$$

$$P(\xi \in V(K)) = 0 \quad \text{if} \quad \sup_{\chi_n} \text{tr}(C(\chi_n)K(\chi_n)^{-1}) = \infty \quad (4)$$


Note: when $K = C$, then $\text{tr}(CK^{-1}) = n$, hence the realizations never belong to the RKHS of the covariance.

¹M. Driscoll, The Reproducing Kernel Hilbert Space Structure of the Sample Paths of Gaussian Processes, 1973

Alternative conditions

If $V_\gamma(C) \subset V_\gamma(K)$, then there exists a self-adjoint bounded linear operator G , s.t.

$$GK(\cdot, x) = C(\cdot, x), \quad \forall x \in \gamma. \quad (5)$$



Conditions for the realizations to belong to an RKHS (II)

1 $V_\gamma(C) \subset V_\gamma(K)$

2

$$P(\xi \in V(K)) = 1 \quad \text{if} \quad \text{tr}(G) < \infty \quad (6)$$

$$P(\xi \in V(K)) = 0 \quad \text{if} \quad \text{tr}(G) = \infty \quad (7)$$

Gaussian random fields over \mathbb{R}^2 - revisited

- 1 if $\sigma_1 \geq \sigma_0$, then $V_{\mathbb{R}^2}(C) \subset V_{\mathbb{R}^2}(K)$
- 2 the action of the G operator is

$$G[f] = \text{const} \int_{\mathbb{R}^2} e^{-\frac{\|x-y\|_2^2}{2\sigma_1^2 - 2\sigma_0^2}} f(x) dx \quad (8)$$

$\text{tr}(G) = \infty$, therefore the realizations of ξ are not in $V_{\mathbb{R}^2}(K)$

Curve-constrained Gaussian random vector fields

Idea:

- define a Gaussian random field over a curve γ
- still need a vector field defined over \mathbb{R}^2
- extend to \mathbb{R}^2 by interpolation

RKHS - review

Restriction of an RKHS to a set

$V_\gamma(K)$ consists of functions in $V_{\mathbb{R}^2}(K)$ restricted to γ with a norm

$$\|f\|_{V_\gamma(K)} = \inf_{g \in V(K), \text{ s.t. } g|_\gamma = f} \|g\|_{V(K)}. \quad (9)$$

Note: there is an isometry between functions in $V_\gamma(K)$ and the subspace of functions in $V_{\mathbb{R}^2}(K)$ which minimize the above norm.

We can extend functions in $V_\gamma(K)$ to be defined over \mathbb{R}^2 .

From finite dimension to infinite dimension

Finite dimensional random fields:

$$\xi_n = \sum_{k=1}^n K(\cdot, x_k) \alpha_k, \quad x_k \in \gamma, \quad \alpha_k \sim \mathcal{N}(0, \Sigma) \quad (10)$$

Covariance:

$$\begin{aligned} C_n(x, y) &= \mathbb{E}[\xi_n(x) \xi_n(y)^T] = \mathbb{E}[K(x, \chi_n) \alpha (K(y, \chi_n) \alpha)^T] = \\ &= K(x, \chi_n) \mathbb{E}[\alpha \alpha^T] K(\chi_n, y) = K(x, \chi_n) \Sigma K(\chi_n, y) \end{aligned} \quad (11)$$

How do we extend to infinite dimensions?

Consistency

Need to select Σ so that ξ_n is consistent with projections on lower dimensional subspaces.

A random field $\xi_m \in V_{\chi_m}(K)$ is *consistent* with the random field $\xi_n \in V_{\chi_n}(K)$, where $\chi_n \subset \chi_m$, if its orthogonal projection $\bar{\xi}_m$ onto $V(\chi_n)$ satisfies

$$\text{Cov}(\bar{\xi}_m) = \text{Cov}(\xi_n). \quad (12)$$

Consistency

Need to select Σ so that ξ_n is consistent with projections on lower dimensional subspaces.

A random field $\xi_m \in V_{\chi_m}(K)$ is *consistent* with the random field $\xi_n \in V_{\chi_n}(K)$, where $\chi_n \subset \chi_m$, if its orthogonal projection $\bar{\xi}_m$ onto $V(\chi_n)$ satisfies

$$\text{Cov}(\bar{\xi}_m) = \text{Cov}(\xi_n). \quad (12)$$

Covariance of the coefficients:

$$\Sigma(\chi_n) = K(\chi_n)^{-1} C(\chi_n) K(\chi_n)^{-1}. \quad (13)$$

Consistency

Need to select Σ so that ξ_n is consistent with projections on lower dimensional subspaces.

A random field $\xi_m \in V_{\chi_m}(K)$ is *consistent* with the random field $\xi_n \in V_{\chi_n}(K)$, where $\chi_n \subset \chi_m$, if its orthogonal projection $\bar{\xi}_m$ onto $V(\chi_n)$ satisfies

$$\text{Cov}(\bar{\xi}_m) = \text{Cov}(\xi_n). \quad (12)$$

Covariance of the coefficients:

$$\Sigma(\chi_n) = K(\chi_n)^{-1} C(\chi_n) K(\chi_n)^{-1}. \quad (13)$$

Covariance of the finite dimensional random field:

$$C_n(x, y) = K(x, \chi_n) K(\chi_n)^{-1} C(\chi_n) K(\chi_n)^{-1} K(\chi_n, y). \quad (14)$$

Consistency

Need to select Σ so that ξ_n is consistent with projections on lower dimensional subspaces.

A random field $\xi_m \in V_{\chi_m}(K)$ is *consistent* with the random field $\xi_n \in V_{\chi_n}(K)$, where $\chi_n \subset \chi_m$, if its orthogonal projection $\bar{\xi}_m$ onto $V(\chi_n)$ satisfies

$$\text{Cov}(\bar{\xi}_m) = \text{Cov}(\xi_n). \quad (12)$$

Covariance of the coefficients:

$$\Sigma(\chi_n) = K(\chi_n)^{-1} C(\chi_n) K(\chi_n)^{-1}. \quad (13)$$

Covariance of the finite dimensional random field:

$$C_n(x, y) = K(x, \chi_n) K(\chi_n)^{-1} C(\chi_n) K(\chi_n)^{-1} K(\chi_n, y). \quad (14)$$

Covariance of the infinite dimensional random field (when $\chi_n \rightarrow \gamma$):

$$C_\gamma(x, y) = \langle \pi_{V_\gamma(K)}(K(\cdot, x)), G \pi_{V_\gamma(K)}(K(\cdot, y)) \rangle_{V_\gamma(K)}. \quad (15)$$

Consistency

Need to select Σ so that ξ_n is consistent with projections on lower dimensional subspaces.

A random field $\xi_m \in V_{\chi_m}(K)$ is *consistent* with the random field $\xi_n \in V_{\chi_n}(K)$, where $\chi_n \subset \chi_m$, if its orthogonal projection $\bar{\xi}_m$ onto $V(\chi_n)$ satisfies

$$\text{Cov}(\bar{\xi}_m) = \text{Cov}(\xi_n). \quad (12)$$

Covariance of the coefficients:

$$\Sigma(\chi_n) = K(\chi_n)^{-1} C(\chi_n) K(\chi_n)^{-1}. \quad (13)$$

Covariance of the finite dimensional random field:

$$C_n(x, y) = K(x, \chi_n) K(\chi_n)^{-1} C(\chi_n) K(\chi_n)^{-1} K(\chi_n, y). \quad (14)$$


Covariance of the infinite dimensional random field (when $\chi_n \rightarrow \gamma$):

$$C_\gamma(x, y) = \langle \pi_{V_\gamma(K)}(K(\cdot, x)), G \pi_{V_\gamma(K)}(K(\cdot, y)) \rangle_{V_\gamma(K)}. \quad (15)$$

Note: when $x, y \in \gamma$, $C_\gamma(x, y) = C(x, y)$.

Question

Let ξ_γ be a centered GRF with covariance $C_\gamma(x, y)$.




Q: Do the realizations of ξ_γ belong to $V_\gamma(K)$?

Example: circle

Let γ be a circle.

Steps to calculate the trace:

- 1 pick an orthonormal basis for $V_\gamma(K)$
 - eigenfunctions of $K[f] = \int_\gamma K(x, y)f(x)dx$
- 2 calculate explicitly: spherical harmonics!
- 3 sum the ratio of eigenvalues $\lambda_k(K), \lambda_k(C)$




Example: circle

Let γ be a circle.

Steps to calculate the trace:

- 1 pick an orthonormal basis for $V_\gamma(K)$
- eigenfunctions of $K[f] = \int_\gamma K(x, y)f(x)dx$
- 2 calculate explicitly: spherical harmonics!
- 3 sum the ratio of eigenvalues $\lambda_k(K)$, $\lambda_k(C)$



Trace formula

$$\text{tr}(G) = \sum_{k=0}^{\infty} \frac{\lambda_k(C)}{\lambda_k(K)} = (e^{2/\sigma_1^2 - 2/\sigma_0^2}) \sum_{k=0}^{\infty} \underbrace{\left(\frac{\sigma_0}{\sigma_1}\right)^{2k}}_{<1} \underbrace{\frac{\sum_{l=0}^{\infty} \frac{(1/\sigma_1)^{2l}}{l!(k+l)!}}{\sum_{l=0}^{\infty} \frac{(1/\sigma_0)^{2l}}{l!(k+l)!}}}_{<1} \leq \infty \quad (16)$$

Example: circle

Let γ be a circle.

Steps to calculate the trace:

- 1 pick an orthonormal basis for $V_\gamma(K)$
 - eigenfunctions of $K[f] = \int_\gamma K(x, y)f(x)dx$
- 2 calculate explicitly: spherical harmonics!
- 3 sum the ratio of eigenvalues $\lambda_k(K)$, $\lambda_k(C)$



Trace formula


$$\text{tr}(G) = \sum_{k=0}^{\infty} \frac{\lambda_k(C)}{\lambda_k(K)} = (e^{2/\sigma_1^2 - 2/\sigma_0^2}) \sum_{k=0}^{\infty} \underbrace{\left(\frac{\sigma_0}{\sigma_1}\right)^{2k}}_{<1} \underbrace{\frac{\sum_{l=0}^{\infty} \frac{(1/\sigma_1)^{2l}}{l!(k+l)!}}{\sum_{l=0}^{\infty} \frac{(1/\sigma_0)^{2l}}{l!(k+l)!}}}_{<1} \leq \infty \quad (16)$$

Q: Do the realizations of ξ_γ belong to $V_\gamma(K)$?

A: Yes.


Example: polygonal curve

Let γ be a polygonal curve.



Example: polygonal curve


Let γ be a polygonal curve.



If γ is a line segment, then $V_\gamma = V_{\mathbb{R}^1}$

Example: polygonal curve

Let γ be a polygonal curve.




If γ is a line segment, then $V_\gamma = V_{\mathbb{R}^1}$

Q: Do the realizations of ξ_γ belong to $V_\gamma(K)$?

A: No.


Example: disc

Let γ be a disc.



Example: disc

Let γ be a disc.



Q: Do the realizations of ξ_γ belong to $V_\gamma(K)$?

A: No.

Question

Let ξ_γ be a centered GRF with covariance $\bar{C}_\gamma(x, y)$.



Q: Do the realizations of ξ_γ belong to $V_\gamma(K)$?

Conclusion

- We have defined a model for curve-constrained Gaussian random vector fields.
- We need to establish for what curves the realizations of those random fields belong to the RKHS of interest.
- We can explore other kernels and covariances.