Post-GWAS Analysis

Gene and Functional Annotation

Training Material
Valentina Tsartsianidou
Postdoctoral Researcher
School of Biology, AUTH

22.05.2026

Summarize GWAS steps - 1st Lecture



GWAS results - Association between SNPs and daily sheep grazing duration

Manhattan plot - Genome-wide level





Chromosomes of interest – 3, 7, 11, 14, 19

SNPs identified on chr3 will be used for the functional annotation

Post-GWAS analysis - SNP Annotation

Annotation: adding information or explanation to something

In genomics, **annotation:** adding biological meaning to DNA sequences or genetic markers

SNP annotation: Identification of candidate genes localized within, upstream, or downstream of the associated SNPs within a defined distance (bp)

Post-GWAS analysis - SNP Annotation

We link variants to nearby genes and their functions to biologically interpret GWAS results and prioritize the genomic markers

- **Genomic context**

- Marker position: exonic, intronic, intergenic, UTR, regulatory element, nearest gene

- **Predicted Consequence of variant change**

- synonymous, missense, nonsense mutations

Consequences (most severe)



Consequence type	Count
intron_variant	1
upstream_gene_variant	2
downstream_gene_variant	1

Genome Browsers to explore genomic regions closely located to the identified markers

NCBI Genome Data Viewer



https://www.ncbi.nlm.nih.gov/gdv/browser/genome/?id=GCF_000298735.2

UCSC Genome Browser



https://genome.ucsc.edu/cgi-bin/hgTracks?db=oviAri4&lastVirtModeType=default&lastVirtModeExtraState=&virtModeType=default&virtMode=0&nonVirtPosition=&position=chr3%3A130065882%2D160065881&hgsid=3912947233_JH7X2OATijgk95vyEagS6lQe7dUL

Ensembl Genome Browser

https://www.ensembl.org/Ovis_aries/Location/View?r=3:145487390-145490781;db=core

However, investigating markers individually is challenging in case of multiple significantly associated SNPs

Systematic workflows is needed for large-scale datasets

Workflow Summary



1. LiftOver Genome Coordinates between different assemblies

widely used options to convert genome coordinates in another assembly

Latest | Open | Jobs | Tutorials | Tags | About | FAQ

Biostars BIOINFORMATICS EXPLAINED — Community | Planet | New Post

The Biostar Handbook
A practical guide to modern bioinformatics
CLICK HERE

Tool:Converting Genome Coordinates From One Genome Version To Another (Ucsc Liftover, Ncbi Remap, Ensembl Api)

195

Some recent posts reminded me that it might be useful for us to review the options for converting between genome coordinate systems.

This comes up in several contexts. Probably the most common is that you have some coordinates for a particular version of a reference genome and you want to determine the corresponding coordinates on a different version of the reference genome for that species. For example, you have a bed file with exon coordinates for human build GRC37 (hg19) and wish to update to GRCh38. By the way, for a nice summary of genome versions and their release names refer to the Assembly Releases and Versions FAQ

Or perhaps you have coordinates of a gene and wish to determine the corresponding coordinates in another species. For example, you have coordinates of a gene in human GRCh38 and wish to determine corresponding coordinates in mouse mm10.

13.1 years ago
Malachi Griffith 20k

Finally you may wish to convert coordinates between coordinate systems within a single assembly. For example, you have the coordinates of a series of exons and you want to determine the position of these exons with respect to the transcript, gene, contig, or entire chromosome.

There are now several well known tools that can help you with these kinds of tasks:

1. **UCSC liftOver**. This tool is available through a simple web interface or it can be downloaded as a standalone executable. To use the executable you will also need to download the appropriate chain file. Each chain file describes conversions between a pair of genome assemblies. Liftover can be used through Galaxy as well. There is a python implementation of liftover called pyliftover that does conversion of point coordinates only.

2. **NCBI Remap**. This tool is conceptually similar to liftOver in that in manages conversions between a pair of genome assemblies but it uses different methods to achieve these mappings. It is also available through a simple web interface or you can use the API for NCBI Remap.

3. **The Ensembl API**. The final example I described above (converting between coordinate systems within a single genome assembly) can be accomplished with the Ensembl core API. Many examples are provided within the installation, overview, tutorial and documentation sections of the Ensembl API project. In particular, refer to these sections of the tutorial: 'Coordinates', 'Coordinate systems', 'Transform', and 'Transfer'.

4. **Assembly Converter**. Ensembl also offers their own simple web interface for coordinate conversions called the Assembly Converter.

5. **Bioconductor rtracklayer** package. For R users, Bioconductor has an implementation of UCSC liftOver in the rtracklayer package. To see documentation on how to use it, open an R session and run the following commands.

```
source("http://bioconductor.org/biocLite.R")
biocLite("rtracklayer")
library(rtracklayer)
?liftOver
```

6. **CrossMap**. A standalone open source program for convenient conversion of genome coordinates (or annotation files) between different assemblies. It supports most commonly used file formats including SAM/BAM, Wiggle/BigWig, BED, GFF/GTF, VCF. CrossMap is designed to liftover genome coordinates between assemblies. It's not a program for aligning sequences to reference genome. Not recommended for converting genome coordinates between species.

7. Flo. A liftover pipeline for different reference genome builds of the same species. It describes the process as follows: "align the new assembly with the old one, process the alignment data to define how a coordinate or coordinate range on the old assembly should be transformed to the new assembly, transform the coordinates."

8. Picard Liftover VCF. Lifts over a VCF file from one reference build to another. This tool adjusts the coordinates of variants within a VCF file to match a new reference. The tool is based on the UCSC liftOver and uses a UCSC chain file to guide its operation.

https://www.biostars.org/p/65558/#391077

CrossMap
Installed in Aristotle cluster in python virtual environment

1. LiftOver Genome Coordinates between different assemblies

Input file: BED, VCF, GFF

File format →

chr3	189803626	189803626	OAR3_204624914.1
chr3	145065882	145065882	OAR3_155103999.1
chr3	131220126	131220126	OAR3_140271376.1
chr3	181785624	181785624	OAR3_195793605_X.1

Chain file

text file with pairwise alignment between different assemblies in continuous segment blocks defining also the gaps in both target and query genomic sequences

↓

Chain file format

alignment score | Target chr | Chr size | strand | Region in target genome | Query chr | Chr size | strand | Region in query genome | Chain ID

```
chain 4900 chrY 58368225 + 25985406 25985566 chr5 151006098 - 43549808 43549970 2
    16        0        2
    60        4        0
    10        0        4
    703
```

Gap in target | Gap in query

separate alignment blocks separated due to presence of gaps or differences



More info: https://crossmap.sourceforge.net/

Chain files for sheep genome provided by UCSC Genomics Institute

↓

https://groups.google.com/a/soe.ucsc.edu/g/genome/c/2BSf_4qAyZY

1. LiftOver Genome Coordinates between different assemblies

How do we trace the genome built that our data correspond to?

1. Use a variant ID (rs ID) in a genome browser that is based on a specific genome built of the studied species

2. If the position does not match the map file, use online liftover tool to determine the assembly of your data



Our data correspond to the *Ovis aries* genome assembly Oar_v4.0 → Texel breed - 2015

Conversion of genome coordinates using ARS-UI_Ramb_v2.0 in-depth functional annotation of the sheep genome (https://academic.oup.com/gigascience/article/doi/10.1093/gigascience/giab096/6521876) → Rambouillet breed - 2022

2. Annotate significantly associated SNPs

Ensembl VEP (Variant Effect Predictor) – Variant Annotation (SNPs, insertions, deletions, structural variants)

1 Input file: a list with the identified variants

VEP default input →

chromosome
start position
end position
allele
strand

https://www.ensembl.org/info/docs/tools/vep/vep_formats.html#default

Daily grazing duration

3 190036206 190036206 A/G
3 145252109 145252109 A/G
3 131380348 131380348 G/A
3 181785624 181785624 G/A
7 62298200 62298200 A/C
7 4217464 4217464 A/G
11 26623188 26623188 A/C
11 29347981 29347981 A/G
11 18248852 18248852 A/G
11 49829927 49829927 C/A
14 39858010 39858010 A/G
14 39795104 39795104 A/G
19 10373068 10373068 G/A

Altitude difference

15 71858906 71858906 G/A
15 63232034 63232034 G/A
15 3499482 3499482 G/A
7 7334440 7334440 C/A
3 18569142 18569142 A/C



https://www.ensembl.org/Multi/Tools/VEP?db=core

2. Annotate significantly associated SNPs

Ensembl VEP (Variant Effect Predictor) – Variant Annotation

2

-Cache file = Ensembl annotation source: **refers to a specific genome built**

- File with all transcripts, regulatory features and variant data for a species
- **It should be compatible to the VEP version installed**
  https://www.ensembl.org/info/docs/tools/vep/script/vep_cache.html#cache

Examples and use cases
Other information

On this page
Caches
GFF/GTF files
FASTA files
Databases
Cache - technical information

Search documentation… Go

- Cache - a downloadable file containing all transcript models, regulatory features and variant data for a species
- GFF or GTF - use transcript models defined in a tabix-indexed GFF or GTF file
  - Requires a FASTA file in --offline mode or if the desired species or assembly is not part of the Ensembl species list.
- Database - connect to a MySQL database server hosting Ensembl databases

Data from VCF, BED and bigWig files can also be incorporated by Ensembl VEP's Custom annotation feature.

Using a cache is the most efficient way to use Ensembl VEP; we would encourage you to use a cache wherever possible. Caches are easy to download and set up using the installer. Follow the tutorial for a simple guide.

Caches

Using a cache (--cache) is the fastest and most efficient way to use Ensembl VEP, as in most cases only a single initial network connection is made and most data is read from local disk. Use offline mode to eliminate all network connections for speed and/or privacy.

⚠ Cache version

We strongly recommend that you download/use the cache version which corresponds to your Ensembl VEP installation, i.e. cache version **115** should be used with the Ensembl VEP tool version **115**.

This is mainly due to the fact that the cache (data content and structure) is generated every Ensembl release, regarding the data and API updates for this release, therefore the cache data format might differ between versions (and be incompatible with a newer version of the Ensembl VEP tool).

Downloading caches

Cache files are created for every species for each Ensembl release. They can be automatically downloaded and configured using INSTALL.pl.

If interested in RefSeq transcripts you may download an alternate cache file (e.g. homo_sapiens_refseq), or a merged file of RefSeq and Ensembl transcripts (eg homo_sapiens_merged); remember to specify --refseq or --merged when running Ensembl VEP to use the relevant cache. See documentation for full details.

Manually downloading caches

It is also simple to download and set up caches without using the installer. By default, Ensembl VEP searches for caches in $HOME/.vep; to use a different directory when running Ensembl VEP, use --dir_cache.

Indexed cache (https://ftp.ensembl.org/pub/release-115/variation/indexed_vep_cache/)

Essential for human and other species with large sets of variant data - requires Bio::DB::HTS (setup by INSTALL.pl) or tabix, e.g.:

```
cd $HOME/.vep
curl -O https://ftp.ensembl.org/pub/release-115/variation/indexed_vep_cache/homo_sapiens_vep_115_GRCh38.tar.gz
tar xzf homo_sapiens_vep_115_GRCh38.tar.gz
```

FTP directories with indexed cache data:

Ensembl:	Vertebrates
Ensembl Genomes:	Bacteria | Fungi | Metazoa | Plants | Protists

Please use the web portal with 1 core per session to install and run VEP!

2. Annotate significantly associated SNPs

Example Output – VEP

Gene Ensembl ID
Consequence
CDS position
Distance from SNP

```
## Codons : Reference and variant codon sequence
## Existing_variation : Identifier(s) of co-located known variants
## Extra column keys:
## IMPACT : Subjective impact classification of consequence type
## DISTANCE : Shortest distance from variant to transcript
## STRAND : Strand of the feature (1/-1)
## FLAGS : Transcript quality flags
## VEP command-line: vep --cache --database 0 --distance 500000 --input_file input_vep.Ensembl --output_file output_vep.Ensembl --species ovis_aries
#Uploaded_variation     Location        Allele  Gene    Feature Feature_type    Consequence     cDNA_position   CDS_position    Protein_position        Amino_acids     Codons  Existing_variation      Extra
3_190834971_A/G 3:190834971     G       ENSOARG00020039732      ENSOART00020005659      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=409525;STRAND=-1
3_190834971_A/G 3:190834971     G       ENSOARG00020003767      ENSOART00020006508      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=319939;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020003767      ENSOART00020006599      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=320099;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020003767      ENSOART00020006708      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=320098;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020003767      ENSOART00020006726      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=319936;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020003767      ENSOART00020006747      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=319939;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020004466      ENSOART00020006859      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=265242;STRAND=-1
3_190834971_A/G 3:190834971     G       ENSOARG00020004542      ENSOART00020007341      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=43693;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020004542      ENSOART00020007365      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=43693;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020004542      ENSOART00020007387      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=43693;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020004542      ENSOART00020007433      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=55413;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020004542      ENSOART00020007461      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=43687;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020003767      ENSOART00020007507      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=319936;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020003767      ENSOART00020043009      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=320096;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020004542      ENSOART00020044492      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=42871;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020004542      ENSOART00020045070      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=42871;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020030797      ENSOART00020049895      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=291786;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020034889      ENSOART00020052086      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=226257;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020004466      ENSOART00020053005      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=278348;STRAND=-1
3_190834971_A/G 3:190834971     G       ENSOARG00020004466      ENSOART00020059129      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=216518;STRAND=-1
3_190834971_A/G 3:190834971     G       ENSOARG00020004466      ENSOART00020060262      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=278348;STRAND=-1
3_190834971_A/G 3:190834971     G       ENSOARG00020004466      ENSOART00020060954      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=249387;STRAND=-1
3_190834971_A/G 3:190834971     G       ENSOARG00020040754      ENSOART00020062327      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=417103;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020040754      ENSOART00020063707      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=417111;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020004542      ENSOART00020066162      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=42871;STRAND=1
3_190834971_A/G 3:190834971     G       ENSOARG00020039732      ENSOART00020067972      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=409676;STRAND=-1
3_190834971_A/G 3:190834971     G       ENSOARG00020004466      ENSOART00020068667      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=227631;STRAND=-1
3_190834971_A/G 3:190834971     G       ENSOARG00020036459      ENSOART00020070106      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=31970;STRAND=-1
3_190834971_A/G 3:190834971     G       ENSOARG00020004466      ENSOART00020070693      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=278348;STRAND=-1
3_190834971_A/G 3:190834971     G       ENSOARG00020004466      ENSOART00020071286      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=216518;STRAND=-1
3_145489075_A/G 3:145489075     G       ENSOARG00020000431      ENSOART00020000557      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=28833;STRAND=-1
3_145489075_A/G 3:145489075     G       ENSOARG00020000431      ENSOART00020000571      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=29663;STRAND=-1
3_145489075_A/G 3:145489075     G       ENSOARG00020000431      ENSOART00020059714      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=28833;STRAND=-1
3_145489075_A/G 3:145489075     G       ENSOARG00020000431      ENSOART00020076609      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=28833;STRAND=-1
3_131527909_G/A 3:131527909     A       ENSOARG00020014046      ENSOART00020011577      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=288466;STRAND=-1
3_131527909_G/A 3:131527909     A       ENSOARG00020013894      ENSOART00020021288      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=305238;STRAND=-1
3_131527909_G/A 3:131527909     A       ENSOARG00020013894      ENSOART00020021355      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=305169;STRAND=-1
3_131527909_G/A 3:131527909     A       ENSOARG00020014046      ENSOART00020021482      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=258124;STRAND=-1
3_131527909_G/A 3:131527909     A       ENSOARG00020014046      ENSOART00020021576      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=266310;STRAND=-1
3_131527909_G/A 3:131527909     A       ENSOARG00020014046      ENSOART00020021657      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=258124;STRAND=-1
3_131527909_G/A 3:131527909     A       ENSOARG00020014249      ENSOART00020021831      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=145132;STRAND=-1
3_131527909_G/A 3:131527909     A       ENSOARG00020014442      ENSOART00020022076      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=65519;STRAND=1
3_131527909_G/A 3:131527909     A       ENSOARG00020014571      ENSOART00020022295      Transcript      downstream_gene_variant -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=61279;STRAND=1
3_131527909_G/A 3:131527909     A       ENSOARG00020014635      ENSOART00020023247      Transcript      upstream_gene_variant   -       -       -       -       -       -       IMPACT=MODIFIER;DISTANCE=88214;STRAND=1
3_131527909_G/A 3:131527909     A       ENSOARG00020014635      ENSOART00020023352      Transcript      upstream_gene_variant                                                   IMPACT=MODIFIER;DISTANCE=88214;STRAND=1
```

3. Functional Annotation

Process of determining gene functions, pathways associated with specific sequences/genes

Gene Ontology (GO) = the widest annotation source of gene products

standardized system to classify gene functions and their products across species in a consistent way with structured vocabulary

GO Has Three Main Categories-Sources:

- **Biological Process (BP)**
  What the gene product does in a wide biological context.
  e.g., cell division, immune response, signal transduction.
- **Molecular Function (MF)**
  The basic activity of the gene product at the molecular level. e.g., ATP binding, enzyme activity, receptor binding.
- **Cellular Component (CC)**
  Where in the cell the gene product is active.
  e.g., nucleus, mitochondrion, plasma membrane.

Types of functional annotation (GO Evidence Codes)

How the gene function was determined

Code	Meaning	
EXP	Inferred from Experiment	experiment-based
IDA	Inferred from Direct Assay	
IEA	Inferred from Electronic Annotation	predicted
ISS	Inferred from Sequence or Structural Similarity	
TAS	Traceable Author Statement	Literature-based
IMP	Inferred from Mutant Phenotype	

3. Functional Enrichment Annotation

definition

A statistical test to determine whether certain functions/pathways are overrepresented in a defined set of genes compared to what is expected by chance

steps

- Gene list as an input
- Select a reference genome including all annotated genes
- Test each annotation term (GO term) for enrichment
- Correct for multiple testing (e.g., Bonferroni, FDR)

3. Functional Enrichment Analysis

g:Profiler2 (R package) - toolset for functional enrichment analysis and visualization

https://pmc.ncbi.nlm.nih.gov/articles/PMC7859841/pdf/f1000research-9-29655.pdf

- Utilizes data from multiple reliable databases GO, KEGG, Reactome etc. for functional annotation across various species
- Input: **gene list** obtained from VEP tool
- Format: Ensembl gene IDs (preferred), Gene symbols
- **Select the organism**

Main functions
gost(): functional profiling
gostplot(): plot the enrichment results
gconvert(): naming conventions between genes, proteins etc across databases and species
gorth(): convert a gene list from a source organism to a target one

https://biit.cs.ut.ee/gprofiler/page/r

3. Functional Enrichment Analysis

Utilization of gprofiler2 (R package)

Install and load gprofiler2 library in RStudio

1. Example input (single query)

```
</> R
gostres <- gost(query = c("X:1000:1000000", "rs17396340", "GO:0005005", "ENSG00000156103", "NLRP1"), organism = "hsapiens")
```

2. Functional annotation from GWAS-derived gene lists (VEP tool results for two SNPs)

Create a named list object in R including two gene lists (Ensembl gene IDs) that correspond to two SNP markers

```
</> R
gene_lists <- list(…)
```

Run functional enrichment analysis - find the argument for multiple gene lists (search in R documentation), optional parameters for significance, threshold, correction)

```
</> R
gostres <- gost(query = gene_lists, organism = "oarambouillet", …)
```

3. Functional Enrichment Results

1. Example input (single query)

</> **R**

```
head(gostres$results)
```

```
 query significant      p_value term_size query_size intersection_size precision
1 query_1        TRUE 4.998799e-02         1          3                 1 0.3333333
2 query_1        TRUE 2.122758e-30        51         20                14 0.7000000
3 query_1        TRUE 4.197869e-18       238         20                13 0.6500000
4 query_1        TRUE 4.437953e-18       239         20                13 0.6500000
5 query_1        TRUE 2.113213e-14       629         20                14 0.7000000
6 query_1        TRUE 2.633096e-14       463         20                13 0.6500000
      recall    term_id source
1 1.00000000 CORUM:6180  CORUM
2 0.27450980 GO:0048013  GO:BP
3 0.05462185 GO:0007411  GO:BP
4 0.05439331 GO:0097485  GO:BP
5 0.02225755 GO:0007169  GO:BP
6 0.02807775 GO:0007409  GO:BP
                                                   term_name effective_domain_size
1                                     PPP2R1A-PPP2R3B complex                 3383
2                                ephrin receptor signaling pathway               21026
3                                                 axon guidance               21026
4                                       neuron projection guidance               21026
5 cell surface receptor protein tyrosine kinase signaling pathway               21026
6                                                 axonogenesis               21026
  source_order                          parents
1         1952                      CORUM:0000000
2        12302                         GO:0007169
3         2892              GO:0007409, GO:0097485
4        18430              GO:0031175, GO:0048812
5         2720                         GO:0007167
6         2891 GO:0048667, GO:0048812, GO:0061564
```

[Explore the query result fields \https://biit.cs.ut.ee/gprofiler/page/apis](https://biit.cs.ut.ee/gprofiler/page/apis)

3. Functional Enrichment Results

2. Functional annotation from GWAS-derived gene lists (VEP tool results for two SNPs)

</> **R**

```
head(gostres$results)
```

```
 query significant      p_value term_size query_size intersection_size precision      recall    term_id source
1 Region1        TRUE 0.0009082825       173         34                 6 0.1764706 0.034682081 GO:0009952  GO:BP
2 Region1        TRUE 0.0164523122       324         34                 6 0.1764706 0.018518519 GO:0003002  GO:BP
3 Region1        TRUE 0.0170941552       398         34                 6 0.1764706 0.015075377 GO:0001501  GO:BP
4 Region1        TRUE 0.0170941552       353         34                 6 0.1764706 0.016997167 GO:0007389  GO:BP
5 Region1        TRUE 0.0170941552      3830         34                18 0.5294118 0.004699739 GO:0010556  GO:BP
6 Region1        TRUE 0.0170941552      3776         34                18 0.5294118 0.004766949 GO:0010468  GO:BP
                                         term_name effective_domain_size source_order                           parents
1         anterior/posterior pattern specification                 16882         3629                         GO:0003002
2                                  regionalization                 16882         1407                         GO:0007389
3                      skeletal system development                 16882          306                         GO:0048731
4                    pattern specification process                 16882         2753             GO:0007275, GO:0032501
5 regulation of macromolecule biosynthetic process                 16882         4023 GO:0009059, GO:0009889, GO:0060255
6                    regulation of gene expression                 16882         3977             GO:0010467, GO:0010556
```

Multi_query=FALSE

What is it enriched in each gene list?

```
 query significant    p_value term_size query_size intersection_size  precision      recall    term_id source
28 Region1        TRUE 0.04588738        72         29                 2 0.06896552 0.027777778 GO:0003730  GO:MF
29 Region1        TRUE 0.04588738       990         29                 6 0.20689655 0.006060606 GO:0008270  GO:MF
30 Region2        TRUE 0.03452436         2         12                 1 0.08333333 0.500000000 GO:0052655  GO:MF
31 Region2        TRUE 0.03452436         2         12                 1 0.08333333 0.500000000 GO:0004084  GO:MF
32 Region2        TRUE 0.03452436         2         12                 1 0.08333333 0.500000000 GO:0052654  GO:MF
33 Region2        TRUE 0.03452436         2         12                 1 0.08333333 0.500000000 GO:0052656  GO:MF
                                         term_name effective_domain_size source_order    parents
28                              mRNA 3'-UTR binding                 16157          339 GO:0003729
29                                 zinc ion binding                 16157         1600 GO:0046914
30     L-valine-2-oxoglutarate transaminase activity                 16157         7464 GO:0004084
31   branched-chain-amino-acid transaminase activity                 16157          553 GO:0008483
32    L-leucine-2-oxoglutarate transaminase activity                 16157         7463 GO:0004084
33 L-isoleucine-2-oxoglutarate transaminase activity                 16157         7465 GO:0004084
```

Explore the query result fields \https://biit.cs.ut.ee/gprofiler/page/apis

3. Functional Enrichment Results

2. Functional annotation from GWAS-derived gene lists (VEP tool results for two SNPs)

</> **R**

```
head(gostres$results)
```

```
 term_id                   p_values significant term_size query_sizes intersection_sizes source
1 GO:0009952 0.0009082825, 1.0000000000 TRUE, FALSE       173      34, -1              6, -1  GO:BP
2 GO:0003700     0.01021126, 1.00000000 TRUE, FALSE       860      29, -1              8, -1  GO:MF
3 GO:0140110     0.01021126, 1.00000000 TRUE, FALSE      1206      29, -1              9, -1  GO:MF
4 GO:0000981     0.01021126, 1.00000000 TRUE, FALSE       760      29, -1              7, -1  GO:MF
5 GO:0000977     0.01021126, 1.00000000 TRUE, FALSE       726      29, -1              7, -1  GO:MF
6 GO:0000976     0.01047532, 0.52532478 TRUE, FALSE       815      29, 12               7, 1  GO:MF
                                                                    term_name effective_domain_size
1                                       anterior/posterior pattern specification                 16882
2                                      DNA-binding transcription factor activity                 16157
3                                               transcription regulator activity                 16157
4           DNA-binding transcription factor activity, RNA polymerase II-specific                 16157
5 RNA polymerase II transcription regulatory region sequence-specific DNA binding                 16157
6                                     transcription cis-regulatory region binding                 16157
  source_order                parents
1         3629             GO:0003002
2          326             GO:0140110
3         9201             GO:0003674
4          117             GO:0003700
5          114             GO:0000976
6          113 GO:0001067, GO:1990837
```

Multi_query=TRUE

Explore the query result fields
\https://biit.cs.ut.ee/gprofiler/page/apis

We compare the gene lists term by term to check how are lists differ for the same GO terms

3. Functional Enrichment Vizualization

Utilization of gprofiler2 (R package)

1. Explore functional enrichment results through an interactive Manhattan plot

```
</>  R

gostplot(gost_output, capped=TRUE, interactive=TRUE)
```

2. Select specific GO terms for final visualization, set interactive=FALSE and then proceed with the following command

3. Visualize enrichment results selecting specific GO terms from both annotated gene lists using the interactive Manhattan plot

```
</>  R

publish_gostplot(gostplot_output, highlight_terms= ...)
```

Functional Enrichment Results

Manhattan plot of enriched GO terms using 2 gene lists (chr3) as an input with multi_query=TRUE



id	source	term_id	term_name	term_size		p_value Region1		p_value Region2
1	GO:MF	GO:0051020	GTPase binding	201		1.0e+00		4.6e-02
2	GO:BP	GO:0001501	skeletal system development	398		3.0e-02		2.8e-01
3	GO:BP	GO:0009952	anterior/posterior pattern specification	171		9.9e-04		1.0e+00

g:Profiler (biit.cs.ut.ee/gprofiler)

Functional Enrichment - Vizualization

Usage of alternative R packages for visualization purposes using the output from gprofiler2 gost()

clusterProfiler
enrichplot
DOSE
ggplot2

Specific columns are required from the data frame produced by gost(), such as

query,
source,
term_id,
term_name,
p_value,
query_size,
intersection_size,
term_size,
effective_domain_size,
intersection

Functional Enrichment Results

Bar plot of enriched GO terms showing number of candidate genes in each biological process detected



barplot()

Functional Enrichment Results

Dot plot of enriched GO terms showing number of candidate genes in each biological process detected



enrichplot()

Use the link to access the material for annotation workflow

https://github.com/valentinatsar/annotation/tree/main