

Introduction

Methods

A Framework for Adversarial Collaboration

Methods of Individual Simulation Studies

Studies by Kriegmair

The methodological setup of my individual simulation studies follows the structure we established for our *adversarial simulation* framework to facilitate stepwise collaboration. In the initial phase of our case study, I independently conducted two separate simulation studies without my collaborator's involvement with the goal to conceptually replicate the findings regarding SAM compared to standard SEM estimation of Rosseel & Loh (2022) and Dhaene & Rosseel (2023). However, there are several differences in the design and setup of the studies compared to the original studies as outlined below.

Aims, objectives and research questions

Both studies aimed to evaluate the performance of vanilla SEM (with maximum likelihood) compared to global SAM (gSAM), local SAM with maximum likelihood (lSAM-ML), and local SAM with unweighted least squares (lSAM-ULS) under various conditions. The two research questions we jointly established prior to conducting the studies served as general basis for both studies:

1. How do SAM and traditional SEM methods (including ML and ULS) compare in terms of bias, Mean Squared Error (MSE), and convergence rates in small to moderate samples?
2. What is the impact of model misspecifications, such as residual correlations and cross-loadings, on the performance of SAM compared to traditional SEM methods?

Population Models and Data Generation Mechanisms

Study 1

Data were generated based on a 5-factor population structural model with 3 indicators for each factor. Four different models were simulated (see figure 1-4). In line with Rosseel & Loh (2022) this model design was chosen to represent a realistic model with sufficient complexity to pose a challenge for the estimation methods, especially in the presence of misspecifications:

- Model 1.1: Correctly specified model.
- Model 1.2: Misspecified with cross-loadings in the population model that are ignored in the estimation model (model 1.1)
- Model 1.3: Misspecified with correlated residuals and a reversed structural path between the third and fourth latent factors in the population model that are ignored in the estimation model (model 1.1)
- Model 1.4: Misspecified with a bidirectional structural relation between factors 3 and 4 specified as only one directional

Factor loadings were fixed across all reliability conditions, with the first indicator of each factor serving as the scaling indicator ($\lambda = 1.0$), and the other two indicators having loadings of 0.7. Indicator reliability levels were manipulated by adjusting the measurement error variances in the Θ matrix. Specifically, the a reliability value was set at different levels (low = 0.3, moderate = 0.5 or high = 0.7) to compute the respective error variances on the diagonal of Θ : $\Theta^* = \text{Var}(\eta)\Lambda^T \times \frac{1}{r-1}$.


Deviation from Rosseel & Loh (2022):

To investigate additioanl possible and realisitic scenarios beyond the ones studied by Rosseel & Loh (2022) model 1.3 included a combination of measuremnt and structural misspecifications as opposed to only measurement misspecifications to introduce an even more severely misspecified model under which SAM methods might perform even better than traditional SEM. Further, model 1.4 included a (not estimated) bidirectional structural relation between factors 3 and 4 as opposed to the unidirectional reversed one. For all models, the population-level values of the structural parameters were set to 0.1.

Study 2:

Figure 1.

Population Model Variations of Study 1



Note. Error terms are not explicitly shown in the figure. Dashed lines represent relations omitted in the estimation model present in the population model.”

Data were generated based on a 5-factor population structural model with 3 indicators for each factor with loadings set to 1, 0.9 and 0.8 for each factor and reliability modulated like in study 1. Regression weights were set to either 0.183 and 0.224 (low) or 0.365 and 0.447 (medium). This should represent varying variance explained (R^2) by the endogenous factors set at low ($R^2 = 0.1$) or medium ($R^2 = 0.4$). Note however that the computation of this was a simplification and does not accurately result in said R^2 values. The aim here was only to generally modulate between lower and higher regression weights. The population models resulted in the following model types with varying misspecification in the estimation model:

- Model 2.1: structural misspecification with falsely specified paths in the estimation model absent in the population model (Figure 5)
- Model 2.2: correlated residuals and a factor cross-loading in either the exogenous (Model 2.2-exo), endogenous (Model 2.2-endo) part of the model or both (Model 2.2-both) with falsely specified paths in the estimation model absent in the population model (see figure 5-6).

Deviation from Dhaene & Rosseel (2023):

To enable the analysis of the impact of falsely specified paths in the estimation model that are not present in the population model and how well the different methods recover these non-existing relations both population models included several such misspecifications in addition to the measurement misspecifications evaluated by Dhaene & Rosseel (2023).

Experimental Design of simulation procedures


Study 1

Study 1 varied three main conditions: (1) sample sizes of small ($N = 100$), moderate ($N = 400$), and large ($N = 6400$); (2) Indicator reliability of low ($= 0.3$), moderate (0.5), high ($= 0.7$); (3) Model specifications: correctly specified model and misspecified with not specified cross loadings in the population model (see figure 2), misspecified with not-specified correlated residuals and a reversed structural path between the the third and the fourth latent factor in the population model (see figure 3) and a recursive structural relation between factor 3 and 4 in the population specified as only one directional (see figure 4).

Study 2

Figure 2.

Population Model Variations of Study 2



Note. Error terms are not explicitly shown in the figure. Dotted paths represent relations specified in the estimation model not present in the population model. For Model 2.2, orange lines represent misspecifications in the exogenous part of the model, and green lines represent misspecifications in the endogenous part of the model. These types of misspecifications result in different realizations of model 2.2 when they are modulated as factors in study 2 but are subsumed under one model here.

Study 2 varied five conditions: (1) sample sizes: small ($N = 100$), medium ($N = 400$), and large ($N = 6400$). (2) Variance explained by endogenous factors: low ($R^2 = 0.1$) and medium ($R^2 = 0.4$). (3) Indicator reliability: low (0.3), moderate (0.5), and high (0.7). (4) Model misspecifications: varying the population model by omitting a residual covariance and a factor cross-loading in different parts of the model. (5) Number of measurement blocks: separate measurement model per latent variable ($b = 5$) and joint measurement model for all exogenous variables ($b = 3$) for the local SAM condition (lSAM-ML).

Method Selection

Both studies compared the performance of four estimation methods: Vanilla SEM with maximum likelihood (ML), Global SAM with maximum likelihood (gSAM), Local SAM with maximum likelihood (lSAM-ML), Local SAM with unweighted least squares (lSAM-ULS).

Performance Measures

For both studies convergence rates were tracked via lavaan’s built-in function that indicates convergence. Further, improper solutions, converged models that showed negative variances (as the only type of improper solution present), were tracked via lavaan warning messages. Next of all converged and proper solutions bias ($\bar{T} - \theta$), and RMSE ($\sqrt{\frac{1}{K} \sum_{k=1}^K (T_k - \theta)^2}$) where T_k is the estimated parameter, \bar{T} the mean of the estimated parameters and θ the true parameter value, and K is the number of replications computed. For comparability across varying regression weights for study 2, relative bias ($\frac{\bar{T} - \theta}{\theta}$) and relative RMSE ($\sqrt{\frac{(\bar{T} - \theta)^2 + S_T^2}{\theta^2}}$) were computed. Monte Carlo standard errors (MCSE) were computed for bias and RMSE as well as relative bias and relative RMSE: $\sqrt{\frac{S_T^2}{K}}$ and $\sqrt{\frac{S_T^2}{K\theta^2}}$ for bias and relative bias, and $\sqrt{\frac{K-1}{K} \sum_{j=1}^K (\text{RMSE}_{(j)} - \text{RMSE})^2}$ and $\sqrt{\frac{K-1}{K} \sum_{j=1}^K (r\text{RMSE}_{(j)} - r\text{RMSE})^2}$ for RMSE and relative RMSE.

Software

All analyses were conducted in R Core Team (2023). Simulation and estimation was done using Rosseel (2012). To ensure reproducibility and avoid synchronization in parallelized a pre-generated list of seeds was used for all replications. For further details and a complete

list of libraries and dependencies, visit <https://github.com/valentinkm/AdversarialSimulation>.

Analysis and Interpretation plan

Similar to the studies by Rosseel & Loh (2022) and Dhaene & Rosseel (2023) results were interpreted by descriptively comparing the performance measures of the different estimation methods under varying sample sizes, indicator reliability levels, and model misspecifications without predetermined cut-off values or critical distances. Performance metric values were aggregated across all parameters excluding the misspecified parameters (present in the population but not in the estimation model).

Studies by Kosanke

Current working directory: `/home/rstudio/VK/thesis`

Does LK/thesis.qmd exist? `TRUE`

Methods

The structure of this section closely aligns to our agreed upon structure of simulation studies in Table 1.

In a first step, I published a simulation protocol containing all the planned analysis to be replicated from the original paper by Robitzsch (2022). This protocol can be accessed here: https://github.com/lkosanke/AdversarialSimulation/blob/main/LK/simulation_protocol.pdf.

Aims, objectives and research questions

For my individual study, I replicated parts of Robitzsch (2022) that were relevant to our two substantive research questions. Overall, I conducted 6 simulation studies.

Population Models and Data Generation Mechanisms

The most important details with regards to the population models and data-generating mechanisms are visible in Table 7. With regards to the population models, all factors in all studies loaded onto 3 indicators each. I chose the population values to align with the original paper

by Robitzsch (2022). The multivariate normally distributed data was generated parametrically, based on a specified population model. All simulations were conducted using seeds to allow for the reproducibility of results.

For more details on the exact values of each study, see the simulation scripts in the Github repository.

Study	Model	Correct model included?	Unmodelled RC	Unmodelled CL	N Sizes	φ/β	λ
Study 1	2-factor-CFA	Yes	1 and 2, both pos. and neg.	x	7	$\varphi = 0.6$	Fixed
Study 1b	2-factor-CFA	Yes	x	x	2	$\varphi = 0.2 - 0.8$	Varied
Study 2	2-factor-CFA	x	x	1 and 2, both pos. and neg.	7	$\varphi = 0.6$	Fixed
Study 3	2-factor-CFA	x	1, pos.	1, pos.	7	$\varphi = 0.6$	Fixed
Study 4	5-factors	Yes	20, all pos.	5, all pos.	7	$\beta = 0.1$	Fixed
Study 4a	5-factors	x	20, all pos.	5, all pos.	7	$\beta = 0.1 - 0.4$	Fixed

Note. Φ : factor correlation, N: sample size, λ : factor loading, σ : residual variance, τ : factor variance, RC: residual correlations, CL: cross-loadings, CFA: confirmatory factor analysis, β : regression coefficient between factors.

Overview of Simulation Studies conducted by Kosanke

Experimental Design of simulation procedures

Overall, 3 different types of factors were varied that can be deduced from Table 7 and are detailed again in the simulation scripts provided.

Firstly, I varied the sample size in all studies, ranging from $N = 50$ to 100.000. I included a smaller sample size $N=50$ for all studies, to be able to answer our substantive research questions in more detail. Study 1b explicitly investigated the small sample bias of LSAM estimation in low sample sizes. Thus, only $N=50$ and $N=100$ were present in this study.

Additionally, I varied the amount of misspecification in all studies, either via different numbers of unmodelled residual correlations, cross-loadings, or both.

Thirdly, in Studies 1b and 4a, I varied the population values for three model parameters (ϕ , β and/ or λ).

Besides studies 1 and 2, I implemented full factorial designs. In Studies 1 and 2 I omitted conditions where both one positive and one negative value would be present. I hypothesize that this was done in Robitzsch (2022) to avoid cancellation of biases, but the authors did not give reasoning for this decision themselves.

In Studies 4 and 4a I investigated the differential performance of the estimators in a model that included a non-saturated structural model (i.e. regressions between some of the factors). These studies were replications not only of the paper by Robitzsch (2022), but of the first paper on the SAM approach by Rosseel & Loh (2022). In contrast to the other studies, studies 4 and 4a differed in the way the misspecification variation was labelled in Robitzsch (2022). Instead of varying a factor misspecification as in the previous study, they varied 3 different data-generating mechanisms (DGM's) as a whole. Thus the conditions are labelled differently: DGM 1 contained no misspecification. DGM 2 contained 5 cross-loadings in the data-generating model, that were not modelled in the estimated models. DGM 3 contained 20 residual correlations that were not modelled in the models. I extended them to investigate the interaction of β and N for the 5-factor regression model, as this again was of interest for our substantial research questions. Additionally, I omitted the inclusion of DGM 1 in Study 4a, as it neither contained misspecification (which is central to our research question), nor did it lead to interesting results in the original study.

Method Selection

In terms of estimation methods, I used constrained SEM maximum-likelihood (SEM-ML) and unweighted-least-squares estimation (SEM-ULS), so that loadings and variance parameters were given the constraints that they had to be positive and larger than 0.01. Additionally, I implemented local-SAM (LSAM) and global-SAM (GSAM) estimation, in both maximum-likelihood (LSAM-ML/ GSAM-ML), and unweighted-least-squares estimation (GSAM-ML/ GSAM-ULS) contexts. Exceptions were studies 1b, 4 and 4a, where only LSAM was investigated, as results did not really differ between the two different SAM-methods (Robitzsch, 2022).

Performance Measures

I calculated the bias and RMSE of the estimated factor correlations in all studies, as well as the standard deviation of the one factor correlation present in Studies 1,2 and 3. For the type of bias calculated, I oriented on Robitzsch (2022), besides in Study 1b. Thus, I calculated average relative bias in Studies 1, 2 and 3, and average absolute bias in Studies 1b, 4 and 4a. In Study 1b, I took the absolute value to see if negative and positive biases canceled each other out in the original study for conditions with lower phi values. In addition to what was done in Robitzsch (2022), I calculated confidence intervals for the bias estimates, but omitted them in the results tables for presentation purposes. The exact computation of the performance measures is detailed in the simulation scripts and results.pdf file in my sub-folder of the Github repository.

I did not include a detailed mechanism to capture model convergence as detailed in the first substantive research question. As Robitzsch (2022) argued in their paper, and was shown already in other simulations, using constrained maximum likelihood estimation should resolve convergence issues of classical maximum likelihood estimation in smaller samples (Lüdtke et al., 2021; Ulitzsch et al., 2023). I did include, however, a mechanism to track the total number of warnings for each estimation and compare it to the total number of estimations as a sanity check.

Software

All analyses were conducted in R (R Core Team, 2023). I used the packages lavaan, purrr, tidyverse, furrr to conduct the simulations, as well as knitr and kableExtra for presenting the results (Rosseel, 2012; Vaughan & Dancho, 2022; Wickham et al., 2019; Wickham & Henry, 2023; Xie, 2024; Zhu et al., 2024) .

Analysis and Interpretation plan

For the interpretation of results, I oriented on cut-offs that were used in the original paper by Robitzsch (2022). For bias, I interpreted differences of 0.05 or higher as substantial. For SD, I explicitly mentioned percentage reductions of more or equal to 5%. For RMSE, the same interpretation was used for differences of 0.03 or higher. The simulation was repeated 1500 times for each Study.

Methods of “Joint” Simulation Study

After collaborating based on the individually conducted studies and the respective results, we did not jointly arrive at the conclusion that conducting a collaborative simulation study as planned was warranted.

However I identified several reasons for setting up another simulation. Firstly, to test and evaluate the viability and technical feasibility of AC for simulation studies, setting up a study based on the individual studies, their results and with Kossman can provide valuable...

Aims, objectives and research questions

Following our framework for collaboration the research questions for the joint study remains the same as specified prior to the individual studies.

Population Models and Data Generation Mechanisms


As in study 1 and 2 data for study 3 was generated based on a 5-factor population structural model with 3 indicators for each factor. Factor loadings and indicator reliability was computed in the same way as in study 1. Two different population models were simulated that resulted in misspecifications of either omitted crossloadings (model 3.1) or omitted correlated residuals (model 3.2). The population-level values of the structural parameters were set to 0.1. Reliability levels were manipulated as in Study 1. The omitted crossloadings (see figure 7) could either be all positive or negative and were set to be 10% lower in absolute values than the factor loadings. Correlated residuals were also either all positive or all negative and were set to not exceed a factor of 0.6 of the residual variances of the indicators.

Experimental Design of simulation procedures**

The joint study varied three conditions: (1) sample sizes of very small ($N = 50$), small ($N = 100$) or moderate ($N = 400$). (2) Indicator reliability of low ($= 0.3$), moderate (0.5) or high ($= 0.7$); (3) Model misspecifications with not-specified cross loadings in the

Figure 3.

Population Model Variations for Study 3



Note. Note. Error terms are not explicitly shown in the figure. Dashed lines represent relations omitted in the estimation model present in the population model. Unspecified crossloadings and correlated residuals could be either positive or negative resulting in 2 modulations of model 3.1 and 3.2 in the study.

population model that were positive or negative (see figure) or not-specified correlated residuals in the population model that were positive or negative (see figure 8).

Method Selection

To address the question ...

Performance Measures

Software

To fully evaluate the effect of bound SEM on convergence convergence rate and rate of proper solutions were tracked condition wise (Kriegmair's individual studies).

Analysis and Interpretation plan

The analysis was conducted largely in the same way as in the individual studies

Results


Results of Individual Simulation Studies

Results of Kriegmair's Simulation Studies

As shown in Figure 4 there were no convergence issues for all SAM methods (gSAM, ISAM ML and ULS) with a convergence rate of 100% and no improper solutions across all conditions even in small samples with low reliability. Standard SEM however showed severe convergence issues in small samples with low to moderate reliability with a convergence rate of as low as 50% and 50% improper solutions in the cross loading misspecification condition as the most challenging condition.

Figure 4.

Convergence Rate and Rate of Proper Solutions in Study 1



Note. Convergence and proper solutions (in parentheses) rates across sample sizes (N), reliability (r), and model misspecifications for global SAM (gSAM), local SAM with Maximum Likelihood (ISAM-ML), Unweighted Least Squares (ISAM-ULS), and SEM.

Convergence rates in study 2 were consistent with this with 100% convergence rates for all SAM methods and as low as 60% for standard SEM with exogenous measurement misspecifications posing more challenges than endogenous misspecifications (see Figure A1 in Appendix B).

Results of Kosanke's Simulation Studies

Results of the “Joint” Simulation Study

Results of the Adversarial Collaboration

Discussion

Discussing the substantial results

Discussing the Adversarial Collaboration

Idea: living simulations

References

- Dhaene, S., & Rosseel, Y. (2023). An evaluation of non-iterative estimators in the structural after measurement (SAM) approach to structural equation modeling (SEM). *Structural Equation Modeling: A Multidisciplinary Journal*, 30(6), 926–940. <https://doi.org/10.1080/10705511.2023.2220135>
- Lüdtke, O., Ulitzsch, E., & Robitzsch, A. (2021). A comparison of penalized maximum likelihood estimation and markov chain monte carlo techniques for estimating confirmatory factor analysis models with small sample sizes. *Frontiers in Psychology*, 12. <https://doi.org/10.3389/fpsyg.2021.615162>
- R Core Team. (2023). *R: A language and environment for statistical computing* [Computer software]. R Foundation for Statistical Computing. <https://www.R-project.org/>


- Robitzsch, A. (2022). Comparing the robustness of the structural after measurement (SAM) approach to structural equation modeling (SEM) against local model misspecifications with alternative estimation approaches. *Stats*, 5(3), 631–672. <https://doi.org/10.3390/stats5030039>
- Rosseel, Y. (2012). Lavaan: An r package for structural equation modeling. *Journal of Statistical Software*, 48, 1–36. <https://doi.org/10.18637/jss.v048.i02>
- Rosseel, Y., & Loh, W. W. (2022). A structural after measurement approach to structural equation modeling. *Psychological Methods*, No Pagination Specified–No Pagination Specified. <https://doi.org/10.1037/met0000503>
- Ulitzsch, E., Lüdtke, O., & Robitzsch, A. (2023). Alleviating estimation problems in small sample structural equation modeling—a comparison of constrained maximum likelihood, bayesian estimation, and fixed reliability approaches. *Psychological Methods*, 28(3), 527–557. <https://doi.org/10.1037/met0000435>
- Vaughan, D., & Dancho, M. (2022). *Furrr: Apply mapping functions in parallel using futures*. <https://furrr.futureverse.org/>
- Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., ... Yutani, H. (2019). Welcome to the tidyverse. *Journal of Open Source Software*, 4(43), 1686. <https://doi.org/10.21105/joss.01686>
- Wickham, H., & Henry, L. (2023). *Purrr: Functional programming tools*. <https://purrr.tidyverse.org/>
- Xie, Y. (2024). *Knitr: A general-purpose package for dynamic report generation in r*. <https://yihui.org/knitr/>
- Zhu, H., Travison, T., Tsai, T., Beasley, W., Xie, Y., Yu, G., Laurent, S., Shepherd, R., Sidi, Y., Salzer, B., Gui, G., Fan, Y., Murdoch, D., Arel-Bundock, V., & Evans, B. (2024). *kableExtra: Construct Complex Table with 'kable' and Pipe Syntax*. <https://cran.r-project.org/web/packages/kableExtra/index.html>

Appendix A: Simulation Protocol

Appendix B: Supplementary Figures

Figure A1.

Convergence Rate and Rate of Proper Solutions in Study 2



Note. Convergence and proper solutions (in parentheses) rates across sample sizes (N), reliability (r), and model misspecification location for global SAM (gSAM), local SAM with Maximum Likelihood (ISAM-ML), Unweighted Least Squares (ISAM-ULS), and SEM.

Appendix C: Detailed Error and Warning Messages

In the following, all different warning and error messages raised during the studies are listed (see Table C1) and shown how often they occurred under various fitting conditions (see Table C2).

Table C1.

List of Unique Warnings and Errors

ID	Message
1	lavaan WARNING: some estimated ov variances are negative
2	lavaan WARNING: the optimizer warns that a solution has NOT been found!
3	lavaan WARNING: the optimizer (NLMINB) claimed the model converged, but not all elements of the gradient are (near) zero; the optimizer may not have found a local solution use <code>check.gradient = FALSE</code> to skip this check.
4	lavaan WARNING: some estimated lv variances are negative
5	lavaan WARNING: some estimated ov variances are negative, lavaan WARNING: some estimated lv variances are negative
6	number of items to replace is not a multiple of replacement length
7	lavaan WARNING: Could not compute standard errors! The information matrix could not be inverted. This may be a symptom that the model is not identified., lavaan WARNING: some estimated ov variances are negative
8	lavaan WARNING: covariance matrix of latent variables is not positive definite; use <code>lavInspect(fit, "cov.lv")</code> to investigate.
9	lavaan WARNING: The variance-covariance matrix of the estimated parameters (vcov) does not appear to be positive definite! The smallest eigenvalue is smaller than or close to zero. This may be a symptom that the model is not identified., lavaan WARNING: some estimated ov variances are negative
10	lavaan WARNING: Could not compute standard errors! The information matrix could not be inverted. This may be a symptom that the model is not identified., lavaan WARNING: some estimated lv variances are negative
11	lavaan WARNING: Could not compute standard errors! The information matrix could not be inverted. This may be a symptom that the model is not identified., lavaan WARNING: some estimated ov variances are negative, lavaan WARNING: some estimated lv variances are negative

Table C1.*List of Unique Warnings and Errors (continued)*

ID	Message
12	lavaan WARNING: some estimated ov variances are negative, lavaan WARNING: covariance matrix of latent variables is not positive definite; use lavInspect(fit, "cov.lv") to investigate.
13	lavaan WARNING: Could not compute standard errors! The information matrix could not be inverted. This may be a symptom that the model is not identified.
14	lavaan WARNING: The variance-covariance matrix of the estimated parameters (vcov) does not appear to be positive definite! The smallest eigenvalue is smaller than or close to zero. This may be a symptom that the model is not identified., lavaan WARNING: covariance matrix of latent variables is not positive definite; use lavInspect(fit, "cov.lv") to investigate.
15	lavaan WARNING: Could not compute standard errors! The information matrix could not be inverted. This may be a symptom that the model is not identified., lavaan WARNING: covariance matrix of latent variables is not positive definite; use lavInspect(fit, "cov.lv") to investigate.
16	lavaan WARNING: the covariance matrix of the residuals of the observed variables (theta) is not positive definite; use lavInspect(fit, "theta") to investigate.
17	lavaan WARNING: The variance-covariance matrix of the estimated parameters (vcov) does not appear to be positive definite! The smallest eigenvalue is smaller than or close to zero. This may be a symptom that the model is not identified.

Note. This table lists all unique warnings and errors encountered during the simulation studies.

Table C2.*Summary of Warnings and Errors by Condition with ID for All Studies*

Study	Model Type	N	Reliability	Method	MessageType	Count	Message ID
Study 1	1.3	100	0.3	SEM	Warning	6860	1
Study 1	1.2	100	0.5	SEM	Warning	3575	1
Study 1	1.3	100	0.5	SEM	Warning	2923	1
Study 1	1.2	100	0.7	SEM	Warning	2903	1
Study 1	1.2	100	0.3	SEM	Warning	2769	1
Study 1	1.1	100	0.3	SEM	Warning	2700	1
Study 1	1.4	100	0.3	SEM	Warning	2577	1
Study 1	1.2	100	0.3	SEM	Warning	2037	2

Table C2.*Summary of Warnings and Errors by Condition with ID for All Studies (continued)*

Study	Model Type	N	Reliability	Method	MessageType	Count	Message ID
Study 1	1.1	100	0.3	SEM	Warning	1258	2
Study 1	1.4	100	0.3	SEM	Warning	1133	2
Study 1	1.2	100	0.3	SEM	Warning	729	3
Study 1	1.3	100	0.3	SEM	Warning	692	3
Study 1	1.3	100	0.7	SEM	Warning	688	1
Study 1	1.3	400	0.3	SEM	Warning	606	1
Study 1	1.3	100	0.3	SEM	Warning	507	2
Study 1	1.2	400	0.3	SEM	Warning	450	1
Study 1	1.2	400	0.5	SEM	Warning	429	1
Study 1	1.2	100	0.3	SEM	Warning	417	4
Study 1	1.2	400	0.7	SEM	Warning	248	1
Study 1	1.1	100	0.3	SEM	Warning	242	3
Study 1	1.4	100	0.3	SEM	Warning	223	3
Study 1	1.2	100	0.5	SEM	Warning	203	2
Study 1	1.1	100	0.5	SEM	Warning	197	1
Study 1	1.4	100	0.5	SEM	Warning	183	1
Study 1	1.2	100	0.3	ISAM- ULS	Warning	150	1
Study 1	1.2	100	0.3	SEM	Warning	146	5
Study 1	1.4	100	0.3	ISAM- ULS	Warning	62	1
Study 1	1.1	100	0.3	ISAM- ULS	Warning	52	1
Study 1	1.2	100	0.3	gSAM	Warning	50	4
Study 1	1.4	100	0.3	SEM	Warning	50	4
Study 1	1.2	100	0.5	SEM	Warning	42	3
Study 1	1.2	100	0.3	ISAM- ULS	Error	38	6
Study 1	1.1	100	0.3	SEM	Warning	29	4
Study 1	1.1	100	0.3	ISAM- ULS	Error	25	6
Study 1	1.4	100	0.3	ISAM- ULS	Error	24	6
Study 1	1.2	100	0.3	SEM	Warning	23	7

Table C2.*Summary of Warnings and Errors by Condition with ID for All Studies (continued)*

Study	Model Type	N	Reliability	Method	MessageType	Count	Message ID
Study 1	1.2	400	0.3	SEM	Warning	15	2
Study 1	1.1	100	0.3	SEM	Warning	14	7
Study 1	1.2	100	0.3	gSAM	Error	14	6
Study 1	1.1	100	0.3	SEM	Warning	12	5
Study 1	1.4	100	0.3	SEM	Warning	11	5
Study 1	1.4	100	0.3	SEM	Warning	9	7
Study 1	1.2	100	0.7	SEM	Warning	7	2
Study 1	1.4	100	0.7	SEM	Warning	7	1
Study 1	1.2	100	0.3	SEM	Warning	5	8
Study 1	1.1	100	0.5	SEM	Warning	4	2
Study 1	1.3	100	0.5	SEM	Warning	4	3
Study 1	1.1	100	0.3	SEM	Warning	3	9
Study 1	1.1	100	0.7	SEM	Warning	3	1
Study 1	1.2	100	0.3	SEM	Warning	3	10
Study 1	1.2	100	0.3	SEM	Warning	3	11
Study 1	1.1	100	0.3	gSAM	Error	2	6
Study 1	1.1	100	0.3	gSAM	Warning	2	4
Study 1	1.1	400	0.3	SEM	Warning	2	1
Study 1	1.2	100	0.3	SEM	Warning	2	12
Study 1	1.2	100	0.3	ISAM- ULS	Warning	2	3
Study 1	1.2	100	0.5	SEM	Warning	2	4
Study 1	1.2	400	0.3	SEM	Warning	2	3
Study 1	1.3	100	0.5	ISAM- ULS	Error	2	6
Study 1	1.3	100	0.5	ISAM- ULS	Warning	2	1
Study 1	1.4	100	0.3	SEM	Warning	2	9
Study 1	1.4	100	0.3	gSAM	Error	2	6
Study 1	1.4	100	0.3	gSAM	Warning	2	4
Study 1	1.4	100	0.5	ISAM- ULS	Error	2	6
Study 1	1.1	100	0.3	SEM	Warning	1	11

Table C2.*Summary of Warnings and Errors by Condition with ID for All Studies (continued)*

Study	Model Type	N	Reliability	Method	MessageType	Count	Message ID
Study 1	1.1	100	0.3	ISAM- ULS	Warning	1	3
Study 1	1.1	100	0.5	ISAM- ULS	Warning	1	1
Study 1	1.2	100	0.3	SEM	Warning	1	9
Study 1	1.2	100	0.3	gSAM	Warning	1	8
Study 1	1.2	100	0.5	ISAM- ULS	Error	1	6
Study 1	1.2	100	0.5	ISAM- ULS	Warning	1	1
Study 1	1.3	100	0.3	SEM	Warning	1	7
Study 1	1.3	100	0.3	ISAM- ULS	Warning	1	1
Study 1	1.3	100	0.5	SEM	Warning	1	2
Study 1	1.3	400	0.5	SEM	Warning	1	1
Study 1	1.4	100	0.3	SEM	Warning	1	10
Study 1	1.4	100	0.3	ISAM- ULS	Warning	1	3
Study 1	1.4	100	0.3	ISAM- ULS	Warning	1	2
Study 1	1.4	100	0.5	ISAM- ULS	Warning	1	1
Study 2	2.2_both	100	0.3	SEM	Warning	5265	1
Study 2	2.2_exo	100	0.3	SEM	Warning	4622	1
Study 2	2.2_endo	100	0.3	SEM	Warning	3615	1
Study 2	2.2_both	100	0.7	SEM	Warning	2904	1
Study 2	2.2_both	100	0.5	SEM	Warning	2743	1
Study 2	2.1	100	0.3	SEM	Warning	2701	1
Study 2	2.2_endo	100	0.7	SEM	Warning	2336	1
Study 2	2.2_exo	100	0.5	SEM	Warning	1814	1
Study 2	2.2_both	100	0.3	SEM	Warning	1625	2
Study 2	2.2_exo	100	0.3	SEM	Warning	1624	2
Study 2	2.2_endo	100	0.5	SEM	Warning	1252	1
Study 2	2.2_endo	100	0.3	SEM	Warning	1211	2

Table C2.*Summary of Warnings and Errors by Condition with ID for All Studies (continued)*

Study	Model Type	N	Reliability	Method	MessageType	Count	Message ID
Study 2	2.1	100	0.3	SEM	Warning	1121	2
Study 2	2.2_exo	100	0.7	SEM	Warning	702	1
Study 2	2.2_both	100	0.3	SEM	Warning	675	3
Study 2	2.2_both	400	0.7	SEM	Warning	599	1
Study 2	2.2_endo	400	0.7	SEM	Warning	588	1
Study 2	2.2_exo	100	0.3	SEM	Warning	559	3
Study 2	2.2_endo	100	0.3	SEM	Warning	305	3
Study 2	2.2_exo	100	0.3	ISAM- ULS	Warning	286	1
Study 2	2.2_both	100	0.3	SEM	Warning	242	4
Study 2	2.2_both	400	0.3	SEM	Warning	239	1
Study 2	2.1	100	0.3	SEM	Warning	220	3
Study 2	2.2_exo	400	0.3	SEM	Warning	195	1
Study 2	2.2_exo	100	0.3	SEM	Warning	175	4
Study 2	2.1	100	0.5	SEM	Warning	165	1
Study 2	2.2_endo	100	0.3	SEM	Warning	157	4
Study 2	2.1	100	0.3	ISAM- ULS	Warning	138	1
Study 2	2.1	100	0.3	SEM	Warning	132	4
Study 2	2.2_both	400	0.5	SEM	Warning	130	1
Study 2	2.2_both	100	0.3	ISAM- ULS	Warning	128	1
Study 2	2.2_exo	100	0.3	ISAM- ULS	Error	121	6
Study 2	2.2_both	100	0.5	SEM	Warning	105	2
Study 2	2.2_exo	100	0.5	SEM	Warning	93	2
Study 2	2.2_endo	100	0.3	ISAM- ULS	Warning	78	1
Study 2	2.2_exo	400	0.5	SEM	Warning	77	1
Study 2	2.2_both	100	0.3	SEM	Warning	57	5
Study 2	2.1	100	0.3	ISAM- ULS	Error	44	6
Study 2	2.2_exo	100	0.3	SEM	Warning	43	5

Table C2.*Summary of Warnings and Errors by Condition with ID for All Studies (continued)*

Study	Model Type	N	Reliability	Method	MessageType	Count	Message ID
Study 2	2.2_both	100	0.3	ISAM- ULS	Error	41	6
Study 2	2.2_endo	400	0.5	SEM	Warning	40	1
Study 2	2.2_both	100	0.3	gSAM	Warning	31	4
Study 2	2.2_endo	400	0.3	SEM	Warning	29	1
Study 2	2.2_endo	100	0.3	SEM	Warning	26	5
Study 2	2.2_endo	100	0.3	gSAM	Warning	26	4
Study 2	2.2_exo	100	0.3	gSAM	Warning	20	4
Study 2	2.1	100	0.3	gSAM	Warning	18	4
Study 2	2.2_both	100	0.5	SEM	Warning	18	3
Study 2	2.2_both	100	0.5	ISAM- ULS	Warning	17	1
Study 2	2.2_exo	100	0.5	ISAM- ULS	Warning	16	1
Study 2	2.1	100	0.3	SEM	Warning	15	5
Study 2	2.1	100	0.3	SEM	Warning	14	7
Study 2	2.2_exo	100	0.5	SEM	Warning	14	3
Study 2	2.2_endo	100	0.3	ISAM- ULS	Error	14	6
Study 2	2.2_endo	100	0.5	SEM	Warning	13	2
Study 2	2.2_exo	100	0.3	SEM	Warning	10	7
Study 2	2.2_exo	400	0.3	SEM	Warning	10	2
Study 2	2.2_both	100	0.3	SEM	Warning	10	7
Study 2	2.2_both	400	0.3	SEM	Warning	8	2
Study 2	2.2_exo	100	0.3	ISAM- ULS	Warning	7	3
Study 2	2.2_both	100	0.3	gSAM	Error	7	6
Study 2	2.2_exo	100	0.3	SEM	Warning	6	10
Study 2	2.2_exo	100	0.5	ISAM- ULS	Error	5	6
Study 2	2.2_both	100	0.5	ISAM- ULS	Error	5	6
Study 2	2.2_exo	400	0.7	SEM	Warning	4	1
Study 2	2.2_endo	100	0.3	SEM	Warning	4	7

Table C2.*Summary of Warnings and Errors by Condition with ID for All Studies (continued)*

Study	Model Type	N	Reliability	Method	MessageType	Count	Message ID
Study 2	2.1	100	0.3	SEM	Warning	3	10
Study 2	2.1	100	0.5	ISAM- ULS	Warning	3	1
Study 2	2.2_exo	100	0.3	SEM	Warning	3	11
Study 2	2.2_both	100	0.3	SEM	Warning	3	10
Study 2	2.2_both	100	0.3	SEM	Warning	3	11
Study 2	2.2_both	100	0.7	SEM	Warning	3	2
Study 2	2.1	100	0.7	SEM	Warning	2	1
Study 2	2.2_exo	100	0.3	gSAM	Error	2	6
Study 2	2.2_exo	100	0.3	ISAM- ULS	Warning	2	2
Study 2	2.2_exo	100	0.7	SEM	Warning	2	2
Study 2	2.2_exo	400	0.3	SEM	Warning	2	3
Study 2	2.2_endo	100	0.5	SEM	Warning	2	3
Study 2	2.2_both	100	0.3	gSAM	Warning	2	3
Study 2	2.2_both	100	0.5	SEM	Warning	2	4
Study 2	2.1	100	0.3	gSAM	Error	1	6
Study 2	2.1	400	0.3	SEM	Warning	1	1
Study 2	2.2_exo	100	0.3	SEM	Warning	1	13
Study 2	2.2_exo	100	0.3	SEM	Warning	1	12
Study 2	2.2_exo	100	0.5	ISAM- ULS	Warning	1	3
Study 2	2.2_exo	100	0.7	SEM	Warning	1	3
Study 2	2.2_exo	400	0.3	ISAM- ULS	Error	1	6
Study 2	2.2_endo	100	0.3	SEM	Warning	1	9
Study 2	2.2_endo	100	0.3	ISAM- ULS	Warning	1	3
Study 2	2.2_endo	100	0.5	ISAM- ULS	Error	1	6
Study 2	2.2_endo	100	0.5	ISAM- ULS	Warning	1	1
Study 2	2.2_endo	400	0.3	SEM	Warning	1	2

Table C2.*Summary of Warnings and Errors by Condition with ID for All Studies (continued)*

Study	Model Type	N	Reliability	Method	MessageType	Count	Message ID
Study 2	2.2_both	100	0.3	ISAM- ULS	Warning	1	2
Study 2	2.2_both	100	0.3	ISAM_ML	Error	1	6
Study 2	2.2_both	100	0.5	ISAM- ULS	Warning	1	2
Study 2	2.2_both	100	0.7	ISAM- ULS	Warning	1	1
Study 2	2.2_both	400	0.3	SEM	Warning	1	3
Study 2	2.2_both	400	0.3	SEM	Warning	1	4
Study 3	3.1	50	0.3	ISAM- ULS	Warning	248	1
Study 3	3.1	50	0.3	gSAM	Warning	209	4
Study 3	3.1_negative	50	0.3	ISAM- ULS	Warning	208	1
Study 3	3.1_negative	50	0.3	gSAM	Warning	164	4
Study 3	3.2	50	0.3	ISAM- ULS	Warning	82	1
Study 3	3.2_negative	50	0.3	ISAM- ULS	Warning	82	1
Study 3	3.1_negative	50	0.3	ISAM- ULS	Error	79	6
Study 3	3.1	50	0.3	ISAM- ULS	Error	72	6
Study 3	3.2	50	0.3	SEM	Warning	71	8
Study 3	3.2_negative	50	0.3	SEM	Warning	71	8
Study 3	3.1	100	0.3	ISAM- ULS	Warning	62	1
Study 3	3.1_negative	50	0.3	SEM	Warning	61	8
Study 3	3.1	50	0.3	SEM	Warning	53	8
Study 3	3.1_negative	100	0.3	ISAM- ULS	Warning	52	1
Study 3	3.1	50	0.5	ISAM- ULS	Warning	51	1
Study 3	3.1_negative	50	0.3	gSAM	Error	47	6

Table C2.*Summary of Warnings and Errors by Condition with ID for All Studies (continued)*

Study	Model Type	N	Reliability	Method	MessageType	Count	Message ID
Study 3	3.1_negative	50	0.5	ISAM- ULS	Warning	37	1
Study 3	3.1	50	0.3	gSAM	Error	36	6
Study 3	3.2	50	0.3	SEM	Warning	31	14
Study 3	3.2	100	0.3	ISAM- ULS	Warning	31	1
Study 3	3.2_negative	50	0.3	SEM	Warning	31	14
Study 3	3.2_negative	100	0.3	ISAM- ULS	Warning	31	1
Study 3	3.1	100	0.3	ISAM- ULS	Error	28	6
Study 3	3.2	50	0.3	ISAM- ULS	Warning	19	2
Study 3	3.2_negative	50	0.3	ISAM- ULS	Warning	19	2
Study 3	3.1	50	0.5	ISAM- ULS	Error	18	6
Study 3	3.2	50	0.3	ISAM- ULS	Warning	18	3
Study 3	3.1_negative	100	0.3	ISAM- ULS	Error	18	6
Study 3	3.2_negative	50	0.3	ISAM- ULS	Warning	18	3
Study 3	3.1	100	0.3	gSAM	Warning	17	4
Study 3	3.2	50	0.3	ISAM- ULS	Error	17	6
Study 3	3.1_negative	50	0.3	SEM	Warning	17	14
Study 3	3.2_negative	50	0.3	ISAM- ULS	Error	17	6
Study 3	3.1_negative	50	0.5	ISAM- ULS	Error	13	6
Study 3	3.1_negative	100	0.3	gSAM	Warning	12	4
Study 3	3.2	100	0.3	ISAM- ULS	Warning	11	3

Table C2.*Summary of Warnings and Errors by Condition with ID for All Studies (continued)*

Study	Model Type	N	Reliability	Method	MessageType	Count	Message ID
Study 3	3.2_negative	100	0.3	ISAM- ULS	Warning	11	3
Study 3	3.1	50	0.3	SEM	Warning	10	15
Study 3	3.1	50	0.3	gSAM	Warning	10	3
Study 3	3.2	50	0.3	gSAM	Warning	10	4
Study 3	3.1_negative	100	0.3	SEM	Warning	10	8
Study 3	3.2_negative	50	0.3	gSAM	Warning	10	4
Study 3	3.1	50	0.3	gSAM	Warning	9	8
Study 3	3.2	100	0.3	ISAM- ULS	Error	9	6
Study 3	3.2_negative	100	0.3	ISAM- ULS	Error	9	6
Study 3	3.1	50	0.5	gSAM	Warning	8	4
Study 3	3.1	50	0.7	ISAM- ULS	Warning	7	1
Study 3	3.2	100	0.3	ISAM- ULS	Warning	7	2
Study 3	3.2_negative	100	0.3	ISAM- ULS	Warning	7	2
Study 3	3.1	50	0.3	SEM	Warning	6	14
Study 3	3.1	100	0.3	SEM	Warning	6	8
Study 3	3.2	50	0.3	SEM	Warning	6	15
Study 3	3.1_negative	50	0.3	SEM	Warning	6	15
Study 3	3.1_negative	50	0.3	gSAM	Warning	6	8
Study 3	3.1_negative	50	0.7	ISAM- ULS	Warning	6	1
Study 3	3.2_negative	50	0.3	SEM	Warning	6	15
Study 3	3.1	50	0.7	ISAM- ULS	Error	5	6
Study 3	3.2	50	0.5	ISAM- ULS	Warning	5	1
Study 3	3.1_negative	100	0.3	gSAM	Error	5	6
Study 3	3.2_negative	50	0.5	ISAM- ULS	Warning	5	1

Table C2.*Summary of Warnings and Errors by Condition with ID for All Studies (continued)*

Study	Model Type	N	Reliability	Method	MessageType	Count	Message ID
Study 3	3.1	50	0.3	ISAM- ULS	Warning	4	2
Study 3	3.2	50	0.5	ISAM- ULS	Error	4	6
Study 3	3.2	100	0.3	SEM	Warning	4	8
Study 3	3.1_negative	50	0.3	ISAM- ULS	Warning	4	2
Study 3	3.1_negative	50	0.5	SEM	Warning	4	8
Study 3	3.2_negative	50	0.5	ISAM- ULS	Error	4	6
Study 3	3.2_negative	100	0.3	SEM	Warning	4	8
Study 3	3.1	50	0.3	ISAM- ULS	Warning	3	16
Study 3	3.1_negative	50	0.3	gSAM	Warning	3	3
Study 3	3.1_negative	100	0.5	ISAM- ULS	Warning	3	1
Study 3	3.1	50	0.3	ISAM- ULS	Warning	2	3
Study 3	3.1	50	0.5	SEM	Warning	2	8
Study 3	3.1	100	0.3	gSAM	Error	2	6
Study 3	3.1	100	0.5	ISAM- ULS	Error	2	6
Study 3	3.1	250	0.3	ISAM- ULS	Warning	2	1
Study 3	3.2	100	0.3	SEM	Warning	2	14
Study 3	3.1_negative	50	0.3	ISAM- ULS	Warning	2	3
Study 3	3.1_negative	50	0.3	ISAM- ULS	Warning	2	16
Study 3	3.1_negative	50	0.5	gSAM	Warning	2	4
Study 3	3.1_negative	100	0.3	SEM	Warning	2	14
Study 3	3.1_negative	100	0.3	ISAM- ULS	Warning	2	3
Study 3	3.2_negative	100	0.3	SEM	Warning	2	14
Study 3	3.1	50	0.5	gSAM	Error	1	6

Table C2.*Summary of Warnings and Errors by Condition with ID for All Studies (continued)*

Study	Model Type	N	Reliability	Method	MessageType	Count	Message ID
Study 3	3.1	50	0.5	ISAM- ULS	Warning	1	3
Study 3	3.1	50	0.5	ISAM- ULS	Warning	1	2
Study 3	3.1	100	0.3	SEM	Warning	1	14
Study 3	3.1	100	0.3	gSAM	Warning	1	8
Study 3	3.1	100	0.3	ISAM- ULS	Warning	1	3
Study 3	3.1	100	0.5	ISAM- ULS	Warning	1	1
Study 3	3.1	250	0.3	ISAM- ULS	Error	1	6
Study 3	3.2	50	0.3	gSAM	Error	1	6
Study 3	3.2	50	0.3	ISAM- ULS	Warning	1	16
Study 3	3.2	50	0.5	gSAM	Warning	1	4
Study 3	3.2	50	0.5	ISAM- ULS	Warning	1	3
Study 3	3.2	50	0.7	ISAM- ULS	Warning	1	1
Study 3	3.2	250	0.3	ISAM- ULS	Warning	1	3
Study 3	3.2	250	0.3	ISAM- ULS	Warning	1	2
Study 3	3.2	250	0.3	ISAM- ULS	Warning	1	1
Study 3	3.2	400	0.3	ISAM- ULS	Warning	1	1
Study 3	3.1_negative	50	0.5	gSAM	Error	1	6
Study 3	3.1_negative	50	0.5	ISAM- ULS	Warning	1	16
Study 3	3.1_negative	50	0.7	ISAM- ULS	Error	1	6
Study 3	3.1_negative	50	0.7	ISAM- ULS	Warning	1	3

Table C2.*Summary of Warnings and Errors by Condition with ID for All Studies (continued)*

Study	Model Type	N	Reliability	Method	MessageType	Count	Message ID
Study 3	3.1_negative	100	0.3	SEM	Warning	1	13
Study 3	3.1_negative	100	0.3	SEM	Warning	1	17
Study 3	3.1_negative	100	0.3	gSAM	Warning	1	8
Study 3	3.1_negative	250	0.3	ISAM- ULS	Error	1	6
Study 3	3.1_negative	250	0.3	ISAM- ULS	Warning	1	1
Study 3	3.2_negative	50	0.3	gSAM	Error	1	6
Study 3	3.2_negative	50	0.3	ISAM- ULS	Warning	1	16
Study 3	3.2_negative	50	0.5	gSAM	Warning	1	4
Study 3	3.2_negative	50	0.5	ISAM- ULS	Warning	1	3
Study 3	3.2_negative	50	0.7	ISAM- ULS	Warning	1	1
Study 3	3.2_negative	250	0.3	ISAM- ULS	Warning	1	3
Study 3	3.2_negative	250	0.3	ISAM- ULS	Warning	1	2
Study 3	3.2_negative	250	0.3	ISAM- ULS	Warning	1	1
Study 3	3.2_negative	400	0.3	ISAM- ULS	Warning	1	1

Note. This table summarizes the count of warnings and errors for each condition in all three simulation studies with the respective ID number corresponding to Table 1.