

Modeling assumptions of repeated-visits, single-visit, and integrated occupancy models

Valentin Lauret, Hélène Labach, Matthieu Authier, Olivier Gimenez

26/03/2021

Contents

Modeling assumptions	1
Repeated-visits occupancy models	1
Single-visit occupancy	1
Integrated occupancy models	2
Summarize the modeling assumptions in a table	2

Literature cited	3
-------------------------	----------

Supporting information of the article *Using single visits into integrated occupancy models to make the most of existing monitoring programs*.

Modeling assumptions

In this section we aimed to list the modeling assumptions of the Repeated-Visits (RV), Single-Visit (SV), and Integrated occupancy models that we ran in the manuscript.

Repeated-visits occupancy models

Citing, (MacKenzie 2006), there are several critical assumptions for the standard occupancy model, i.e. RV occupancy.

1. Occupancy status at each site does not change over the survey season; that is, sites are “closed” to changes in occupancy
2. The probability of occupancy is constant across sites, or differences in occupancy probability are modeled using covariates
3. The probability of detection is constant across all sites and surveys or is a function of site-survey covariates
4. There is no unmodeled heterogeneity in detection probabilities
5. Detection of species and detection histories at each location are independent

Single-visit occupancy

Developing SV occupancy models, (Lele, Moreno, and Bayne 2012) underlined that SV occupancy relax the closure assumption of sampled sites between visits. Besides, the literature about SV provide some requirements and guidance to a valid application of SV occupancy models. We listed the elements below:

1. Occupancy probability and detection probability depend on covariates

2. At least two independent continuous covariates are used to estimate occupancy probability and detection probability. Shared covariates can result in biased estimates for regression coefficients.
3. Ensure adequate numbers of occurrence. (Peach, Cohen, and Frair 2017), suggested that “estimates of occupancy probability remained unbiased across our scenarios, whereas colonization and extinction estimates became biased as occupancy probability approached extremes (i.e. 0.1 or 0.9).”
4. Prefer nonlinear detection model to provide accurate parameter estimates and to assume a more realistic relationship between detection and effort (Knappe and Korner-Nievergelt 2015) for single-visit occupancy models.

Integrated occupancy models

Combining multiple datasets into occupancy models have been developed previously by (Nichols et al. 2008) in details by estimating occupancy at two the spatial scales. In our integration process, we extended the parametrization of a standard occupancy model detection process to include two different datasets with different detection probabilities. Doing it, we must assume that

- the two monitoring programs are independent, i.e. detection by program 1 does not affect detection probability of program 2.

Subsequently, integrated repeated-visits occupancy models have the modeling assumption of both RV occupancy and of integrated occupancy. Similarly, using integrated single-visit occupancy models cumulate both the assumptions of SV occupancy and of integrated occupancy.

Summarize the modeling assumptions in a table

Modeling assumptions	RV occupancy	SV occupancy
Closure assumption, i.e. site does not change over the survey season	Yes	No
Latent occupancy process	Occupancy prob. constant across sites, or modeled with covariates	Continuous covariate needed, independent from the covariate used for the detection prob.
Detection process	Detection prob. constant across sites, or modeled with covariates	Continuous covariate needed, independent from the covariate used for the occupancy prob.
No unmodeled heterogeneity in detection prob.	Yes	Yes
Detection of species and detection histories at each site are independent	Yes	Yes
Proportion of occurrence over the total number of sites is > 10%	No	Yes
Data integration Detection process	Integrated RV occupancy Monitoring programs must be independent, or dependency must be accounted.	Integrated SV occupancy Monitoring programs must be independent, or dependency must be accounted.

Literature cited

- Knape, Jonas, and Fränzi Korner-Nievergelt. 2015. “Estimates from Non-Replicated Population Surveys Rely on Critical Assumptions.” Edited by Nigel Yoccoz. *Methods in Ecology and Evolution* 6 (3): 298–306. <https://doi.org/10.1111/2041-210X.12329>.
- Lele, S. R., M. Moreno, and E. Bayne. 2012. “Dealing with Detection Error in Site Occupancy Surveys: What Can We Do with a Single Survey?” *Journal of Plant Ecology* 5 (1): 22–31. <https://doi.org/10.1093/jpe/rtr042>.
- MacKenzie, Darryl I., ed. 2006. *Occupancy Estimation and Modeling: Inferring Patterns and Dynamics of Species*. Amsterdam ; Boston: Elsevier.
- Nichols, James D., Larissa L. Bailey, Allan F. O’Connell Jr., Neil W. Talancy, Evan H. Campbell Grant, Andrew T. Gilbert, Elizabeth M. Annand, Thomas P. Husband, and James E. Hines. 2008. “Multi-Scale Occupancy Estimation and Modelling Using Multiple Detection Methods.” *Journal of Applied Ecology* 45 (5): 1321–29. <https://doi.org/10.1111/j.1365-2664.2008.01509.x>.
- Peach, Michelle A., Jonathan B. Cohen, and Jacqueline L. Frair. 2017. “Single-Visit Dynamic Occupancy Models: An Approach to Account for Imperfect Detection with Atlas Data.” *Journal of Applied Ecology* 54 (6): 2033–42. <https://doi.org/10.1111/1365-2664.12925>.