

Quizzes:

Q1: $\psi(x, t) = \int dx' U(x, x', t, t_0) \psi(x', t_0)$ ✓

Q2: $U(x_N, x_0, t_N, t_0) = A \int Dx(t) e^{\frac{i}{\hbar} S(x(t))}$ ✓

Q3: i) $\hat{U}(t, t_0) = e^{-\frac{i}{\hbar} \int_{t_0}^t \hat{H}(t') dt'}$ is unitary.

ii) $U(x_N, x_0, t_N, t_0)$ is the time evolution operator for any path from x_0 to x_N . In the Schrödinger picture we do not look at multiple paths, but the propagator (in S.P.: time evolution operator) is only time dependent: $\vec{U}(t, t_0)$. ✓

Exercise 1:

1.: $e^{\hat{A}} \cdot e^{-\hat{A}} = \left(\sum_{n=0}^{\infty} \frac{\hat{A}^n}{n!} \right) \cdot \left(\sum_{n=0}^{\infty} \frac{(-\hat{A})^n}{n!} \right) = \left(\sum_{i=0}^{\infty} a_i \right) \cdot \left(\sum_{j=0}^{\infty} b_j \right) = \sum_{h=0}^{\infty} \sum_{l=0}^h a_l b_{h-l}$

$$= \sum_{h=0}^{\infty} \sum_{l=0}^h \frac{\hat{A}^l}{l!} \frac{(-\hat{A})^{h-l}}{(h-l)!}$$

$$= \sum_{h=0}^{\infty} \frac{\hat{A}^h}{h!} \sum_{l=0}^h \frac{(-1)^{h-l}}{(h-l)!} \hat{A}^l$$

$$e^{\hat{A}} \cdot e^{-\hat{A}} = \left(\sum_{n=0}^{\infty} \frac{\hat{A}^n}{n!} \right) \cdot \left(\sum_{n=0}^{\infty} \frac{(-\hat{A})^n}{n!} \right)$$

$$= \left(1 + \sum_{n=1}^{\infty} \frac{\hat{A}^n}{n!} \right) \cdot \left(1 + \sum_{n=1}^{\infty} \frac{(-\hat{A})^n}{n!} \right)$$

$$= 1 + \sum_{n=1}^{\infty} \frac{(-\hat{A})^n}{n!} + \sum_{n=1}^{\infty} \frac{\hat{A}^n}{n!} + \left(\sum_{n=1}^{\infty} \frac{\hat{A}^n}{n!} \right) \left(\sum_{n=1}^{\infty} \frac{(-\hat{A})^n}{n!} \right)$$

$$= 1 + \sum_{n=1}^{\infty} \frac{1}{n!} [(-\hat{A})^n + \hat{A}^n] + \sum_{n=1}^{\infty} \sum_{l=0}^{n+1} \frac{\hat{A}^{l+1}}{(l+1)!} \frac{(-\hat{A})^{n-l}}{(n-l)!}$$

$$= 1 + \sum_{n=1}^{\infty} \frac{\hat{A}^n}{n!} [1 + (-1)^n] + \sum_{n=1}^{\infty} \sum_{l=0}^{n+1} \hat{A}^{n+1} \frac{(-1)^{n-l}}{(l+1)!(n-l)!}$$

$$= 1 + \sum_{n=1}^{\infty} \frac{\hat{A}^{n+1}}{(n+1)!} [1 + (-1)^{n+1}] + \sum_{n=1}^{\infty} \hat{A}^{n+1} \sum_{l=0}^{n+1} \frac{(-1)^{n-l}}{(l+1)!(n-l)!}$$

$$\stackrel{!}{=} 1$$

$$\Rightarrow \sum_{n=0}^{\infty} \frac{\hat{A}^{n+1}}{(n+1)!} [1 + (-1)^{n+1}] + \sum_{n=1}^{\infty} \hat{A}^{n+1} \sum_{l=0}^{n+1} \frac{(-1)^{n-l}}{(l+1)!(n-l)!} = 0$$

$$\Rightarrow \frac{1 + (-1)^{n+1}}{(n+1)!} + \sum_{l=0}^{n+1} \frac{(-1)^{n-l}}{(l+1)!(n-l)!} = 0$$

1) Exponentiating Operators

$$e^{\hat{A}} = \sum_{n=0}^{\infty} \frac{1}{n!} \hat{A}^n$$

1) Show: $e^{\hat{A}} e^{-\hat{A}} = 1$

$$\begin{aligned} e^{\hat{A}} e^{-\hat{A}} &= \left(\sum_{n=0}^{\infty} \frac{1}{n!} \hat{A}^n \right) \left(\sum_{m=0}^{\infty} \frac{1}{m!} (-\hat{A})^m \right) \\ &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{1}{n! m!} \hat{A}^n (-\hat{A})^m \\ &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{(-1)^m}{n! m!} \hat{A}^{(n+m)} \quad | \quad p = n+m \\ &= \sum_{p=0}^{\infty} \sum_{n=0}^p \frac{(-1)^{p-n}}{n! (p-n)!} \hat{A}^p \\ &= 1 + \sum_{p=1}^{\infty} \frac{(-\hat{A})^p}{p!} \sum_{n=0}^p (-1)^n \binom{p}{n} \quad \text{with } \binom{p}{n} = \frac{p!}{n! (p-n)!} \quad | \quad 0 = (1-1)^p = \sum_{n=0}^p \binom{p}{n} 1^{p-n} (-1)^n = \sum_{n=0}^p \binom{p}{n} (-1)^n \end{aligned}$$

$$\Rightarrow e^{\hat{A}} e^{-\hat{A}} = 1 + 0 = 1 \quad \square \quad \checkmark$$

4.2 ?

3) Show: $e^{\hat{A}} \hat{B} e^{-\hat{A}} = \sum_{n=0}^{\infty} \frac{1}{n!} [\hat{A}, \hat{B}]_n$

$$\begin{aligned} e^{\hat{A}} \hat{B} e^{-\hat{A}} &= \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \frac{1}{i! j!} \hat{A}^i \hat{B} (-\hat{A})^j \quad | \quad n = i+j \\ &= \sum_{n=0}^{\infty} \frac{1}{n!} \left(\sum_{i=0}^n \frac{n!}{i! (n-i)!} \hat{A}^{(n-i)} \hat{B} (-\hat{A})^i \right) \\ &= \sum_{n=0}^{\infty} \frac{1}{n!} [\hat{A}, \hat{B}]_n \quad \square \end{aligned}$$

$$\text{Show: } [\hat{A}, \hat{B}]_n = \sum_{i=0}^n \frac{n!}{i! (n-i)!} \hat{A}^{n-i} \hat{B} (-\hat{A})^i$$

$$\cdot \boxed{n=0}: [\hat{A}, \hat{B}]_0 = \hat{A}^0 \hat{B} (-\hat{A})^0 = \hat{B}$$

$$\cdot \boxed{n+1}: [\hat{A}, \hat{B}]_{n+1} = [\hat{A}, [\hat{A}, \hat{B}]_n]$$

$$= [\hat{A}, \sum_{i=0}^n \frac{n!}{i! (n-i)!} \hat{A}^{n-i} \hat{B} (-\hat{A})^i]$$

$$= \sum_{i=0}^n \frac{n!}{i! (n-i)!} [\hat{A}, \hat{A}^{n-i} \hat{B} (-\hat{A})^i]$$

$$= \sum_{i=0}^n \frac{n!}{i! (n-i)!} (\hat{A}^{n+1-i} \hat{B} (-\hat{A})^i + \hat{A}^{n-i} \hat{B} (-\hat{A})^{i+1})$$

$$= (\dots)$$

3) Perturbed Harmonic Oscillator

$$2) |A_n| = \delta_{fi} - \frac{i}{\hbar} \int_{t_0}^t dt' \exp[i\omega_k(t-t')] \langle f^n | \hat{H}_{int} | i^{(0)} \rangle$$

$$\cdot \langle f^n | \hat{H}_{int} | i^{(0)} \rangle = \langle f^n | a \hat{x}^p e^{i\gamma t} | i^{(0)} \rangle$$

$$= a e^{\frac{i\gamma t}{\hbar}} \int_{\mathbb{R}} dx x^p \psi_f(x) \psi_i(x)$$

$$= \frac{ab}{\sqrt{\pi}} \frac{e^{\frac{i\gamma t}{\hbar}}}{\sqrt{2} \ell_F} \int_{\mathbb{R}} dx x^p \underbrace{e^{-\frac{b^2 x^2}{2}}}_{f(x)} \underbrace{H_p(bx)}_{g(x)} \quad \left| \quad \psi_n(x) = \left(\frac{b}{\pi} \right)^{\frac{1}{4}} \frac{1}{\sqrt{2^n n!}} H_n(bx) e^{-\frac{b^2 x^2}{2}} \quad \text{where } b = \sqrt{\frac{m\omega}{\hbar}} \right.$$

$$\cdot \text{Assume: } |i\rangle = |n=0\rangle \Rightarrow \psi_i = \left(\frac{b}{\pi} \right)^{\frac{1}{4}} e^{-\frac{b^2 x^2}{2}}$$

$$\cdot f(-x) = (-1)^p f(x)$$

$$\cdot g(-x) = g(x)$$

$$\cdot h(-x) = H_p(-x) = (-1)^p H_p(x) = (-1)^p h(x)$$

$$\left. \begin{aligned} &\left\{ \begin{aligned} &\text{for even } p: P_{ft} \sim \int dx f(x) g(x) h(x) = 0 \quad \text{for even } f \rightarrow \text{accessible for odd } f \\ &\text{for odd } p: P_{ft} \sim \int dx f(x) g(x) h(x) = 0 \quad \text{for odd } f \rightarrow \text{accessible for even } f \end{aligned} \right. \end{aligned} \right\} \quad \checkmark$$

$$3) P_{n,0} = |A_n|^2 = \left| -\frac{i}{\hbar} \int_{t_0}^t dt' \exp[i\omega_k(t-t')] \langle 1 | \hat{H}_1 | 0 \rangle \right|^2$$

$$= \left| -\frac{i}{\hbar} \int_{t_0}^t dt' \exp[i\omega_k(t-t')] \int_{\mathbb{R}} dx a \exp\left[-\frac{b^2 x^2}{2}\right] x^p \psi_0(x) \psi_1(x) \right|^2 \quad \left| \quad \psi_0(x) = \left(\frac{b}{\pi} \right)^{\frac{1}{4}} e^{-\frac{b^2 x^2}{2}} \quad \wedge \quad \psi_1(x) = \left(\frac{b^2}{\pi} \right)^{\frac{1}{4}} b x e^{-\frac{b^2 x^2}{2}} \quad (p=1) \right.$$

$$= \left| -\frac{i}{\hbar} \frac{ab}{\sqrt{\pi}} \int_{t_0}^t dt' \exp[i\omega_k(t-t')] \exp\left[-\frac{b^2 x^2}{2}\right] \int_{\mathbb{R}} dx x^2 \exp[-b^2 x^2] \right|^2 \quad \left| \quad \int_{\mathbb{R}} dx x^2 \exp(-b^2 x^2) = -\frac{2}{\partial b} \int_{\mathbb{R}} dx \exp(-b^2 x^2) = -\frac{2}{\partial b} \sqrt{\frac{\pi}{b^2}} = \frac{\sqrt{\pi}}{b^2} \right.$$

$$= \left| -\frac{i}{\hbar} \frac{a}{b} \exp(-u t_0) \int_{t_0}^t dt' \exp(u t' - v t'^2) \right|^2 \quad \text{where } u = i\omega \quad \wedge \quad v = \gamma^2 \quad | \quad t_0 \rightarrow -\infty \quad \wedge \quad t \rightarrow \infty$$

$$= \left| -\frac{i}{\hbar} \frac{a}{b} \exp(-u t_0) \exp\left(\frac{u^2}{4v}\right) \sqrt{\frac{\pi}{v}} \right|^2$$

$$= \left| -i a \gamma \sqrt{\frac{\pi}{\hbar m \omega}} \exp(-i \omega t_0) \exp\left(-\frac{\gamma^2}{4} \omega^2 \gamma^2\right) \right|^2$$

$$= \frac{a^2 \pi \gamma^2}{\hbar m \omega} \exp\left(-\frac{\gamma^2}{2} \omega^2 \gamma^2\right) \quad \checkmark$$

$$\hookrightarrow \text{for } \gamma \rightarrow 0: \quad \left. \begin{aligned} &i) a = \text{const.} \Rightarrow P_{n,0} \rightarrow 0 \\ &ii) a \sim \frac{1}{\sqrt{\gamma}} \Rightarrow P_{n,0} \rightarrow 0 \end{aligned} \right\} \quad \checkmark$$

Exercise 2:

1.: $c_f(t)$ is the coefficient, that "says how much of" $|f\rangle$, $|\psi(t)\rangle$ has:

$$\text{I: } |\psi(t)\rangle = \sum_n c_n(t) |n\rangle = \sum_{n \neq f} c_n(t) |n\rangle + c_f(t) |f\rangle$$

$$\text{Starting from } |i^{(0)}\rangle = \sum_n c_n(i) |n^{(0)}\rangle = \sum_n \delta_{in} |n^{(0)}\rangle$$

$$|\psi(t_0)\rangle = |i^{(0)}\rangle \leadsto |\psi(t)\rangle = |f^{(0)}\rangle$$

$$\Rightarrow P_{fi}(t) = \underbrace{|\langle f^{(0)} | i^{(0)} \rangle|^2}_{= |c_f(t)|^2} \quad \text{I}$$

2.:

$$\begin{aligned} |\psi(t)\rangle &= \sum_n d_n(t) |\psi_n^{(0)}(t)\rangle \\ &= \sum_n d_n(t) \hat{U}(t, t_0) |\psi_n^{(0)}(t_0)\rangle \\ &= \sum_n d_n(t) \hat{U}(t, t_0) |n^{(0)}\rangle \end{aligned}$$

$$|\psi(t)\rangle = \sum_n c_n(t) |n^{(0)}\rangle, \text{ then:}$$

$$\begin{aligned} c_n(t) &= d_n(t) \hat{U}(t, t_0) \\ &= d_n(t) e^{-\frac{i}{\hbar} \hat{H}_0 (t-t_0)} \quad \checkmark \end{aligned}$$

$$3. \quad E_i^{(0)} |i\rangle = \hat{H}_0 |i\rangle$$

$$\text{What is } \dot{d}_f? = \frac{d(d_f(t))}{dt} \quad \text{yes}$$

$$\langle f^{(0)} | \hat{H}_1(t) | \psi(t) \rangle = \sum_n \langle f^{(0)} | \hat{H}_1(t) \cdot d_n(t) e^{-\frac{i}{\hbar} \hat{H}_0 (t-t_0)} | n^{(0)} \rangle$$

$$\hat{H}_1(t) |\psi(t)\rangle = i\hbar \frac{\partial}{\partial t} |\psi(t)\rangle$$

$$\Rightarrow \hat{H}_1(t) d_n(t) e^{-\frac{i}{\hbar} \hat{H}_0 (t-t_0)} |n^{(0)}\rangle = i\hbar \frac{\partial}{\partial t} (d_n(t) e^{-\frac{i}{\hbar} \hat{H}_0 (t-t_0)} |n^{(0)}\rangle) \quad \text{why?}$$

$$\begin{aligned} \Rightarrow (\hat{H}_1(t) + \hat{H}_0) d_n(t) e^{-\frac{i}{\hbar} \hat{H}_0 (t-t_0)} |n^{(0)}\rangle &= i\hbar e^{-\frac{i}{\hbar} \hat{H}_0 (t-t_0)} (\dot{d}_n(t) |f^{(0)}\rangle + \frac{1}{i\hbar} \hat{H}_0 d_n(t) |n^{(0)}\rangle) \\ &= e^{-\frac{i}{\hbar} \hat{H}_0 (t-t_0)} (i\hbar \dot{d}_n(t) |f^{(0)}\rangle + \hat{H}_0 d_n(t) |n^{(0)}\rangle) \end{aligned}$$

$$\Rightarrow \hat{H}_1(t) d_n(t) e^{-\frac{i}{\hbar} \hat{H}_0 (t-t_0)} |n^{(0)}\rangle = i\hbar \dot{d}_n(t) e^{-\frac{i}{\hbar} \hat{H}_0 (t-t_0)} |f^{(0)}\rangle$$

$$\Rightarrow \hat{H}_1(t) d_n(t) e^{-\frac{i}{\hbar} E_n^{(0)} (t-t_0)} |n^{(0)}\rangle = i\hbar \dot{d}_n(t) e^{-\frac{i}{\hbar} E_f^{(0)} (t-t_0)} |f^{(0)}\rangle$$

$$\Rightarrow i\hbar \dot{d}_n = \sum_n \langle f^{(0)} | \hat{H}_1(t) | n^{(0)} \rangle d_n(t) e^{-i\omega_f (t-t_0)} \quad \text{with } \omega_f = \frac{E_f^{(0)} - E_n^{(0)}}{\hbar} \quad \checkmark \quad \square$$

4:

