

## Targeting interhemispheric inhibition with neuromodulation to enhance stroke rehabilitation



L.J. Boddington, J.N.J. Reynolds\*

Brain Health Research Centre and Brain Research NZ Centre of Research Excellence, Department of Anatomy, University of Otago, New Zealand

### ARTICLE INFO

#### Article history:

Received 26 February 2016

Received in revised form

10 December 2016

Accepted 10 January 2017

Available online 11 January 2017

#### Keywords:

Interhemispheric inhibition

Stroke

Rehabilitation

Neuromodulation

Transcranial magnetic stimulation

Electrical stimulation

### ABSTRACT

**Background/Objectives:** Interhemispheric inhibition in the brain plays a dynamic role in the production of voluntary unimanual actions. In stroke, the interhemispheric imbalance model predicts the presence of asymmetry in interhemispheric inhibition, with excessive inhibition from the contralesional hemisphere limiting maximal recovery. Stimulation methods to reduce this asymmetry in the brain may be promising as a stroke therapy, however determining how to best measure and modulate interhemispheric inhibition and who is likely to benefit, remain important questions.

**Methods:** This review addresses current understanding of interhemispheric inhibition in the healthy and stroke lesioned brain. We present a review of studies that have measured interhemispheric inhibition using different paradigms in the clinic, as well as results from recent animal studies investigating stimulation methods to target abnormal inhibition after stroke.

**Main findings/Discussion:** The degree to which asymmetric interhemispheric inhibition impacts on stroke recovery is controversial, and we consider sources of variation between studies which may contribute to this debate. We suggest that interhemispheric inhibition is not static following stroke in terms of the movement phase in which it is aberrantly engaged. Instead it may be dynamically increased onto perilesional areas during early movement, thus impairing motor initiation. Hence, its effect on stroke recovery may differ between studies depending on the technique and movement phase of eliciting the measurement. Finally, we propose how modulating excitability in the brain through more specific targeting of neural elements underlying interhemispheric inhibition via stimulation type, location and intensity may raise the ceiling of recovery following stroke and enhance functional return.

© 2017 Elsevier Inc. All rights reserved.

### 1. Introduction

Stroke is a leading cause of death and disability in the developed world. Unfortunately many who survive stroke never achieve complete restoration of function. Extensive physical therapy is the current best practice to maximise return of function after stroke, however, within several months many stroke survivors reach a plateau in the gains made with rehabilitation alone, leaving some


degree of residual disability affecting activities of daily life [1]. Non-invasive neuromodulation techniques such as repetitive transcranial magnetic stimulation (rTMS) have been extensively trialled to determine their effectiveness at augmenting motor recovery after stroke. Although individual gains have been reported [2–7], there is no convincing evidence of benefit over and above the ceiling of recovery that is reached after approximately six months of usual rehabilitation [1,8,9]. It is clear that new approaches are required to enhance the effects of physical therapy and elevate the ceiling of recovery, to reduce the burden of stroke on individuals and their families.

Unilateral stroke is classically regarded as leading to impaired inhibition between the cerebral hemispheres. The interhemispheric imbalance model (see Fig. 1) assumes that in the healthy brain inhibition is balanced between the hemispheres, but predicts that after a stroke there is reduced inhibition from the stroke-affected area onto the unaffected (contralesional) hemisphere. This results in increased excitability of the contralesional

**Abbreviations:** TMS, Transcranial magnetic stimulation; rTMS, repetitive transcranial magnetic stimulation; GABA,  $\gamma$ -amino butyric acid; tDCS, transcranial direct current stimulation; TBS, theta-burst stimulation; iTBS, intermittent theta-burst stimulation; cTBS, continuous theta-burst stimulation; S-IHI, Short latency interhemispheric inhibition; L-IHI, Long latency interhemispheric inhibition; iSP, ipsilateral silent period; fMRI, functional magnetic resonance imaging.

\* Corresponding author. Department of Anatomy, University of Otago School of Medical Sciences, PO Box 913, Dunedin 9054, New Zealand.

E-mail address: [john.reynolds@otago.ac.nz](mailto:john.reynolds@otago.ac.nz) (J.N.J. Reynolds).



**Fig. 1.** The interhemispheric imbalance model, which predicts dysfunction after unilateral stroke, as represented by a diagram of the rat brain with unilateral cortical stroke. The interhemispheric imbalance model predicts that each side of the brain inhibits the other equally. After a stroke, the inhibition coming from the stroke-affected hemisphere is decreased (interhemispheric dashed line) alongside a decreased excitability in the peri-lesional tissue. The unaffected hemisphere therefore becomes more excitable and exerts a stronger inhibition onto the peri-lesional tissue. Modulating this zone of inhibition in the peri-infarct tissue appears to be a potential target for stroke therapy.

hemisphere which ultimately leads to excessive interhemispheric inhibition onto ipsilesional cortical areas [10–12]. The connectivity in the residual surviving tissue could conceivably be remapped to recover lost function [13–17], however this excessive imbalanced inhibition is thought to impede neuroplasticity in these areas and limit the gains in motor recovery that can be achieved through rehabilitation [10,18,19]. Recently however, conflicting conclusions have been reported across studies as to whether a true ‘imbalance’ in interhemispheric inhibition exists in the human brain after stroke [10,20–22]. In the sections that follow, we aim to synthesise current understanding of interhemispheric inhibition after stroke and to assess the validity of current neurostimulation methods for targeting and modulating imbalanced interhemispheric inhibition. We conclude with a consideration of novel neuromodulation techniques and suggest how these alternative methods of neuromodulation may be effective in improving stroke recovery.

## 2. Interhemispheric inhibition in the healthy motor cortex

Interhemispheric inhibition in the motor cortex is the mechanism by which each hemisphere inhibits the other during the production of voluntary unimanual movements. It is thought that interhemispheric inhibition underlies the rapid inhibition of the contralateral motor cortex that is present during movement initiation, in order to suppress a ‘mirrored’ movement that may be deleterious to task performance [23–25]. This is mediated via the largest fibre bundle tract in the brain, the corpus callosum [26,27]. Evidence in humans that a transcallosal pathway conveys interhemispheric inhibitory signals despite the predominance of direct excitatory connections between motor areas [23,28–30] was first demonstrated in 1992 by Ferbert et al. The authors pioneered the use of a double transcranial magnetic stimulation (TMS) pulse paradigm using a separate coil positioned over each motor cortex. They reported that stimulating the motor cortex in one hemisphere

inhibited movement-related activity initiated by the opposite motor cortex. A suprathreshold ‘test’ pulse was applied to one hemisphere to elicit a motor-evoked potential, however a ‘conditioning’ pulse applied to the opposite hemisphere at specific intervals prior to the test pulse reduced the amplitude of the test motor-evoked potential. Since then, two distinct phases of interhemispheric inhibition elicited by the double coil method have been described, the first where conditioning to test pulse intervals are short (6–15 ms; S-IHI), and the second where the conditioning to test interval is long (40–50 ms; L-IHI) [31–34], likely representing the engagement of different neuronal mechanisms in the target motor cortex.

A single TMS pulse applied to the motor cortex can also inhibit tonic ipsilateral muscle activity elicited by sustained muscle contraction [35]. This period of inhibition is lacking in patients with agenesis of the corpus callosum [26] indicating that its origin is transcallosal in nature. Many studies have since utilised this period of inhibition as an alternative measure of interhemispheric communication, calling it the ipsilateral silent period (ISP). There is convincing evidence to suggest that interhemispheric inhibition elicited by the double TMS coil and ISP methods activate different neuronal circuitry under particular circumstances [33]. Firstly, L-IHI and ISP are thought to be mediated by GABA<sub>B</sub> receptors [31,36], whereas S-IHI is not influenced by GABA<sub>B</sub> receptor activation and may instead rely on GABA<sub>A</sub> receptors [31]. Muscle activation also modifies the measurement of S-IHI, however has little effect on L-IHI or ISP [25,33]. Furthermore, interhemispheric inhibition elicited by double TMS coil or ISP methods differentially modulates intracortical circuitry [37,38]. Altogether these studies suggest that ISP and short and long latency IHI should be considered as complementary, not equivalent measures of interhemispheric communication [33]. Since ISP measurements require continuous muscle contraction, this measure provides information related to ongoing cortical motor output during sustained isometric activation, in the absence of visible movement [39]. In contrast, interhemispheric inhibition, as measured using the double TMS coil paradigm, appears more related to the initiation of voluntary movement, since it is normally released during movement execution [10]. Since the purpose of interhemispheric inhibition is thought to be preventing unwanted bimanual movements, the double TMS coil method would therefore appear to be more suitable for interrogating the mechanisms relevant to movement initiation. The experimental ease of conducting the ISP measurement however means this method currently dominates the literature. Keeping in mind the differences in the underlying circuitry, we suggest that future studies should compare both measures before making broad claims regarding the influence that interhemispheric inhibition has on either the healthy or stroke-affected brain.

## 3. Modulation and mechanism of interhemispheric inhibition in the healthy brain

In the healthy brain, interhemispheric inhibition can be modulated using common rTMS paradigms. Short-latency interhemispheric inhibition can be directly modulated by the application of low and high frequency rTMS [40–42], quadripulse TMS [43], theta-burst stimulation (TBS) [44], or bilateral transcranial direct current stimulation (tDCS) via external electrodes [45,46]. Similarly, rTMS applied to one hemisphere has also been shown to alter cortical excitability in the unstimulated hemisphere [47–50], presumably through modulation of interhemispheric communication however the underlying mechanisms remain unknown. In order to investigate the cellular mechanisms underlying interhemispheric inhibition and its modulation by TBS, Barry et al. (2014) recently developed an *in vivo* animal model in healthy adult rats using sharp

electrode intracellular recording [51]. By recording monosynaptic potentials from inside individual pyramidal neurons in response to electrical stimulation of the ipsilateral cerebral cortex, they found that interhemispheric inhibition can be induced in the motor cortical circuitry by focal electrical stimulation of the contralateral motor cortex (see Fig. 2A). The protocols for eliciting interhemispheric inhibition were applied with the limb muscles at rest, and were modelled on those used in human double coil TMS studies [10,35,52,53]. Importantly, interhemispheric inhibition was only observed when the intensity of the conditioning electrical stimulation was set below the threshold to elicit an excitatory post-synaptic potential within the recorded neuron. Thus very low-intensity contralateral electrical stimulation seemed to preferentially target a low-threshold neural circuit which may underlie interhemispheric inhibition. Further work is required to identify the neural elements activated at low threshold by this stimulation, and to determine whether the inhibition is mono- or disynaptic. Interestingly, in the auditory system, there is evidence in other mammals of inhibition between homologous cortical areas, at latencies consistent with a direct transcallosal inhibitory circuit [54].

#### 4. Interhemispheric inhibition in the stroke-affected brain


The disruption of blood flow during a stroke prevents cells in the ischaemic area from being able to maintain ionic gradients across cell membranes and consequently leads to excitotoxic cell death. The resulting region of cell death is surrounded by the peri-infarct or penumbral tissue which displays heightened plasticity following stroke, thought to be related to remapping lost functions into the surviving tissue [55–60]. Remote electrophysiological changes in ipsilesional and contralesional hemispheres are also common [12,26,61–64] and targeting these may be beneficial at enhancing recovery after stroke.

Studies using animal models have identified that over the first week following stroke or cortical injury, excitability in the

ipsilesional hemisphere is reduced [62] and the contralesional hemisphere becomes hyper-excitable [61]. Rapid increases in contralesional excitability (within an hour following stroke onset) [65] may represent a release of interhemispheric inhibition as a result of acute damage to callosal circuitry [30]. Over time, with or without intervention, excitability in the contralesional hemisphere normalises, which appears to track with improvements in recovery [66,67]. In human stroke patients, similar changes occur where increased intracortical excitability is observed in the contralesional hemisphere in both the acute [12,68] and subacute [69,70] phases after stroke. Increased contralesional excitability has also been associated with a reduction of inhibition emanating from the stroke-affected tissue [64]. Furthermore patients with severe impairment also retain abnormal contralesional activation [71], whereas those that show substantial recovery over time show a normalisation of brain activity to activation predominantly in the ipsilesional hemisphere [72–76]. This suggests that up-regulating the ipsilesional hemisphere or down-regulating the contralesional hemisphere may be beneficial after stroke [77,78].

Evidence suggesting that stroke can also directly alter interhemispheric inhibition was first presented by Boroojerdi et al. (1996) and later supported by Shimizu et al. (2002). Reductions in interhemispheric inhibition characterized by both the double coil TMS and ISP methods were noted in stroke patients with mixed cortical-subcortical lesions but not in those with purely subcortical lesions, further suggesting that interhemispheric inhibition is predominantly mediated by circuits in the cortex [69,79].

Since the functional role of interhemispheric inhibition in the healthy brain is likely to be associated with the initiation and performance of voluntary movement, Murase et al. [10], hypothesised that meaningful abnormalities in interhemispheric inhibition after stroke would be better detected when active movements of a paretic limb were performed. Indeed in stroke patients, an abnormally strong inhibitory drive from contralesional to ipsilesional hemisphere was seen close to presentation of the go signal to



**Fig. 2. (A)** Rodent model of interhemispheric inhibition. Electrical test pulses elicit post-synaptic potentials (shown) recorded *in vivo* in the motor cortex of a rat. When an electrical conditioning stimulus (CS; orange arrows) in the opposite motor cortex, precedes an electrical test pulse (purple arrows) applied in the recorded motor cortex, at specific inter-stimulus intervals (ISI), interhemispheric inhibition is seen as a reduction in the amplitude, and maximal slope of the post-synaptic potential (seen in red). In Barry et al. (2014) this interhemispheric inhibition was abolished in healthy rats by the application of low intensity electrical iTBS to the same conditioning electrode. **(B)** Electrical iTBS applied to the unaffected hemisphere after stroke enhances stroke recovery in rats. Rats with focal lesions in the primary motor cortex received either sham stimulation or electrical iTBS or cTBS over a number of days after stroke. Application of iTBS enhanced recovery on a grid-walking task compared to both cTBS or sham stimulation, suggesting that this may be through modulation of interhemispheric inhibition. Figures reprinted from Experimental Neurology, Barry et al. (2014), Vol 261, 258–266; with permission from Elsevier. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

commence voluntary movement in their paretic hand, suggesting that abnormal interhemispheric inhibition was present during movement preparation. A greater imbalance of interhemispheric inhibition between contralesional and ipsilesional hemispheres, also correlated with poorer performance [10,21], thus it has been proposed that increased inhibition in the ipsilesional hemisphere may limit maximal functional recovery after stroke and contribute to residual motor disability [10,21]. These findings have been fundamental in shaping the interhemispheric imbalance model and many neuromodulatory therapies have been based on this premise of an asymmetric interhemispheric inhibition [80].

Evidence in favor of the interhemispheric imbalance model is apparent in two rare case studies where the symptoms associated with a single hemisphere stroke were alleviated by a second stroke that occurred in the opposite hemisphere [18,19]. Vuilleumier et al. 1996 described a patient initially presenting with a right-sided parietal infarct resulting in severe unilateral neglect. Notably the symptoms of neglect abruptly disappeared after the patient experienced a second stroke in the left frontal lobe. More recently, Sauerbrei & Liepert (2012) reported a similar case of a patient initially presenting with severe right-sided hemiparesis after a left-hemisphere haemorrhage. Remarkably the hemiparesis disappeared almost completely after a later right-parietal infarct. In both of these reported cases, the investigators state that the rapid time course of recovery seen in their patients cannot entirely be explained by the brain's natural ability to spontaneously recover, and they speculate that rebalancing of interhemispheric rivalry played a key role in their patients' recovery. Of note there have also been reports of worsening function after a second contralesional stroke [13,15,16]. However, all were due to subcortical infarcts, which are less likely to affect interhemispheric inhibition than a subsequent cortical infarct [69,79].

The presence of altered interhemispheric inhibition after stroke is not universally agreed, with different groups reporting variable results. In some studies, no asymmetry in interhemispheric inhibition from the unaffected hemisphere was detected [20,81,82]. Others have demonstrated: a greater inhibition from the ipsilesional onto contralesional hemispheres [83], that the contralesional hemisphere does not over-inhibit the ipsilesional hemisphere [84], and that reducing contralesional inhibition results in poorer performance [85], indeed all opposing the interhemispheric inhibition model. A common factor linking all of these studies is that the measurement of interhemispheric inhibition using the double TMS coil method or iSP was performed at rest or during sustained isometric contraction, respectively.

Functional connectivity and dynamic causal modelling analyses of fMRI data suggest that there are striking differences in interhemispheric connectivity during active movements of the paretic limb compared with the limb at rest [78,86,87]. During active movements in stroke patients (typically repetitive fist closures), there is a stronger inhibitory influence of the contralesional motor cortex onto the ipsilesional motor cortex [78,86] and a disinhibition of the contralesional hemisphere from the ipsilesional hemisphere [87]. This disruption to functional connectivity originating contralesionally also predicts poorer motor performance [78,86,88,89]. In comparison, in stroke patients at rest there is normal connectivity between motor areas compared with healthy controls [86]. These findings collectively suggest that interhemispheric inhibition is dynamically influenced by movement and is likely to be more prominent and aberrantly engaged in the peri-lesional areas during the critical movement preparation and early execution periods as opposed to when the affected limb is completely at rest or isometrically contracted. To verify this empirically, measurement of iSPs during sustained muscle activation or interhemispheric inhibition either at rest or during sustained isometric contraction may

not be capturing an abnormally enhanced interhemispheric inhibition, which may be impairing the processes of initiating a useful movement after chronic stroke. It therefore is likely that the best approach by which to detect asymmetric interhemispheric inhibition after stroke is by applying the double coil method during production of a voluntary movement, where the active functional role of such inhibition is to prevent bimanual movements, and not when the limb is at rest.

**Restoration of symmetry in interhemispheric connections and cortical excitability after stroke does not always predict functional improvement [90,91].** Those with large infarcts may instead show significant remapping of cortical representations to remote areas not normally involved in the lost function, a process known as vicariation [92,93]. The recently proposed 'Bimodal Balance-Recovery' model suggests that neither the interhemispheric imbalance model, nor the vicariation model, can predict recovery in all patients [94]. It proposes that if there is a high structural reserve, such as if some residual function remains due to partial retention of the integrity of the neural circuitry of the corticospinal tract, then the interhemispheric imbalance model will dominate and in these cases it may be beneficial to utilise therapies for rebalancing interhemispheric inhibition [95]. **When structural reserve is low, vicariation will dominate, with more distant, even contralesional, areas recruited to take over lost function [93,94],** which may explain why severely impaired patients often retain abnormal contralesional activation [96–98]. Thus in these cases, interhemispheric rebalancing may not be beneficial and these patients may benefit from facilitation of contralesional activity instead [95]. Utilising information about lesion size and corticospinal tract integrity will likely help in the stratification of patients towards therapies that are likely to be of most benefit for each individual [99]. Plow et al. (2016) suggest that neuromodulation techniques that boost plasticity based on the Bimodal Balance-Recovery model will result in better outcomes for patients, and are aiming to determine specific neurophysiological markers and cut-offs that may help to identify which patients will benefit from particular neuromodulation therapies. Clarification of these markers will certainly assist in determining the influence that interhemispheric imbalance has on functional recovery after stroke.

## 5. Current methods of neuromodulation to target interhemispheric inhibition

Understanding who is likely to benefit from 'rebalancing' therapies is an important first step towards raising the ceiling of maximum functional return that is currently reached through rehabilitation and spontaneous recovery. The next step for stroke survivors exhibiting a degree of interhemispheric imbalance amenable to therapy, is knowing how best to modulate excitability in the brain to accelerate and augment functional recovery. Neuromodulation through electrical or magnetic stimulation has offered much promise as techniques to augment stroke rehabilitation, however little is known of the effects of these therapies on brain circuits. Repetitive TMS or transcranial direct current stimulation (tDCS) are two non-invasive stimulation techniques that have been trialled clinically with the aim of modulating brain excitability in order to improve function following stroke [98,100]. The use of these stimulation techniques alongside or to 'prime' standard physical rehabilitation has become a large area of interest in the field, however there is still little consensus on the optimal stimulation paradigms or neuroanatomical targets to best maximise the benefits of rehabilitation.

**Commonly, non-invasive brain stimulation is used over the ipsilesional hemisphere to enhance excitability in the peri-infarct tissue.** Anodal (excitatory) tDCS over the affected hemisphere of



chronic stroke patients has been shown to transiently improve hand motor function on the Jebsen Taylor hand function test [2]. Similarly ipsilesional application of rTMS, either alone or combined with physical rehabilitation, has also been shown to enhance upper limb function with effects persisting from ten days to several months after the end of treatment [3–7,101]. Not all have seen success with this approach however. High frequency rTMS applied ipsilesionally alongside constraint-induced therapy did not enhance motor function [9] and has been shown to differentially affect cortical and subcortical stroke where it enhanced dexterity after subcortical stroke, but worsened dexterity after cortical stroke [102].

Another approach that has been utilised with some success is to down-regulate the excitability in the unaffected hemisphere, where excessive interhemispheric inhibition is said to originate [103,104], or to target interhemispheric inhibition directly [11]. Low frequency rTMS applied to the contralesional hemisphere can enhance various measures of paretic upper limb performance [7,105–111]. Additionally contralesional low-frequency rTMS can enhance L-IHI [11], and reduce the extent and duration of interhemispheric inhibition from the contra-to ipsilesional hemispheres [106,112,113] which correlates with rTMS-induced improvements in motor function [106,113]. Forms of TBS [114] have also commonly been applied using rTMS. Both intermittent theta-burst stimulation (iTBS) applied to the ipsilesional motor cortex [5,115], or continuous theta-burst stimulation (cTBS) applied to the contralesional motor cortex have been shown to improve paretic hand performance during a task, although contralesional cTBS has also been associated with reduced overall limb function [5].

It is important to note here that there are inconsistencies between studies investigating the effects of rTMS on motor function, or on physiological measures such as interhemispheric inhibition [8,116,117]. Not all studies demonstrate that rTMS can influence interhemispheric inhibition [52,118,119]. Indeed the effects of rTMS and TBS specifically have also been frequently shown to be highly variable between individuals [120,121]. Analysis of a large sample of participants has revealed that changing the orientation of the TMS coil delivering the stimulation produces different effects between individuals, possibly due to activation of different neural elements in each participant [122]. Alongside inter-individual variability, various studies have shown that the same TMS pulse can facilitate or suppress consequent cortical responses depending on the timing of the pulse relative to prior spontaneous or evoked brain activity [123,124]. Similarly, repeated application of the same tDCS protocols can induce opposing effects on motor cortex excitability depending on the time interval between the application of the different sets of stimulation [125]. These differences in prior brain-state and subsequent differences in homeostatic plasticity may therefore account for some of the inconsistencies between, or even within studies.

The vast parameter space of single or multiple stimulation patterns and sites, combined with the variability of neural responses in healthy individuals as a consequence of prior brain-states [123,124] make understanding the effects of rTMS in healthy individuals alone a daunting task. This is yet further amplified when attempting to use rTMS paradigms in stroke patients, who on their own represent a highly variable and heterogeneous population. This highlights a critical need for studies investigating the cellular mechanisms engaged by rTMS and how this may differ in the stroke-affected brain [126]. This should be followed by systematic investigations of how to optimise stimulation parameters for individuals before these approaches find their way into routine clinical practice for the treatment of stroke.

## 6. Novel methods of neuromodulation to target interhemispheric inhibition

Considering the inconsistencies of rTMS protocols, motor recovery following stroke might be enhanced using a therapy that does not attempt to grossly rebalance hemispheric excitability, but rather one that targets specific functional circuitry related to interhemispheric inhibition and movement activation. One approach is to drive passive movements of an affected limb via a device that is mechanically coupled to the movements generated in the other limb [127], essentially physiologically 'priming' specific motor circuits [53] instead of unfocused global hemispheric priming using rTMS. This approach enhanced the rate at which patients reached their peak motor recovery post stroke, and the course of improvement correlated with progressive increases in interhemispheric inhibition from the peri-infarct area to the contralesional hemisphere [128]. In healthy individuals, passive mechanical coupling also leads to reduced interhemispheric inhibition from the active onto the passive hemisphere [53], and a short-lasting ( $\approx 30$  min) disinhibition of cortical areas representing the passive limb [53,129]. Interestingly this effect was most pronounced when symmetrical (mirror) movements were induced compared to asymmetrical movements. Collectively, this evidence suggests that focussed disinhibition may involve plasticity in transcallosal circuits normally specialised in preventing mirror movements. This short period of disinhibition may present a brief therapeutic window in which accelerated rehabilitation gains can be achieved, despite ongoing peri-lesional tonic inhibition. We propose that effective augmentation of stroke rehabilitation in cases where interhemispheric inhibition is dynamically enhanced requires targeted neuromodulation of the specific neural elements underlying interhemispheric inhibition.

Whilst not yet sufficiently developed for translation into human use, viral targeting and optogenetic modulation of specific interhemispheric circuits might present a novel way of specifically modulating the neural elements underlying abnormal interhemispheric inhibition after stroke [130]. The benefit of optogenetic stimulation is in the ability to selectively target particular neuronal projections, from within a specific brain nucleus. In comparison, the cellular effects of TMS or tDCS are much less focal, activating mixed populations of neurons and producing a range of effects that may ultimately cancel. Rodent studies have shown that precise activation of specific neuronal subtypes using optogenetics with patterns that mimic physiologically normal brain activity can enhance function in neurological disorders better than simple repetitive stimulation protocols [131]. If specific excitatory or inhibitory transcallosal circuitry were found to be abnormally active in animal models of stroke, theoretically the same circuitry could be identified and targeted in humans in order to optogenetically activate or inhibit this abnormal activity. The benefits of human translation of these optogenetic techniques are being discussed [132] and therefore may eventually become a powerful tool in the neuromodulation toolbox available to improve function in neurological conditions.

In the meantime, electrical stimulation using implanted electrodes may represent the most focal approach for brain stimulation of specific neural areas. In an animal model of interhemispheric inhibition, low-intensity TBS paradigms seemed to specifically target neural elements underlying interhemispheric inhibition. Specifically, electrical iTBS applied to the hemisphere contralateral to the recorded neuron via an implanted electrode, abolishes interhemispheric inhibition from the stimulated to the recorded hemisphere [51]. Low intensity electrical stimulation is known to activate low-threshold cortical neurons [37], thus in this context the targeted neural circuits might be those involved in inhibition

onto the recorded hemisphere. In a rat stroke model, iTBS applied at similar low intensities via an implanted contralesional electrode accelerated motor recovery (see Fig. 2B). Following three weeks of this electrical stimulation, limb function was 40% better than that measured in sham-stimulated rats [51]. Interestingly, low-intensity cTBS tended to impede recovery, raising the possibility that electrical cTBS, which potentiated monosynaptic connections from the stimulated cortex onto pyramidal cells in the contralateral motor cortex, may have also up-regulated inhibitory circuits (monosynaptic or polysynaptic) underlying interhemispheric inhibition, raising the threshold for activation of pyramidal motor neurons. Invasive stimulation of this nature may facilitate modulation of specific neuronal circuitry, in a more targeted manner than can be achieved using tDCS or TMS.

Invasive cortical electrical stimulation has been previously trialled in human stroke patients in the Northstar Neuroscience EVEREST Phase III trial. This trial attempted to enhance excitability in the surviving tissue in order to augment recovery from stroke and showed very promising Phase I and II results [133]. The Phase III trial failed to reach its primary efficacy end-point, however it was noted that extradural stimulating electrodes were not placed over viable tissue in many cases [134–137]. Post-hoc analysis revealed that in patients with stimulating electrodes placed over viable cortical tissue, 69% displayed a significant increase on the upper extremity Fugl-Meyer and Arm Motor Ability Tests at 4, 12 and 24-week assessment points compared with controls [138]. Whilst overall the EVEREST trial did not meet its primary end-point, this should not rule out more invasive stimulation methods as a potential option for some patients. Application of electrical stimulation to the contralesional hemisphere, where tissue still remains viable, may represent a method to dynamically modulate interhemispheric inhibition and normalise interhemispheric asymmetry when it is at its most disadvantageous following stroke, such as close to movement initiation [10]. An implanted electrode could facilitate longer or even continuous application of stimulation throughout activities of daily living, circumventing limitations of TMS, which currently requires costly external equipment, and application by a skilled operator.

## 7. Conclusions

Interhemispheric imbalance is one of the phenomena that could be targeted by neuromodulation following stroke to maximise recovery, and yet in only a proportion of patients the restoration of symmetry in interhemispheric communication appears to facilitate accelerated or enhanced recovery. Although previous studies have proven variable, we believe that future investigations should be designed with the assessment of interhemispheric inhibition made in the context of movement initiation in order to obtain a more meaningful picture of the negative impact of asymmetry on functional outcomes. We propose that interhemispheric inhibition in stroke recovery is a dynamic phenomenon strongly related to movement initiation, rather than one that remains static and persistently altered following stroke.

Modulating interhemispheric communication through invasive or non-invasive brain stimulation methods holds promise for augmenting rehabilitation therapy. However it is important to gain a greater understanding of the cellular mechanisms engaged by these treatments, since this will ultimately allow therapies to be more specifically targeted to the pathology affecting individuals. Well-designed animal experiments are tremendously valuable in this regard. Precise modulation of disrupted circuitry using more focal or specific neuronal targeting (including implanted electrical or optogenetic technology) may form the next generation of therapies for enhancing or accelerating recovery. Overall, modulation of

interhemispheric inhibition after stroke alongside conventional rehabilitation does remain a potential avenue for therapy to allow the ceiling of stroke recovery to be raised.

## Conflict of interest statement

The Authors declare no conflicts of interest.

## Acknowledgements

Laura Boddington was supported by a W & B Miller Neurological Foundation of New Zealand Scholarship (partially funded by the HB Williams Turanga Trust), a University of Otago Doctoral Scholarship and a University of Otago Postgraduate Publishing Bursary. John Reynolds was supported by a Rutherford Discovery Fellowship from the Royal Society of New Zealand. We thank Natalie Matheson and Rose Melchers for their helpful comments on the manuscript.

## References

- [1] Schaechter JD. Motor rehabilitation and brain plasticity after hemiparetic stroke. *Prog Neurobiol* 2004 May;73(1):61–72.
- [2] Hummel F, Celnik P, Giraux P, Flöel A, Wu W-H, Gerloff C, et al. Effects of non-invasive cortical stimulation on skilled motor function in chronic stroke. *Brain* 2005 Mar;128(Pt 3):490–9.
- [3] Khedr EM, Abdel-Fadeil MR, Farghali A, Qaid M. Role of 1 and 3 Hz repetitive transcranial magnetic stimulation on motor function recovery after acute ischaemic stroke. *Eur J Neurol* 2009 Dec;16(12):1323–30.
- [4] Khedr EM, Etraby AE, Hemeda M, Nasef AM, Razek AAE. Long-term effect of repetitive transcranial magnetic stimulation on motor function recovery after acute ischemic stroke. *Acta Neurol Scand* 2009 Dec 8;121(1):30–7. Blackwell Publishing Ltd.
- [5] Ackerley SJ, Stinear CM, Barber PA, Byblow WD. Combining theta burst stimulation with training after subcortical stroke. *Stroke* 2010 Jul;41(7):1568–72.
- [6] Chang WH, Kim Y-H, Bang OY, Kim ST, Park YH, Lee PKW. Long-term effects of rTMS on motor recovery in patients after subacute stroke. *J Rehabil Med* 2010 Sep;42(8):758–64.
- [7] Emara TH, Moustafa RR, ElNahas NM, ElGanzoury AM, Abdo TA, Mohamed SA, et al. Repetitive transcranial magnetic stimulation at 1Hz and 5Hz produces sustained improvement in motor function and disability after ischaemic stroke. *Eur J Neurol* 2010 Apr 8;17(9):1203–9.
- [8] Talelli P, Wallace A, Dileone M, Hoad D, Cheeran B, Oliver R, et al. Theta burst stimulation in the rehabilitation of the upper limb: a semirandomized, placebo-controlled trial in chronic stroke patients. *Neurorehabil Neural Repair* 2012 Oct;26(8):976–87.
- [9] Malcolm MP, Triggs WJ, Light KE, Gonzalez Rothi LJ, Wu S, Reid K, et al. Repetitive transcranial magnetic stimulation as an adjunct to constraint-induced therapy. *Am J Phys Med Rehabil* 2007 Sep;86(9):707–15.
- [10] Murase N, Duque J, Mazzocchio R, Cohen LG. Influence of interhemispheric interactions on motor function in chronic stroke. *Ann Neurol* 2004;55(3):400–9.
- [11] Kirton A, deVeber G, Gunraj C, Chen R. Cortical excitability and interhemispheric inhibition after subcortical pediatric stroke: plastic organization and effects of rTMS. *Clin Neurophysiol* 2010 Nov;121(11):1922–9.
- [12] Liepert J, Hamzei F, Weiller C. Motor cortex disinhibition of the unaffected hemisphere after acute stroke. *Muscle Nerve* 2000 Nov;23(11):1761–3.
- [13] Ago T, Kitazono T, Ooboshi H, Takada J. Deterioration of pre-existing hemiparesis brought about by subsequent ipsilateral lacunar infarction. *J Neurol* 2003;247(8):1152–3.
- [14] Nudo RJ. Mechanisms for recovery of motor function following cortical damage. *Curr Opin Neurobiol* 2006 Dec;16(6):638–44.
- [15] Song Y-M, Lee J-Y, Park J-M, Yoon B-W, Roh J-K. Ipsilateral hemiparesis caused by a corona radiata infarct after a previous stroke on the opposite side. *Arch Neurol* 2005 May;62(5):809–11.
- [16] Kobayashi Z, Akaza M, Endo H, Numasawa Y, Tomimitsu H, Shintani S. Deterioration of pre-existing hemiparesis due to an ipsilateral internal capsule infarction after a contralateral stroke. *J Neurol Sci* 2015 Jul 15;354(1–2):140–1.
- [17] Nudo RJ, Wise BM, SiFuentes F, Milliken GW. Neural substrates for the effects of rehabilitative training on motor recovery after ischemic infarct. *Science* 1996 Jun 21;272(5269):1791–4.
- [18] Vuilleumier P, Hester D, Assal G, Regli F. Unilateral spatial neglect recovery after sequential strokes. *Neurology* 1996 Jan 1;46(1):184–9.
- [19] Sauerbrei R, Liepert J. Support of the concept of interhemispheric rivalry by two consecutive strokes occurring in both hemispheres: a case study. *J Neurol* 2012 Nov;259(11):2484–5.

- [20] Stinear CM, Petoe MA, Byblow WD. Primary motor cortex excitability during recovery after stroke: implications for neuromodulation. *Brain Stimul* 2015 Nov;8(6):1183–90.
- [21] Duque J, Hummel F, Celnik P, Murase N, Mazzocchio R, Cohen LG. Transcallosal inhibition in chronic subcortical stroke. *NeuroImage* 2005 Dec;28(4):940–6.
- [22] Borich MR, Wheaton LA, Brodie SM, Lakhani B. Evaluating interhemispheric cortical responses to transcranial magnetic stimulation in chronic stroke: a TMS-EEG investigation. *Neuroscience* 2016;618:25–30.
- [23] Beaulé V, Tremblay S, Théoret H. Interhemispheric control of unilateral movement. *Neural Plast* 2012;2012:627816.
- [24] Mayston MJ, Harrison LM, Stephens JA. A neurophysiological study of mirror movements in adults and children. *Ann Neurol* 1999 May;45(5):583–94.
- [25] Hinder MR, Schmidt MW, Garry MI, Summers JJ. Unilateral contractions modulate interhemispheric inhibition most strongly and most adaptively in the homologous muscle of the contralateral limb. *Exp Brain Res* 2010 Sep;205(3):423–33.
- [26] Meyer BU, Röricht S, Gräfin von Einsiedel H, Kruggel F, Weindl A. Inhibitory and excitatory interhemispheric transfers between motor cortical areas in normal humans and patients with abnormalities of the corpus callosum. *Brain* 1995 Apr;118:429–40.
- [27] Di Lazzaro V, Oliviero A, Profice P, Insola A. Direct demonstration of interhemispheric inhibition of the human motor cortex produced by transcranial magnetic stimulation. *Exp Brain Res* 1999;124:520–4.
- [28] Porter LL, White EL. Synaptic connections of callosal projection neurons in the vibrissal region of mouse primary motor cortex: an electron microscopic/horseradish peroxidase study. *J Comp Neurol* 1986 Jun 22;248(4):573–87.
- [29] Conti F, Manzoni T. The neurotransmitters and postsynaptic actions of callosally projecting neurons. *Behav Brain Res* 1994;64:37–53.
- [30] Dancause N, Touvykine B, Mansoori BK. Inhibition of the contralesional hemisphere after stroke: reviewing a few of the building blocks with a focus on animal models. *Prog Brain Res* 2015;218:361–87. Elsevier.
- [31] Irlbacher K, Brocke J, Mechow JV, Brandt SA. Effects of GABA(A) and GABA(B) agonists on interhemispheric inhibition in man. *Clin Neurophysiol* 2007 Feb;118(2):308–16.
- [32] Ni Z, Gunraj C, Nelson AJ, Yeh I-J, Castillo G, Hoque T, et al. Two phases of interhemispheric inhibition between motor related cortical areas and the primary motor cortex in human. *Cereb Cortex* 2009 Jul;19(7):1654–65.
- [33] Chen R, Yung D, Li J-Y. Organization of ipsilateral excitatory and inhibitory pathways in the human motor cortex. *J Neurophysiol* 2003 Mar;89(3):1256–64.
- [34] Kukawadia S, Wagle-Shukla A, Morgante F, Gunraj C, Chen R. Interactions between long latency afferent inhibition and interhemispheric inhibitions in the human motor cortex. *J Physiol (Lond)* 2005 Mar 8;563(3):915–24.
- [35] Ferbert A, Priori A, Rothwell JC, Day BL, Colebatch JG, Marsden CD. Interhemispheric inhibition of the human motor cortex. *J Physiol (Lond)* 1992;453:525–46.
- [36] Werhahn KJ, Kunesch E, Noachtar S, Benecke R, Classen J. Differential effects on motorcortical inhibition induced by blockade of GABA uptake in humans. *J Physiol (Lond)* 1999 Jun 1;517(2):591–7.
- [37] Daskalakis ZJ, Christensen BK, Fitzgerald PB, Roshan L, Chen R. The mechanisms of interhemispheric inhibition in the human motor cortex. *J Physiol (Lond)* 2004 Jul 22;543(1):317–26.
- [38] Avanzino L, Teo JTH, Rothwell JC. Intracortical circuits modulate transcallosal inhibition in humans. *J Physiol (Lond)* 2007 Aug 15;583(Pt 1):99–114.
- [39] Giovannelli F, Borgheresi A, Balestrieri F, Zaccara G, Viggiano MP, Cincotta M, et al. Modulation of interhemispheric inhibition by volitional motor activity: an ipsilateral silent period study. *J Physiol (Lond)* 2009 Nov 15;587(Pt 22):5393–410.
- [40] Pal PK, Hanajima R, Gunraj CA, Li J-Y, Wagle-Shukla A, Morgante F, et al. Effect of low-frequency repetitive transcranial magnetic stimulation on interhemispheric inhibition. *J Neurophysiol* 2005 Sep;94(3):1668–75.
- [41] Gilio F, Rizzo V, Siebner HR, Rothwell JC. Effects on the right motor hand-area excitability produced by low-frequency rTMS over human contralateral homologous cortex. *J Physiol (Lond)* 2004 Jul 16;551(2):563–73.
- [42] Cincotta M, Giovannelli F, Borgheresi A, Balestrieri F, Zaccara G, Inghilleri M, et al. Modulatory effects of high-frequency repetitive transcranial magnetic stimulation on the ipsilateral silent period. *Exp Brain Res* 2006 Jun;171(4):490–6.
- [43] Tsutsumi R, Hanajima R, Terao Y, Shirota Y, Ohminami S, Shimizu T, et al. Effects of the motor cortical quadripulse transcranial magnetic stimulation (QPS) on the contralateral motor cortex and interhemispheric interactions. *J Neurophysiol* 2014 Jan;111(1):26–35. American Physiological Society.
- [44] Premji A, Zapallow C, Tsang P, Tang R, Jacobs M, Nelson AJ. Influence of area 5 on interhemispheric inhibition. *NeuroReport* 2011 Dec 21;22(18):974–8.
- [45] Tazoe T, Endoh T, Kitamura T, Ogata T. Polarity specific effects of transcranial direct current stimulation on interhemispheric inhibition. *PLoS One* 2014;9(12):e114244.
- [46] Williams JA, Pascual-Leone A, Fregni F. Interhemispheric modulation induced by cortical stimulation and motor training. *Phys Ther* 2010 Mar 1;90(3):398–410. American Physical Therapy Association.
- [47] Gersler A, Bäumer T, Weiller C, Münchau A, Liepert J. Interhemispheric effects of high and low frequency rTMS in healthy humans. *Clin Neurophysiol* 2003 Oct;114(10):1800–7.
- [48] Schambra HM, Sawaki L, Cohen LG. Modulation of excitability of human motor cortex (M1) by 1 Hz transcranial magnetic stimulation of the contralateral M1. *Clin Neurophysiol* Jan 2003;114:130–3.
- [49] Kobayashi M, Hutchinson S, Théoret H, Schlaug G, Pascual-Leone A. Repetitive TMS of the motor cortex improves ipsilateral sequential simple finger movements. *Neurology* 2004 Jan 13;62(1):91–8.
- [50] Plewnia C, Lotze M, Gerloff C. Disinhibition of the contralateral motor cortex by low-frequency rTMS. *NeuroReport* 2003 Mar 24;14(4):609.
- [51] Barry MD, Boddington LJ, Igelström KM, Gray JP, Shemmell J, Tseng KY, et al. Utility of intracerebral theta burst electrical stimulation to attenuate interhemispheric inhibition and to promote motor recovery after cortical injury in an animal model. *Exp Neurol* 2014 Nov;261:258–66.
- [52] Suppa A, Ortu E, Zafar N, Deriu F, Paulus W, Berardelli A, et al. Theta burst stimulation induces after-effects on contralateral primary motor cortex excitability in humans. *J Physiol (Lond)* 2008 Sep 15;586(18):4489–500.
- [53] Byblow WD, Stinear CM, Smith M-C, Bjerre L, Flakager BK, McCambridge AB. Mirror symmetric bimanual movement priming can increase corticomotor excitability and enhance motor learning. *PLoS One* 2012;7(3):1–14.
- [54] Kitzes LM, Doherty D. Influence of callosal activity on units in the auditory cortex of ferret (*Mustela putorius*). *J Neurophysiol* 1994 May 1;71(5):1740–51. American Physiological Society.
- [55] Carmichael T, Chesselet M-F. Synchronous neuronal activity is a signal for axonal sprouting after cortical lesions in the adult. *J Neurosci* 2002 Jul 15;22(14):6062–70.
- [56] Carmichael T, Archibeque I, Luke L, Nolan T, Momiy J, Li S. Growth-associated gene expression after stroke: evidence for a growth-promoting region in peri-infarct cortex. *Exp Neurol* 2005 Jun;193(2):291–311.
- [57] Nudo RJ, Milliken GW. Reorganization of movement representations in primary motor cortex following focal ischemic infarcts in adult squirrel monkeys. *J Neurophysiol Am Physiol Soc* 1996;75(5):2144–9.
- [58] Calabresi P, Centonze D, Pisani A, Cupini LM. Synaptic plasticity in the ischaemic brain. *Lancet Neurol* 2003 Oct;2(10):622–9.
- [59] Schiene K, Bruehl C, Zilles K, QM, Hagemann G, Kraemer M, et al. Neuronal hyperexcitability and reduction of GABA<sub>A</sub>-receptor expression in the surround of cerebral photothrombosis. *J Cereb Blood Flow Metab* 1996 Sep;16:906–14.
- [60] Hagemann G, Redecker C, Neumann-Haefelin T, Freund H-J, Witte OW. Increased long-term potentiation in the surround of experimentally induced focal cortical infarction. *Ann Neurol* 1998 Aug;44(2):255–8.
- [61] Buchkremer-Ratzmann I, Witte OW. Extended brain disinhibition following small photothrombotic lesions in rat frontal cortex. *NeuroReport* 1997 Jan;8(2):519–22.
- [62] Neumann-Haefelin T, Witte OW. Periinfarct and Remote Excitability Changes after transient middle cerebral artery occlusion. *J Cereb Blood Flow Metab* 2000 Jan;20(1):45–52.
- [63] Witte OW, Bidmon HJ, Schiene K, Redecker C, Hagemann G. Functional differentiation of multiple perilesional zones after focal cerebral ischemia. *J Cereb Blood Flow Metab* 2000 Aug;20(8):1149–65.
- [64] Niehaus L, Bajbouj M, Meyer B-U. Impact of interhemispheric inhibition on excitability of the non-lesioned motor cortex after acute stroke. In: Paulus W, Tergau F, Nitsche MA, Rothwell JC, Ziemann U, Hallett M, editors. Transcranial magnetic stimulation and transcranial direct current stimulation [Internet]. Elsevier; 2003. p. 181–6.
- [65] Mohajerani MH, Aminoltejeri K. Targeted mini-strokes produce changes in interhemispheric sensory signal processing that are indicative of disinhibition within minutes. 2011.
- [66] Mansoori BK, Jean-Charles L, Touvykine B, Liu A, Quessy S, Dancause N. Acute inactivation of the contralesional hemisphere for longer durations improves recovery after cortical injury. *Exp Neurol* 2014 Apr;254:18–28.
- [67] Jablonka JA, Burnat K, Witte OW, Kossut M. Remapping of the somatosensory cortex after a photothrombotic stroke: dynamics of the compensatory reorganization. *Neuroscience* 2010 Jan 13;165(1):90–100.
- [68] Manganotti P, Patuzzo S, Cortese F, Palermo A, Smania N, Fiaschi A. Motor disinhibition in affected and unaffected hemisphere in the early period of recovery after stroke. *Clin Neurophysiol* 2002 Jun;113(6):936–43.
- [69] Shimizu T, Hosaki A, Hino T, Sato M, Komori K, Hirai S, et al. Motor cortical disinhibition in the unaffected hemisphere after unilateral cortical stroke. *Brain* 2002 Aug;125(Pt 8):1896–907.
- [70] Bütefisch CM, Netz J, Wessling M, Seitz RJ, Hömberg V. Remote changes in cortical excitability after stroke. *Brain Oxford University Press* 2003 Feb;vol. 126(Pt 2):470–81.
- [71] Nelles G, Spiekramann G, Jueptner M, Leonhardt G, Müller S, Gerhard H, et al. Evolution of functional reorganization in hemiplegic stroke: a serial positron emission tomographic activation study. *Ann Neurol* 1999 Dec;46(6):901–9.
- [72] Marshall RS, Perera GM, Lazar RM, Krakauer JW, Constantine RC, DeLaPaz RL. Evolution of cortical activation during recovery from corticospinal tract infarction. *Stroke* 2000 Mar;31(3):656–61.
- [73] Ward NS, Brown MM, Thompson AJ, Frackowiak RSJ. Neural correlates of motor recovery after stroke: a longitudinal fMRI study. *Brain* 2003 Nov;126(Pt 11):2476–96.
- [74] Rehme AK, Eickhoff SB, Rottschy C, Fink GR, Grefkes C. Activation likelihood estimation meta-analysis of motor-related neural activity after stroke. *NeuroImage* 2012 Feb 1;59(3):2771–82.



- [75] Liepert J, Miltner W, Bauder H, Sommer M. Motor cortex plasticity during constraint-induced movement therapy in stroke patients. *Neurosci Lett* 1998;250:5–8.
- [76] Harris-Love ML, Chan E, Dromerick AW, Cohen LG. Neural substrates of motor recovery in severely impaired stroke patients with hand paralysis. *Neurorehabil Neural Repair* 2016 May;30(4):328–38. SAGE Publications.
- [77] Di Lazzaro V, Profice P, Pilato F, Capone F, Ranieri F, Pasqualetti P, et al. Motor cortex plasticity predicts recovery in acute stroke. *Cereb Cortex* 2010 Jul;20(7):1523–8.
- [78] Rehme AK, Eickhoff SB, Wang LE, Fink GR, Grefkes C. Dynamic causal modeling of cortical activity from the acute to the chronic stage after stroke. *NeuroImage* 2011 Apr 1;55(3):1147–58. Elsevier Inc.
- [79] Boroojerdi B, Diefenbach K, Ferbert A. Transcallosal inhibition in cortical and subcortical cerebral vascular lesions. *J Neurol Sci* 1996 Dec;144(1–2):160–70.
- [80] Nowak DA, Grefkes C, Ameli M, Fink GR. Interhemispheric competition after stroke: brain stimulation to enhance recovery of function of the affected hand. *Neurorehabil Neural Repair* 2009 Sep;23(7):641–56.
- [81] Butefisch CM, We ling M, Netz J, Seitz RJ, Homberg V. Relationship between interhemispheric inhibition and motor cortex excitability in subacute stroke patients. *Neurorehabil Neural Repair* 2007 Apr 4;22(1):4–21.
- [82] Cassidy JM, Chu H, Anderson DC, Krach LE, Snow L, Kimberley TJ, et al. A comparison of primed low-frequency repetitive transcranial magnetic stimulation treatments in chronic stroke. *Brain Stimul* 2015 Nov;8(6):1074–84.
- [83] Borich MR, Neva JL, Boyd LA. Evaluation of differences in brain neurophysiology and morphometry associated with hand function in individuals with chronic stroke. *Restor Neurol Neurosci* 2015;33(1):31–42. IOS Press.
- [84] Dimyan MA, Perez MA, Auh S, Tarula E, Wilson M, Cohen LG. Nonparetic arm force does not overinhibit the paretic arm in chronic poststroke hemiparesis. *Arch Phys Med Rehabil* 2014 May;95(5):849–56.
- [85] Mang CS, Borich MR, Brodie SM, Brown KE, Snow NJ, Wadden KP, et al. Diffusion imaging and transcranial magnetic stimulation assessment of transcallosal pathways in chronic stroke. *Clin Neurophysiol* 2015 Oct 1;126(10):1959–71. International Federation of Clinical Neurophysiology.
- [86] Grefkes C, Nowak DA, Eickhoff SB, Dafotakis M, Küst J, Karbe H, et al. Cortical connectivity after subcortical stroke assessed with functional magnetic resonance imaging. *Ann Neurol* 2008 Feb;63(2):236–46.
- [87] Volz LJ, Sarfeld A-S, Diekhoff S, Rehme AK, Pool E-M, Eickhoff SB, et al. Motor cortex excitability and connectivity in chronic stroke: a multimodal model of functional reorganization. *Brain Struct Funct* 2015 Mar;220(2):1093–107.
- [88] Carter AR, Astafiev SV, Lang CE, Connor LT, Rengachary J, Strube MJ, et al. Resting interhemispheric functional magnetic resonance imaging connectivity predicts performance after stroke. *Ann Neurol* 2010 Mar;67(3):365–75.
- [89] Zheng X, Sun L, Yin D, Jia J, Zhao Z, Jiang Y, et al. The plasticity of intrinsic functional connectivity patterns associated with rehabilitation intervention in chronic stroke patients. *Neuroradiology* 2016 Apr;58(4):417–27.
- [90] Favre I, Zeffiro TA, Detante O, Krainik A, Hommel M, Jaillard A. Upper limb recovery after stroke is associated with ipsilesional primary motor cortical activation: a meta-analysis. *Stroke* 2014 Apr;45(4):1077–83. Lippincott Williams & Wilkins.
- [91] Cramer SC, Nelles G, Benson RR, Kaplan JD, Parker RA, Kwong KK, et al. A functional MRI study of subjects recovered from hemiparetic stroke. *Stroke* 1997 Dec;28(12):2518–27.
- [92] Frost SB, Barbay S, Friel KM, Plautz EJ, Nudo RJ. Reorganization of remote cortical regions after ischemic brain injury: a potential substrate for stroke recovery. *J Neurophysiol* 2003 Jun;89(6):3205–14.
- [93] Slavin MD, Laurence S, Stein DG. Another look at vicariation. In: Finger S, Levere TE, Almli CR, Stein DG, editors. *Brain injury and recovery*. Boston, MA: Springer US; 1988. p. 165–79.
- [94] Di Pino G, Pellegrino G, Assenza G, Capone F, Ferreri F, Formica D, et al. Modulation of brain plasticity in stroke: a novel model for neuro-rehabilitation. *Nat Rev Neurol* 2014 Oct;10(10):597–608.
- [95] Plow EB, Sankarasubramanian V, Cunningham DA, Potter-Baker K, Varnerin N, Cohen LG, et al. Models to tailor brain stimulation therapies in stroke. *Neural Plast* 2016;2016(4):1–17.
- [96] Bradnam LV, Stinear CM, Barber PA, Byblow WD. Contralesional hemisphere control of the proximal paretic upper limb following stroke. *Cereb Cortex* 2012 Oct 4;22(11):2662–71.
- [97] Bestmann S, Swayne O, Blankenburg F, Ruff CC, Teo J, Weiskopf N, et al. The role of contralesional dorsal premotor cortex after stroke as studied with concurrent TMS-fMRI. *J Neurosci* 2010 Sep 8;30(36):11926–37. Society of Neuroscience.
- [98] Schlaug G, Renga V, Nair D. Transcranial direct current stimulation in stroke recovery 2008 Dec;vol. 65(12):1571–6.
- [99] Stinear CM, Barber PA, Petoe M, Anwar S, Byblow WD. The PREP algorithm predicts potential for upper limb recovery after stroke. *Brain* 2012 Aug;135(Pt 8):2527–35.
- [100] Lefaucheur J-P, André-Obadia N, Antal A, Ayache SS, Baeken C, Benninger DH, et al. Evidence-based guidelines on the therapeutic use of repetitive transcranial magnetic stimulation (rTMS). *Clin Neurophysiol* 2014 Nov;125(11):2150–206.
- [101] Ackerley SJ, Byblow WD, Barber PA, MacDonald H, McIntyre-Robinson A, Stinear CM. Primed physical therapy enhances recovery of upper limb function in chronic stroke patients. *Neurorehabil Neural Repair* 2016 May;30(4):339–48.
- [102] Ameli M, Grefkes C, Kemper F, Riegg FP, Rehme AK, Karbe H, et al. Differential effects of high-frequency repetitive transcranial magnetic stimulation over ipsilesional primary motor cortex in cortical and subcortical middle cerebral artery stroke. *Ann Neurol* 2009 Sep;66(3):298–309.
- [103] Hsu WY, Cheng CH, Liao KK, Lee IH, Lin YY. Effects of repetitive transcranial magnetic stimulation on motor functions in patients with stroke: a meta-analysis. *Stroke* 2012;43:1849–57.
- [104] Tang I-N. The effect of repetitive transcranial magnetic stimulation on upper extremity motor function in stroke patients: a meta-analytical review. *J Food Drug Anal* 2012;20:1–5.
- [105] Mansur CG, Fregni F, Boggio PS, Riberto M, Gallucci-Neto J, Santos CM, et al. A sham stimulation-controlled trial of rTMS of the unaffected hemisphere in stroke patients. *Neurology* 2005 May 24;64(10):1802–4.
- [106] Takeuchi N, Chuma T, Matsuo Y, Watanabe I, Ikoma K. Repetitive transcranial magnetic stimulation of contralesional primary motor cortex improves hand function after stroke. *Stroke* 2005 Dec;36(12):2681–6.
- [107] Liepert J, Zittel S, Weiller C. Improvement of dexterity by single session low-frequency repetitive transcranial magnetic stimulation over the contralesional motor cortex in acute stroke: a double-blind placebo-controlled crossover trial. *Restor Neurol Neurosci* 2007;25(5–6):461–5.
- [108] Nowak DA, Grefkes C, Dafotakis M, Eickhoff S, Küst J, Karbe H, et al. Effects of low-frequency repetitive transcranial magnetic stimulation of the contralesional primary motor cortex on movement kinematics and neural activity in subcortical stroke. *Arch Neurol* 2008 Jun;65(6):741–7. American Medical Association.
- [109] Dafotakis M, Grefkes C, Eickhoff SB, Karbe H, Fink GR, Nowak DA. Effects of rTMS on grip force control following subcortical stroke. *Exp Neurol* 2008 Jun;211(2):407–12.
- [110] Takeuchi N, Tada T, Toshima M, Chuma T, Matsuo Y, Ikoma K. Inhibition of the unaffected motor cortex by 1 Hz repetitive transcranial magnetic stimulation enhances motor performance and training effect of the paretic hand in patients with chronic stroke. *J Rehabil Med* 2008 Apr;40(4):298–303.
- [111] Demirtas-Tatlidede A, Alonso-Alonso M, Shetty RP, Ronen I, Pascual-Leone A, Fregni F. Long-term effects of contralesional rTMS in severe stroke: safety, cortical excitability, and relationship with transcallosal motor fibers. *NeuroRehabilitation* 2015;36(1):51–9.
- [112] Takeuchi N, Tada T, Matsuo Y, Ikoma K. Low-frequency repetitive TMS plus anodal transcranial DCS prevents transient decline in bimanual movement induced by contralesional inhibitory rTMS after stroke. *Neurorehabil Neural Repair* 2012 Oct;26(8):988–98. SAGE Publications.
- [113] Avenanti A, Coccia M, Ladavas E, Provinciali L, Ceravolo MG. Low-frequency rTMS promotes use-dependent motor plasticity in chronic stroke: a randomized trial. *Neurology* 2012 Jan 24;78(4):256–64. Lippincott Williams & Wilkins.
- [114] Huang Y-Z, Edwards MJ, Rounis E, Bhatia KP, Rothwell JC. Theta burst stimulation of the human motor cortex. *Neuron* 2005 Jan 20;45(2):201–6.
- [115] Talelli P, Greenwood RJ, Rothwell JC. Exploring Theta Burst Stimulation as an intervention to improve motor recovery in chronic stroke. *Clin Neurophysiol* 2007 Feb;118(2):333–42.
- [116] Theilig S, Podubecka J, Bösl K, Wiederer R, Nowak DA. Functional neuromuscular stimulation to improve severe hand dysfunction after stroke: does inhibitory rTMS enhance therapeutic efficiency? *Exp Neurol* 2011 Jul;230(1):149–55.
- [117] Seniów J, Bilik M, Leśniak M, Waldowski K, Iwański S, Członkowska A. Transcranial magnetic stimulation combined with physiotherapy in rehabilitation of poststroke hemiparesis: a randomized, double-blind, placebo-controlled study. *Neurorehabil Neural Repair* 2012 Nov;26(9):1072–9. SAGE Publications.
- [118] Di Lazzaro V, Pilato F, Dileone M, Profice P, Oliviero A, Mazzone P, et al. The physiological basis of the effects of intermittent theta burst stimulation of the human motor cortex. *J Physiol (Lond)* 2008 Aug 15;586(16):3871–9.
- [119] Lüdemann-Podubecká J, Bösl K, Nowak DA. Inhibition of the contralesional dorsal premotor cortex improves motor function of the affected hand following stroke. *Eur J Neurol* 2016 Jan 23;23(4):823–30.
- [120] Brownjohn PW, Reynolds JNJ, Matheson N, Fox J, Shemmell JBH. The effects of individualized theta burst stimulation on the excitability of the human motor system. *Brain Stimul* 2014 Mar;7(2):260–8.
- [121] Maeda F, Keenan JP, Tormos JM, Topka H, Pascual-Leone A. Interindividual variability of the modulatory effects of repetitive transcranial magnetic stimulation on cortical excitability. *Exp Brain Res* 2000 Aug;133(4):425–30.
- [122] Hamada M, Murase N, Hasan A, Balaratnam M, Rothwell JC. The role of interneuron networks in driving human motor cortical plasticity. *Cereb Cortex* 2013 Jul;23(7):1593–605.
- [123] Pasley BN, Allen EA, Freeman RD. State-dependent variability of neuronal responses to transcranial magnetic stimulation of the visual cortex. *Neuron* 2009 Apr 30;62(2):291–303.
- [124] Moliadze V, Zhao Y, Eysel U, Funke K. Effect of transcranial magnetic stimulation on single-unit activity in the cat primary visual cortex. *J Physiol (Lond)* 2003 Dec 1;553(Pt 2):665–79.
- [125] Fricke K, Seeber AA, Thiruganasambandam N, Paulus W, Nitsche MA, Rothwell JC. Time course of the induction of homeostatic plasticity generated



- by repeated transcranial direct current stimulation of the human motor cortex. *J Neurophysiol* 2011 Mar;105(3):1141–9.
- [126] Matheson NA, Shemmell JBH, De Ridder D, Reynolds JNJ. Understanding the effects of repetitive transcranial magnetic stimulation on neuronal circuits. *Front Neural Circuits* 2016;10:67.
- [127] Stinear JW, Byblow WD. Rhythmic bilateral movement training modulates corticomotor excitability and enhances upper limb motricity poststroke: a pilot study. *J Clin Neurophysiol* 2004 Mar;21(2):124–31.
- [128] Stinear CM, Petoe MA, Anwar S, Barber PA, Byblow WD. Bilateral priming accelerates recovery of upper limb function after stroke: a randomized controlled trial. *Stroke* 2014 Jan;45(1):205–10.
- [129] Stinear JW, Byblow WD. Disinhibition in the human motor cortex is enhanced by synchronous upper limb movements. *J Physiol (Lond)* [Internet] 2004 Jul 22;543(1):307–16.
- [130] Cheng MY, Aswendt M, Steinberg GK. Optogenetic approaches to target specific neural circuits in post-stroke recovery. 2015 Dec 23.
- [131] Seeger-Armbruster S, Bosch-Bouju C, Little STC, Smither RA, Hughes SM, Hyland BI, et al. Patterned, but not tonic, optogenetic stimulation in motor thalamus improves reaching in acute drug-induced Parkinsonian rats. *J Neurosci* 2015 Jan 21;35(3):1211–6. Society for Neuroscience.
- [132] Chow BY, Boyden ES. Optogenetics and translational medicine. *Sci Transl Med Am Assoc Adv Sci* 2013 Mar 20;5(177):177ps5.
- [133] Harvey RL, Winstein CJ. Everest Trial Group. Design for the everest randomized trial of cortical stimulation and rehabilitation for arm function following stroke. *Neurorehabil Neural Repair* 2009 Jan;23(1):32–44.
- [134] Plow EB, Carey JR, Nudo RJ, Pascual-Leone A. Invasive cortical stimulation to promote recovery of function after stroke: a critical appraisal. *Stroke* 2009 May;40(5):1926–31.
- [135] Brown JA, Lutsep HL, Weinand M, Cramer SC. Motor cortex stimulation for the enhancement of recovery from stroke: a prospective, multicenter safety study. *Neurosurgery* 2006 Mar;58(3):464–73.
- [136] Huang M, Harvey RL, Stoykov ME, Ruland S, Weinand M, Lowry D, et al. Cortical stimulation for upper limb recovery following ischemic stroke: a small phase II pilot study of a fully implanted stimulator. *Top Stroke Rehabil* 2008 Mar;15(2):160–72.
- [137] Levy R, Ruland S, Weinand M, Lowry D, Dafer R, Bakay R. Cortical stimulation for the rehabilitation of patients with hemiparetic stroke: a multicenter feasibility study of safety and efficacy. 2008 Apr;vol. 108(4):707–714.
- [138] Levy RM, Harvey RL, Kissela BM, Winstein CJ, Lutsep HL, Parrish TB, et al. Epidural electrical stimulation for stroke rehabilitation: results of the prospective, multicenter, randomized, single-blinded everest trial. *Neurorehabil Neural Repair* 2016 Feb;30(2):107–19.