Oblig 1b: Terningdropp Analysis

Your Name

Date of Submission

1 Introduction

Briefly describe the objective of the assignment and the methods used.

2 Assignment Questions

2.1 Question 2a: First 5 Measurements Regression

```
\# Linear Regression on the first 5 measurements in R \operatorname{lm\_first5} \leftarrow \operatorname{lm}(\operatorname{Lengde} \ \tilde{}\ \operatorname{Dropp},\ \operatorname{data=df}[1:5,\ ])
```

Calculated Regression Line: Include your manual calculation or calculator output here.

2.2 Question 2b: Scatter Plot of Data Points

Scatter plot of the Dropp vs Lengde data (placeholder image).

2.3 Question 2c: Regression Line Plot

Scatter plot with regression line (placeholder image).

2.4 Question 2d: Regression Analysis Entire Dataset

```
\# Linear Regression on the entire dataset in R lm_-full \leftarrow lm(Lengde \ ^\circ Dropp, \ data=df)
```

2.5 Question 2e: Sum of Squared Residuals (SSe)

2.5.1 First 5 Measurements

```
ssr_first5 <- sum(residuals(lm_first5)^2)
```

SSR for First 5 Measurements:

 $SSR = < calculated_value >$

Scatterplot av Dropp vs Lengde




Figure 1: Scatter plot of Dropp vs Lengde

2.5.2 Entire Dataset

 $ssr_full \leftarrow sum(residuals(lm_full)^2)$

SSR for Entire Dataset:

 $SSR = < calculated_value >$

2.6 Question 2f: Standard Error (se)

2.6.1 First 5 Measurements

 $se_first5 \leftarrow sqrt(ssr_first5 / lm_first5 df.residual)$

Standard Error for First 5 Measurements:

 $SE = < calculated_value >$

2.6.2 Entire Dataset

 $se_full \leftarrow sqrt(ssr_full / lm_full df.residual)$

Standard Error for Entire Dataset:

 $SE = < calculated_value >$

3 Conclusion

Summarize your findings and observations from the assignment.




Figure 2: Scatter plot with regression line