

Notes on a design of a simple spatial sampling
method (S3M) for assessing coverage of health
and nutrition programmes in Liberia

Valid International

2018-06-19

Contents

Simple Spatial Sampling Method (S3M)	5
1 Introduction	7
2 The survey sample	9
3 The first stage sample	11
3.1 Step 1: Find a map	12
4 The second stage sample	15
5 Analysis	17
6 liberiaS3M	19
6.1 Example	19

Simple Spatial Sampling Method (S3M)



Chapter 1

Introduction

The Simple Spatial Survey Method (S3M) was developed from the CSAS coverage survey method as a response to the widespread adoption of community management of acute malnutrition (CMAM) by ministries of health. Large-scale programs need a large-scale survey method and S3M was developed to meet that need.

S3M was designed to :

- Be simple enough for MoH, NGO, and UNO personnel without specialist statistical training to perform.
- Provide a general survey method. S3M can be used to survey and map :
- Need for and coverage of selective-entry programs such as CMAM and TSFP as well as universal programs such as EPI, GMP, GFD (general ration), and “blanket” SFP over wide areas.
- Levels of indicators such as those for IYCF, WASH, and period prevalence / cumulative prevalence of ARI, fever, and diarrhoea over wide areas.

This document concentrates on using S3M to assess the need for and coverage of a variety of selective-entry feeding programs. The indicators discussed in this manual are:

- Therapeutic feeding (OTP and TSFP) programs :
- Prevalence of SAM and coverage of treatment of SAM in children aged between 6 and 59 months.
- Prevalence of MAM and coverage of treatment of MAM in children aged between 6 and 59 months.
- Prevalence of MAM and treatment of MAM and in pregnant and lactating women (PLWs).
- Food-based prevention of malnutrition (FBPM) programs :

- Prevalence of need for and coverage of food-based prevention of malnutrition in younger children at risk of developing MAM and SAM.
- Prevalence of need for and coverage of food-based prevention of malnutrition in pregnant and lactating women (PLWs) at risk of developing MAM and SAM.
- Coverage of screening for all of the above programs.
- Coverage of Behaviour Change Communication (BCC) programs focussing on maternal and child health and nutrition to all principal carers of children (usually their mothers) and all PLWs.

Chapter 2

The survey sample

The survey method described here uses a two-stage sample:

- **First-stage:** We take an even (or near-even) spatial sample of communities from all of the communities in the survey area.
- **Second-stage:** We take a sample of eligible individuals from each of the communities identified in the first stage of sampling.

Two-stage sampling is used in many survey methods. A typical example of a survey method that uses a two- stage sample is the SMART method that is commonly used for nutritional anthropometry surveys.

The main difference between the sample taken in S3M based surveys and in SMART type surveys is that S3M based samples used a spatial sample in the first stage whereas SMART type surveys use a proportional to population size (PPS) sample.

The advantages of using a spatial first stage sample is that such a sample allows us to identify where (and why) coverage is good, and where (and why) coverage is poor. This information is essential to improving program coverage and ensuring equitable access to services.

A spatial sample can be used to produce equivalent results to a traditional proportional to population size (PPS) sample as is used in (e.g.) SMART type surveys using a weighted analysis. This means that a spatial sample can be made to act as a PPS sample. A PPS type sample cannot, however, be made to act as a spatial sample.

Chapter 3

The first stage sample

3.1 Step 1: Find a map




Figure 3.1: Small scale map of Liberia showing counties and settlements

The first step in an S3M survey is to find a map of the survey area.

Try to find a map showing the locations of all towns and villages in the survey area. You may need to update the map to take into account migration and displacement.

If you are surveying a very large area then you will find it useful to have:

- A small scale map of the entire survey area. This map does not need to show the location of all towns and villages in survey area.
- A collection of larger scale maps showing the locations of all towns and villages. This collection of maps should cover the entire survey area.

The small scale map will be useful for identifying initial sampling locations.

The large scale maps will be useful for identifying the precise locations of sampling points and for selecting the communities to be sampled.

Chapter 4

The second stage sample

Chapter 5

Analysis

Chapter 6

liberiaS3M

The goal of liberiaS3M is to ...

6.1 Example

This is a basic example which shows you how to solve a common problem:

```
## basic example code
```