Your routine

Helps you schedule and keep track of the exercises your physio prescribed. Set by your physio at your visit.

Date: _____ Physio: _____ Stage: _____

Days: ☐ Sun ☐ Mon ☐ Tue ☐ Wed ☐ Thu ☐ Fri ☐ Sat

	Exercise	Sets	Reps / sec
Mobility			
☐	Ankle alphabet (per side)		
☐	Ankle pumps		
☐	Knee to wall (per side)		
☐	Calf stretch (per side)		
Strength			
☐	Band: foot out (per side)		
☐	Band: foot in (per side)		
☐	Band: toes up (per side)		
☐	Band: toes down (per side)		
☐	Heel raises, both legs		
☐	Heel raises, one leg (per side)		
Balance			
☐	Single-leg balance (per side)		
☐	Single-leg balance, eyes closed (per side)		
☐	Single-leg reach (per side)		
Hops			
☐	Small hops, both legs		
☐	Side-to-side hops		
☐	Hop and hold (per side)		

Notes from the visit:

Next visit: _____

Stop and get checked if any of these apply

- You can't put weight on your foot or take 4 steps
- Your ankle or foot looks deformed or out of place
- Your foot is numb, tingling, cold or pale
- You felt or heard a "pop" at the back of your heel, or can't push off your toes
- Your calf is painful, swollen, warm or red
- Pain or swelling is getting worse instead of better

Don't do the exercises. Contact your physio or go to urgent care.

Emergency **911**

Video link pending

Exercise videos

Scan with your phone camera

Tip: set a phone alarm for your exercise days.

Exercises

Ankle alphabet	Sit with your leg out and your foot off the floor. Trace the letters A to Z in the air with your big toe, moving only at the ankle.
Ankle pumps	Lying down or sitting with your leg up, point your toes away, then pull them toward you. Move slowly and smoothly.
Knee to wall	Stand facing a wall with your foot a few centimeters away. Keep your heel down and bend your knee toward the wall. Move your foot back a little as it gets easier.
Calf stretch	Hands on a wall, one foot back with the knee straight and heel down. Lean forward until you feel a gentle stretch in the calf. Hold.
Band: foot out	Sit with a resistance band around your forefoot, anchored to the inside. Turn the sole of your foot outward against the band, then return slowly.
Band: foot in	Sit with a resistance band around your forefoot, anchored to the outside. Turn the sole of your foot inward against the band, then return slowly.
Band: toes up	Sit with a band around the top of your foot, anchored in front of you. Pull your toes toward you against the band, then return slowly.
Band: toes down	Sit with your leg out and a band around the ball of your foot, holding the ends. Push your toes away against the band, then return slowly.
Heel raises, both legs	Stand holding a wall or chair. Rise onto the balls of both feet, pause, and lower slowly.
Heel raises, one leg	Stand on one leg, lightly holding a wall. Rise onto the ball of your foot, pause, and lower slowly over 3 seconds.
Single-leg balance	Stand on one leg near a wall or counter, knee slightly bent. Hold your balance. Touch the wall only if you need to.
Single-leg balance, eyes closed	Same as single-leg balance, next to a wall, with your eyes closed. Open your eyes or touch the wall whenever you need to.
Single-leg reach	Stand on one leg. With the other foot, reach forward, to the side and behind, lightly tapping the floor each time, then return to the center.

Small hops, both legs	Hop gently in place on both feet, landing softly on the balls of your feet with knees slightly bent.
Side-to-side hops	Hop side to side over a line on both feet, landing softly. Progress to one leg when your physio or coach says it's fine.
Hop and hold	Hop forward from one leg and land on the same leg. Hold the landing still for 2 seconds, knee slightly bent over your toes.

This app is not a diagnosis or treatment, and no one monitors what you log. If your ankle gets worse, contact your physio or go to urgent care.