

Homework

1 Basic operation and their notation

Exercise 1.1: Inner/outer products in Dirac notation

$$\begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 \end{pmatrix} \begin{pmatrix} 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 11 \end{pmatrix}$$

The last one in Dirac notation :

$$\begin{aligned} & (\langle 0| + 2\langle 1|) \times (3|0\rangle + 4|1\rangle) \\ &= 3\langle 0|0\rangle + 4\langle 0|1\rangle + 6\langle 1|0\rangle + 8\langle 1|1\rangle \\ &= 3 + 8 \\ &= 11 \end{aligned}$$

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 \\ 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 3 \\ 4 \end{pmatrix} \begin{pmatrix} 1 & 2 \end{pmatrix} = \begin{pmatrix} 3 & 6 \\ 4 & 8 \end{pmatrix}$$

The last two in Dirac notation :

$$\begin{aligned} & (3|0\rangle + 4|1\rangle) \times (\langle 0| + 2\langle 1|) \\ &= 3|0\rangle\langle 0| + 6|0\rangle\langle 1| + 4|1\rangle\langle 0| + 8|1\rangle\langle 1| \end{aligned}$$

Exercise 1.2: Matrix products in Dirac notation

$$\begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 3 \\ 2 & 4 \end{pmatrix} \begin{pmatrix} 5 \\ 6 \end{pmatrix} = \begin{pmatrix} 23 \\ 34 \end{pmatrix}$$

The last one in Dirac notation :

$$\begin{aligned} & (|0\rangle\langle 0| + 3|0\rangle\langle 1| + 2|1\rangle\langle 0| + 4|1\rangle\langle 1|) \times (5|0\rangle + 6|1\rangle) \\ &= 5|0\rangle\langle 0|0\rangle + 6|0\rangle\langle 0|1\rangle + 15|0\rangle\langle 1|0\rangle + 18|0\rangle\langle 1|1\rangle + \\ & \quad 10|1\rangle\langle 0|0\rangle + 12|1\rangle\langle 0|1\rangle + 20|1\rangle\langle 1|0\rangle + 24|1\rangle\langle 1|1\rangle \\ &= 5\langle 0|0\rangle|0\rangle + 18\langle 1|1\rangle|0\rangle + 10\langle 0|0\rangle|1\rangle + 24\langle 1|1\rangle|1\rangle \\ &= 5|0\rangle + 18|0\rangle + 10|1\rangle + 24|1\rangle \\ &= 23|0\rangle + 24|1\rangle \end{aligned}$$

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 3 \\ 2 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 3 \\ 2 & 4 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

The last one in Dirac notation :

$$\begin{aligned} & (1/\sqrt{2}|0\rangle\langle 0| + 1/\sqrt{2}|0\rangle\langle 1| + 1/\sqrt{2}|1\rangle\langle 0| - 1/\sqrt{2}|1\rangle\langle 1|)^2 \\ &= 1/2|0\rangle\langle 0|0\rangle\langle 0| + 1/2|0\rangle\langle 0|0\rangle\langle 1| + 1/2|0\rangle\langle 1|1\rangle\langle 0| - 1/2|0\rangle\langle 1|1\rangle\langle 1| \\ & \quad 1/2|1\rangle\langle 0|0\rangle\langle 1| + 1/2|1\rangle\langle 0|0\rangle\langle 0| - 1/2|1\rangle\langle 1|1\rangle\langle 0| + 1/2|1\rangle\langle 1|1\rangle\langle 1| \\ &= 1/2|0\rangle\langle 0| + 1/2|0\rangle\langle 0| + 1/2|1\rangle\langle 1| + 1/2|1\rangle\langle 1| \\ &= |0\rangle\langle 0| + |1\rangle\langle 1| \end{aligned}$$

Exercise 1.3: Tensor products in Dirac/Coecke notation

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \quad \begin{pmatrix} 0 \\ 1 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix} \quad \begin{pmatrix} 1 \\ 2 \end{pmatrix} \otimes \begin{pmatrix} 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 3 \\ 4 \\ 6 \\ 8 \end{pmatrix}$$

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 1 \end{pmatrix} \otimes \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix}$$

The last two in Dirac notation :

$$(|0\rangle + 2|1\rangle) \otimes (3|0\rangle + 4|1\rangle) = 3|00\rangle + 4|01\rangle + 6|10\rangle + 8|11\rangle$$

$$|0\rangle \otimes |0\rangle + |1\rangle \otimes |1\rangle = |00\rangle + |11\rangle$$

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} \otimes \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} = \begin{pmatrix} 1/2 & 1/2 & 1/2 & 1/2 \\ 1/2 & -1/2 & 1/2 & -1/2 \\ 1/2 & 1/2 & -1/2 & -1/2 \\ 1/2 & -1/2 & -1/2 & 1/2 \end{pmatrix}$$

In Dirac notation :

$$\begin{aligned} & (|0\rangle\langle 0| + |1\rangle\langle 1|) \otimes (|0\rangle\langle 0| + |1\rangle\langle 1|) \\ &= |0\rangle\langle 0| \otimes |0\rangle\langle 0| + |0\rangle\langle 0| \otimes |1\rangle\langle 1| + |1\rangle\langle 1| \otimes |0\rangle\langle 0| + |1\rangle\langle 1| \otimes |1\rangle\langle 1| \\ &= |0\rangle\langle 0| + |1\rangle\langle 1| + |2\rangle\langle 2| + |3\rangle\langle 3| \end{aligned}$$

$$\begin{aligned} & (|0\rangle\langle 0|) \otimes (|0\rangle\langle 0| + |1\rangle\langle 1|) \\ &= |0\rangle\langle 0| \otimes |0\rangle\langle 0| + |0\rangle\langle 0| \otimes |1\rangle\langle 1| \\ &= |0\rangle\langle 0| + |1\rangle\langle 1| \end{aligned}$$

$$\begin{aligned} & 1/\sqrt{2}(|0\rangle\langle 0| + |0\rangle\langle 1| + |1\rangle\langle 0| - |1\rangle\langle 1|) \otimes 1/\sqrt{2}(|0\rangle\langle 0| + |1\rangle\langle 0| + |1\rangle\langle 0| - |1\rangle\langle 1|) \\ &= 1/2(|0\rangle\langle 0| + |0\rangle\langle 1| + |0\rangle\langle 2| + |0\rangle\langle 3| + \\ & \quad |1\rangle\langle 0| - |1\rangle\langle 1| + |1\rangle\langle 2| - |1\rangle\langle 3| + \\ & \quad |2\rangle\langle 0| + |2\rangle\langle 1| - |2\rangle\langle 2| - |2\rangle\langle 3| + \\ & \quad |3\rangle\langle 0| - |3\rangle\langle 1| - |3\rangle\langle 2| + |3\rangle\langle 3|) \end{aligned}$$

We want to prove $(A \otimes B)(C \otimes D) = (AC) \otimes (BD)$

- Dirac's notation :
- Coecke's notation :

Exercice 1.4: Dagger in Dirac/Coecke notation

$$\begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix}^\dagger = \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} \quad \begin{pmatrix} 1 & 3i \\ 2 & 4i \end{pmatrix}^\dagger = \begin{pmatrix} 1 & 2 \\ -3i & -4i \end{pmatrix}$$

In Dirac notation:

$$\begin{aligned} & 1/\sqrt{2}(|0\rangle\langle 0| + |0\rangle\langle 1| + |1\rangle\langle 0| - |1\rangle\langle 1|)^\dagger \\ &= 1/\sqrt{2}(|0\rangle\langle 0| + |1\rangle\langle 0| + |0\rangle\langle 1| - |1\rangle\langle 1|) \end{aligned}$$

$$\begin{aligned} & (|0\rangle\langle 0| + 3i|0\rangle\langle 1| + 2|1\rangle\langle 0| + 4i|1\rangle\langle 1|)^\dagger \\ &= (|0\rangle\langle 0| - 3i|1\rangle\langle 0| + 2|0\rangle\langle 1| - 4i|1\rangle\langle 1|) \end{aligned}$$

Exercice 1.5: Gates in Dirac notations

$$H = 1/\sqrt{2}(|0\rangle\langle 0| + |0\rangle\langle 1| + |1\rangle\langle 0| - |1\rangle\langle 1|)$$

$$CNot = |0\rangle\langle 0| + |1\rangle\langle 1| + |3\rangle\langle 2| + |2\rangle\langle 3|$$

$$T = |0\rangle\langle 0| + e^{\frac{i\pi}{4}}|1\rangle\langle 1|$$

Proof that are unitary matrix :

- $H : H^\dagger H = Id_1$ already do in Exercice-1.2
- $CNot$:

$$\begin{aligned} CNot^\dagger CNot &= (|0\rangle\langle 0| + |1\rangle\langle 1| + |3\rangle\langle 2| + |2\rangle\langle 3|)(|0\rangle\langle 0| + |1\rangle\langle 1| + |2\rangle\langle 3| + |3\rangle\langle 2|) \\ &= (|0\rangle\langle 0| + |1\rangle\langle 1| + |2\rangle\langle 2| + |3\rangle\langle 3|) \\ &= Id_4 \end{aligned}$$

- T :

$$\begin{aligned} (|0\rangle\langle 0| + e^{\frac{i\pi}{4}}|1\rangle\langle 1|)^2 &= |0\rangle\langle 0|0\rangle\langle 0| + e^{\frac{i\pi}{4}}|0\rangle\langle 0|1\rangle\langle 1| + e^{\frac{i\pi}{4}}(|1\rangle\langle 1|0\rangle\langle 0|) + e^{\frac{i\pi}{2}}(|1\rangle\langle 1|1\rangle\langle 1|) \\ &= |0\rangle\langle 0| + |1\rangle\langle 1| \end{aligned}$$

Exercice 1.6: Pauli matrices in Dirac/Coecke notation

- For all $i, k \in [0, 3]$ we want to show $\sigma_i \sigma_j = \delta_{ij} I + i \sum_k \epsilon_{ijk} \sigma_k$
 - If $i = j$ then $\sigma_i \sigma_j = I$ and for all k we have $\epsilon_{ijk} = 0$.
So we have $\delta_{ij} I = I = \sigma_i \sigma_j$

- For all $i, k \in [0, 3]$ we want to show $[\sigma_i, \sigma_j] = 2i \sum_k \epsilon_{ijk}$

$$\begin{aligned}
[\sigma_i, \sigma_j] &= \sigma_i \sigma_j - \sigma_j \sigma_i \\
&= (\delta_{ij} I + i \sum_k \epsilon_{ijk} \sigma_k) - (\delta_{ji} I + i \sum_k \epsilon_{jik} \sigma_k) \\
&= i \sum_k \epsilon_{ijk} \sigma_k - i \sum_k \epsilon_{jik} \sigma_k \\
&= i \sum_k \epsilon_{ijk} \sigma_k + i \sum_k \epsilon_{ijk} \sigma_k \\
&= 2i \sum_k \epsilon_{ijk} \sigma_k
\end{aligned}$$

- For all $i, k \in [0, 3]$ with $i \neq j$ we want to show $\{\sigma_i, \sigma_j\} = 0$

$$\begin{aligned}
\{\sigma_i, \sigma_j\} &= \sigma_i \sigma_j + \sigma_j \sigma_i \\
&= \delta_{ij} I + i \sum_k \epsilon_{ijk} \sigma_k + \delta_{ji} I - i \sum_k \epsilon_{ijk} \sigma_k & \epsilon_{jik} = -\epsilon_{ijk} \\
&= 2\delta_{ij} I \\
&= 0 & i \neq j
\end{aligned}$$

2 Postulates on pure states

Exercise 2.1: Evolutions

Let $|\psi\rangle = |0\rangle \otimes |0\rangle$ the initial state of two qubits. We want to compute $CNot(H \otimes I)|\psi\rangle$.

$$\begin{aligned}
CNot(H \otimes I)(|0\rangle \otimes |0\rangle) &= CNot((|0\rangle + |1\rangle)/\sqrt{2} \otimes (|0\rangle + |1\rangle)/\sqrt{2}) \\
&= \frac{1}{2} CNot(|00\rangle + |01\rangle + |10\rangle + |11\rangle) \\
&= \frac{1}{2} (|00\rangle + |01\rangle + |11\rangle + |10\rangle) := |\psi'\rangle
\end{aligned}$$

$$\begin{aligned}
(H \otimes I)CNot|\psi'\rangle &= (H \otimes I) \frac{1}{2} (|00\rangle + |01\rangle + |10\rangle + |11\rangle) \\
&= |00\rangle & H^2|\psi\rangle = |\psi\rangle
\end{aligned}$$

1. $T^4 H |0\rangle$
2. $HT^4 H |0\rangle$
3. $CNot(H|0\rangle \otimes HT^4 H|0\rangle)$
- 4.
5. $(H \otimes (H^2))(|0\rangle \otimes |0\rangle)$

Exercise 2.2: Measuring in another basis

- orthogonal :

$$\begin{aligned}\langle +|- \rangle &= \left(\frac{1}{\sqrt{2}}\right)^2 + \frac{1}{\sqrt{2}} \times \frac{-1}{\sqrt{2}} \\ &= \left(\frac{1}{\sqrt{2}}\right)^2 - \left(\frac{1}{\sqrt{2}}\right)^2 \\ &= 0\end{aligned}$$

- norm one :

$$\begin{aligned}\langle ++ \rangle &= \left(\frac{1}{\sqrt{2}}\right)^2 + \left(\frac{1}{\sqrt{2}}\right)^2 & \langle -- \rangle &= \left(\frac{1}{\sqrt{2}}\right)^2 + \left(\frac{-1}{\sqrt{2}}\right)^2 \\ &= \frac{1}{2} + \frac{1}{2} & &= \frac{1}{2} + \frac{1}{2} \\ &= 1 & &= 1\end{aligned}$$

- generate \mathbb{C}^2

Let $c = \alpha|0\rangle + \beta|1\rangle \in \mathbb{C}^2$ we want to find x and y such that $x|+\rangle + y|-\rangle = c$.

$$\begin{aligned}x|+\rangle + y|-\rangle &= \frac{x}{\sqrt{2}}(|0\rangle + |1\rangle) + \frac{y}{\sqrt{2}}(|0\rangle - |1\rangle) \\ &= \left(\frac{x+y}{\sqrt{2}}\right)|0\rangle + \left(\frac{x-y}{\sqrt{2}}\right)|1\rangle\end{aligned}$$

So we have this system :

$$\begin{aligned}\begin{cases} (x+y)/\sqrt{2} &= \alpha \\ (x-y)/\sqrt{2} &= \beta \end{cases} &\Rightarrow \begin{cases} x+y &= \sqrt{2}\alpha \\ x-y &= \sqrt{2}\beta \end{cases} \\ &\Rightarrow \begin{cases} 2x &= \sqrt{2}(\alpha + \beta) \\ x-y &= \sqrt{2}\beta \end{cases} \\ &\Rightarrow \begin{cases} x &= \frac{\sqrt{2}}{2}(\alpha + \beta) \\ -x+y &= -\sqrt{2}\beta \end{cases} \\ &\Rightarrow \begin{cases} x &= \frac{\sqrt{2}}{2}(\alpha + \beta) \\ y &= \frac{\sqrt{2}}{2}(\alpha - \beta) \end{cases}\end{aligned}$$

$B = \{|0\rangle, |1\rangle\}$ is another o.n.b of \mathbb{C}^2

We need to show that $\sum_{M \in \mathcal{M}_{\pm}} M^{\dagger} M = 1$

$$\begin{aligned}\sum_{M \in \mathcal{M}_{\pm}} &= (|+\rangle\langle +|)^{\dagger}(|+\rangle\langle +|) + (|-\rangle\langle -|)^{\dagger}(|-\rangle\langle -|) \\ &= (|+\rangle\langle +| + |-\rangle\langle -|) \\ &= |+\rangle\langle +| + |-\rangle\langle -| \\ &= \frac{1}{2}(|0\rangle\langle 0| + |0\rangle\langle 1| + |1\rangle\langle 0| + |1\rangle\langle 1|) + \frac{1}{2}(|0\rangle\langle 0| - |0\rangle\langle 1| - |1\rangle\langle 0| + |1\rangle\langle 1|) \\ &= |0\rangle\langle 0| + |1\rangle\langle 1| \\ &= 1\end{aligned}$$

So \mathcal{M}_{\pm} is a valid measurement.

We have $|\psi\rangle = \frac{1}{3}|0\rangle + \frac{\sqrt{8}}{3}|1\rangle$

- For $|+\rangle\langle+|$

$$\begin{aligned} p(|+\rangle) &= \langle\psi|(|+\rangle\langle+|)^{\dagger}|+\rangle\langle+||\psi\rangle \\ &= \langle\psi||+\rangle\langle+||\psi\rangle \\ &= \langle\psi|+\rangle\langle+|\psi\rangle \\ &= \left(\frac{1}{3} \times \frac{1}{\sqrt{2}} + \frac{\sqrt{8}}{3} \times \frac{1}{\sqrt{2}}\right)^2 \\ &= \frac{1}{2} \times \left(\frac{1+\sqrt{8}}{3}\right)^2 = \frac{1}{2} \times \frac{(1+\sqrt{8})^2}{9} = \frac{(1+\sqrt{8})^2}{18} \end{aligned}$$

- For $|-\rangle\langle-|$

Exercice 2.3: Measuring to distinguish

We defined:

$$M_0 = \begin{pmatrix} 1 & 0 \\ -1 & 0 \end{pmatrix} \quad M_+ = \begin{pmatrix} 1 & -1 \\ 2 & 2 \end{pmatrix} \quad M_f = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$$

3 Some mathematics

Exercice 3.1: Spectral theorems complements

Let D and D' two diagonal matrices and U a unitary matrix.

- Let $A = UDU^{\dagger}$ and $B = UD'U^{\dagger}$

$$\begin{aligned} AB &= UDU^{\dagger}UD'U^{\dagger} \\ &= UDD'U^{\dagger} \\ &= UD'DU^{\dagger} \\ &= UD'U^{\dagger}UDU^{\dagger} \\ &= BA \end{aligned} \quad \begin{array}{l} U \text{ is a unitary matrix} \\ D \text{ and } D' \text{ are diagonal} \end{array}$$

- Let $M = UDU^{\dagger}$

Exercice 3.2: Isometry versus unitary versus involution

- Matrix 2×2 unitary that is not an involution :

$$\begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix}$$

- Matrix $m \times n$ unitary that is not an involution :

4 On the nature of quantum information

Exercice 4.1: Hadamard

- $a = 0$

$$\begin{aligned} H|0\rangle &= \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) \\ &= \frac{1}{\sqrt{2}}(|0\rangle + (-1)^0|1\rangle) \end{aligned}$$

- $a = 1$

$$\begin{aligned} H|1\rangle &= \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle) \\ &= \frac{1}{\sqrt{2}}(|0\rangle + (-1)^1|1\rangle) \end{aligned}$$

Exercice 4.2: Who controls whom?

We define :

$$NotC = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

$$NotC|00\rangle = |00\rangle$$

$$NotC|01\rangle = |11\rangle$$

$$NotC|10\rangle = |10\rangle$$

$$NotC|11\rangle = |01\rangle$$

We want to proof $(H \otimes H)CNot(H \otimes H) = NotC$:

$$\begin{aligned} (H \otimes H)(|x\rangle \otimes |y\rangle) &= (H|x\rangle \otimes H|y\rangle) \\ &= \frac{1}{\sqrt{2}}(|0\rangle + (-1)^x|1\rangle) \otimes \frac{1}{\sqrt{2}}(|0\rangle + (-1)^y|1\rangle) \\ &= \frac{1}{2}(|00\rangle + (-1)^x|10\rangle + (-1)^y|01\rangle + (-1)^{x+y}|11\rangle) \end{aligned}$$

We apply the operator $CNot$:

$$\begin{aligned} &CNot\left(\frac{1}{2}(|00\rangle + (-1)^x|10\rangle + (-1)^y|01\rangle + (-1)^{x+y}|11\rangle)\right) \\ &= \frac{1}{2}(|00\rangle + (-1)^x|11\rangle + (-1)^y|01\rangle + (-1)^{x+y}|10\rangle) \\ &= \frac{1}{\sqrt{2}}(|0\rangle + (-1)^{x+y}|1\rangle) \otimes \left(\frac{1}{\sqrt{2}}(|0\rangle + (-1)^y|1\rangle)\right) \\ &= H|x \oplus y\rangle \otimes H|y\rangle \end{aligned} \quad (-1)^{x+y} = (-1)^{x \oplus y} \quad (x \oplus y \in \{0, 1\})$$

Finlay we apply $(H \otimes H)$:

$$\begin{aligned} HH|x \oplus y\rangle \otimes HH|y\rangle &= |x \oplus y\rangle \otimes |y\rangle \\ &= NotC(|x\rangle \otimes |y\rangle) \end{aligned}$$

5 Protocols

6 Quantum error correction

7 Bell