

The value of observational causal inference for medical decision making

Wouter van Amsterdam


Department of Data Science and Biostatistics
Julius Center for Health Sciences and Primary Care
University Medical Center Utrecht, the Netherlands

Medical decisions optimize for patient outcomes under interventions

Medical decisions are interventions (treatments) that change the outcome of a patient


Medical decisions optimize for patient outcomes under interventions

Need expected outcome under intervention (prediction under intervention)




Medical decisions optimize for patient outcomes under interventions

Need expected outcome under intervention (prediction under intervention) *conditional on characteristics*



Medical decisions optimize for patient outcomes under interventions

Need expected outcome under intervention (prediction under intervention)



How to estimate treatment effects?

Treatment effect estimation requires unconfoundedness

Which generally does not hold in observational data

concurrent vs sequential

Treatment effect estimation requires unconfoundedness

Which generally does not hold in observational data


concurrent vs sequential



overall survival


Treatment effect estimation requires unconfoundedness

Which generally does not hold in observational data



Treatment effect estimation requires unconfoundedness


Which generally does not hold in observational data



$$Y|do(T) \neq Y|T$$

Treatment effect estimation requires unconfoundedness


Which generally does not hold in observational data



$$Y|do(T) \neq Y|T$$
$$Y|do(T), Z = Y|T, Z$$

Treatment effect estimation requires unconfoundedness

Which generally does not hold in observational data




$$Y|do(T) \neq Y|T$$

$$Y|do(T), Z = Y|T, Z$$

$$Y|do(T) = \sum_Z Y|T, Z$$

Treatment effect estimation requires unconfoundedness

Which generally does not hold in observational data




$$Y|do(T) \neq Y|T$$

$$Y|do(T), Z = Y|T, Z$$

$$Y|do(T) \neq \sum_Z Y|T, Z$$


Randomized controlled trials guarantee unconfoundedness

Thus allow for causal inference



Randomized controlled trials guarantee unconfoundedness

Thus allow for causal inference



$$Y|do(T) = Y|T$$

Randomized controlled trials can estimate any interventional parameter

But generally only measure average treatment effect

Randomized controlled trials can estimate any interventional parameter

But generally only measure average treatment effect

Average Treatment Effect (ATE) $\mathbb{E} Y|do(T = A) - Y|do(T = B)$

Randomized controlled trials can estimate any interventional parameter

But generally only measure average treatment effect

Average Treatment Effect (ATE) $\mathbb{E} Y|do(T = A) - Y|do(T = B)$

Average Treatment Effect (risk ratio) $\frac{\mathbb{E} Y|do(T = A)}{\mathbb{E} Y|do(T = B)}$

Randomized controlled trials can estimate any interventional parameter

But generally only measure average treatment effect

Average Treatment Effect (ATE) $\mathbb{E} Y|do(T = A) - Y|do(T = B)$

Average Treatment Effect (risk ratio) $\frac{\mathbb{E} Y|do(T = A)}{\mathbb{E} Y|do(T = B)}$

Conditional Average Treatment Effect $\mathbb{E} Y|do(T = A), X = x - Y|do(T = B), X = x$

Randomized controlled trials can estimate any interventional parameter

But generally only measure average treatment effect *often on a relative scale*


Average Treatment Effect (ATE) $\mathbb{E} Y|do(T = A) - Y|do(T = B)$

Average Treatment Effect (risk ratio) $\frac{\mathbb{E} Y|do(T = A)}{\mathbb{E} Y|do(T = B)}$

Problems with RCTs for medical decision making


Problems with RCTs for medical decision making

RCTs are conducted in non-random samples of the population so transportability assumptions are needed



Problems with RCTs for medical decision making

RCTs are conducted in non-random samples of the population so transportability assumptions are needed



e.g. transportability of risk ratio (RR):


$$RR(\text{pop.} = \text{RCT}) = RR(\text{pop.} = \text{clinical})$$

Trading one unverifiable assumption (unconfoundedness) for another (transportability)

Problems with RCTs for medical decision making

Suspected treatment effect heterogeneity, but RCT too small to measure and lack of equipoise


- ▶ RCT establishes average treatment effect



Problems with RCTs for medical decision making

Suspected treatment effect heterogeneity, but RCT too small to measure and lack of equipoise


- ▶ RCT establishes average treatment effect
- ▶ Treatment effect heterogeneity assumed



Problems with RCTs for medical decision making

Suspected treatment effect heterogeneity, but RCT too small to measure and lack of equipoise


- ▶ RCT establishes average treatment effect
- ▶ Treatment effect heterogeneity assumed
- ▶ RCT too small to test this heterogeneity (e.g. subgroup tests)



Problems with RCTs for medical decision making

Suspected treatment effect heterogeneity, but RCT too small to measure and lack of equipoise


- ▶ RCT establishes average treatment effect
- ▶ Treatment effect heterogeneity assumed
- ▶ RCT too small to test this heterogeneity (e.g. subgroup tests)
- ▶ Cannot do a new trial because of lack of equipoise



Problems with RCTs for medical decision making

Suspected treatment effect heterogeneity, but RCT too small to measure and lack of equipoise


- ▶ RCT establishes average treatment effect
- ▶ Treatment effect heterogeneity assumed
- ▶ RCT too small to test this heterogeneity (e.g. subgroup tests)
- ▶ Cannot do a new trial because of lack of equipoise
- ▶ Need preliminary evidence



Problems with RCTs for medical decision making

Suspected treatment effect heterogeneity, but RCT too small to measure and lack of equipoise

- ▶ RCT establishes average treatment effect
- ▶ Treatment effect heterogeneity assumed
- ▶ RCT too small to test this heterogeneity (e.g. subgroup tests)
- ▶ Cannot do a new trial because of lack of equipoise
- ▶ Need preliminary evidence **from observational studies** (W. A. C. v. Amsterdam et al. 2022)



Problems with RCTs for medical decision making

Treatments in trials may differ from standard clinical care


Problems with RCTs for medical decision making

Treatments in trials may differ from standard clinical care



Problems with RCTs for medical decision making


Treatments in trials may differ from standard clinical care



- ▶ Patients don't take medications as prescribed **except when monitored in a trial**

Problems with RCTs for medical decision making


Treatments in trials may differ from standard clinical care



- ▶ Patients don't take medications as prescribed **except when monitored in a trial**
- ▶ Trading one unverifiable assumption (unconfoundedness) for another (consistency)

Problems with RCTs for medical decision making


RCTs solve confounding but have their own limitations




Transportability

Problems with RCTs for medical decision making

RCTs solve confounding but have their own limitations




Transportability




Heterogeneity

Problems with RCTs for medical decision making


RCTs solve confounding but have their own limitations



Transportability



Heterogeneity



Consistency

How can observational causal inference help medical decision making?

How observational causal inference can support medical decision making


Constant relative treatment effect + varying untreated risk = varying treatment effect

- ▶ Transportability assumption needed: most often on *relative scale*

How observational causal inference can support medical decision making

Constant relative treatment effect + varying untreated risk = varying treatment effect


- ▶ Transportability assumption needed: most often on *relative scale*
- ▶ Variance in expected outcome under *no treatment* ($\mathbb{E} Y|do(T = A), X$),



How observational causal inference can support medical decision making

Constant relative treatment effect + varying untreated risk = varying treatment effect


- ▶ Transportability assumption needed: most often on *relative scale*
- ▶ Variance in expected outcome under *no treatment* ($\mathbb{E} Y|do(T = A), X$),
 - ▶ implies differing effectiveness on absolute probability scale



How observational causal inference can support medical decision making

Constant relative treatment effect + varying untreated risk = varying treatment effect

- ▶ Transportability assumption needed: most often on *relative scale*
- ▶ Variance in expected outcome under *no treatment* ($\mathbb{E} Y|do(T = A), X$),
 - ▶ implies differing effectiveness on absolute probability scale
 - ▶ **thus varying treatment recommendations**



How observational causal inference can support medical decision making


Transportability: Need good models of untreated risk

- ▶ Widely used paradigm (cardiovascular risk, chemotherapy in breast cancer, ...)

How observational causal inference can support medical decision making

Transportability: Need good models of untreated risk


- ▶ Widely used paradigm (cardiovascular risk, chemotherapy in breast cancer, ...)
- ▶ Untreated risk is a quantity of the interventional distribution (i.e. *causal*)



How observational causal inference can support medical decision making

Transportability: Need good models of untreated risk


- ▶ Widely used paradigm (cardiovascular risk, chemotherapy in breast cancer, ...)
- ▶ Untreated risk is a quantity of the interventional distribution (i.e. *causal*)
- ▶ Current risk-models: mix of treated / untreated patients (W. v. Amsterdam et al. 2022),



How observational causal inference can support medical decision making

Transportability: Need good models of untreated risk


- ▶ Widely used paradigm (cardiovascular risk, chemotherapy in breast cancer, ...)
- ▶ Untreated risk is a quantity of the interventional distribution (i.e. *causal*)
- ▶ Current risk-models: mix of treated / untreated patients (W. v. Amsterdam et al. 2022),
- ▶ or ungrounded methods (Candido dos Reis et al. 2017; Xu et al. 2021).



How observational causal inference can support medical decision making

Transportability: Need good models of untreated risk

- ▶ Widely used paradigm (cardiovascular risk, chemotherapy in breast cancer, ...)
- ▶ Untreated risk is a quantity of the interventional distribution (i.e. *causal*)
- ▶ Current risk-models: mix of treated / untreated patients (W. v. Amsterdam et al. 2022),
- ▶ or ungrounded methods (Candido dos Reis et al. 2017; Xu et al. 2021).
- ▶ Need better 'causal' methods (Amsterdam et al 2023)



How observational causal inference can support medical decision making

Rare outcomes and statistical efficiency

- ▶ Need large studies for rare events (e.g. COVID-19 vaccinations)

How observational causal inference can support medical decision making

Rare outcomes and statistical efficiency

- ▶ Need large studies for rare events (e.g. COVID-19 vaccinations)
- ▶ Observational data may improve statistical efficiency of RCTs

How observational causal inference can support medical decision making

Rare outcomes and statistical efficiency

- ▶ Need large studies for rare events (e.g. COVID-19 vaccinations)
- ▶ Observational data may improve statistical efficiency of RCTs
 - ▶ RCT and observational data share* outcome mechanism

*under transportability assumption

How observational causal inference can support medical decision making

Rare outcomes and statistical efficiency

- ▶ Need large studies for rare events (e.g. COVID-19 vaccinations)
- ▶ Observational data may improve statistical efficiency of RCTs
 - ▶ RCT and observational data share* outcome mechanism
 - ▶ Need less randomized patients when using both data sources (Rosenman et al. 2020; Ilse et al. 2023)

*under transportability assumption

Observational Causal Inference Takeaway

- Challenges

Observational Causal Inference Takeaway

- ▶ Challenges
 - ▶ Unobserved confounding

Observational Causal Inference Takeaway

- ▶ Challenges
 - ▶ Unobserved confounding
- ▶ Opportunities

Observational Causal Inference Takeaway

- ▶ Challenges
 - ▶ Unobserved confounding
- ▶ Opportunities
 - ▶ Right population (to be treated)

Observational Causal Inference Takeaway

- ▶ Challenges
 - ▶ Unobserved confounding
- ▶ Opportunities
 - ▶ Right population (to be treated)
 - ▶ Right treatments (as given in standard clinical care)

Observational Causal Inference Takeaway

- ▶ Challenges
 - ▶ Unobserved confounding
- ▶ Opportunities
 - ▶ Right population (to be treated)
 - ▶ Right treatments (as given in standard clinical care)
 - ▶ Bigger data, allows estimating:

Observational Causal Inference Takeaway

- ▶ Challenges
 - ▶ Unobserved confounding
- ▶ Opportunities
 - ▶ Right population (to be treated)
 - ▶ Right treatments (as given in standard clinical care)
 - ▶ Bigger data, allows estimating:
 - ▶ Heterogeneity

Observational Causal Inference Takeaway


- ▶ Challenges
 - ▶ Unobserved confounding
- ▶ Opportunities
 - ▶ Right population (to be treated)
 - ▶ Right treatments (as given in standard clinical care)
 - ▶ Bigger data, allows estimating:
 - ▶ Heterogeneity
 - ▶ Rare events

Observational Causal Inference and RCTs interplay


Observational studies

RCT


Observational Causal Inference and RCTs interplay




Observational Causal Inference and RCTs interplay



Observational Causal Inference and RCTs interplay







Observational causal inference and RCTs for medical decision making



Must not abandon RCTs,
observational causal inference can be of great value for medical decision making

References

-  Amsterdam, W. van et al. (2022). "Decision making in cancer: causal questions require causal answers". In: *submitted*.
-  Amsterdam, Wouter A. C. van et al. (Apr. 2022). "Individual treatment effect estimation in the presence of unobserved confounding using proxies: a cohort study in stage III non-small cell lung cancer". en. In: *Scientific Reports* 12.1. Number: 1 Publisher: Nature Publishing Group, p. 5848. ISSN: 2045-2322. DOI: 10.1038/s41598-022-09775-9. URL: <https://www.nature.com/articles/s41598-022-09775-9> (visited on 04/12/2022).
-  Amsterdam et al, Wouter A. C. van (Apr. 2023). *Conditional average treatment effect estimation with treatment offset models*. arXiv:2204.13975 [stat]. URL: <http://arxiv.org/abs/2204.13975> (visited on 06/29/2022).
-  Candido dos Reis, Francisco J. et al. (Dec. 2017). "An updated PREDICT breast cancer prognostication and treatment benefit prediction model with independent validation". en. In: *Breast Cancer Research* 19.1. 80 citations (Crossref) [2021-08-06], p. 58. ISSN: 1465-542X. DOI: 10/gbhgpq. URL: <http://breast-cancer-research.com>