
Individual studies within monotonic−treatment meta−analyses

RR = p_t / p_c; SR = (1 − p_t) / (1 − p_c). Page 1/9 | MA rows 1−6 of 54

● Finite △ Infinite ratio ▽ Infinite x × Undefined ratio (0/0)


Screened treatment effect — Decreases the probability of Y



Likely/certain monotonicity; known screened direction; at least 5 study records per MA. No continuity correction.
Nonfinite ratios: top edge; infinite x: right edge; no connecting lines.
Black dashed: matched−scale mean (1 − m) and opposite−scale prediction; m estimated with equal weights from finite ratios.
Curves span observed nonsingular baselines. * Estimate outside [0, 1], not clipped. Source records retained.

Individual studies within monotonic−treatment meta−analyses

RR = p_t / p_c; SR = (1 − p_t) / (1 − p_c). Page 2/9 | MA rows 7−12 of 54




Likely/certain monotonicity; known screened direction; at least 5 study records per MA. No continuity correction.
Nonfinite ratios: top edge; infinite x: right edge; no connecting lines.
Black dashed: matched−scale mean (1 − m) and opposite−scale prediction; m estimated with equal weights from finite ratios.
Curves span observed nonsingular baselines. * Estimate outside [0, 1], not clipped. Source records retained.

Individual studies within monotonic−treatment meta−analyses

RR = p_t / p_c; SR = (1 − p_t) / (1 − p_c). Page 3/9 | MA rows 13−18 of 54

● Finite △ Infinite ratio ▽ Infinite x × Undefined ratio (0/0)

Screened treatment effect — Decreases the probability of Y




Likely/certain monotonicity; known screened direction; at least 5 study records per MA. No continuity correction.
Nonfinite ratios: top edge; infinite x: right edge; no connecting lines.
Black dashed: matched−scale mean (1 − m) and opposite−scale prediction; m estimated with equal weights from finite ratios.
Curves span observed nonsingular baselines. * Estimate outside [0, 1], not clipped. Source records retained.

Individual studies within monotonic−treatment meta−analyses

RR = p_t / p_c; SR = (1 − p_t) / (1 − p_c). Page 4/9 | MA rows 19−24 of 54

● Finite △ Infinite ratio ▽ Infinite x × Undefined ratio (0/0)

Screened treatment effect — Decreases the probability of Y




Likely/certain monotonicity; known screened direction; at least 5 study records per MA. No continuity correction.
Nonfinite ratios: top edge; infinite x: right edge; no connecting lines.
Black dashed: matched−scale mean (1 − m) and opposite−scale prediction; m estimated with equal weights from finite ratios.
Curves span observed nonsingular baselines. * Estimate outside [0, 1], not clipped. Source records retained.

Individual studies within monotonic−treatment meta−analyses

RR = p_t / p_c; SR = (1 − p_t) / (1 − p_c). Page 5/9 | MA rows 25−30 of 54

Screened treatment effect — Decreases the probability of Y

● Finite △ Infinite ratio ▽ Infinite x




Likely/certain monotonicity; known screened direction; at least 5 study records per MA. No continuity correction.
Nonfinite ratios: top edge; infinite x: right edge; no connecting lines.
Black dashed: matched−scale mean (1 − m) and opposite−scale prediction; m estimated with equal weights from finite ratios.
Curves span observed nonsingular baselines. * Estimate outside [0, 1], not clipped. Source records retained.

Individual studies within monotonic−treatment meta−analyses

RR = p_t / p_c; SR = (1 − p_t) / (1 − p_c). Page 6/9 | MA rows 31−36 of 54

● Finite △ Infinite ratio ▽ Infinite x × Undefined ratio (0/0)


Screened treatment effect — Decreases the probability of Y — Increases the probability of Y



Likely/certain monotonicity; known screened direction; at least 5 study records per MA. No continuity correction.
Nonfinite ratios: top edge; infinite x: right edge; no connecting lines.
Black dashed: matched−scale mean (1 − m) and opposite−scale prediction; m estimated with equal weights from finite ratios.
Curves span observed nonsingular baselines. * Estimate outside [0, 1], not clipped. Source records retained.

Individual studies within monotonic−treatment meta−analyses


RR = p_t / p_c; SR = (1 − p_t) / (1 − p_c). Page 7/9 | MA rows 37−42 of 54



Likely/certain monotonicity; known screened direction; at least 5 study records per MA. No continuity correction.
Nonfinite ratios: top edge; infinite x: right edge; no connecting lines.
Black dashed: matched−scale mean (1 − m) and opposite−scale prediction; m estimated with equal weights from finite ratios.
Curves span observed nonsingular baselines. * Estimate outside [0, 1], not clipped. Source records retained.

Individual studies within monotonic−treatment meta−analyses

RR = p_t / p_c; SR = (1 − p_t) / (1 − p_c). Page 8/9 | MA rows 43−48 of 54




Likely/certain monotonicity; known screened direction; at least 5 study records per MA. No continuity correction.
Nonfinite ratios: top edge; infinite x: right edge; no connecting lines.
Black dashed: matched−scale mean (1 − m) and opposite−scale prediction; m estimated with equal weights from finite ratios.
Curves span observed nonsingular baselines. * Estimate outside [0, 1], not clipped. Source records retained.

Individual studies within monotonic−treatment meta−analyses

RR = p_t / p_c; SR = (1 − p_t) / (1 − p_c). Page 9/9 | MA rows 49−54 of 54

Screened treatment effect — Increases the probability of Y

● Finite △ Infinite ratio ▽ Infinite x × Undefined ratio (0/0)



Likely/certain monotonicity; known screened direction; at least 5 study records per MA. No continuity correction.
Nonfinite ratios: top edge; infinite x: right edge; no connecting lines.
Black dashed: matched−scale mean (1 − m) and opposite−scale prediction; m estimated with equal weights from finite ratios.
Curves span observed nonsingular baselines. * Estimate outside [0, 1], not clipped. Source records retained.