

Homework 6, Due: Thursday, 12/13

This assignment is due on **Thursday, December 13**, by 11:59 PM. Your assignment should be well-organized, typed (or neatly written and scanned) and saved as a .pdf for submission on Canvas. You must show all of your work to receive full credit. For problems requiring the use of MATLAB code, remember to also submit your .m-files on Canvas as a part of your completed assignment. Your code should be appropriately commented to receive full credit.

Problems

- 1 (6 points) Consider the initial value problem

$$y' = \frac{2}{t}y + t^2 e^t, \quad 1 \leq t \leq 2, \quad y(1) = 0.$$

Use Theorem 5.4 to show that the initial value problem has a unique solution.

- 2 Consider Euler's method (Algorithm 5.1, pg. 267 of the text) for solving the initial value problem

$$y'(t) = f(t, y), \quad a \leq t \leq b, \quad y(a) = \alpha.$$

- (a) (6 points) Write a MATLAB **function** implementing Euler's method. Name your function **eulers.m**. List the input and output variables of your function, and insert comments to describe what each line of the code does.
- (b) (6 points) Use your function from part (a) to approximate the solution to

$$y' = \frac{2 - 2ty}{t^2 + 1}, \quad 0 \leq t \leq 1, \quad y(0) = 1$$

with $h = 0.5$, $h = 0.25$, and $h = 0.1$.

- (c) (6 points) Compare your approximations from part (b) to the true solution

$$y(t) = \frac{2t + 1}{t^2 + 1}$$

by computing the absolute error at each mesh point t_i for each h . Make a plot of the absolute errors in each case, and discuss your results.

Note: For any of the above problems for which you use MATLAB to help you solve, you must submit your code/.m-files as part of your work. Your code must run in order to receive full credit. If you include any plots, make sure that each has a title, axis labels, and readable font size, and include the final version of your plots as well as the code used to generate them.