

Genotyping information for *Tg(Cited-Cre)/-ERT2, -EGFP1Mdca*

Scott Boyle, Andrew Misfeldt, Kelly J. Chandler, Karen K. Deal, E. Michelle Southard-Smith, Douglas P. Mortlock, H. Scott Baldwin, Mark de Caestecker, Fate mapping using Cited1-CreERT2 mice demonstrates that the cap mesenchyme contains self-renewing progenitor cells and gives rise exclusively to nephronic epithelia, *Developmental Biology*, Volume 313, Issue 1, 2008, Pages 234-245, ISSN 0012-1606, <https://doi.org/10.1016/j.ydbio.2007.10.014>.
(<https://www.sciencedirect.com/science/article/pii/S0012160607014571>)

Strain description:

- This strain contains an unmapped transgene insertion.
- Zygosity may not be assessed using standard PCR-based genotyping approaches.
- Copy number analysis qPCR or ddPCR assays will be required for zygosity assessment.
Note that the original article suggests the transgene copy number = 4.

Automated genotyping:

A Transnetyx assay is available for this strain that targets the eGFP sequence.

Standard genotyping protocol from original publication:

Founders and subsequent offspring were genotyped using the following primer sets;

Cre: 5' GGCGCGGCAACACCATTTTT; 3' TCCGGGCTGCCACGACCAA

Transgene copy number was estimated using TaqMan based real-time PCR using genomic DNA as described ([Chandler et al., 2007](#)). Briefly, the chloramphenicol cassette within the BAC vector was amplified using a custom primer set from Applied Biosystems (5': GCA CAA GTT TTA TCC GGC CTT TAT T; 3': GTC TTT CAT TGC CAT ACG GAA CTC; and an internal FAM-labeled probe CCG CCT GAT GAA TGC). To estimate copy number, triplicate Δ CT values for the chloramphenicol cassette relative to a genomic DNA control, *Jun* kinase, were compared to the Δ CT values of known quantities of BAC DNA prepared to estimate copy number equivalents.