


## Author's Proof

Before checking your proof, **please read the instructions below**.

- Carefully read the entire proof and mark all corrections in the appropriate place, using the Adobe Reader commenting tools ([Adobe Help](#)). Do not use the Edit tool, as direct edits to the text will be ignored; annotate your corrections instead.
- Provide your corrections in a single PDF file or post your comments in the Production Forum, making sure to reference the relevant query/line number. Upload or post all your corrections directly in the Production Forum to avoid any comments being missed.
- We do not accept corrections in the form of edited manuscripts or via email.
- Do not provide scanned or handwritten corrections.
- Before you submit your corrections, make sure that you have checked your proof carefully as once you approve it, you won't be able to make any further corrections.
- To ensure timely publication of your article, please submit your corrections within 48 hours. After submitting, do not email or query asking for confirmation of receipt.




Do you need help? Visit our [Production Help Center](#) for more information. If you can't find an answer to your question, contact your Production team directly by posting in the Production Forum.

**NOTE FOR CHINESE-SPEAKING AUTHORS:** If you'd like to see a Chinese translation, click on the  symbol next to each query. **Only respond in English** as non-English responses will not be considered. Translated instructions for providing corrections can be found [here](#).

### Quick checklist

- ☐ **Author names** - Complete, accurate and consistent with your previous publications.
- ☐ **Affiliations** - Complete and accurate. Follow this style when applicable: Department, Institute, University, City, Country.
- ☐ **Tables** - Make sure our formatting style has not changed the meaning/alignment of your Tables.
- ☐ **Figures** - Make sure we are using the latest versions.
- ☐ **Funding and Acknowledgments** - List all relevant funders and acknowledgments.
- ☐ **Conflict of interest** - Ensure any relevant conflicts are declared.
- ☐ **Supplementary files** - Ensure the latest files are published and that no line numbers and tracked changes are visible.  
Also, the supplementary files should be cited in the article body text.
- ☐ **Queries** - You must reply to **all of the typesetter's queries below** in order for production to proceed.
- ☐ **Content** - Read all content carefully and ensure any necessary corrections are made, then **upload them** to the Production Forum.

## Author queries form

Query no.	Details required	Author's response
<b>Q1</b>	The citation and surnames of all of the authors have been highlighted. Check that they are correct and consistent with your previous publications, and correct them if needed. Please note that this may affect the indexing of your article in repositories such as PubMed. If adding/removing authors, or changing the order of this list, please provide us with a signed <a href="#">Authorship Change form</a> . 	Please change the second author's name to: Cameron A. MacKenzie
<b>Q2</b>	Confirm whether the article title is correct. 	Yes
<b>Q3</b>	Confirm that all author affiliations are correctly listed. Per our style guidelines, affiliations are listed sequentially and follow author order. Requests for non-sequential affiliation listing will not be fulfilled. Note that affiliations should reflect those at the time during which the work was undertaken. If adding new affiliations, specify if these should be listed as a present address instead of a regular affiliation. 	These are correct

Query no.	Details required	Author's response
<b>Q4</b>	Confirm that the email address in your correspondence section is accurate. Any changes to corresponding authors requires individual confirmation from all original and added/removed corresponding authors. 🌐	The corresponding author should be changed to Cameron A. MacKenzie. Email address: camacken@iastate.edu
<b>Q5</b>	Confirm that the keywords are correct, and keep them to a maximum of eight and a minimum of five. (Note: a keyword can be made up of one or more words.) 🌐	Please remove the first instance of the keywords 'decision analysis'.
<b>Q6</b>	If you decide to use previously published and/or copyrighted figures in your article, please keep in mind that it is your responsibility as the author to obtain the appropriate permissions and licenses to reproduce them, and to follow any citation instructions requested by third-party rights holders. If obtaining the reproduction rights involves the payment of a fee, these charges are to be paid by the authors. 🌐	Understood.
<b>Q7</b>	Check if the section headers (i.e., section leveling) have been correctly captured. 🌐	They are correct.
<b>Q8</b>	If applicable, check your taxonomy and/or nomenclature carefully to avoid any errors that may invalidate it. 🌐	They appear to be correct.
<b>Q9</b>	Ensure that all the figures, tables, and captions are correct, and that all figures are of the highest quality/resolution. If necessary, you may upload improved figures to the Production Forum. Please note that figures and tables must be cited sequentially, per the author guidelines. 🌐	Please see comment on Table 1, Table 2. The figures got put in the wrong order somehow - please see the replacement instruction comments in each figure (1-4) and some figure caption changes.
<b>Q10</b>	Check that all equations and special characters are displayed correctly. 🌐	They appear to be correct.
<b>Q11</b>	Confirm that the Data availability statement is accurate. Note that this statement may have been amended to adhere to our Publication Ethics guidelines. 🌐	Please add the text as indicated in the comment in that section.
<b>Q12</b>	Confirm that the details in the "Author Contributions" section are correct. We have added the sentence "All authors contributed to the article and approved the submitted version." 🌐	Confirmed.
<b>Q13</b>	Ensure all grant numbers and funding information are included and accurate (after publication it is not possible to change this information). All funders should be credited, and all grant numbers should be correctly included in this section. 🌐	Confirmed.

Query no.	Details required	Author's response
<b>Q14</b>	<p>Ensure that any supplementary material is correctly published at this link: <a href="https://www.frontiersin.org/articles/10.3389/fsufs.2023.1040927/full#supplementary-material">https://www.frontiersin.org/articles/10.3389/fsufs.2023.1040927/full#supplementary-material</a></p> <p>If the link does not work, you can check the file(s) directly in the production forum; the published supplementary files appear in green. Please make sure all Supplementary files are cited. Please also provide captions for these files, if relevant. If you have any corrections, please provide new files and republish them in the forum. Frontiers will deposit ALL supplementary files to FigShare and they will receive a DOI. Notify us of any previously deposited material. If the Supplementary Material files contain identifiable or copyright images, please keep in mind that it is your responsibility, as the author, to ensure you have permission to use the images in the article. Please check this <a href="#">link</a> for information on author responsibilities and the publication images. 🌐</p>	The material is correctly published.
<b>Q15</b>	<p>Confirm if the text included in the Conflict of interest statement is correct. Please do not suggest edits to the wording of the final sentence, as this is standard for Frontiers' journal style. 🌐</p>	Confirmed.
<b>Q16</b>	<p>Note that the city name for affiliation 1 has been inserted. Confirm that the city is correct. 🌐</p>	It is correct.
<b>Q17</b>	<p>Note that the keywords "Iowa, Maize" has been repeated twice in the Keyword section, so we have repeated the duplicate. Please confirm if this is fine.</p>	I do not see this duplication. The keywords 'decision analysis' were repeated, and I requested the duplicate be eliminated (see Q5).
<b>Q18</b>	<p>The following reference citations has been changed to match reference list. Please check if correct.</p> <p>"FAO, 2022" was changed to "Food and Agriculture Organization (FAO), 2020"</p> <p>"Quinn et al., 2022" was changed to "Quinn et al., 2021"</p>	<p>Good catch, thank you.</p> <p>Thapa, R., Mirsky, S. B., &amp; Tully, K. L. (2018). Cover crops reduce nitrate leaching in agroecosystems: A global meta-analysis. <i>Journal of environmental quality</i>, 47(6), 1400-1411.</p>
<b>Q19</b>	<p>Please provide complete details for references "Thapa et al., 2018." 🌐</p>	
<b>Q20</b>	<p>Cite the following references inside the text. 🌐</p> <p>"Chatterjee et al., 2020; Da Silva et al., 2021; Martinez-Feria et al., 2019; Pantoja et al., 2015; Park et al., 2022; Roesch-McNally et al., 2018."</p>	<p>Chatterjee et al. 2020 = inside Table 1; Da Silva et al. 2021 = should be Araldi-Da-Silva et al. 2022, added to Table 1; ; Martinez-Feria et al. 2019 = appears inside Table 1; Pantoja et al. 2015 = added in line 491; Park et al. 2022 = added in line 145; Roesch-McNally et al. 2018 = eliminate from citations</p>
<b>Q21</b>	<p>Provide the publisher location for "Da Silva et al., 2021; Hardaker et al., 2015." 🌐</p>	<p>The Da Silva et al. 2021 citation was incorrect, correction added in reference section. Information for Hardaker added in ref section.</p>
<b>Q22</b>	<p>Provide the publisher name and location for "Howard and Abbas, 2010." 🌐</p>	<p>The date on the citation was incorrect - I corrected it in the text and added the publisher info in the ref section</p>
<b>Q23</b>	<p>Provide doi for the reference "Kaspar et al., 2001."</p>	<p>I could not find a doi. I emailed the journal and have not heard back</p>
<b>Q24</b>	<p>provide volume number and page range for the reference "Park et al., 2022."</p>	<p>The doi was added in the reference section.</p>
<b>Q25</b>	<p>The abstract should ideally be structured according to the IMRaD format (Introduction, Methods, Results and Discussion). Provide a structured abstract if possible.</p>	<p>A restructured abstract will be uploaded.</p>



## OPEN ACCESS

## EDITED BY

Jacob Jungers,  
University of Minnesota Twin Cities,  
United States

## REVIEWED BY

Antonio DiTommaso,  
Cornell University, United States  
Atique ur Rehman,  
Bahauddin Zakariya University, Pakistan

## \*CORRESPONDENCE

Gina A. Nichols  
✉ virginia.nichols@gmail.com

## SPECIALTY SECTION

This article was submitted to  
Agroecology and Ecosystem Services,  
a section of the journal  
Frontiers in Sustainable Food Systems

RECEIVED 09 September 2022

ACCEPTED 27 February 2023

PUBLISHED xx xx 2023

## CITATION

Nichols GA and MacKenzie C (2023) Identifying  
research priorities through decision analysis: A  
case study for cover crops.  
*Front. Sustain. Food Syst.* 7:1040927.  
doi: 10.3389/fsufs.2023.1040927

## COPYRIGHT

© 2023 Nichols and MacKenzie. This is an  
open-access article distributed under the terms  
of the [Creative Commons Attribution License](#)  
(CC BY). The use, distribution or reproduction  
in other forums is permitted, provided the  
original author(s) and the copyright owner(s)  
are credited and that the original publication in  
this journal is cited, in accordance with  
accepted academic practice. No use,  
distribution or reproduction is permitted which  
does not comply with these terms.

# Identifying research priorities through decision analysis: A case study for cover crops

Gina A. Nichols<sup>1,2\*</sup> and Cameron MacKenzie<sup>3</sup>

<sup>1</sup>Department of Plant Sciences, University of California, Davis, Davis, CA, United States, <sup>2</sup>Department of Agronomy, Iowa State University, Ames, IA, United States, <sup>3</sup>Department of Industrial and Manufacturing Systems Engineering, Iowa State University, Ames, IA, United States

Midwestern maize (*Zea mays* L.)-based systems, planting an over-wintering cover crop such as rye (*Secale cereale* L.) following fall harvests of summer crops maintains continuous soil cover, offering numerous environmental advantages. However, while adoption of cover crops has increased over the past decade, on a landscape-scale it remains low. Identifying where agronomic research could be most impactful in increasing adoption is therefore a useful exercise. Decision analysis (DA) is a tool for clarifying decision trade-offs, quantifying risk, and identifying optimal decisions. Several fields regularly utilize DA frameworks including the military, industrial engineering, business strategy, and economics, but it is not yet widely applied in agriculture. Here we apply DA to a maize-soybean [*Glycine max* (L.) Merr.] rotation using publicly available weather, management, and economic data from central Iowa. In this region, planting a cover crop following maize (preceding soybean) poses less risk to the producer compared to planting following soybean, meaning it may be a more palatable entry point for producers. Furthermore, the risk of reduced maize yields when planting <14 days following rye termination substantially contributes to the overall risk cover crops pose to producers, but also has significant potential to be addressed through agronomic research. In addition to identifying research priorities, DA provided clarity to a complex problem, was performed using publicly available data, and by incorporating risk it better estimated true costs to the producer compared to using input costs alone. We believe DA is a valuable and underutilized tool in agronomy and could aid in increasing adoption of cover crops in the Midwest.

## KEYWORDS

decision analysis, cover crop, soybean, risk, decision analysis (DA), Iowa (USA), maize (*Zea mays* L.)

## 1. Introduction

Many cropping systems in the United States (US) have undergone simplifications, now being composed of only a few, often annual, crops (Aguilar et al., 2015; Hijmans et al., 2016; Crossley et al., 2021). These systems frequently leave the soil fallow for some period of time, presenting notable environmental challenges including but not limited to increased risk of soil erosion and an increased potential for nutrient loss (Mitsch et al., 2001; Hatfield et al., 2009; Syswerda et al., 2012). The notion of “continuous living cover” has been used to encourage creative solutions to these issues by focusing cropping system re-design on eliminating these environmentally-challenging fallow periods. Planting cover crops to reduce fallow periods is one such tactic that could at least partially address many of the environmental problems presented by annual cropping systems.

The US produces approximately one-third of the world's maize (*Zea mays* L.) and soybean [*Glycine max* (L.) Merr.] [Food and Agriculture Organization (FAO), 2020], with five states in the Midwestern region contributing over half of that production [Feyereisen et al., 2006]. It follows that large amounts of agricultural land in the Midwestern US are dedicated to cropping systems that grow only maize and soybean [Boryan et al., 2011; USDA National Agricultural Statistics Service Cropland Data Layer (USDA NASS CDL), 2021]. Utilizing over-wintering cover crops in these systems has been shown to reduce soil erosion and nitrate leaching [Kaspar et al., 2007, 2012; Chen et al., 2022], is associated with a reduction in crop insurance losses due to drought, excess heat, and excess moisture [Aglasan and Rejesus, 2021], and possibly offer numerous other context-specific benefits such as increased soil infiltration rates, higher soil water-holding capacity, or increased soil organic matter content [Moore et al., 2014; Basche and DeLonge, 2017; Krupek et al., 2022; Nichols et al., 2022]. However, the Midwestern maize-soybean systems present challenges to cover crop adoption. In some regions of the US, cover crop adoption on annual cropland is above 25% and growing [Hamilton et al., 2017]. Meanwhile, states comprising the Midwestern US exhibit some of the lowest adoption rates, with most states well below 10% adoption [Hamilton et al., 2017; Rundquist and Carlson, 2017; Seifert et al., 2018].

Low adoption rates within the Midwest have been the subject of numerous studies, and it is clearly a complex issue involving economics, climate constraints, field operations, management, equipment, culture, and technical knowledge [Lee et al., 2018; Church et al., 2020; Nichols et al., 2020a; Thompson et al., 2021; Yoder et al., 2021]. One barrier we believe merits more attention is that of risk. Risk incorporates two components, uncertainty and negative consequences, and is frequently measured with probabilities describing the potential severity of consequences [Kaplan and Garrick, 1981; Bedford and Cooke, 2001; Hubbard, 2020]. Cover crops present both direct, and indirect risks. Managerially, maize and soybean are both are planted in the late spring (April, May) and harvested in the fall (September, October, November). Producers typically fit over-wintering cover crops into these systems by planting a cover crop in the fall after the cash crop harvest, and terminating the cover crop in the spring before the next cash crop is planted [Sustainable Agriculture Research and Education (SARE), 2020]. Therefore, both the planting and termination of an over-wintering cover crop such as rye (*Secale cereale* L.) can conflict with cash crop management. As such, using a cover crop requires complex decision-making that balances risk and rewards in uncertain conditions. While perceived risks associated with cover cropping are often cited as barriers to adoption [Arbuckle and Roesch-McNally, 2015], quantifying those risks in economic terms is challenging (e.g., Bergtold et al., 2019; Plastina et al., 2020). Furthermore, while lists of cover crop research priorities have been proposed (e.g., Carlson and Stockwell, 2013; Basche and Roesch-McNally, 2017), a tool for ranking priorities would be useful. By quantifying the risk associated with each decision point for producers, research priorities can be set to address points posing the highest risk. The use of risk as a ranking tool would also help researchers and funding organizations assess how resources can be used most impactfully. Furthermore, understanding how uncertainties around weather conditions elevate risks of profit loss is important for understanding

both the mechanisms for delivering incentives, and the amount producers may require for meaningful participation.

Decision analysis is an interdisciplinary tool that can be applied to analyze decision-making under uncertain conditions (Howard, 1988; Howard and Abbas, 2010; Clemen and Reilly, 2013). It can leverage both quantitative information and expert knowledge, incorporate different degrees of risk aversion, and through sensitivity analyses can allow exploration of the decision space [Cegan et al., 2017; Shackelford et al., 2019]. It is a recognized tool for coping with risk in agriculture [Hardaker et al., 2015] and has been applied to a range of agronomic-related topics including agroforestry adoption risks, nitrate pollution loading, cover crop species selection, optimal cropping system choices, and promoting sustainable agricultural practices [Almasri and Kaluarachchi, 2005; Gandorfer et al., 2011; Ramírez-García et al., 2015; Talukder et al., 2017; Do et al., 2020]. However, to our knowledge decision analysis frameworks have had limited application regarding management decisions related to cover crops in the maize/soybean systems of the Midwestern US. Therefore, the objectives of this study were two-fold:

- 1) Provide a case study using publicly available data to demonstrate the process and utility of applying decision analysis to cover crop systems.
- 2) Use a basic analysis to suggest research priorities for cover crops in Central Iowa.

We chose to use Central Iowa as a case study because it has large areas in maize/soybean systems that are broadly representative of the US Midwest [USDA National Agricultural Statistics Service Cropland Data Layer (USDA NASS CDL), 2021], and currently demonstrates a moderate amount of cover crop adoption [Rundquist and Carlson, 2017]. Furthermore, Iowa's land grant institution, Iowa State University, as well as the United States Department of Agriculture (USDA) National Laboratory for Agriculture and the Environment (NLAE) are located in Central Iowa and support a strong infrastructure for publicly funded agronomic research trials in this region that provide rich sources of public data.

## 2. Methods and materials

### 2.1. Decision set

We used cereal rye (*Secale cereale* L.) as our "model" over-wintering cover crop because it is the most used cover crop in Iowa and is one of the most widely used cover crop species in the Midwest [Singer, 2008]. Assuming a producer has both the maize and soybean phase of a maize-soybean rotation growing at a given time, there are two scenarios for cover crop integration, each including three decision alternatives with unique benefits and challenges (Table 1). Concomitant benefits and challenges of each decision alternative highlights the need to use a quantitative approach to decision optimization, which can be achieved using decision analysis frameworks.

**TABLE 1** Two scenarios each including three decision alternatives related to cover cropping in a maize/soybean rotation with various benefits and challenges associated with each alternative.

Decision alternative	Description	Benefits	Challenges
<b>In fields with a soybean crop</b>			
1	Do not plant a cover crop following soybean harvest	No added costs or risks due to cover crop	Low residue from soybean crop leaves soil vulnerable to erosion (Dickey et al., 1985) Soil nitrogen is likely to be lost from the field in the spring to leaching (Qi et al., 2008) Low residue contributes minimally to non-chemical weed control
2	Plant a cover crop, plan to terminate early April	Soybeans are harvested earlier in the fall compared to maize [USDA National Agricultural Statistics Service (USDA NASS), 2022], allowing for earlier cover crop planting which increases likelihood of successful establishment and more cover crop growth (Chatterjee et al., 2020; Nichols et al., 2020b) Cover crop residue reduces soil erosion following soybeans (Kaspar et al., 2001) Cover crop residue may provide weed control following soybeans (Nelson and Bennett, 2018) Cover crop growth can uptake soil nitrate thus mitigating nutrient pollution (Qi et al., 2008; Kaspar et al., 2012; Martinez-Feria et al., 2019)	Cover crop may indirectly reduce subsequent maize yields by competing for workable field days and delaying maize planting, which often results in lower maize yields (Baum et al., 2019) Planting maize less than two weeks following cover crop termination may result in reduced yields, but the effect is unpredictable (Johnson et al., 1998; Acharya et al., 2017, 2020)
3	Plant a cover crop, plan to terminate late April	Enhances cover crop benefits due to more cover crop growth and biomass	Increases chances of delayed maize planting, and thus reduced maize yields
<b>In fields with a maize crop</b>			
4	Do not plant a cover crop following maize harvest	No added costs or risks due to cover crop	Soil nitrogen is likely to be lost from the field in the spring to leaching (Qi et al., 2008)
5	Plant a cover crop, plan to terminate early April	Maize can leave large nitrate reserves in the soil at harvest, and cover crop growth can uptake the nitrate thus mitigating nutrient pollution (Qi et al., 2008; Kaspar et al., 2012; Martinez-Feria et al., 2019) Soybean planting dates are less sensitive to planting dates compared to maize (Kessler et al., 2020)	Timely fall cover crop planting can be difficult following maize harvest Maize is harvested in late fall, and late-planted cover crops can result in low spring cover crop biomass (Chatterjee et al., 2020; Nichols et al., 2020b), and therefore minimal benefits, if terminated in early April
6	Plant a cover crop, plan to terminate late April	Enhances cover crop benefits due to more cover crop growth and biomass	Larger amounts of cover crop biomass may be more difficult to terminate uniformly

## 2.2. Decision structure

The decision set was translated into decision models with known states, uncertainties, and values, each described below.

### 2.2.1. Fall weather uncertainties

Cover crops are most often planted following cash crop harvests [Sustainable Agriculture Research and Education (SARE), 2020]. Soybean crops in Central Iowa are harvested in September or

October, and maize in October or November [USDA National Agricultural Statistics Service (USDA NASS), 2022]. Planting cover crops into standing crops before harvest can increase the probability of establishment (Wilson et al., 2014) but requires specialized equipment that is not yet widely available. We therefore assume cover crop planting occurs after cash crop harvest.

Seeds require precipitation to germinate, and heat units to establish such that the plants emerge and survive the winter. Failure of a cover crop to germinate or establish in the fall results in wasted seed, wasted fuel, and possible weed problems the following



spring. While the amount of precipitation needed for rye to germinate depends on soil moisture conditions at planting, crop advisors and producers often assume 1.27 cm (0.5 inches) is needed (Sarrantonio, 1994), which is consistent with field studies (Fisher et al., 2011; Wilson et al., 2013) and simulation model assumptions (Feyereisen et al., 2006; Marcillo et al., 2019). While we assumed 1.27 cm was needed for our baseline analysis, this assumption was tested through a sensitivity analysis (see Section 2.3.2).

Growing degree days (GDDs) represent an estimation of the number of heat units accumulated above a threshold temperature specific to a crop. For rye the threshold is 0 or 1°C (Feyereisen et al., 2006). We acknowledge the number of GDDs required for rye to successfully over-winter will depend on several additional factors including soil texture and snow cover. A study in Minnesota suggested rye required at least 100 GDDs in the fall to produce biomass in the spring (Kantar and Porter, 2014). We therefore estimated rye requires 100 GDDs to successfully establish before winter, but tested the sensitivity of this assumption (see Section 2.3.2).

To estimate the probability of successful rye establishment, we used 30 years of historical weather data (1988–2019) collected at the AMES-8-WSW station from the Iowa Environmental Mesonet (IEM) (2022). We chose this dataset because it had previously undergone an extensive quality check (Archontoulis et al., 2020). Using 30 years of weather data, we calculated (i) the probability the site received 1.27 cm of rainfall during an allotted timeframe, and (ii) the probability of achieving 100 GDDs in the allotted timeframe. The timeframes differed by decision alternative to account for the generally earlier harvest dates for soybean compared to maize [USDA National Agricultural Statistics Service (USDA NASS), 2022]. The precipitation timeframes were 15-Oct through 30-Nov and 1-Nov through 30-Nov for rye following soybean and rye following maize, respectively. The GDD accumulation timeframes were 15-Oct through 1-Dec and 1-Nov through 1-Dec for rye following soybean and rye following maize, respectively. We chose to calculate the precipitation and GDD probabilities separately rather than as a joint probability to aid in assessing how breeding efforts could increase changes of establishment. We recognize this model for establishment is a simplification of the complex interactions between weather, soil, and management considerations. While more sophisticated modeling approaches have been utilized for predicting cover crop establishment (Baker and Griffis, 2009; Marcillo et al., 2019; Nichols et al., 2020b), they require specialized skillsets and a significant time commitment. Our goal in this exercise is to demonstrate how insights can be obtained using publicly available data and approachable methodologies.

## 2.2.2. Spring weather uncertainties

Iowa has a humid continental climate wherein a significant amount of precipitation occurs during the spring months. In addition to the direct constraints on management that precipitation exerts, performing field operations in wet soils can result in undesirable outcomes including long-term soil compaction and equipment malfunctions. The USDA National Agricultural Statistics Service (NASS) surveys producers to determine the

number of days suitable for fieldwork (workable-field day; WFD) for each week throughout the year [USDA National Agricultural Statistics Service (USDA NASS), 2018]. A “suitable” day is defined as one in which weather and field conditions allow producers to work in fields the majority of a given day. Determining whether a day is a “suitable” is subjective, but provides valuable information about the progress and constraints of agricultural production on a landscape level.

Historical data shows that in Iowa, the number of WFDs during the spring can severely restrict field activities (Urban et al., 2015; Edwards, 2020). To comply with governmental crop insurance cost-share policies, cover crops must be terminated before the cash crop is planted [Bergtold et al., 2019; USDA Risk Management Agency (USDA RMA), 2019]. Therefore, the presence of a living cover crop that must be terminated before the cash crop can be planted can potentially add to the spring workload for a producer. While this depends on whether producers typically have a pre-plant or pre-emergent herbicide pass, the operation is much less crucial when the goal is simply to eliminate weeds around cash crop planting compared to killing a live cover crop to comply with federal crop insurance requirements. To account for the increased importance of timely cover crop termination, in this exercise we assumed cover crop termination requires an additional set of field working-days compared to systems without a cover crop. However, because many producers do a pre-plant or pre-emergent herbicide pass in systems without cover crops, we did not assume extra herbicide or fuel costs associated with terminating the cover crop. In short, we assumed producers who plant a rye cover crop require two more spring WFDs than those who do not. This fact introduces an important component of risk that is often not accounted for explicitly in economic analyses.

The decision of cover crop termination timing will also affect WFDs, and therefore may indirectly affect cash crop yields. If a producer has WFDs in early April, the producer must choose whether to utilize them to terminate the cover crop, or wait in order to accrue more benefits from prolonged cover crop growth (Table 1). Societal-level benefits such as reduced nitrate leaching, as well as farm-level benefits such as the potential to off-set weed control costs, increase as spring cover crop termination dates are delayed and cover crop biomass increases (Finney et al., 2016; Thapa et al., 2018; Nichols et al., 2020b). However, the by choosing not to utilize early April WFDs, the producer risks not having sufficient WFDs in late April to terminate the cover crop or plant the cash crop, resulting in delayed cash crop planting and a possible concomitant reduction in yields. Therefore, understanding the uncertainty around WFDs in the spring is an important component in assessing optimal decision alternatives.

In this analysis we only include the uncertainties associated with WFDs. In years with very low spring precipitation, delaying cover crop termination can also result in decreased cash crop yields due to the cover crop’s use of stored soil water needed for cash crop production. While this risk is possible, due to climatic patterns it is not common in Central Iowa (Daigh et al., 2014; Martinez-Feria et al., 2016). Therefore, the risk of cover crops inducing drought-related yield reductions in the following cash crop is not considered in this exercise.

Q19

Workable field days are estimated by surveying farmers about how many days in the previous week were field-workable. The data is therefore reported as a number of days within a 7-day calendar period, with this period being inconsistent between years. For the purposes of this exercise, we chose to take the total WFDs over the 7-day reporting period and divide the total by seven to assign a number of WFDs to each calendar day the reporting week included. We then created five spring categories (early April, late April, early May, and late May, June). Workable field day values were then summed within these spring calendar categories. More details, including R code, concerning this procedure can be found in [Supplementary material](#). We assumed cover crop termination would require two WFDs within a spring category, and cash crop planting would likewise require two WFDs. Therefore, cover crop termination and cash crop planting within a given window would require four WFDs. The probability of two and four WFDs being reported in a given spring category was calculated using 30 years of historical data (1988–2019).

### 2.2.3. Subsequent maize yield uncertainties

On average, winter cover crops such as rye have been shown to have a neutral effect on subsequent maize and soybean yields ([Marcillo and Miguez, 2017](#)). However, numerous studies have shown that under certain conditions, planting maize <10–14 days following cover crop termination can result in lower maize yields ([Johnson et al., 1998](#); [Acharya et al., 2017](#); [Hirsh et al., 2021](#); [Quinn et al., 2021](#)). We assumed a producer would plant their maize crop as early as possible, regardless of the penalty that would be incurred due to the <14 day window. We made this assumption because conversations with producers confirmed that while they were aware there may be a yield penalty from a small termination-planting window, it was inconsistent and may not occur at all, and they were therefore more concerned with timely maize planting. We therefore assumed if there four WFDs in a given spring category, the producer would plant maize but there would be a 50% chance of a 10% decrease in maize yield. We acknowledge that in our scenarios, the 10% yield penalty from the small termination-planting window is larger than the penalty incurred for delaying planting until late May, but we believe our decision structure captures the uncertainty currently associated with whether that yield penalty will be incurred. Soybeans are not impacted by the time between rye termination and soybean planting ([Acharya et al., 2020](#)), so no yield penalty was assigned in those circumstances.

### 2.2.4. Value

The main contributors to decision value were estimated using partial budgets and included the costs from planting a cover crop, the savings from planting a cover crop, and the income from the subsequent cash crop. Extension publications, farming group publications, and peer-reviewed literature were used to guide each estimation. Sensitivity analyses were performed on assumed values (Section 2.3.2), and instances where conclusions were overly sensitive to assumptions were noted.

To estimate the direct costs associated with planting and terminating a cover crop we used Iowa State University's "Economics of Cover Crops" decision tool ([Iowa State University](#)

[Extension, 2018](#)). While these prices will fluctuate depending on the price of fuel and labor, we feel they are sufficiently representative for this exercise ([Table 2](#)).

In order to account for the effect of cover cropping on income from crop yields, we needed to estimate the net revenue a producer expects per unit crop yield. The net revenue from a crop will depend on producer costs of production as well as market prices, both of which vary significantly across years. To overcome this variability, we looked at production costs ([Iowa State University Extension, 2022](#)) and market prices [[USDA National Agricultural Statistics Service \(USDA NASS\), 2022](#)] from 2013 to 2021, calculated the net revenue per unit crop yield for each year, then took the year with the maximum net revenue for each crop ([Figure 1](#)). By calculating the net revenue in this manner, when a rye cover crop negatively impacted cash crop yields our analyses represented the highest potential costs of those effects. All prices and calculations are available in [Supplementary material](#).

Maize was assumed to have a maximum yield of 10.7 dry Mg ha<sup>-1</sup> (200 bu ac<sup>-1</sup>) and soybean a yield of 1.4 dry Mg ha<sup>-1</sup> (60 bu ac<sup>-1</sup>), which are representative of the state average yields in Iowa [[USDA National Agricultural Statistics Service \(USDA NASS\), 2022](#)]. Maize yield is sensitive to planting date, with later planting dates being associated with lower yields ([Kucharik, 2008](#); [Baum et al., 2019](#)). We therefore assume a graduated yield penalty increasing 5–20% as maize planting occurs past April ([Supplementary Table S2](#)). In summary, the decision of whether to terminate the cover crop early or late impacts the available WFDs ([Table 3](#)), which impact whether the producer incurs a termination-planting penalty or a late-planting penalty, both of which impact the value of the decision.

Soybean yields are less sensitive to planting dates compared to maize ([Kessler et al., 2020](#)) and therefore was assumed to have a less severe graduated penalty as planting was delayed (5–10%; [Table 2](#)).

When the cover crop was followed by a maize crop (decision alternatives 1–3), we assumed herbicide costs were equal in the cover crop and no-cover alternatives (\$205 ha<sup>-1</sup>). When the cover crop was followed by a soybean crop (decision alternatives 4–6), we utilized information from on-farm experiments showing producers reduced herbicide costs due to the mulch provided by a late-terminated cover crop. Therefore, in the decision alternative where the cover crop was terminated in late April or later followed by soybean planting (decision alternative 6), a \$37 ha<sup>-1</sup> savings in herbicides was applied ([Nelson and Bennett, 2018](#)).

There are currently no payments available to farmers in Iowa for the societal benefits reaped from delaying cover crop termination. However, other areas in the US have implemented payment structures that reward late termination due to the societal benefits gained from late termination ([Maryland Department of Agriculture, 2022](#)), so the potential for this payment in decision alternatives 3 and 6 was included in sensitivity analyses.

## 2.3. Decision analysis

### 2.3.1. Building decision trees

Decisions can be visualized and modeled using decision tree notation ([Howard and Abbas, 2010](#); [Clemen and Reilly, 2013](#)).



TABLE 2 Summary of economic assumptions for each scenario, the assumed relative cash crop yield indicated in parentheses.

	No cover crop system	Cover crop system	
		14+ day gap	<14 day gap <sup>a,b</sup>
Cover crop			
Cover crop seed	–	\$20 ha <sup>-1</sup>	\$20 ha <sup>-1</sup>
Cover crop planting	–	\$32 ha <sup>-1</sup>	\$32 ha <sup>-1</sup>
Cost-shares/insurance discounts with cover crop planting	–	\$12–74 ha <sup>-1</sup>	\$12–74 ha <sup>-1</sup>
Cover crop preceding maize			
Herbicide costs	\$205 ha <sup>-1</sup>	\$205 ha <sup>-1</sup>	\$205 ha <sup>-1</sup>
Maize income (assumed \$2.14 net income per bushel)			
Planted early April	\$1057 ha <sup>-1</sup>	–	\$1057 ha <sup>-1</sup> /\$951 ha <sup>-1</sup> (90%)
Planted late April	\$1057 ha <sup>-1</sup>	\$1057 ha <sup>-1</sup>	\$1057 ha <sup>-1</sup> /\$951 ha <sup>-1</sup> (90%)
Planted early May <sup>c</sup>	\$1004 ha <sup>-1</sup> (95%)	\$1004 ha <sup>-1</sup> (95%)	\$1004 ha <sup>-1</sup> (95%)/\$889 ha <sup>-1</sup> (85%)
Planted late May	\$951 ha <sup>-1</sup> (90%)	\$951 ha <sup>-1</sup> (90%)	\$951 ha <sup>-1</sup> (90%)/\$846 ha <sup>-1</sup> (80%)
Planted June	\$846 ha <sup>-1</sup> (80%)	\$846 ha <sup>-1</sup> (80%)	\$846 ha <sup>-1</sup> (80%)/\$740 ha <sup>-1</sup> (70%)
Cover crop preceding soybean			
Herbicide costs <sup>d</sup>	\$205 ha <sup>-1</sup>	\$168 ha <sup>-1</sup>	\$168 ha <sup>-1</sup>
Soybean income (assumed \$4.06 net income per bushel)			
Planted early April	–	–	–
Planted late April	\$601 ha <sup>-1</sup>	\$601 ha <sup>-1</sup>	\$601 ha <sup>-1</sup>
Planted early May	\$601 ha <sup>-1</sup>	\$601 ha <sup>-1</sup>	\$601 ha <sup>-1</sup>
Planted late May <sup>e</sup>	\$571 ha <sup>-1</sup> (95%)	\$571 ha <sup>-1</sup> (95%)	\$571 ha <sup>-1</sup> (95%)
Planted June	\$541 ha <sup>-1</sup> (90%)	\$541 ha <sup>-1</sup> (90%)	\$541 ha <sup>-1</sup> (90%)

<sup>a</sup>The decision model for rye following soybean includes a 50% chance a <14 day maize yield reduction will not occur (first values listed), and 50% chance the <14 day maize yield reduction will occur (second values listed).

<sup>b</sup>Estimated maize yield reduction due to termination-planting gap are based on Johnson et al. (1998), Hirsh et al. (2021), and Quinn et al. (2021).

<sup>c</sup>Estimated maize yield reduction due to delayed maize planting are based on Kucharik (2008) and Baum et al. (2019).

<sup>d</sup>Estimated reduction in herbicide costs based on Nelson and Bennett (2018).

<sup>e</sup>Estimated soybean yield reduction based on Kessler et al. (2020).

The full decision model is available in [Supplementary material](#) and consists of building out a branch for each unique decision node and uncertainty outcome with probabilities, then assigning a value to each branch. We assume a risk-neutral decision maker which means that the decision maker should choose the alternative that maximizes his or her expected value. A square in the decision tree represents a choice between two or more alternatives, and a circle represents an uncertainty where each branch stemming from the uncertainty is assigned a probability. The first decision for the producer is whether or not to plant a cover crop in the fall (Figure 1). If the producers choose to plant a cover crop, there is an uncertainty about whether or not sufficient precipitation occurs followed by a second uncertainty about whether or not sufficient GDDs are accumulated. If sufficient precipitation and sufficient GDDs occur, the producer makes a second decision about whether or not to terminate in early April (Supplementary Figure S1). This decision is followed by uncertainties in the number of WFDs available in a given time frame, and whether there is a penalty when maize is planted in the same spring category as cover crop termination. The decision tree is solved using a “rollback” procedure starting from the right-hand side of the tree. If a decision

(square) node is encountered, the alternative with the largest expected monetary value is selected. If an uncertainty (circle) node is encountered, the expected monetary value is calculated using the probabilities on the branches as weights. This procedure results in identifying the alternative for a given decision (e.g., whether or not to plant a cover crop in the fall) that maximizes the producer’s expected monetary value.

2.3.2. Sensitivity analyses

Sensitivity analysis on the uncertainty and parameter assumptions can provide insight into the criticality and importance of an assumption or variable to the decision. The sensitivity of outcomes was assessed for the precipitation required for rye germination (ranging from 0 to 3.5 cm in 1 mm increments), the number of GDDs needed for rye to over-winter (ranging from 0 to 300 in 5 GDD increments), the potential relative reduction in maize yields when maize was planted <14 days following rye termination (ranging from 0 to 20% in 5% increments), the incentive payments offered to plant rye (ranging from \$0 to 200




TABLE 3 Summary of probabilities of workable field days (WFDs) in a given timeframe based on 30 years of NASS survey data [USDA National Agricultural Statistics Service (USDA NASS), 2022].

Management window	Probability of two or more workable field days (WFDs)	Probability of four or more WFDs
1-Apr through 15-Apr (early April)	69%	48%
16-Apr through 30-Apr (late April)	71%	37%
1-May through 15-May (early May)	89%	45%
16-May through 31-May (late May)	87%	55%

ha<sup>-1</sup> in \$1 increments), and the incentive payments offered to delay termination of rye (ranging from \$0 to 200 ha<sup>-1</sup> in \$1 increments). Additionally, sensitivity analyses were performed on the assumed revenues and costs associated with each scenario to ensure conclusions were not overly sensitive to these assumptions (see [Supplementary material](#); [Gupta, 2022](#) for details).

2.3.3. Value of information

In our decision model, if a producer has two WFDs within 14 days following cover crop termination, they have a 50% of incurring a 10% maize yield reduction if they choose to plant. This uncertainty is due to research gaps—we do not yet have sufficient information to provide a producer to help them determine whether this reduction will occur. By estimating the value of the decision if the producer knows whether the yield reduction will occur, one

can estimate the “value of perfect information” ([Repo, 1989](#)). This provides an estimate of what that information would be worth to producers, thus allowing researchers to assess how impactful such research would be. We therefore estimated the value of knowing when there would not be a reduction in maize yields when planting <14 days after cover crop termination.

3. Results and discussion

3.1. Optimal decisions

Assuming there is no cost-share available for planting a cover crop and long-term or societal economic benefits are not accounted for, the overall expected monetary value of not planting a cover crop is greater than the expected monetary value of planting a cover crop, regardless of the sequencing scenario ([Figure 2](#)). This analysis shows that in addition to the cost of seed and fuel to plant the cover crop (\$50 ha<sup>-1</sup>), when rye precedes maize there is an additional \$40–70 ha<sup>-1</sup> cost associated with the risk that the spring management of the cover crop will result in reduced maize yields (either through delayed maize planting due to insufficient WFDs or <14 day gap penalties). When rye precedes soybeans, the costs of planting the cover crop and risks of reduced yields due to delayed planting are partially compensated by through reduced herbicide costs. Within the decision sets that include the alternative of planting a cover crop, the value of the decision is always maximized if the cover crop is terminated in early vs. late April.

Many of the benefits reaped from planting cover crops (e.g., reduced soil erosion, reduced nitrate leaching, non-chemical weed control) are directly related to the amount of biomass the cover crop produces ([Finney et al., 2016](#); [Thapa et al., 2018](#); [Nichols et al., 2020b](#)). However, in areas that lack incentives for delaying cover crop termination to allow the cover crop to grow, our analyses show the optimal decision is to terminate the cover crop as soon as possible, even when there might be cost savings from reduced herbicide use (Rye-Soybean scenario in [Figure 2](#)). Notably, the termination decision differential is highest when the cover crop precedes maize, meaning the sequencing where society may benefit the most (higher mitigation of erosion and nitrate leaching, [Table 1](#)) would also require the highest incentives to render late April termination the optimal decision. The US state of Maryland has created a tiered incentive system wherein producers are compensated more for early planting and late termination of cover crops ([Maryland Department of Agriculture, 2022](#)). Our analysis indicates having compensation rates differ by cropping sequence may also be an approach worth considering.

Our analyses also expose a potential moral hazard. If a producer chooses to plant a cover crop preceding a maize crop and receives a cost-share or incentive for doing so, failed cover crop establishment will lead to a better financial result than successful establishment (Rye-Maize scenario in [Figure 2](#)). It is important to provide support for producers as they learn to manage cover crops, and often cover crop establishment is out of a producer’s control, but our analyses demonstrate the complexity in determining the best payment structures, and the need to include the risks the cover crop may pose to the subsequent crop yields.




FIGURE 2

Value of each decision alternative (rounded to the nearest \$10) assuming no incentive/cost share payments for planting a cover crop.


### 3.2. Sensitivity to cost-share/incentives

If there are no cost-shares or incentive programs, the overall expected monetary value of not planting any cover crop is greater than the expected value of planting a cover crop, regardless of the sequencing scenario (in the top panel of Figure 3, this is seen from the “do not plant rye” alternative having a greater value when the cost share or incentive is \$0 on the horizontal axis). However, current incentive programs may be enough to make planting a cover crop preceding a soybean cash crop (“Rye-Soybean”) the optimal decision. If the incentive is greater than \$30 ha<sup>-1</sup>, the expected monetary value of planting rye prior to soybeans is greater than not planting rye.

When a cover crop precedes a maize crop (“Rye-Maize” in top panel of Figure 3), within the current range of incentives the optimal decision is to not plant a cover crop. However, this recommendation is sensitive to the reduction in maize yield due to planting <10 days following cover crop termination (bottom panel of Figure 3). If the potential reduction in yield were eliminated,

the difference between the value of not planting a cover crop and planting a cover crop could be reduced from \$85 ha<sup>-1</sup> to \$60 ha<sup>-1</sup>, bringing the difference into the range of current incentive programs in this area (\$12–74 ha<sup>-1</sup>).

The exact causes of the reduced yield in maize are not yet clear and it is currently not possible to predict when they will manifest (e.g., Patel et al., 2019; Quinn et al., 2021). The value of perfect information is worth \$20–25 depending on the planned cover crop timing, which is roughly equal to the increased value from eliminating the yield penalty. This indicates that research that allows producers to accurately predict when the yield penalty will occur is equally as valuable as eliminating the yield penalty. Potential mechanisms include altered nutrient dynamics, disease pressure, allelopathy, rye stands that are not fully terminated, changes in soil temperature and/or moisture in a rye cover crop system. A meta-analysis of studies may aid in identifying factors that drive the variation in the effect. Our analyses demonstrate that this phenomenon poses a significant risk to producers, and a better understanding of the drivers and identification of ways to predict



when yield declines are likely would greatly reduce the financial risk associated with planting a rye cover crop in these systems.

### 3.3. Sensitivity to weather

On average, Central Iowa received 7.4 and 4.2 cm of rain from 15-Oct and 1-Nov through 30-Nov, respectively. This equated to a high probability (>80%) of the rye cover crop receiving sufficient precipitation for germination (>1.27 cm) in both sequences (Figure 4, Supplementary Table S3). This result was robust against uncertainty in our assumptions; even if rye required almost double the assumed precipitation, the probability of receiving that amount of rainfall did not drop below 80% for either planting scenario (Figure 4). While the probability of accumulating sufficient GDDs (100) was 100% when the rye was planted following soybeans (15-Oct planting date), it dropped to 71% chance of success when planted following maize (1-Nov planting date; Supplementary Table S3). The probability of

establishment was very sensitive to the sequencing (rye following soybeans or rye following maize). For the 1-Nov planting date, the results are very sensitive to the assumed GDDs required for establishment.


These results can be used to guide research efforts. Our analysis demonstrates that in most cases, precipitation is not the limiting factor for cover crop establishment in Central Iowa. Breeding varieties that require less precipitation to germinate would likely involve breeding for smaller seeds, which carries inherent tradeoffs (e.g., Carleton and Cooper, 1972; Mohler et al., 2009). A study done in Minnesota showed precipitation accounts for the highest amount of variation in rye establishment, followed by temperature (Wilson et al., 2013), demonstrating the value of evaluating weather-related risks locally. While our results do not account for how the precipitation is distributed across time and how that may impact germination, our results suggest this area of Iowa can support larger precipitation requirements for cover crops without experiencing a significant reduction in the probability of cover crop germination.

Our results also show when planting after soybean harvest, the cover crop is almost guaranteed to gain 100 GDDs in the fall (Figure 4). Conversely, after maize harvest the probability is very sensitive to how many GDDs are assumed to be needed. Our analyses highlight the need to better understand conditions that lead to successful establishment, particularly in the later months of the year. Additionally, research focused on identifying management tactics that allow for earlier cover crop planting may be most effective in increasing the probability of successful cover crop establishment in Central Iowa. For example, some producers report switching to earlier maturing soybean and maize varieties when adopting cover crops in order to plant the cover crop earlier (Plastina et al., 2020). Some areas have organized blocks of producers who share in aerial seeding costs, and custom seeding equipment/services that allows for seeding into a standing crop are becoming more common. Our analyses indicate these types of activities are well-suited to reducing the risk associated with planting a cover crop in Central Iowa.

In the spring, the number of WFDs presented a great deal of uncertainty (Table 3). Averaged over the entire spring period (1-Apr through 31-May), there was a 79% probability of two or more WFDs in a given 2-week period, and only a 46% probability of four or more WFDs. We assumed two or more WFDs were needed to successfully complete a cover crop termination activity, and two additional WFDs were needed to complete cash crop planting activities. Therefore, producers wishing to terminate and plant within a 2-week period may not have sufficient WFDs to do so. The probability of two or more WFDs was higher in May compared to April, indicating paying producers to delay cover crop termination may also increase the chances the producer can terminate in their planned timeframe.

Our analyses indicate in Central Iowa, there is generally a high probability the fall conditions will foster cover crop establishment, and that the majority of risk occurs due to the potential for the additional management required in the spring to delay cash crop planting. A Midwestern focus group found some producers had been switching to winterkill cover crop varieties because of the difficulties associated with killing the cover crop and planting a cash crop in a timely manner in the spring (Plastina et al., 2020). For





this analysis we assumed the rye cover crop could be terminated at any point, but the stage of rye growth will affect how easy it is to terminate, particularly when using mechanical termination (Creamer and Dabney, 2002; Mirsky et al., 2009). Decision support tools that help producers decide if early termination is the best choice could be beneficial in helping producers manage this risk.

4. Conclusions

Using publicly available data and reasonable assumptions, we were able to gain significant insight into localized priorities for cover crop research. Using historical weather data, NASS surveys on WFDs, extension publications, and a partial budget for cover crop economics we were able to build a single-attribute decision model, and model decision values assuming a risk-neutral producer. Our analysis does not include possible long-term impacts such as the maintenance of productivity, long-term impacts on weeds or insects, or changes in yield stability over time, which could be incorporated in future applications of this framework. We found including only the costs of seed and fuel in cover crop economics underestimates the additional financial risk producers assume due to the extra spring work cover crops might entail in areas with limited numbers of WFDs during that time. We found there is minimal information on the number of GDDs required for a rye cover crop to successfully overwinter, and that this may have a large impact on risks associated with planting cover crops in Central Iowa. In Central Iowa, identifying ways to ensure early cover crop planting and managements that render maize yields less sensitive to rye cover crop termination timing, or that allow that reduction to be more predictable, could significantly help reduce the financial risk of planting cover crops. Furthermore, flat payments for planting cover crops may result in a moral hazard, wherein the decision value for planting a cover crop preceding a maize crop is maximized when the cover crop fails to establish in the fall. Policies that promote tiered payment structures could rectify this while still providing support for producers as they learn to manage cover crops.

Data availability statement

The original contributions presented in the study are included in the article/Supplementary material, further inquiries can be directed to the corresponding author.

Author contributions

GN conceived of and designed the analyses, collected the data, performed the analyses, and wrote the first draft of the manuscript. CM designed the analyses, and edited the manuscript. All authors contributed to the article and approved the submitted version.

Funding

This material is based upon work supported in part by the National Science Foundation (Grant No. DGE-1828942).

Acknowledgments

We acknowledge Dr. Sarah Ryan for organizing a workshop for the NSF DataFEWSion program that sparked this collaboration, Dr. Raj Raman for his helpful feedback, and two reviewers for their insightful comments and suggestions that greatly improved the quality of this manuscript.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of



their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## References

- Acharya, J., Bakker, M. G., Moorman, T. B., Kaspar, T. C., Lenssen, A. W., and Robertson, A. E. (2017). Time interval between cover crop termination and planting influences corn seedling disease, plant growth, and yield. *Plant Dis.* 101, 591–600. doi: 10.1094/PDIS-07-16-0975-RE
- Acharya, J., Moorman, T. B., Kaspar, T. C., Lenssen, A. W., and Robertson, A. E. (2020). Cover crop rotation effects on growth and development, seedling disease, and yield of corn and soybean. *Plant Dis.* 104, 677–687. doi: 10.1094/PDIS-09-19-1904-RE
- Aglasan, S., and Rejesus, R. M. (2021). “An analysis of crop insurance losses, cover crops, and weather in US Crop Production,” in *2021 Agricultural and Applied Economics Association Annual Meeting* (Austin, TX).
- Aguilar, J., Gramig, G. G., Hendrickson, J. R., Archer, D. W., Forcella, F., et al. (2015). Crop species diversity changes in the United States: 1978–2012. *PLoS ONE* 10, e0136580. doi: 10.1371/JOURNAL.PONE.0136580
- Almasri, M. N., and Kaluarachchi, J. J. (2005). Multi-criteria decision analysis for the optimal management of nitrate contamination of aquifers. *J. Environ. Manage.* 74, 365–381. doi: 10.1016/j.jenvman.2004.10.006
- Arbuckle, J. G., and Roesch-McNally, G. (2015). Cover crop adoption in Iowa: The role of perceived practice characteristics. *J. Soil Water Conserv.* 70, 418–429. doi: 10.2489/jswc.70.6.418
- Archontoulis, S. V., Castellano, M. J., Licht, M. A., Nichols, V., Baum, M., Huber, L., et al. (2020). Predicting crop yields and soil-plant nitrogen dynamics in the US Corn Belt. *Crop Sci.* 60, 721–738. doi: 10.1002/csc2.20039
- Baker, J. M., and Griffis, T. J. (2009). Evaluating the potential use of winter cover crops in corn-soybean systems for sustainable co-production of food and fuel. *Agric. For. Meteorol.* 149, 2120–2132. doi: 10.1016/j.agrformet.2009.05.017
- Basche, A., and DeLonge, M. (2017). The impact of continuous living cover on soil hydrologic properties: a meta-analysis. *Soil Sci. Soc. Am. J.* 81, 1179–1190. doi: 10.2136/sssaj2017.03.0077
- Basche, A. D., and Roesch-McNally, G. E. (2017). Research topics to scale up cover crop use: Reflections from innovative Iowa farmers. *J. Soil Water Conserv.* 72, 59A–63A. doi: 10.2489/jswc.72.3.59A
- Baum, M. E., Archontoulis, S. V., and Licht, M. A. (2019). Planting date, hybrid maturity, and weather effects on maize yield and crop stage. *Agron. J.* 111, 303–313. doi: 10.2134/agnonj2018.04.0297
- Bedford, T., and Cooke, R. (2001). *Probabilistic Risk Analysis: Foundations and Methods*. Cambridge: Cambridge University Press. doi: 10.1017/CBO9780511813597
- Bergtold, J. S., Ramsey, S., Maddy, L., and Williams, J. R. (2019). A review of economic considerations for cover crops as a conservation practice. *Renew. Agric. Food Syst.* 34, 62–76. doi: 10.1017/S1742170517000278
- Boryan, C., Yang, Z., Mueller, R., and Craig, M. (2011). Monitoring US agriculture: the US Department of Agriculture, National Agricultural Statistics Service, Cropland Data Layer Program. *Geocarto Int.* 26, 341–358. doi: 10.1080/10106049.2011.562309
- Carleton, A. E., and Cooper, C. S. (1972). Seed size effects upon seedling vigor of three forage legumes 1. *Crop Sci.* 12, 183–186. doi: 10.1002/aecp.13248
- Carlson, S., and Stockwell, R. (2013). Research priorities for advancing adoption of cover crops in agriculture-intensive regions. *J. Agric. Food Syst. Commun. Develop.* 3, 125–129. doi: 10.5304/jafscd.2013.034.017
- Cegan, J. C., Fillion, A. M., Keisler, J. M., and Linkov, I. (2017). Trends and applications of multi-criteria decision analysis in environmental sciences: literature review. *Environ. Syst. Decis.* 37, 123–133. doi: 10.1007/s10669-017-9642-9
- Chatterjee, N., Archontoulis, S. V., Bastidas, A., Proctor, C. A., Elmore, R. W., and Basche, A. D. (2020). Simulating winter rye cover crop production under alternative management in a corn-soybean rotation. *Agron. J.* 112, 4648–4665. doi: 10.1002/aj2.20377
- Chen, L., Rejesus, R. M., Aglasan, S., Hagen, S. C., and Salas, W. (2022). The of cover on soil erosion in the US Midwest. *J. Environ. Manage.* 324. doi: 10.1016/j.jenvman.2022.116168

## Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fsufs.2023.1040927/full#supplementary-material>

- Church, S. P., Lu, J., Ranjan, P., Reimer, A. P., and Prokopy, L. S. (2020). The role of systems thinking in cover crop adoption: implications for conservation communication. *Land Use Policy* 94, 104508. doi: 10.1016/j.landusepol.2020.104508
- Clemen, R. T., and Reilly, T. (2013). *Making hard decisions with DecisionTools*. Beijing: Cengage Learning.
- Creamer, N. G., and Dabney, S. M. (2002). Killing cover crops mechanically: review of recent literature and assessment of new research results. *Am. J. Altern. Agric.* 17, 32–40. doi: 10.1079/AJAA200204
- Crossley, M. S., Burke, K. D., Schoville, S. D., and Radeloff, V. C. (2021). Recent collapse of crop belts and declining diversity of US agriculture since 1840. *Glob. Chang. Biol.* 27, 151–164. doi: 10.1111/gcb.15396
- Da Silva, G. A., Han, S., Candel, Y. R., Mueller, D. S., Helmers, M., Kaspar, T. C., et al. (2021). *Field Study of the Effect of Rye Cover Crop on Soybean Root Disease and Productivity*. PhytoFrontiers.
- Daigh, A. L., Helmers, M. J., Klavivko, E., Zhou, X., Goeken, R., Cavdini, J., et al. (2014). Soil water during the drought of 2012 as affected by rye cover crops in fields in Iowa and Indiana. *J. Soil Water Conserv.* 69, 564–573. doi: 10.2489/jswc.69.6.564
- Dickey, E. C., Shelton, D. P., Jasa, P. J., and Peterson, T. R. (1985). Soil erosion from tillage systems used in soybean and corn residues. *Trans. ASAE* 28, 1124–1130. doi: 10.13031/2013.32399
- Do, H., Luedeling, E., and Whitney, C. (2020). Decision analysis of agroforestry options reveals adoption risks for resource-poor farmers. *Agron. Sustain. Dev.* 40, 1–12. doi: 10.1007/s13593-020-00624-5
- Edwards, W. (2020). *The Number of Days Suitable for Fieldwork in Iowa is Shrinking*. Iowa State University Extension and Outreach. Available online at: <https://www.extension.iastate.edu/agdm/articles/edwards/EdwMar20.html> (accessed September 2, 2022).
- Feyerisen, G. W., Wilson, B. N., Sands, G. R., Strock, J. S., and Porter, P. M. (2006). Potential for a rye cover crop to reduce nitrate loss in southwestern Minnesota. *Agron. J.* 98, 1416–1426. doi: 10.2134/agnonj2005.0134
- Finney, D. M., White, C. M., and Kaye, J. P. (2016). Biomass production and carbon/nitrogen ratio influence ecosystem services from cover crop mixtures. *Agron. J.* 108, 39–52. doi: 10.2134/agnonj15.0182
- Fisher, K. A., Momen, B., and Kratochvil, R. J. (2011). Is broadcasting seed an effective winter cover crop planting method? *Agron. J.* 103, 472–478. doi: 10.2134/agnonj2010.0318
- Food and Agriculture Organization (FAO) (2020). *FAOSTAT Crops and Livestock Products*. Available online at: <https://www.fao.org/faostat/en/#data/QCL> (accessed September 1, 2022).
- Gandorfer, M., Pannell, D., and Meyer-Aurich, A. (2011). Analyzing the effects of risk and uncertainty on optimal tillage and nitrogen fertilizer intensity for field crops in Germany. *Agric. Syst.* 104, 615–622. doi: 10.1016/j.agsy.2011.06.004
- Gupta, S. (2022). *Sensitivity Analysis in Excel. One and Two Variable Data Table*. Available online at: <https://www.wallstreetmojo.com/sensitivity-analysis-in-excel/> (accessed September 1, 2022).
- Hamilton, A. V., Mortensen, D. A., and Allen, M. K. (2017). The state of the cover crop nation and how to set realistic future goals for the popular conservation practice. *J. Soil Water Conserv.* 72, 111A–115A. doi: 10.2489/jswc.72.5.111A
- Hardaker, F. R., Lien, G., Anderson, J. R., and Huirne, R. B. (2015). *Coping With Risk in Agriculture: Applied Decision Analysis*. Cabi. doi: 10.1079/9781780645742.0000
- Hatfield, J. L., McMullen, L. D., and Jones, C. S. (2009). Nitrate-nitrogen patterns in the Raccoon River Basin related to agricultural practices. *J. Soil Water Conserv.* 64, 190–199. doi: 10.2489/jswc.64.3.190
- Hijmans, R. J., Choe, H., and Perlman, J. (2016). Spatiotemporal patterns of field crop diversity in the United States, 1870–2012. *Agric. Environ. Lett.* 1, 160022. doi: 10.2134/aehl2016.05.0022

- Hirsh, S. M., Duiker, S. W., Graybill, J., Nichols, K., and Weil, R. R. (2021). Scavenging and recycling deep soil nitrogen using cover crops on mid-Atlantic, USA farms. *Agric. Ecosyst. Environ.* 309, 107274. doi: 10.1016/j.agee.2020.107274
- Howard, R., and Abbas, A. E. (2010). *Foundations of Decision Analysis*. (2010). (Q22)
- Howard, R. A. (1988). Decision analysis: practice and promise. *Manage. Sci.* 34, 679–695. doi: 10.1287/mnsc.34.6.679
- Hubbard, D. W. (2020). *The Failure of Risk Management: Why It's Broken and How to Fix It*. New York: John Wiley and Sons. doi: 10.1002/9781119521914
- Iowa Environmental Mesonet (IEM) (2022). Available online at: <https://mesonet.agron.iastate.edu> (accessed September 1, 2022).
- Iowa State University Extension (2018). *Economics of Cover Crops Worksheets*. Available online at: <https://www.extension.iastate.edu/agdm/crops/html/a1-91.html> (accessed September 1, 2022).
- Iowa State University Extension (2022). *Estimated Costs of Crop Production*. Available online at: <https://www.extension.iastate.edu/agdm/crops/html/a1-20.html> (accessed September 1, 2022).
- Johnson, T. J., Kaspar, T. C., Kohler, K. A., Corak, S. J., and Logsdon, S. D. (1998). Oat and rye overseeded into soybean as fall cover crops in the upper Midwest. *J. Soil Water Conserv.* 53, 276–279.
- Kantar, M., and Porter, P. (2014). Relationship between planting date, growing degree days and the winter rye (*Secale cereale* L.) variety “Rymin” in Minnesota. *Crop Manage.* 13, 1–9. doi: 10.2134/CM-2013-0096-RS
- Kaplan, S., and Garrick, B. J. (1981). On the quantitative definition of risk. *Risk Anal.* 1, 11–27. doi: 10.1111/j.1539-6924.1981.tb01350.x
- Kaspar, T. C., Jaynes, D. B., Parkin, T. B., and Moorman, T. B. (2007). Rye cover crop and gamagrass strip effects on NO<sub>3</sub> concentration and load in tile drainage. *J. Environ. Qual.* 36, 1503–1511. doi: 10.2134/jeq2006.0468
- Kaspar, T. C., Jaynes, D. B., Parkin, T. B., Moorman, T. B., and Singer, J. W. (2012). Effectiveness of oat and rye cover crops in reducing nitrate losses in drainage water. *Agric. Water Manage.* 110, 25–33. doi: 10.1016/j.agwat.2012.03.010
- Kaspar, T. C., Radke, J. K., and Lafen, J. (2001). Small grain cover crops and wheel traffic effects on infiltration, runoff, and erosion. *J. Soil Water Conserv.* 56, 160–164. (Q23)
- Kessler, A., Archontoulis, S. V., and Licht, M. A. (2020). Soybean yield and crop stage response to planting date and cultivar maturity in Iowa, USA. *Agron. J.* 112, 382–394. doi: 10.1002/aj2.20053
- Krupek, F. S., Mizero, S. M., Redfearn, D., and Basche, A. (2022). Assessing how cover crops close the soil health gap in on-farm experiments. *Agric. Environ. Lett.* 7, e20088. doi: 10.1002/acl2.20088
- Kucharik, C. J. (2008). Contribution of planting date trends to increased maize yields in the central United States. *Agron. J.* 100, 328–336. doi: 10.2134/agronj2007.0145
- Lee, D., Arbuckle, J. G., Zhu, Z., and Nowatzke, L. (2018). Conditional causal mediation analysis of factors associated with cover crop adoption in Iowa, USA. *Water Resour. Res.* 54, 9566–9584. doi: 10.1029/2017WR022385
- Marcillo, G. S., Carlson, S., Filbert, M., Kaspar, T., Plastina, A., and Miguez, F. E. (2019). Maize system impacts of cover crop management decisions: a simulation analysis of rye biomass response to planting populations in Iowa, USA. *Agric. Syst.* 176, 102651. doi: 10.1016/j.agsy.2019.102651
- Marcillo, G. S., and Miguez, F. E. (2017). Corn yield response to winter cover crops: an updated meta-analysis. *J. Soil Water Conserv.* 72, 226–239. doi: 10.2489/jswc.72.3.226
- Martinez-Feria, R., Nichols, V., Basso, B., and Archontoulis, S. (2019). Can multi-strategy management stabilize nitrate leaching under increasing rainfall? *Environ. Res. Lett.* 14, 124079. doi: 10.1088/1748-9326/ab5ca8
- Martinez-Feria, R. A., Dietzel, R., Liebman, M., Helmers, M. J., and Archontoulis, S. V. (2016). Rye cover crop effects on maize: a system-level analysis. *Field Crops Res.* 196, 145–159. doi: 10.1016/j.fcr.2016.06.016
- Maryland Department of Agriculture (2022). *Maryland's 2022–2023 Cover Crop Program*. Available online at: [https://mda.maryland.gov/resource\\_conservation/Pages/cover\\_crop.aspx](https://mda.maryland.gov/resource_conservation/Pages/cover_crop.aspx) (accessed September 1, 2022).
- Mirsky, S. B., Curran, W. S., Mortensen, D. A., Ryan, M. R., and Shumway, D. L. (2009). Control of cereal rye with a roller/crimper as influenced by cover crop phenology. *Agron. J.* 101, 1589–1596. doi: 10.2134/agronj2009.0130
- Mitsch, W. J., Day, J. W., Gilliam, J. W., Groffman, P. M., Hey, D. L., Randall, G. W., et al. (2001). Reducing nitrogen loading to the Gulf of Mexico from the Mississippi River Basin: strategies to counter a persistent ecological problem: ecotechnology-the use of natural ecosystems to solve environmental problems-should be a part of efforts to shrink the zone of hypoxia in the Gulf of Mexico. *Bioscience* 51, 373–388. doi: 10.1641/0006-3568(2001)051(0373:RNLTG)2.0.CO;2
- Mohler, C. L., Liebman, M., and Staver, C. P. (2009). *Weed Life History: Identifying Vulnerabilities. Ecological Management of Agricultural Weeds*. Cambridge: Cambridge University Press, 40–98. doi: 10.1017/CBO9780511541810
- Moore, E. B., Wiedenhoef, M. H., Kaspar, T. C., and Cambardella, C. A. (2014). Rye cover crop effects on soil quality in no-till corn silage-soybean cropping systems. *Soil Sci. Soc. Am. J.* 78, 968–976. doi: 10.2136/sssaj2013.09.0401
- Nelson, H., and Bennett, S. (2018). *Cereal Rye Cover Crop for Reducing Herbicides in Soybeans. Practical Farmers of Iowa Cooperators' Program*. Available online at: <https://practicalfarmers.org/research/cereal-rye-cover-crop-for-reducing-herbicides-in-soybeans/> (accessed February 2022).
- Nichols, V., English, L., Carlson, S., Gailans, S., and Liebman, M. (2020a). Effects of long-term cover cropping on weed seedbanks. *Front. Agron.* 2, 591091. doi: 10.3389/fagro.2020.591091
- Nichols, V., Martinez-Feria, R., Weisberger, D., Carlson, S., Basso, B., and Basche, A. (2020b). Cover crops and weed suppression in the US Midwest: a meta-analysis and modeling study. *Agricultural and Environmental Letters* 5, e20022. doi: 10.1002/acl2.20022
- Nichols, V. A., Moore, E. B., Gailans, S., Kaspar, T. C., and Liebman, M. (2022). Site-specific effects of winter cover crops on soil water storage. *Agrosyst. Geosci. Environ.* 5, e20238. doi: 10.1002/agg2.20238
- Pantoja, J. L., Woli, K. P., Sawyer, J. E., and Barker, D. W. (2015). Corn nitrogen fertilization requirement and corn-soybean productivity with a rye cover crop. *Soil Sci. Soc. Am. J.* 79, 1482–1495. doi: 10.2136/sssaj2015.02.0084
- Park, B., Rejesus, R. M., Aglasan, S., Che, Y., Hagen, S. C., and Salas, W. (2022). Payments from agricultural conservation programs and cover crop adoption. *Appl. Econ. Perspect. Policy.* (Q20)
- Patel, S., Sawyer, J. E., and Lundvall, J. P. (2019). Can management practices enhance corn productivity in a rye cover crop system? *Agron. J.* 111, 3161–3171. doi: 10.2134/agronj2019.03.0158 (Q24)
- Plastina, A., Liu, F., Miguez, F., and Carlson, S. (2020). Cover crops use in Midwestern US agriculture: perceived benefits and net returns. *Renew. Agric. Food Syst.* 35, 38–48. doi: 10.1017/S1742170518000194
- Qi, Z., Helmers, M. J., and Lawlor, P. A. (2008). “Effect of different land covers on nitrate-nitrogen leaching and nitrogen uptake in Iowa,” in *2008 Providence, Rhode Island, June 29-July 2, 2008* (p. 1). American Society of Agricultural and Biological Engineers.
- Quinn, D. J., Poffenberger, H. J., Leuthold, S. J., and Lee, C. D. (2021). Corn response to in-furrow fertilizer and fungicide across rye cover crop termination timings. *Agron. J.* 113, 3384–3398. doi: 10.1002/aj2.20723
- Ramirez-Garcia, J., Carrillo, J. M., Ruiz, M., Alonso-Ayuso, M., and Quemada, M. (2015). Multicriteria decision analysis applied to cover crop species and cultivars selection. *Field Crops Res.* 175, 106–115. doi: 10.1016/j.fcr.2015.02.008
- Repo, A. J. (1989). The value of information: approaches in economics, accounting, and management science. *J. Am. Soc. Inform. Sci.* 40, 68–85. doi: 10.1002/(SICI)1097-4571(198903)40:2<68::AID-ASIS2>3.0.CO;2-J
- Roesch-McNally, G. E., Basche, A. D., Arbuckle, J. G., Tyndall, J. C., Miguez, F. E., Bowman, T., et al. (2018). The trouble with cover crops: Farmers' experiences with overcoming barriers to adoption. *Renew. Agric. Food Syst.* 33, 322–333. doi: 10.1017/S1742170517000096 (Q20)
- Rundquist, S., and Carlson, S. (2017). *Mapping Cover Crops on Corn and Soybeans in Illinois, Indiana and Iowa, 2015–2016*. Environmental Working Group. Available online at: <https://www.ewg.org/research/mapping-cover-crops-corn-and-soybeans-illinois-indiana-and-iowa-2015-2016> (accessed September 1, 2022).
- Sarrantonio, M. (1994). *Northeast Cover Crop Handbook*. Kutztown, PA: Soil Health Series. Rodale Institute.
- Seifert, C. A., Azzari, G., and Lobell, D. B. (2018). Satellite detection of cover crops and their effects on crop yield in the Midwestern United States. *Environ. Res. Lett.* 13, 064033. doi: 10.1088/1748-9326/aac4c8
- Shackelford, G. E., Kelsey, R., Sutherland, W. J., Kennedy, C. M., Wood, S. A., Gennet, S., et al. (2019). Evidence synthesis as the basis for decision analysis: a method of selecting the best agricultural practices for multiple ecosystem services. *Front. Sustain. Food Systems* 3, 83. doi: 10.3389/fsufs.2019.00083
- Singer, J. W. (2008). Corn belt assessment of cover crop management and preferences. *Agron. J.* 100, 1670–1672. doi: 10.2134/agronj2008.0151
- Sustainable Agriculture Research and Education (SARE) (2020). *National Cover Crop Surveys*. Available online at: <https://www.sare.org/publications/cover-crops/national-cover-crop-surveys/> (accessed September 2, 2022).
- Syswerda, S. P., Basso, B., Hamilton, S. K., Tausig, J. B., and Robertson, G. P. (2012). Long-term nitrate loss along an agricultural intensity gradient in the Upper Midwest USA. *Agric. Ecosyst. Environ.* 149, 10–19. doi: 10.1016/j.agee.2011.12.007
- Talukder, B., Blay-Palmer, A., Hipel, K. W., and vanLoon, G. W. (2017). Elimination method of multi-criteria decision analysis (MCDA): a simple methodological approach for assessing agricultural sustainability. *Sustainability* 9, 287. doi: 10.3390/su9020287
- Thompson, N. M., Reeling, C. J., Fleckenstein, M. R., Prokopy, L. S., and Armstrong, S. D. (2021). Examining intensity of conservation practice adoption: evidence from cover crop use on US Midwest farms. *Food Policy* 101, 102054. doi: 10.1016/j.foodpol.2021.102054

Urban, D. W., Roberts, M. J., Schlenker, W., and Lobell, D. B. (2015). The effects of extremely wet planting conditions on maize and soybean yields. *Clim. Change* 130, 247–260. doi: 10.1007/s10584-015-1362-x

USDA National Agricultural Statistics Service (USDA NASS) (2018). *National Crop Progress—Terms and Definitions*. Available online at: [https://www.nass.usda.gov/Publications/National\\_Crop\\_Progress/Terms\\_and\\_Definitions/index.php](https://www.nass.usda.gov/Publications/National_Crop_Progress/Terms_and_Definitions/index.php) (accessed September 1, 2022).

USDA National Agricultural Statistics Service (USDA NASS) (2022). *Quick Stats*. Available online at: <https://quickstats.nass.usda.gov> (accessed September 1, 2022).

USDA National Agricultural Statistics Service Cropland Data Layer (USDA NASS CDL) (2021). *Published Corn Data Layer*. Washington, DC: USDA-NASS. Available online at: <https://nassgeodata.gmu.edu/CropScape/> (accessed August 10, 2021).

USDA Risk Management Agency (USDA RMA) (2019). *2020 Cover Crops Insurance and NRCS Cover Crop Termination Guidelines*. Available online at: <https://www.rma.usda.gov/en/News-Room/Frequently-Asked-Questions/2020-Cover-Crops-Insurance-and-NRCS-Cover-Crop-Termination-Guidelines> (accessed September 1, 2022).

Wilson, M. L., Allan, D. L., and Baker, J. M. (2014). Aerially seeding cover crops in the northern US Corn Belt: limitations, future research needs, and alternative practices. *J. Soil Water Conserv.* 69, 67A–72A. doi: 10.2489/jswc.69.3.67A

Wilson, M. L., Baker, J. M., and Allan, D. L. (2013). Factors affecting successful establishment of aerially seeded winter rye. *Agron. J.* 105, 1868–1877. doi: 10.2134/agronj2013.0133

Yoder, L., Houser, M., Bruce, A., Sullivan, A., and Farmer, J. (2021). Are climate risks encouraging cover crop adoption among farmers in the southern Wabash River Basin? *Land Use Policy* 102, 105268. doi: 10.1016/j.landusepol.2020.105268