

Identifying research priorities through decision analysis: a case study for cover crops

1 **Gina A. Nichols^{1,2*}, Cameron MacKenzie³**

2 ¹Department of Plant Sciences, University of California, Davis, CA, USA

3 ²Department of Agronomy, Iowa State University, Ames, IA, USA

4 ³Department of Industrial and Manufacturing Systems Engineering, Iowa State University, Ames, IA,
5 USA

6

7 *** Correspondence:**

8 Corresponding author

9 virginia.nichols@gmail.com

10 **Keywords: decision analysis, cover crop, Iowa, maize, soybean, risk**

11 **Abstract**

12 In Midwestern maize (*Zea-mays* L.)-based systems, planting an over-wintering cover crop such as
13 rye (*Secale cereale* L.) following fall harvests of summer crops maintains continuous soil cover,
14 offering numerous environmental advantages. However, while adoption of cover crops has increased
15 over the past decade, on a landscape-scale it remains low. Identifying where agronomic research
16 could be most impactful in increasing adoption is therefore a useful exercise. Decision analysis (DA)
17 is a tool for clarifying decision trade-offs, quantifying risk, and identifying optimal decisions. Several
18 fields regularly utilize DA frameworks including the military, industrial engineering, business
19 strategy, and economics, but it is not yet widely applied in agriculture. Here we apply DA to a maize-
20 soybean [*Glycine max* (L.) Merr.] rotation using publicly available weather, management, and
21 economic data from central Iowa. In this region, planting a cover crop following maize (preceding
22 soybean) poses less risk to the producer compared to planting following soybean, meaning it may be
23 a more palatable entry point for producers. Furthermore, the risk of reduced maize yields when
24 planting less than 14 days following rye termination substantially contributes to the overall risk cover
25 crops pose to producers, but also has significant potential to be addressed through agronomic
26 research. In addition to identifying research priorities, DA provided clarity to a complex problem,
27 was performed using publicly available data, and by incorporating risk it better estimated true costs
28 to the producer compared to using input costs alone. We believe DA is a valuable and underutilized
29 tool in agronomy and could aid in increasing adoption of cover crops in the Midwest.

30 **1 Introduction**

31 Many cropping systems in the United States (US) have undergone simplifications, now being
32 composed of only a few, often annual, crops (Aguilar et al. 2015; Hijmans et al. 2016; Crossley et al.
33 2021). These systems frequently leave the soil fallow for some period of time, presenting notable
34 environmental challenges including but not limited to increased risk of soil erosion and an increased

potential for nutrient loss (Mitsch et al., 2001; Hatfield et al., 2009, Syswerda et al., 2012). The notion of ‘continuous living cover’ has been used to encourage creative solutions to these issues by focusing cropping system re-design on eliminating these environmentally-challenging fallow periods. Planting cover crops to reduce fallow periods is one such tactic that could at least partially address many of the environmental problems presented by annual cropping systems.

The US produces approximately one-third of the world’s maize (*Zea mays* L.) and soybean [*Glycine max* (L.) Merr.] (FAO, 2022), with five states in the Midwestern region contributing over half of that production (USDA-NASS, 2022). It follows that large amounts of agricultural land in the Midwestern US are dedicated to cropping systems that grow only maize and soybean (Boryan et al., 2011; USDA-NASS Cropland Data Layer, 2021). Utilizing over-wintering cover crops in these systems has been shown to reduce soil erosion and nitrate leaching (Kaspar et al., 2007; Kaspar et al. 2012; La Chen et al., 2022), is associated with a reduction in crop insurance losses due to drought, excess heat, and excess moisture (Aglasan and Rejesus, 2021), and possibly offer numerous other context-specific benefits such as increased soil infiltration rates, higher soil water-holding capacity, or increased soil organic matter content (Moore et al., 2014; Basche et al., 2017; Nichols et al., 2022; Krupek et al., 2022). However, the Midwestern maize-soybean systems present challenges to cover crop adoption. In some regions of the US, cover crop adoption on annual cropland is above 25% and growing (Hamilton et al., 2017). Meanwhile, states comprising the Midwestern US exhibit some of the lowest adoption rates, with most states well below 10% adoption (Hamilton et al., 2017; Rundquist and Carlson, 2017; Seifert et al., 2018).

Low adoption rates within the Midwest have been the subject of numerous studies, and it is clearly a complex issue involving economics, climate constraints, field operations, management, equipment, culture, and technical knowledge (Lee et al., 2018, Church et al., 2020, Nichols et al., 2020a, Thompson et al., 2021; Yoder et al., 2021). One barrier we believe merits more attention is that of risk. Risk incorporates two components, uncertainty and negative consequences, and is frequently measured with probabilities describing the potential severity of consequences (Kapland and Garrick, 1981; Bedford and Cooke, 2001; Hubbard 2020). Cover crops present both direct, and indirect risks. Managerially, maize and soybean are both are planted in the late spring (April, May) and harvested in the fall (September, October, November). Producers typically fit over-wintering cover crops into these systems by planting a cover crop in the fall after the cash crop harvest, and terminating the cover crop in the spring before the next cash crop is planted (SARE 2020). Therefore, both the

planting and termination of an over-wintering cover crop such as rye (*Secale cereale* L.) can conflict with cash crop management. As such, using a cover crop requires complex decision-making that balances risk and rewards in uncertain conditions. While perceived risks associated with cover cropping are often cited as barriers to adoption (Arbuckle and Roesch-McNally, 2015), quantifying those risks in economic terms is challenging (e.g., Bergtold et al., 2019; Plastina et al., 2020). Furthermore, while lists of cover crop research priorities have been proposed (e.g., Carlson and Stockwell, 2013; Basche and Roesch-McNally, 2017), a tool for ranking priorities would be useful. By quantifying the risk associated with each decision point for producers, research priorities can be set to address points posing the highest risk. The use of risk as a ranking tool would also help researchers and funding organizations assess how resources can be used most impactfully. Furthermore, understanding how uncertainties around weather conditions elevate risks of profit loss is important for understanding both the mechanisms for delivering incentives, and the amount producers may require for meaningful participation.

Decision analysis is an interdisciplinary tool that can be applied to analyze decision-making under uncertain conditions (Howard, 1988; Howard and Abbas, 2010; Clemen and Reilly, 2013). It can leverage both quantitative information and expert knowledge, incorporate different degrees of risk aversion, and through sensitivity analyses can allow exploration of the decision space (Cegan et al., 2017; Shackelford et al. 2019). It is a recognized tool for coping with risk in agriculture (Hardaker et al. 2015) and has been applied to a range of agronomic-related topics including agroforestry adoption risks, nitrate pollution loading, cover crop species selection, optimal cropping system choices, and promoting sustainable agricultural practices (Almasri et al., 2005; Gandorfer et al., 2011; Ramirez-Garcia et al., 2015; Talukder et al., 2017; Do et al., 2020). However, to our knowledge decision analysis frameworks have had limited application regarding management decisions related to cover crops in the maize/soybean systems of the Midwestern US. Therefore, the objectives of this study were two-fold:

- 1) Provide a case study using publicly available data to demonstrate the process and utility of applying decision analysis to cover crop systems
- 2) Use a basic analysis to suggest research priorities for cover crops in Central Iowa

We chose to use Central Iowa as a case study because it has large areas in maize/soybean systems that are broadly representative of the US Midwest (USDA NASS CDL, 2021), and currently

demonstrates a moderate amount of cover crop adoption (Rundquist and Carlson, 2017). Furthermore, Iowa's land grant institution, Iowa State University, as well as the United States Department of Agriculture (USDA) National Laboratory for Agriculture and the Environment (NLAE) are located in Central Iowa and support a strong infrastructure for publicly funded agronomic research trials in this region that provide rich sources of public data.

2 Methods and Materials

2.1 Decision set

We used cereal rye (*Secale cereale* L.) as our 'model' over-wintering cover crop because it is the most used cover crop in Iowa and is one of the most widely used cover crop species in the Midwest (Singer 2008). Assuming a producer has both the maize and soybean phase of a maize-soybean rotation growing at a given time, there are two scenarios for cover crop integration, each including three decision alternatives with unique benefits and challenges (**Table 1**). Concomitant benefits and challenges of each decision alternative highlights the need to use a quantitative approach to decision optimization, which can be achieved using decision analysis frameworks.

2.2 Decision structure

The decision set was translated into decision models with known states, uncertainties, and values, each described below.

2.2.1 Fall weather uncertainties

Cover crops are most often planted following cash crop harvests (SARE 2020). Soybean crops in Central Iowa are harvested in September or October, and maize in October or November (USDA NASS 2022). Planting cover crops into standing crops before harvest can increase the probability of establishment (Wilson et al. 2014) but requires specialized equipment that is not yet widely available. We therefore assume cover crop planting occurs after cash crop harvest.

Seeds require precipitation to germinate, and heat units to establish such that the plants emerge and survive the winter. Failure of a cover crop to germinate or establish in the fall results in wasted seed, wasted fuel, and possible weed problems the following spring. While the amount of precipitation needed for rye to germinate depends on soil moisture conditions at planting, crop advisors and producers often assume 1.27 cm (0.5 inches) is needed (Sarrantonio1994), which is consistent with field studies (Fisher et al., 2011; Wilson et al. 2013) and simulation model assumptions (Feyereisen

et al. 2006; Marcillo et al. 2019). While we assumed 1.27 cm was needed for our baseline analysis, this assumption was tested through a sensitivity analysis (see Section 2.3.2).

Growing degree days (GDDs) represent an estimation of the number of heat units accumulated above a threshold temperature specific to a crop. For rye the threshold is 0 or 1 deg Celsius (Feyereisen et al. 2006). We acknowledge the number of GDDs required for rye to successfully over-winter will depend on several additional factors including soil texture and snow cover. A study in Minnesota suggested rye required at least 100 GDDs in the fall to produce biomass in the spring (Kantar and Porter 2014). We therefore estimated rye requires 100 GDDs to successfully establish before winter, but tested the sensitivity of this assumption (see Section 2.3.2).

To estimate the probability of successful rye establishment, we used 30 years of historical weather data (1988-2019) collected at the AMES-8-WSW station from the Iowa Environmental Mesonet (IEM 2022). We chose this dataset because it had previously undergone an extensive quality check (Archontoulis et al. 2020). Using 30 years of weather data, we calculated (i) the probability the site received 1.27 cm of rainfall during an allotted timeframe, and (ii) the probability of achieving 100 GDDs in the allotted timeframe. The timeframes differed by decision alternative to account for the generally earlier harvest dates for soybean compared to maize (USDA NASS 2022). The precipitation timeframes were 15-Oct through 30-Nov and 1-Nov through 30-Nov for rye following soybean and rye following maize, respectively. The GDD accumulation timeframes were 15-Oct through 1-Dec and 1-Nov through 1-Dec for rye following soybean and rye following maize, respectively. We chose to calculate the precipitation and GDD probabilities separately rather than as a joint probability to aid in assessing how breeding efforts could increase changes of establishment. We recognize this model for establishment is a simplification of the complex interactions between weather, soil, and management considerations. While more sophisticated modelling approaches have been utilized for predicting cover crop establishment (Baker et al. 2009, Marcillo et al. 2019; Nichols et al. 2020b), they require specialized skillsets and a significant time commitment. Our goal in this exercise is to demonstrate how insights can be obtained using publicly available data and approachable methodologies.

2.2.2 Spring weather uncertainties

Iowa has a humid continental climate wherein a significant amount of precipitation occurs during the spring months. In addition to the direct constraints on management that precipitation exerts, performing field operations in wet soils can result in undesirable outcomes including long-term soil

compaction and equipment malfunctions. The USDA National Agricultural Statistics Service (NASS) surveys producers to determine the number of days suitable for fieldwork (workable-field day; WFD) for each week throughout the year (USDA NASS 2018). A ‘suitable’ day is defined as one in which weather and field conditions allow producers to work in fields the majority of a given day. Determining whether a day is a ‘suitable’ is subjective, but provides valuable information about the progress and constraints of agricultural production on a landscape level.

Historical data shows that in Iowa, the number of WFDs during the spring can severely restrict field activities (Urban et al 2015; Edwards 2020). To comply with governmental crop insurance cost-share policies, cover crops must be terminated before the cash crop is planted (Bergtold et al. 2019; USDA RMA 2019). Therefore, the presence of a living cover crop that must be terminated before the cash crop can be planted can potentially add to the spring workload for a producer. While this depends on whether producers typically have a pre-plant or pre-emergent herbicide pass, the operation is much less crucial when the goal is simply to eliminate weeds around cash crop planting compared to killing a live cover crop to comply with federal crop insurance requirements. To account for the increased importance of timely cover crop termination, in this exercise we assumed cover crop termination requires an additional set of field working-days compared to systems without a cover crop. However, because many producers do a pre-plant or pre-emergent herbicide pass in systems without cover crops, we did not assume extra herbicide or fuel costs associated with terminating the cover crop. In short, we assumed producers who plant a rye cover crop require two more spring WFDs than those who do not. This fact introduces an important component of risk that is often not accounted for explicitly in economic analyses.

The decision of cover crop termination timing will also affect WFDs, and therefore may indirectly affect cash crop yields. If a producer has WFDs in early April, the producer must choose whether to utilize them to terminate the cover crop, or wait in order to accrue more benefits from prolonged cover crop growth (**Table 1**). Societal-level benefits such as reduced nitrate leaching, as well as farm-level benefits such as the potential to off-set weed control costs, increase as spring cover crop termination dates are delayed and cover crop biomass increases (Finney et al. 2016; Thapa et al. 2018; Nichols et al. 2020b). However, by choosing not to utilize early April WFDs, the producer risks not having sufficient WFDs in late April to terminate the cover crop or plant the cash crop, resulting in delayed cash crop planting and a possible concomitant reduction in yields. Therefore,

186 understanding the uncertainty around WFDs in the spring is an important component in assessing
187 optimal decision alternatives.

188 In this analysis we only include the uncertainties associated with WFDs. In years with very low
189 spring precipitation, delaying cover crop termination can also result in decreased cash crop yields due
190 to the cover crop's use of stored soil water needed for cash crop production. While this risk is
191 possible, due to climatic patterns it is not common in Central Iowa (Daigh et al. 2014; Martinez-Feria
192 et al. 2016). Therefore, the risk of cover crops inducing drought-related yield reductions in the
193 following cash crop is not considered in this exercise.

194 Workable field days are estimated by surveying farmers about how many days in the previous week
195 were field-workable. The data is therefore reported as a number of days within a seven-day calendar
196 period, with this period being inconsistent between years. For the purposes of this exercise, we chose
197 to take the total WFDs over the seven-day reporting period and divide the total by seven to assign a
198 number of WFDs to each calendar day the reporting week included. We then created five spring
199 categories (early April, late April, early May, late May, June). Workable field day values were then
200 summed within these spring calendar categories. More details, including R code, concerning this
201 procedure can be found in supplemental material. We assumed cover crop termination would require
202 two WFDs within a spring category, and cash crop planting would likewise require two WFDs.
203 Therefore, cover crop termination and cash crop planting within a given window would require four
204 WFDs. The probability of two and four WFDs being reported in a given spring category was
205 calculated using 30 years of historical data (1988-2019).

206 **2.2.3 Subsequent maize yield uncertainties**

207 On average, winter cover crops such as rye have been shown to have a neutral effect on subsequent
208 maize and soybean yields (Marcillo and Miguez, 2017). However, numerous studies have shown that
209 under certain conditions, planting maize less than 10-14 days following cover crop termination can
210 result in lower maize yields (Johnson et al. 1998, Acharya et al. 2017, Hirsh et al. 2021, Quinn et al.
211 2022). We assumed a producer would plant their maize crop as early as possible, regardless of the
212 penalty that would be incurred due to the <14 day window. We made this assumption because
213 conversations with producers confirmed that while they were aware there may be a yield penalty
214 from a small termination-planting window, it was inconsistent and may not occur at all, and they
215 were therefore more concerned with timely maize planting. We therefore assumed if there four
216 WFDs in a given spring category, the producer would plant maize but there would be a 50% chance

of a 10% decrease in maize yield. We acknowledge that in our scenarios, the 10% yield penalty from the small termination-planting window is larger than the penalty incurred for delaying planting until late May, but we believe our decision structure captures the uncertainty currently associated with whether that yield penalty will be incurred. Soybeans are not impacted by the time between rye termination and soybean planting (Acharya et al. 2020), so no yield penalty was assigned in those circumstances.

2.2.4 Value

The main contributors to decision value were estimated using partial budgets and included the costs from planting a cover crop, the savings from planting a cover crop, and the income from the subsequent cash crop. Extension publications, farming group publications, and peer-reviewed literature were used to guide each estimation. Sensitivity analyses were performed on assumed values (Section 2.3.2), and instances where conclusions were overly sensitive to assumptions were noted.

To estimate the direct costs associated with planting and terminating a cover crop we used Iowa State University's 'Economics of Cover Crops' decision tool (Iowa State University Extension, 2018). While these prices will fluctuate depending on the price of fuel and labor, we feel they are sufficiently representative for this exercise (**Table 2**).

In order to account for the effect of cover cropping on income from crop yields, we needed to estimate the net revenue a producer expects per unit crop yield. The net revenue from a crop will depend on producer costs of production as well as market prices, both of which vary significantly across years. To overcome this variability, we looked at production costs (Iowa State University Extension, 2022) and market prices (USDA NASS 2022) from 2013-2021, calculated the net revenue per unit crop yield for each year, then took the year with the maximum net revenue for each crop (**Figure 1**). By calculating the net revenue in this manner, when a rye cover crop negatively impacted cash crop yields our analyses represented the highest potential costs of those effects. All prices and calculations are available in supplemental material.

Maize was assumed to have a maximum yield of 10.7 dry Mg ha⁻¹ (200 bu ac⁻¹) and soybean a yield of 1.4 dry Mg ha⁻¹ (60 bu ac⁻¹), which are representative of the state average yields in Iowa (USDA NASS 2022). Maize yield is sensitive to planting date, with later planting dates being associated with lower yields (Kucharik et al. 2008; Baum et al. 2019). We therefore assume a graduated yield penalty increasing 5-20% as maize planting occurs past April (**Table S2**). In summary, the decision of

whether to terminate the cover crop early or late impacts the available WFDs (**Table 3**), which impact whether the producer incurs a termination-planting penalty or a late-planting penalty, both of which impact the value of the decision.

Soybean yields are less sensitive to planting dates compared to maize (Kessler et al. 2020) and therefore was assumed to have a less severe graduated penalty as planting was delayed (5-10%; **Table 2**).

When the cover crop was followed by a maize crop (decision alternatives 1-3), we assumed herbicide costs were equal in the cover crop and no-cover alternatives (\$205 ha⁻¹). When the cover crop was followed by a soybean crop (decision alternatives 4-6), we utilized information from on-farm experiments showing producers reduced herbicide costs due to the mulch provided by a late-terminated cover crop. Therefore, in the decision alternative where the cover crop was terminated in late April or later followed by soybean planting (decision alternative 6), a \$37 ha⁻¹ savings in herbicides was applied (Nelson and Bennett, 2018).

There are currently no payments available to farmers in Iowa for the societal benefits reaped from delaying cover crop termination. However, other areas in the US have implemented payment structures that reward late termination due to the societal benefits gained from late termination (Maryland Department of Agriculture, 2022), so the potential for this payment in decision alternatives 3 and 6 was included in sensitivity analyses.

2.3 Decision analysis

2.3.1 Building decision trees

Decisions can be visualized and modelled using decision tree notation (Howard and Abbas, 2010; Clemen and Reilly, 2013). The full decision model is available in supplemental files and consists of building out a branch for each unique decision node and uncertainty outcome with probabilities, then assigning a value to each branch. We assume a risk-neutral decision maker which means that the decision maker should choose the alternative that maximizes his or her expected value. A square in the decision tree represents a choice between two or more alternatives, and a circle represents an uncertainty where each branch stemming from the uncertainty is assigned a probability. The first decision for the producer is whether or not to plant a cover crop in the fall (**Figure 1**). If the producers choose to plant a cover crop, there is an uncertainty about whether or not sufficient precipitation occurs followed by a second uncertainty about whether or not sufficient GDDs are

accumulated. If sufficient precipitation and sufficient GDDs occur, the producer makes a second decision about whether or not to terminate in early April (**Figure S1**). This decision is followed by uncertainties in the number of WFDs available in a given time frame, and whether there is a penalty when maize is planted in the same spring category as cover crop termination. The decision tree is solved using a “rollback” procedure starting from the right-hand side of the tree. If a decision (square) node is encountered, the alternative with the largest expected monetary value is selected. If an uncertainty (circle) node is encountered, the expected monetary value is calculated using the probabilities on the branches as weights. This procedure results in identifying the alternative for a given decision (e.g., whether or not to plant a cover crop in the fall) that maximizes the producer’s expected monetary value.

2.3.2 Sensitivity analyses

Sensitivity analysis on the uncertainty and parameter assumptions can provide insight into the criticality and importance of an assumption or variable to the decision. The sensitivity of outcomes was assessed for the precipitation required for rye germination (ranging from 0-3.5 cm in 1 mm increments), the number of GDDs needed for rye to over-winter (ranging from 0-300 in 5 GDD increments), the potential relative reduction in maize yields when maize was planted less than 14 days following rye termination (ranging from 0-20% in 5% increments), the incentive payments offered to plant rye (ranging from \$0-200 ha⁻¹ in \$1 increments), and the incentive payments offered to delay termination of rye (ranging from \$0-200 ha⁻¹ in \$1 increments). Additionally, sensitivity analyses were performed on the assumed revenues and costs associated with each scenario to ensure conclusions were not overly sensitive to these assumptions (see supplemental files and Gupta 2022 for details).

2.3.3 Value of information

In our decision model, if a producer has two WFDs within 14 days following cover crop termination, they have a 50% of incurring a 10% maize yield reduction if they choose to plant. This uncertainty is due to research gaps – we do not yet have sufficient information to provide a producer to help them determine whether this reduction will occur. By estimating the value of the decision if the producer knows whether the yield reduction will occur, one can estimate the ‘value of perfect information’ (Repo 1989). This provides an estimate of what that information would be worth to producers, thus allowing researchers to assess how impactful such research would be. We therefore estimated the value of knowing when there would not be a reduction in maize yields when planting <14 days after cover crop termination.

3 Results and Discussion

3.1 Optimal decisions

Assuming there is no cost-share available for planting a cover crop and long-term or societal economic benefits are not accounted for, the overall expected monetary value of not planting a cover crop is greater than the expected monetary value of planting a cover crop, regardless of the sequencing scenario (**Figure 2**). This analysis shows that in addition to the cost of seed and fuel to plant the cover crop (\$50 ha⁻¹), when rye precedes maize there is an additional \$40-70 ha⁻¹ cost associated with the risk that the spring management of the cover crop will result in reduced maize yields (either through delayed maize planting due to insufficient WFDs or <14 day gap penalties). When rye precedes soybeans, the costs of planting the cover crop and risks of reduced yields due to delayed planting are partially compensated by through reduced herbicide costs. Within the decision sets that include the alternative of planting a cover crop, the value of the decision is always maximized if the cover crop is terminated in early versus late April.

Many of the benefits reaped from planting cover crops (e.g., reduced soil erosion, reduced nitrate leaching, non-chemical weed control) are directly related to the amount of biomass the cover crop produces (Finney et al. 2016; Thapa et al. 2018; Nichols et al. 2020b). However, in areas that lack incentives for delaying cover crop termination to allow the cover crop to grow, our analyses show the optimal decision is to terminate the cover crop as soon as possible, even when there might be cost savings from reduced herbicide use (Rye-Soybean scenario in **Figure 2**). Notably, the termination decision differential is highest when the cover crop precedes maize, meaning the sequencing where society may benefit the most (higher mitigation of erosion and nitrate leaching, **Table 1**) would also require the highest incentives to render late April termination the optimal decision. The US state of Maryland has created a tiered incentive system wherein producers are compensated more for early planting and late termination of cover crops (Maryland Department of Agriculture, 2022). Our analysis indicates having compensation rates differ by cropping sequence may also be an approach worth considering.

Our analyses also expose a potential moral hazard. If a producer chooses to plant a cover crop preceding a maize crop and receives a cost-share or incentive for doing so, failed cover crop establishment will lead to a better financial result than successful establishment (Rye-Maize scenario

in **Figure 2**). It is important to provide support for producers as they learn to manage cover crops, and often cover crop establishment is out of a producer's control, but our analyses demonstrate the complexity in determining the best payment structures, and the need to include the risks the cover crop may pose to the subsequent crop yields.

3.2 Sensitivity to cost-share/incentives

If there are no cost-shares or incentive programs, the overall expected monetary value of not planting any cover crop is greater than the expected value of planting a cover crop, regardless of the sequencing scenario (in the top panel of **Figure 3**, this is seen from the 'do not plant rye' alternative having a greater value when the cost share or incentive is \$0 on the horizontal axis). However, current incentive programs may be enough to make planting a cover crop preceding a soybean cash crop ('Rye-Soybean') the optimal decision. If the incentive is greater than \$30 ha⁻¹, the expected monetary value of planting rye prior to soybeans is greater than not planting rye.

When a cover crop precedes a maize crop ('Rye-Maize' in top panel of **Figure 3**), within the current range of incentives the optimal decision is to not plant a cover crop. However, this recommendation is sensitive to the reduction in maize yield due to planting less than 10 days following cover crop termination (bottom panel of **Figure 3**). If the potential reduction in yield were eliminated, the difference between the value of not planting a cover crop and planting a cover crop could be reduced from \$85 ha⁻¹ to \$60 ha⁻¹, bringing the difference into the range of current incentive programs in this area (\$12-74 ha⁻¹).

The exact causes of the reduced yield in maize are not yet clear and it is currently not possible to predict when they will manifest (e.g., Patel et al. 2019; Quinn et al. 2021). The value of perfect information is worth \$20-25 depending on the planned cover crop timing, which is roughly equal to the increased value from eliminating the yield penalty. This indicates that research that allows producers to accurately predict when the yield penalty will occur is equally as valuable as eliminating the yield penalty. Potential mechanisms include altered nutrient dynamics, disease pressure, allelopathy, rye stands that are not fully terminated, changes in soil temperature and/or moisture in a rye cover crop system. A meta-analysis of studies may aid in identifying factors that drive the variation in the effect. Our analyses demonstrate that this phenomenon poses a significant risk to producers, and a better understanding of the drivers and identification of ways to predict when yield declines are likely would greatly reduce the financial risk associated with planting a rye cover crop in these systems.

3.3 Sensitivity to weather

On average, Central Iowa received 7.4 and 4.2 cm of rain from 15-Oct and 1-Nov through 30-Nov, respectively. This equated to a high probability (>80%) of the rye cover crop receiving sufficient precipitation for germination (>1.27 cm) in both sequences (**Figure 4, Table S3**). This result was robust against uncertainty in our assumptions; even if rye required almost double the assumed precipitation, the probability of receiving that amount of rainfall did not drop below 80% for either planting scenario (**Figure 4**). While the probability of accumulating sufficient GDDs (100) was 100% when the rye was planted following soybeans (15-Oct planting date), it dropped to 71% chance of success when planted following maize (1-Nov planting date; **Table S3**). The probability of establishment was very sensitive to the sequencing (rye following soybeans or rye following maize). For the 1-Nov planting date, the results are very sensitive to the assumed GDDs required for establishment.

These results can be used to guide research efforts. Our analysis demonstrates that in most cases, precipitation is not the limiting factor for cover crop establishment in Central Iowa. Breeding varieties that require less precipitation to germinate would likely involve breeding for smaller seeds, which carries inherent tradeoffs (e.g., Carleton and Cooper 1972, Mohler et al. 2009). A study done in Minnesota showed precipitation accounts for the highest amount of variation in rye establishment, followed by temperature (Wilson et al. 2013), demonstrating the value of evaluating weather-related risks locally. While our results do not account for how the precipitation is distributed across time and how that may impact germination, our results suggest this area of Iowa can support larger precipitation requirements for cover crops without experiencing a significant reduction in the probability of cover crop germination.

Our results also show when planting after soybean harvest, the cover crop is almost guaranteed to gain 100 GDDs in the fall (**Figure 4**). Conversely, after maize harvest the probability is very sensitive to how many GDDs are assumed to be needed. Our analyses highlight the need to better understand conditions that lead to successful establishment, particularly in the later months of the year. Additionally, research focused on identifying management tactics that allow for earlier cover crop planting may be most effective in increasing the probability of successful cover crop establishment in Central Iowa. For example, some producers report switching to earlier maturing soybean and maize varieties when adopting cover crops in order to plant the cover crop earlier (Plastina et al., 2020). Some areas have organized blocks of producers who share in aerial seeding

costs, and custom seeding equipment/services that allows for seeding into a standing crop are becoming more common. Our analyses indicate these types of activities are well-suited to reducing the risk associated with planting a cover crop in Central Iowa.

In the spring, the number of WFDs presented a great deal of uncertainty (**Table 3**). Averaged over the entire spring period (1-Apr through 31-May), there was a 79% probability of two or more WFDs in a given two-week period, and only a 46% probability of four or more WFDs. We assumed two or more WFDs were needed to successfully complete a cover crop termination activity, and two additional WFDs were needed to complete cash crop planting activities. Therefore, producers wishing to terminate and plant within a two-week period may not have sufficient WFDs to do so. The probability of two or more WFDs was higher in May compared to April, indicating paying producers to delay cover crop termination may also increase the chances the producer can terminate in their planned timeframe.

Our analyses indicate in Central Iowa, there is generally a high probability the fall conditions will foster cover crop establishment, and that the majority of risk occurs due to the potential for the additional management required in the spring to delay cash crop planting. A Midwestern focus group found some producers had been switching to winterkill cover crop varieties because of the difficulties associated with killing the cover crop and planting a cash crop in a timely manner in the spring (Plastina et al., 2020). For this analysis we assumed the rye cover crop could be terminated at any point, but the stage of rye growth will affect how easy it is to terminate, particularly when using mechanical termination (Creamer et al., 2002; Mirsky et al., 2009). Decision support tools that help producers decide if early termination is the best choice could be beneficial in helping producers manage this risk.

4 Conclusions

Using publicly available data and reasonable assumptions, we were able to gain significant insight into localized priorities for cover crop research. Using historical weather data, NASS surveys on WFDs, extension publications, and a partial budget for cover crop economics we were able to build a single-attribute decision model, and model decision values assuming a risk-neutral producer. Our analysis does not include possible long-term impacts such as the maintenance of productivity, long-term impacts on weeds or insects, or changes in yield stability over time, which could be incorporated in future applications of this framework. We found including only the costs of seed and fuel in cover crop economics underestimates the additional financial risk producers assume due to the

extra spring work cover crops might entail in areas with limited numbers of WFDs during that time. We found there is minimal information on the number of GDDs required for a rye cover crop to successfully overwinter, and that this may have a large impact on risks associated with planting cover crops in Central Iowa. In Central Iowa, identifying ways to ensure early cover crop planting and managements that render maize yields less sensitive to rye cover crop termination timing, or that allow that reduction to be more predictable, could significantly help reduce the financial risk of planting cover crops. Furthermore, flat payments for planting cover crops may result in a moral hazard, wherein the decision value for planting a cover crop preceding a maize crop is maximized when the cover crop fails to establish in the fall. Policies that promote tiered payment structures could rectify this while still providing support for producers as they learn to manage cover crops.

5 Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

6 Author Contributions

GN conceived of and designed the analyses, collected the data, performed the analyses, and wrote the first draft of the manuscript. CM designed the analyses, and edited the manuscript.

7 Funding

This material is based upon work supported in part by the National Science Foundation (Grant no. DGE-1828942).

8 Acknowledgments

We acknowledge Dr. Sarah Ryan for organizing a workshop for the NSF DataFEWSion program that sparked this collaboration, Dr. Raj Raman for his helpful feedback, and two reviewers for their insightful comments and suggestions that greatly improved the quality of this manuscript.

9 References

- Acharya, J., Bakker, M. G., Moorman, T. B., Kaspar, T. C., Lenssen, A. W., & Robertson, A. E. (2017). Time interval between cover crop termination and planting influences corn seedling disease, plant growth, and yield. *Plant disease*, 101(4), 591-600.
- Acharya, J., Moorman, T. B., Kaspar, T. C., Lenssen, A. W., and Robertson, A. E. (2020). Cover crop rotation effects on growth and development, seedling disease, and yield of corn and soybean. *Plant disease*, 104(3), 677-687.
- Aglasan, S., and Rejesus, R. M. (2021). An Analysis of Crop Insurance Losses, Cover Crops, and Weather in US Crop Production. e 2021 Agricultural & Applied Economics Association Annual Meeting, Austin, TX, August 1 – August 3

- 465 Aguilar, J., Gramig, G. G., Hendrickson, J. R., Archer, D. W., Forcella, F. et al. (2015). Crop Species
466 Diversity Changes in the United States: 1978–2012. PLOS ONE 10(8): e0136580. doi:
467 10.1371/JOURNAL.PONE.0136580.
- 468 Almasri, M. N., and Kaluarachchi, J. J. (2005). Multi-criteria decision analysis for the optimal
469 management of nitrate contamination of aquifers. *Journal of Environmental management*. 74(4):365-
470 381.
- 471 Arbuckle, J. G., & Roesch-McNally, G. (2015). Cover crop adoption in Iowa: The role of perceived
472 practice characteristics. *Journal of Soil and Water Conservation*, 70(6), 418-429.
- 473 Archontoulis, S.V., Castellano, M.J., Licht, M.A., Nichols, V., Baum, M., Huber, I., Martinez-Feria,
474 R., Puntel, L., Ordóñez, R.A., Iqbal, J. and Wright, E.E. (2020). Predicting crop yields and soil-plant
475 nitrogen dynamics in the US Corn Belt. *Crop Science*, 60(2), pp.721-738.
- 476 Baker, J.M. and Griffis, T.J. (2009). Evaluating the potential use of winter cover crops in corn–
477 soybean systems for sustainable co-production of food and fuel. *Agricultural and forest meteorology*,
478 149(12), pp.2120-2132.
- 479 Basche, A., & DeLonge, M. (2017). The impact of continuous living cover on soil hydrologic
480 properties: A meta-analysis. *Soil Science Society of America Journal*, 81(5), 1179-1190.
- 481 Basche, A. D., & Roesch-McNally, G. E. (2017). Research topics to scale up cover crop use:
482 Reflections from innovative Iowa farmers. *Journal of Soil and Water Conservation*, 72(3), 59A-63A.
- 483 Baum, M. E., S. V. Archontoulis, and M. A. Licht. (2019). Planting date, hybrid maturity, and
484 weather effects on maize yield and crop stage. *Agronomy Journal* 111(1):303-313.
- 485 Bedford, T., and Cooke, R. (2001). Probabilistic risk analysis: foundations and methods. Cambridge
486 University Press.
- 487 Bergtold, J. S., Ramsey S., Maddy, L., and Williams, J. R. (2019). A review of economic
488 considerations for cover crops as a conservation practice." *Renewable Agriculture and Food Systems*
489 34(1): 62-76.
- 490 Boryan, C., Yang, Z., Mueller, R. and Craig, M. (2011). Monitoring US agriculture: the US
491 Department of Agriculture, National Agricultural Statistics Service, Cropland Data Layer Program.
492 <http://dx.doi.org/10.1080/10106049.2011.562309> 26(5): 341–358. doi:
493 10.1080/10106049.2011.562309.
- 494 Carlson, S., & Stockwell, R. (2013). Research priorities for advancing adoption of cover crops in
495 agriculture-intensive regions. *Journal of Agriculture, Food Systems, and Community Development*,
496 3(4), 125-129.
- 497 Carleton, A. E., and Cooper, C. S. (1972). Seed size effects upon seedling vigor of three forage
498 legumes 1." *Crop Science* 12(2): 183-186. <https://doi.org/10.1002/aepp.13248>
- 499 Cegan, J. C., Fillion, A. M., Keisler, J. M., and Linkov, I. (2017). Trends and applications of multi-
500 criteria decision analysis in environmental sciences: literature review. *Environment Systems and*
501 *Decisions*, 37(2), 123-133.

- Chatterjee, N., Archontoulis, S. V., Bastidas, A., Proctor, C. A., Elmore, R. W., and Basche, A. D. (2020). Simulating winter rye cover crop production under alternative management in a corn-soybean rotation. *Agronomy Journal*, 112(6), 4648-4665.
- Church, S. P., Lu, J., Ranjan, P., Reimer, A. P., & Prokopy, L. S. (2020). The role of systems thinking in cover crop adoption: Implications for conservation communication. *Land use policy*, 94, 104508.
- Clemen, R. T., and Reilly, T. (2013). *Making hard decisions with DecisionTools*. Cengage Learning.
- Creamer, N. G., & Dabney, S. M. (2002). Killing cover crops mechanically: Review of recent literature and assessment of new research results. *American Journal of Alternative Agriculture*, 17(1), 32-40.
- Crossley, M.S., Burke, K. D., Schoville, S. D. and Radeloff, V. C. (2021). Recent collapse of crop belts and declining diversity of US agriculture since 1840. *Global Change Biology* 27(1): 151–164. doi: 10.1111/GCB.15396.
- Daigh, A. L., Helmers, M. J., Kladvko, E., Zhou, X., Goeken, R., Cavdini, J., ... and Sawyer, J. (2014). Soil water during the drought of 2012 as affected by rye cover crops in fields in Iowa and Indiana. *Journal of Soil and Water Conservation*, 69(6), 564-573.
- Da Silva, G. A., Han, G., Kandel, Y. R., Mueller, D. S., Helmers, M., Kaspar, T. C., & Leandro, L. F. (2021). Field Studies on the Effect of Rye Cover Crop on Soybean Root Disease and Productivity. *PhytoFrontiers*, (ja).
- Dickey, E. C., Shelton, D. P., Jasa, P. J., & Peterson, T. R. (1985). Soil erosion from tillage systems used in soybean and corn residues. *Transactions of the ASAE*, 28(4), 1124-1130.
- Do, H., Luedeling, E., & Whitney, C. (2020). Decision analysis of agroforestry options reveals adoption risks for resource-poor farmers. *Agronomy for Sustainable Development*, 40(3), 1-12.
- Edwards, W. (2020). The number of days suitable for fieldwork in Iowa is shrinking. Iowa State University Extension and Outreach. <https://www.extension.iastate.edu/agdm/articles/edwards/EdwMar20.html> [Accessed September 2, 2022]
- Feyereisen, G. W., Wilson, B. N., Sands, G. R., Strock, J. S., & Porter, P. M. (2006). Potential for a rye cover crop to reduce nitrate loss in southwestern Minnesota. *Agronomy Journal*, 98(6), 1416-1426.
- Finney, D. M., White, C. M., & Kaye, J. P. (2016). Biomass production and carbon/nitrogen ratio influence ecosystem services from cover crop mixtures. *Agronomy Journal*, 108(1), 39-52.
- Fisher, K. A., Momen, B., & Kratochvil, R. J. (2011). Is broadcasting seed an effective winter cover crop planting method?. *Agronomy Journal*, 103(2), 472-478.
- Food and Agriculture Organization (FAO). (2020). FAOSTAT Crops and livestock products. <https://www.fao.org/faostat/en/#data/QCL> [Accessed September 1, 2022]

- 538 Gandorfer, M., Pannell, D., and Meyer-Aurich, A. (2011). Analyzing the effects of risk and
539 uncertainty on optimal tillage and nitrogen fertilizer intensity for field crops in Germany.
540 *Agricultural Systems* 104(8): 615-622.
- 541 Guputa, S. (2022). Sensitivity Analysis in Excel | One & Two Variable Data Table.
542 <https://www.wallstreetmojo.com/sensitivity-analysis-in-excel/> [Accessed September 1, 2022]
- 543 Hamilton, A. V., Mortensen, D. A., & Allen, M. K. (2017). The state of the cover crop nation and
544 how to set realistic future goals for the popular conservation practice. *Journal of Soil and Water*
545 *Conservation*, 72(5), 111A-115A.
- 546 Hardaker, J. B., Lien, G., Anderson, J. R., & Huirne, R. B. (2015). *Coping with risk in agriculture:*
547 *Applied decision analysis*. Cabi.
- 548 Hatfield, J. L., McMullen, L. D., & Jones, C. S. (2009). Nitrate-nitrogen patterns in the Raccoon
549 River Basin related to agricultural practices. *Journal of soil and water conservation*, 64(3), 190-199.
- 550 Hijmans, R.J., Choe, H., and Perlman, J. (2016). Spatiotemporal Patterns of Field Crop Diversity in
551 the United States, 1870–2012. *Agricultural & Environmental Letters* 1(1): 160022. doi:
552 10.2134/AEL2016.05.0022.
- 553 Hirsh, S. M., Duiker, S. W., Graybill, J., Nichols, K., & Weil, R. R. (2021). Scavenging and
554 recycling deep soil nitrogen using cover crops on mid-Atlantic, USA farms. *Agriculture, Ecosystems*
555 *& Environment*, 309, 107274.
- 556 Howard, R. A. (1988). Decision analysis: Practice and promise. *Management science*, 34(6), 679-
557 695.
- 558 Howard, R., and Abbas, A. E. (2010). *Foundations of Decision Analysis* (manuscript).
- 559 Hubbard, D. W. (2020). *The failure of risk management: Why it's broken and how to fix it*. John
560 Wiley & Sons.
- 561 Iowa Environmental Mesonet (IEM). (2022). <https://mesonet.agron.iastate.edu> [Accessed September
562 1, 2022]
- 563 Iowa State University Extension. (2018). *Economics of Cover Crops worksheets*.
564 <https://www.extension.iastate.edu/agdm/crops/html/a1-91.html> [Accessed September 1, 2022]
- 565 Iowa State University Extension. (2022). *Estimated Costs of Crop Production*.
566 <https://www.extension.iastate.edu/agdm/crops/html/a1-20.html> [Accessed September 1, 2022]
- 567 Johnson, T. J., Kaspar, T. C., Kohler, K. A., Corak, S. J., and Logsdon, S. D. (1998). Oat and rye
568 overseeded into soybean as fall cover crops in the upper Midwest. *Journal of Soil and Water*
569 *Conservation*, 53(3), 276-279.
- 570 Kantar, M., & Porter, P. (2014). Relationship between planting date, growing degree days and the
571 winter rye (*Secale cereale* L.) variety “Rymin” in Minnesota. *Crop Management*, 13(1), 1-9.

- 572 Kaplan, S., and Garrick, B. J. (1981). On the quantitative definition of risk. *Risk analysis*, 1(1), 11-
573 27.
- 574 Kaspar, T. C., Radke, J. K., & Laflen, J. M. (2001). Small grain cover crops and wheel traffic effects
575 on infiltration, runoff, and erosion. *Journal of Soil and Water Conservation*, 56(2), 160-164.
- 576 Kaspar, T.C., Jaynes, D.B., Parkin, T.B. and Moorman, T.B., (2007). Rye cover crop and gamagrass
577 strip effects on NO₃ concentration and load in tile drainage. *Journal of environmental quality*, 36(5),
578 pp.1503-1511.
- 579 Kaspar, T. C., Jaynes, D. B., Parkin, T. B., Moorman, T. B., and Singer, J. W. (2012). Effectiveness
580 of oat and rye cover crops in reducing nitrate losses in drainage water. *Agricultural Water*
581 *Management*, 110, 25-33.
- 582 Kessler, A., Archontoulis, S. V., & Licht, M. A. (2020). Soybean yield and crop stage response to
583 planting date and cultivar maturity in Iowa, USA. *Agronomy Journal*, 112(1), 382-394.
- 584 Krupek, F. S., Mizero, S. M., Redfearn, D., and Basche, A. (2022). Assessing how cover crops close
585 the soil health gap in on-farm experiments. *Agricultural & Environmental Letters*, 7(2), e20088.
- 586 Kucharik, C. J. (2008). Contribution of planting date trends to increased maize yields in the central
587 United States. *Agronomy Journal*, 100(2), 328-336.
- 588 Chen, L., Rejesus, R. M., Aglasan, S., Hagen, S. C., & Salas, W. (2022). The of cover on soil erosion
589 in the US Midwest. *Journal of Environmental Management*, 324.
- 590 Lee, D., Arbuckle, J. G., Zhu, Z., & Nowatzke, L. (2018). Conditional causal mediation analysis of
591 factors associated with cover crop adoption in Iowa, USA. *Water Resources Research*, 54(11), 9566-
592 9584.
- 593 Marcillo, G. S., & Miguez, F. E. (2017). Corn yield response to winter cover crops: An updated
594 meta-analysis. *Journal of Soil and Water Conservation*, 72(3), 226-239.
- 595 Marcillo, G. S., Carlson, S., Filbert, M., Kaspar, T., Plastina, A., & Miguez, F. E. (2019). Maize
596 system impacts of cover crop management decisions: A simulation analysis of rye biomass response
597 to planting populations in Iowa, USA. *Agricultural Systems*, 176, 102651.
- 598 Martinez-Feria, R. A., Dietzel, R., Liebman, M., Helmers, M. J., & Archontoulis, S. V. (2016). Rye
599 cover crop effects on maize: A system-level analysis. *Field Crops Research*, 196, 145-159.
- 600 Martinez-Feria, R., Nichols, V., Basso, B., & Archontoulis, S. (2019). Can multi-strategy
601 management stabilize nitrate leaching under increasing rainfall?. *Environmental Research Letters*,
602 14(12), 124079.
- 603 Maryland Department of Agriculture (2022). Maryland's 2022-2023 Cover Crop Program.
604 https://mda.maryland.gov/resource_conservation/Pages/cover_crop.aspx [Accessed September 1,
605 2022]

- 606 Mirsky, S. B., Curran, W. S., Mortensen, D. A., Ryan, M. R., & Shumway, D. L. (2009). Control of
607 cereal rye with a roller/crimper as influenced by cover crop phenology. *Agronomy Journal*, 101(6),
608 1589-1596.
- 609 Mitsch, W. J., Day, J. W., Gilliam, J. W., Groffman, P. M., Hey, D. L., Randall, G. W., & Wang, N.
610 (2001). Reducing Nitrogen Loading to the Gulf of Mexico from the Mississippi River Basin:
611 Strategies to Counter a Persistent Ecological Problem: Ecotechnology—the use of natural ecosystems
612 to solve environmental problems—should be a part of efforts to shrink the zone of hypoxia in the
613 Gulf of Mexico. *BioScience*, 51(5), 373-388.
- 614 Mohler, C. L., Liebman, M. and Staver, C. P. (2009). Weed life history: identifying
615 vulnerabilities. *Ecological management of agricultural weeds*: 40-98. Cambridge University Press.
616 <https://doi.org/10.1017/CBO9780511541810>
- 617 Moore, E. B., Wiedenhoef, M. H., Kaspar, T. C., & Cambardella, C. A. (2014). Rye cover crop
618 effects on soil quality in no-till corn silage–soybean cropping systems. *Soil Science Society of
619 America Journal*, 78(3), 968-976.
- 620 Nelson, H. and S. Bennett. (2018). Cereal Rye Cover Crop for Reducing Herbicides in Soybeans.
621 Practical Farmers of Iowa Cooperators' Program. [https://practicalfarmers.org/research/cereal-rye-](https://practicalfarmers.org/research/cereal-rye-cover-crop-for-reducing-herbicides-in-soybeans/)
622 [cover-crop-for-reducing-herbicides-in-soybeans/](https://practicalfarmers.org/research/cereal-rye-cover-crop-for-reducing-herbicides-in-soybeans/) [Accessed February 2022]
- 623 Nichols, V., English, L., Carlson, S., Gailans, S., and Liebman, M. (2020a). Effects of long-term
624 cover cropping on weed seedbanks. *Frontiers in Agronomy*, 2, 591091.
- 625 Nichols, V., Martinez-Feria, R., Weisberger, D., Carlson, S., Basso, B., & Basche, A. (2020b). Cover
626 crops and weed suppression in the US Midwest: A meta-analysis and modeling study. *Agricultural &
627 Environmental Letters*, 5(1), e20022.
- 628 Nichols, V. A., Moore, E. B., Gailans, S., Kaspar, T. C., and Liebman, M. (2022). Site-specific
629 effects of winter cover crops on soil water storage. *Agrosystems, Geosciences & Environment*, 5(1),
630 e20238.
- 631 Pantoja, J. L., Woli, K. P., Sawyer, J. E., & Barker, D. W. (2015). Corn nitrogen fertilization
632 requirement and corn–soybean productivity with a rye cover crop. *Soil Science Society of America
633 Journal*, 79(5), 1482-1495.
- 634 Park, B., Rejesus, R. M., Aglasan, S., Che, Y., Hagen, S. C., & Salas, W. (2022). Payments from
635 agricultural conservation programs and cover crop adoption. *Applied Economic Perspectives and
636 Policy*.
- 637 Patel, S., Sawyer, J. E., & Lundvall, J. P. (2019). Can management practices enhance corn
638 productivity in a rye cover crop system?. *Agronomy Journal*, 111(6), 3161-3171.
- 639 Plastina, A., Liu, F., Miguez, F., & Carlson, S. (2020). Cover crops use in Midwestern US
640 agriculture: perceived benefits and net returns. *Renewable Agriculture and Food Systems*, 35(1), 38-
641 48.

- 642 Qi, Z., Helmers, M. J., & Lawlor, P. A. (2008). Effect of different land covers on nitrate-nitrogen
643 leaching and nitrogen uptake in Iowa. In 2008 Providence, Rhode Island, June 29–July 2, 2008 (p. 1).
644 American Society of Agricultural and Biological Engineers.
- 645 Quinn, D. J., Poffenbarger, H. J., Leuthold, S. J., & Lee, C. D. (2021). Corn response to in-furrow
646 fertilizer and fungicide across rye cover crop termination timings. *Agronomy Journal*, 113(4), 3384-
647 3398.
- 648 Ramírez-García, J., Carrillo, J. M., Ruiz, M., Alonso-Ayuso, M., & Quemada, M. (2015).
649 Multicriteria decision analysis applied to cover crop species and cultivars selection. *Field Crops*
650 *Research*, 175, 106-115.
- 651 Repo, A. J. (1989). The value of information: Approaches in economics, accounting, and
652 management science. *Journal of the American Society for Information Science*, 40(2), 68-85.
- 653 Roesch-McNally, G. E., Basche, A. D., Arbuckle, J. G., Tyndall, J. C., Miguez, F. E., Bowman, T., &
654 Clay, R. (2018). The trouble with cover crops: Farmers' experiences with overcoming barriers to
655 adoption. *Renewable Agriculture and Food Systems*, 33(4), 322-333.
- 656 Rundquist, S. and Carlson, S. (2017). Mapping cover crops on corn and soybeans in Illinois, Indiana
657 and Iowa, 2015-2016. Environmental Working Group. [https://www.ewg.org/research/mapping-](https://www.ewg.org/research/mapping-cover-crops-corn-and-soybeans-illinois-indiana-and-iowa-2015-2016)
658 [cover-crops-corn-and-soybeans-illinois-indiana-and-iowa-2015-2016](https://www.ewg.org/research/mapping-cover-crops-corn-and-soybeans-illinois-indiana-and-iowa-2015-2016) [Accessed September 1, 2022]
- 659 Sustainable Agriculture Research and Education (SARE). (2020). National Cover Crop Surveys.
660 <https://www.sare.org/publications/cover-crops/national-cover-crop-surveys/> [Accessed September 2,
661 2022]
- 662 Sarrantonio, M. (1994). Northeast Cover Crop Handbook. Soil Health Series. Rodale Institute,
663 Kutztown, PA
- 664 Seifert, C. A., Azzari, G., and Lobell, D. B. (2018). Satellite detection of cover crops and their effects
665 on crop yield in the Midwestern United States. *Environmental Research Letters*, 13(6), 064033.
- 666 Shackelford, G. E., Kelsey, R., Sutherland, W. J., Kennedy, C. M., Wood, S. A., Gennet, S., ... &
667 Dicks, L. V. (2019). Evidence synthesis as the basis for decision analysis: a method of selecting the
668 best agricultural practices for multiple ecosystem services. *Frontiers in sustainable food systems*, 83.
- 669 Singer, J. W. (2008). Corn belt assessment of cover crop management and preferences. *Agronomy*
670 *Journal*, 100(6), 1670-1672.
- 671 Syswerda, S. P., Basso, B., Hamilton, S. K., Tausig, J. B., and Robertson, G. P. (2012). Long-term
672 nitrate loss along an agricultural intensity gradient in the Upper Midwest USA. *Agriculture,*
673 *Ecosystems & Environment*, 149, 10-19.
- 674 Thompson, N. M., Reeling, C. J., Fleckenstein, M. R., Prokopy, L. S., and Armstrong, S. D. (2021).
675 Examining intensity of conservation practice adoption: Evidence from cover crop use on US
676 Midwest farms. *Food Policy*, 101, 102054.

- 677 Talukder, B., Blay-Palmer, A., Hipel, K. W., and vanLoon, G. W. (2017). Elimination method of
 678 multi-criteria decision analysis (MCDA): A simple methodological approach for assessing
 679 agricultural sustainability. *Sustainability* 9:287. doi:10.3390/su9020287
- 680 Urban, D. W., Roberts, M. J., Schlenker, W., and Lobell, D. B. (2015). The effects of extremely wet
 681 planting conditions on maize and soybean yields. *Climatic Change* 130(2):247-260.
- 682 USDA National Agricultural Statistics Service (USDA NASS). (2018). National Crop Progress –
 683 Terms and Definitions.
 684 https://www.nass.usda.gov/Publications/National_Crop_Progress/Terms_and_Definitions/index.php
 685 [Accessed September 1, 2022]
- 686 USDA National Agricultural Statistics Service (USDA NASS). (2022). Quick Stats.
 687 <https://quickstats.nass.usda.gov> [Accessed September 1, 2022]
- 688 USDA National Agricultural Statistics Service Cropland Data Layer (USDA NASS CDL). (2021).
 689 Published corn data layer [Online]. USDA-NASS, Washington, DC.
 690 <https://nassgeodata.gmu.edu/CropScape/> [Accessed August 10, 2021]
- 691 USDA Risk Management Agency (USDA RMA). (2019). 2020 Cover Crops Insurance and NRCS
 692 Cover Crop Termination Guidelines. [https://www.rma.usda.gov/en/News-Room/Frequently-Asked-](https://www.rma.usda.gov/en/News-Room/Frequently-Asked-Questions/2020-Cover-Crops-Insurance-and-NRCS-Cover-Crop-Termination-Guidelines)
 693 [Questions/2020-Cover-Crops-Insurance-and-NRCS-Cover-Crop-Termination-Guidelines](https://www.rma.usda.gov/en/News-Room/Frequently-Asked-Questions/2020-Cover-Crops-Insurance-and-NRCS-Cover-Crop-Termination-Guidelines) [Accessed
 694 September 1, 2022]
- 695 Wilson, M. L., Baker, J. M., and Allan, D. L. (2013). Factors affecting successful establishment of
 696 aerially seeded winter rye. *Agronomy Journal*, 105(6), 1868-1877.
- 697 Wilson, M. L., Allan, D. L. and Baker, J. M. (2014). Aerially seeding cover crops in the northern US
 698 Corn Belt: Limitations, future research needs, and alternative practices. *Journal of Soil and Water*
 699 *Conservation* 69(3): 67A-72A.
- 700 Yoder, L., Houser, M., Bruce, A., Sullivan, A., and Farmer, J. (2021). Are climate risks encouraging
 701 cover crop adoption among farmers in the southern Wabash River Basin?. *Land Use Policy*, 102,
 702 105268.

703

704 1 Data Availability Statement

705 The datasets and R code generated for this study can be found in the Github repository
 706 (https://github.com/vanichols/Nichols_Frontiers_CoverCropRisk).

707 2 Figure legends and tables

708 **Figure 1** Decision tree visualization for planting a cover crop following a soybean crop. The first
 709 decision (light blue square) is whether or not to plant a cover crop. If the producer chooses to plant a
 710 cover crop, there is uncertainty about precipitation and growing degree days (GDDs); if the cover
 711 crop is successfully established the producer will have to decide whether to terminate the cover crop
 712 in early April, or to wait until late-April. Each decision branch has a monetary value.

Figure 2. Value of each decision alternative (rounded to the nearest \$10) assuming no incentive/cost share payments for planting a cover crop.

Figure 3. (Top) Planting a rye cover crop (green line) required \$85 ha⁻¹ and \$30 cost shares/incentives, respectively for a soybean-rye-maize and maize-rye-soybean scenario to make decision values equal to not planting a cover crop (gold line); gray box represents range of current incentive values. (Bottom) Current estimates show maize yields can be reduced by approximately 10% when maize is planted <14 days after terminating a cover crop; if agronomic research efforts were able to eliminate this yield reduction the difference in decision values would be within the range of current incentive programs.

Figure 4 Sensitivity of outcomes to assumption of fall weather for rye planted following maize (dark blue; 1-Nov planting date) and following soybeans (pink; 15-Oct planting date). (Top) The probability of receiving sufficient precipitation is not sensitive to sequencing nor the assumed amount required for germination. (Bottom) The probability of accumulating sufficient GDDs for rye establishment is very sensitive to both the sequencing and the number of GDDs required

Table 1 Two scenarios each including three decision alternatives related to cover cropping in a maize/soybean rotation with various benefits and challenges associated with each alternative.

Decision alternative	Description	Benefits	Challenges
<i>In fields with a soybean crop</i>			
1	Do not plant a cover crop following soybean harvest	No added costs or risks due to cover crop	Low residue from soybean crop leaves soil vulnerable to erosion (Dickey et al, 1985) Soil nitrogen is likely to be lost from the field in the spring to leaching (Qi et al., 2008) Low residue contributes minimally to non-chemical weed control
2	Plant a cover crop, plan to terminate early April	Soybeans are harvested earlier in the fall compared to maize (USDA NASS 2022), allowing for earlier cover crop planting which increases likelihood of successful establishment and more cover crop growth	Cover crop may indirectly reduce subsequent maize yields by competing for workable field days and delaying maize planting, which often results in lower maize yields (Baum et al. 2019)

		(Chatterjee et al. 2020; Nichols et al. 2020b)	Planting maize less than two weeks following cover crop termination may result in reduced yields, but the effect is unpredictable (Johnson et al. 1998; Archarya et al. 2017; Acharya et al. 2020)
		Cover crop residue reduces soil erosion following soybeans (Kaspar et al. 2001)	
		Cover crop residue may provide weed control following soybeans (Nelson and Bennett 2018)	
		Cover crop growth can uptake soil nitrate thus mitigating nutrient pollution (Qi et al., 2008; Kaspar et al. 2012; Martinez-Feria et al. 2019)	
3	Plant a cover crop, plan to terminate late April	Enhances cover crop benefits due to more cover crop growth and biomass	Increases chances of delayed maize planting, and thus reduced maize yields
<i>In fields with a maize crop</i>			
4	Do not plant a cover crop following maize harvest	No added costs or risks due to cover crop	Soil nitrogen is likely to be lost from the field in the spring to leaching (Qi et al. 2008)
5	Plant a cover crop, plan to terminate early April	Maize can leave large nitrate reserves in the soil at harvest, and cover crop growth can uptake the nitrate thus mitigating nutrient pollution (Qi et al., 2008; Kaspar et al. 2012; Martinez-Feria et al. 2019)	Timely fall cover crop planting can be difficult following maize harvest
		Soybean planting dates are less sensitive to planting dates compared to maize (Kessler et al. 2020)	Maize is harvested in late fall, and late-planted cover crops can result in low spring cover crop biomass (Chatterjee et al. 2020; Nichols et al. 2020b), and therefore minimal benefits, if terminated in early April
6	Plant a cover crop, plan to terminate late April	Enhances cover crop benefits due to more cover crop growth and biomass	Larger amounts of cover crop biomass may be more difficult to terminate uniformly

730 **Table 2.** Summary of economic assumptions for each scenario, the assumed relative cash crop yield
 731 indicated in parentheses

No cover crop system		Cover crop system	
		14+ day gap	<14 day gap ^{1,2}
Cover crop			
Cover crop seed	-	\$20 ha ⁻¹	\$20 ha ⁻¹
Cover crop planting	-	\$32 ha ⁻¹	\$32 ha ⁻¹
Cost-shares/insurance	-	\$12-74 ha ⁻¹	\$12-74 ha ⁻¹
Cover crop preceding maize			
Herbicide costs	\$205 ha ⁻¹	\$205 ha ⁻¹	\$205 ha ⁻¹
Maize income (assumed \$2.14 net income per bushel)			
Planted early April	\$1057 ha ⁻¹	-	\$1057 ha ⁻¹ / \$951 ha ⁻¹ (90%)
Planted late April	\$1057 ha ⁻¹	\$1057 ha ⁻¹	\$1057 ha ⁻¹ / \$951 ha ⁻¹ (90%)
Planted early May ²	\$1004 ha ⁻¹ (95%)	\$1004 ha ⁻¹ (95%)	\$1004 ha ⁻¹ (95%)/ \$889 ha ⁻¹ (85%)
Planted late May	\$951 ha ⁻¹ (90%)	\$951 ha ⁻¹ (90%)	\$951 ha ⁻¹ (90%)/ \$846 ha ⁻¹ (80%)
Planted June	\$846 ha ⁻¹ (80%)	\$846 ha ⁻¹ (80%)	\$846 ha ⁻¹ (80%)/ \$740 ha ⁻¹ (70%)
Cover crop preceding soybean			
Herbicide costs ³	\$205 ha ⁻¹	\$168 ha ⁻¹	\$168 ha ⁻¹
Soybean income (assumed \$4.06 net income per bushel)			
Planted early April	-	-	-
Planted late April	\$601 ha ⁻¹	\$601 ha ⁻¹	\$601 ha ⁻¹
Planted early May	\$601 ha ⁻¹	\$601 ha ⁻¹	\$601 ha ⁻¹
Planted late May ⁴	\$571 ha ⁻¹ (95%)	\$571 ha ⁻¹ (95%)	\$571 ha ⁻¹ (95%)
Planted June	\$541 ha ⁻¹ (90%)	\$541 ha ⁻¹ (90%)	\$541 ha ⁻¹ (90%)

¹The decision model for rye following soybean includes a 50% chance a <14 day maize yield reduction will not occur (first values listed), and 50% chance the <14 day maize yield reduction will occur (second values listed)

²Estimated maize yield reduction due to termination-planting gap are based on Johnson et al., 1998; Hirsh et al., 2021; Quinn et al., 2022

²Estimated maize yield reduction due to delayed maize planting are based on Kucharik et al., 2008 and Baum et al., 2019

Table 3. Summary of probabilities of workable field days (WFDs) in a given timeframe based on 30 years of NASS survey data (USDA NASS 2022)

<i>Management Window</i>	<i>Probability of two or more workable field days (WFDs)</i>	<i>Probability of four or more WFDs</i>
1-Apr through 15-Apr (early April)	69%	48%
16-Apr through 30-Apr (late April)	71%	37%
1-May through 15-May (early May)	89%	45%
16-May through 31-May (late May)	87%	55%