

Dicamba-Tolerant Soybean Combined Cover Crop to Control Palmer amaranth

Author(s): Garret B. Montgomery, Angela T. McClure, Robert M. Hayes, Forbes R. Walker, Scott A. Senseman and Lawrence E. Steckel

Source: Weed Technology, 32(2):109-115.

Published By: Weed Science Society of America

URL: <http://www.bioone.org/doi/full/10.1017/wet.2017.96>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

Weed Management-Major Crops

Cite this article: Montgomery GB, McClure AT, Hayes RM, Walker FR, Senseman SA and Steckel LE (2018) Dicamba-Tolerant Soybean Combined Cover Crop to Control Palmer amaranth. *Weed Technol* 32:109–115. doi: 10.1017/wet.2017.96

Received: 29 March 2017
Accepted: 22 September 2017

Associate Editor:

Kevin Bradley, University of Missouri

Nomenclature:

Glyphosate; dicamba; fomesafen; Palmer amaranth, *Amaranthus palmeri* (S.) Wats.; hairy vetch, *Vicia villosa* Roth.; wheat, *Triticum aestivum* L.; soybean, *Glycine max* (L.) Merr.

Key words:

Cover crops; cultural weed control; resistance management

Author for correspondence:

Lawrence E. Steckel, Department of Plant Sciences, University of Tennessee, Jackson, TN 38301. (E-mail: lsteckel@utk.edu)

Dicamba-Tolerant Soybean Combined Cover Crop to Control Palmer amaranth

Garret B. Montgomery¹, Angela T. McClure², Robert M. Hayes³,
Forbes R. Walker⁴, Scott A. Senseman⁵ and Lawrence E. Steckel³

¹Graduate Research Assistant, Department of Plant Sciences, University of Tennessee, Jackson, TN, USA, ²Associate Professor, Department of Plant Sciences, University of Tennessee, Jackson, TN, USA, ³Professor, Department of Plant Sciences, University of Tennessee, Jackson, TN, USA, ⁴Associate Professor, Department of Biosystems Engineering and Soil Science, 2506 E.J. Chapman Dr., Knoxville, TN, USA and ⁵Professor and Head of Department of Plant Sciences, University of Tennessee, Knoxville, TN, USA

Abstract

A study was conducted to evaluate the response of glyphosate- and dicamba-tolerant (GDT) soybean and weed control from cover crop different termination intervals before and after soybean planting. Cover crop biomass was highest when terminated at planting, decreased with the 7- and 14-d preplant (DPP) and day-after-planting (DAP) timings, and again at the 14 DPP and DAP timings. Glyphosate + dicamba provided total control of cover crops by 21 DAP. Cover crop termination timing did not influence soybean population or yield. Palmer amaranth control at the 21 and 28 d after termination (DAT) was 97% to 99%. Differences in Palmer amaranth control were not detected among herbicide programs or termination intervals at the end of season rating, and all treatments provided $\geq 97\%$ control. Although differences in Palmer amaranth control were not apparent at the end of the season, the delay in cover crop affected the number of days until 10-cm Palmer amaranth was present. When utilizing a wheat + hairy vetch cover crop in DGT soybeans, producers should delay cover crop termination until 11 to 14 DPP and make at least one POST application of glyphosate + dicamba + an additional herbicide mode of action (MOA) to maximize Palmer amaranth control and soybean yields.

Introduction

Introduction of glyphosate-resistant (GR) soybean (1996), cotton (1997), and corn (1998) into the US marketplace (Roundup Ready[®], Monsanto, St. Louis, MO) soon drastically changed the approach to weed control (Burke et al. 2005; Corbett et al. 2004; Culpepper 2006; Norsworthy et al. 2012). Over-reliance on glyphosate caused a shift in the overall weed spectrum through extreme selection pressure, and GR biotypes of key weed species, such as horseweed [*Conyza canadensis* (L.) Cronq.] and Palmer amaranth, have become common in the major agronomic areas of the United States (Culpepper 2006; Norsworthy et al. 2008; VanGessel 2001). Palmer amaranth was first confirmed GR in Georgia in 2004 and is currently documented as GR in most of the major US agronomic states (Culpepper 2006; Heap 2016). Since that time, Palmer amaranth has become one of the most economically damaging weeds in the United States and dominates in-season weed management decisions where present (Price et al. 2011; Webster and Sosnoskie 2010). The aggressive growth rate, stature, and inherent survival abilities of this weed make it extremely competitive with agronomic crops (Klingaman and Oliver 1994; Bond and Oliver 2006; Culpepper et al. 2006). For example, Klingaman and Oliver (1994) reported soybean yield percentage reductions at the following Palmer amaranth densities (in plants m⁻¹): 0.33 (17%), 0.66 (27%), 1 (32%), 2 (48%), 3.33 (64%), and 10 (68%).

Growers planted approximately 4.4 million ha of soybeans in Tennessee in 2016, making it an important crop in the state (Anonymous 2012). In 2014, 94% of the soybean hectarage in the United States was planted with herbicide-resistant soybean cultivars (USDA NASS 2014). The majority of soybeans sown in Tennessee between the late 1990s and 2015 were glyphosate-tolerant (L. Steckel, personal communication). In the past, soybean producers in Tennessee and the Mid-South have heavily relied on protoporphyrinogen IX oxidase (PPO) herbicides for control of GR Palmer amaranth (Miller and Norsworthy 2016). However, the confirmation and spread of PPO-resistant Palmer amaranth resulted in growers shifting to more glufosinate-resistant soybean 2016 (Heap 2016; L. Steckel, personal communication). The loss of PPO herbicides eliminated effective POST herbicide options for controlling Palmer amaranth in GR soybean. In response to the increased incidence of weed resistance to glyphosate and other herbicides, seed companies have developed soybean cultivars with resistance to multiple herbicides such as glyphosate + dicamba (GDT).

Dicamba has been widely used for over 40 yr and is an effective herbicide for the control of most broadleaf weed species (Mueller et al. 2013; Shaner 2014). Dicamba is an auxin-mimicking herbicide that controls GR Palmer amaranth and other broadleaf weeds alone or in combination with other herbicides (Cahoon et al. 2015; Merchant et al. 2013; Samples et al. 2013; York et al. 2012, 2015). Inman et al. (2016) reported that glyphosate plus dicamba significantly decreased the frequency and total Palmer amaranth density when compared to glyphosate alone. Crow et al. (2016) reported that dicamba + diflufenzopyr provided similar to or greater control of large (>20 cm) Palmer amaranth plants when compared to other single-herbicide mode-of-action (MOA) treatments in corn.

Additionally, producers have begun utilizing other management practices such as cover crops to combat multiple herbicide-resistant weed species. No-till crop production is prevalent in Tennessee, where 71% of the corn (*Zea mays* L.), cotton (*Gossypium hirsutum* L.), soybean, and wheat hectares were produced in a no-till environment in 2014 (Kenerson 2014). No-tillage methods limit the cultural control methods available to producers for combating difficult herbicide-resistant weeds, such as Palmer amaranth (Price et al. 2011). However, cover crops can be implemented into no-till systems to increase the sustainability of weed control programs (Mirsky et al. 2013; Wiggins et al. 2016a, b). Cereal rye (*Secale cereale* L.) and winter wheat are two common winter annual grass species used for cover cropping systems in the southeastern United States (Norsworthy et al. 2011; Wiggins et al. 2016b). Winter wheat is an appealing cover crop for many producers, because it is economical; moreover, many producers already have experience growing it as a cash crop. However, winter wheat should be managed differently if it is being grown as a cover crop (SARE 2007). Crimson clover (*Trifolium incarnatum* L.) and hairy vetch are two winter-annual legume species that have been researched extensively as cover crops (Norsworthy et al. 2010; Reddy 2001). Annual legumes have been shown to reduce pressure on some winter- and summer-annual weeds similarly to winter-annual grass species (Fisk et al. 2001; Isik et al. 2009). However, utilizing a cereal + legume cover crop combination can create a synergistic effect increasing cover crop biomass (Wiggins et al. 2016a; Zotarelli et al. 2009). The quantity of biomass present at a given time by a cover crop directly contributes to the amount of weed suppression that can be achieved (Mirsky et al. 2013; Norsworthy et al. 2011; Ryan et al. 2011a; Wiggins et al. 2016b).

Cover crop termination is one of the most influential factors in determining the amount of weed suppression (Mirsky et al. 2013). Delaying cover crop termination until at or near planting of the cash crop allows the cover crop a longer growing season, thus producing more biomass (Ashford and Reeves 2003; Mirsky et al. 2009, 2011; Wortman et al. 2012). Although chemical termination of cereal cover crop species such as cereal rye or wheat can easily be accomplished with glyphosate (Ashford and Reeves 2003; Price et al. 2009), termination of legume cover crops with herbicides labeled for at or near cash crop planting can be more challenging (Davis 2010; Fisk et al. 2001; Wiggins et al. 2016a, b). However, auxin herbicides such as 2,4-D or dicamba have proven to be effective herbicides to control legume cover crop species (Curran et al. 2015; McCurdy et al. 2013). White and Worsham (1990) reported that dicamba provided 97% control of hairy vetch prior to planting corn in North Carolina.

Cereal + legume cover crop mixtures are becoming more common in areas where producers are utilizing cover crops to

combat difficult to control weed species. The advent of GDT soybean technologies could provide producers with an effective herbicide option for terminating such cover crop mixtures near soybean planting. The ability to delay termination of cover crop mixtures used in combination with new herbicide-tolerant crop technologies could provide producers an effective means to increase sustainability of weed control systems for soybeans in no-till environments. Therefore, we conducted research to evaluate glyphosate + dicamba for terminating a wheat + hairy vetch cover crop at different timings before and after planting GDT soybean in Tennessee.

Materials and Methods

Field studies to evaluate wheat + hairy vetch termination and in-crop weed control with fomesafen and dicamba in GDT soybean systems were conducted in the growing seasons of 2015 and 2016 at the West Tennessee Research and Education Center, in Jackson, TN (35.633° N, 88.856° W). The experimental site was planted to soybeans in each of the previous site years, and both cover crops and soybeans were sown into long-term no-till environments common to western Tennessee.

Wheat and hairy vetch cover crops (seeded at 67 and 22 kg ha⁻¹, respectively) were drilled in September and October of 2014 and 2015, respectively, using a no-till drill, and allowed to over-winter (Table 1). An experimental, proprietary, late-four maturity group GDT soybean variety (Monsanto Co., St. Louis, MO) was sown on May 11, 2015 and May 16, 2016. Soybeans were planted 3 cm deep at a population of 346,000 seed ha⁻¹ into the existing cover crop residue with a no-tillage planter. Individual plots consisted of two 76-cm rows that were 9.1 m long. Treatments were replicated four times in a split-plot design within a randomized complete block. The whole plot was subjected to termination timing and consisted of 14 d prior to planting (DPP), 7 DPP, at planting, 7 d after planting (DAP), and 14 DAP. The subplot was POST herbicide and consisted of a premix of glyphosate + fomesafen (1,120 and 280 g ae ha⁻¹, respectively) (Flexstar® GT 3.5 herbicide label, Syngenta Crop Protection, LLC, Greensboro, NC) or a proprietary premix of glyphosate + the diglycolamine salt of dicamba (1,120 and 560 g ae ha⁻¹, respectively) applied when Palmer amaranth for that termination interval reached 10 cm in height. The research site was infested with nearly 100% GR Palmer amaranth (L. Steckel, unpublished data), so these treatments are referred to as fomesafen and dicamba, respectively. Termination dates and cover crop growth stage at each termination timing are presented in Table 1 for wheat and hairy vetch (Zadok et al. 1974). POST herbicide treatments were applied using a CO₂-pressurized backpack sprayer calibrated to deliver 140 L ha⁻¹ equipped with TTI110025 nozzles (TTI Turbo TeeJet Spray Tips, TeeJet Technologies, Wheaton, IL), in accordance with currently proposed nozzle requirements for using dicamba on GDT soybeans (Anonymous 2017).

Cover crop biomass was collected from 0.5-m² quadrants at each termination timing, and samples were then dried in a forced-air oven at 60 C for 48 hr. Soybean stand was counted and the count averaged over two randomly selected sections of 0.5 m per row in each plot; this count was then converted to plants ha⁻¹. Cover crop control was visually estimated on a scale of 0 to 100%, where 0 represented no injury or control and 100 represented complete plant death at 7, 14, and 21 DAP. Control ratings for cover crops did not begin until after planting, because the authors'

Table 1. Wheat and hairy vetch growth stages at 14, 7, and 0 d before and 7 and 14 d after soybean planting termination intervals at the West Tennessee Research and Education Center in Jackson, TN, in 2015 and 2016.

Termination interval ^a	Wheat ^b		Hairy vetch		Application dates		Days to POST ^c	
	2015	2016	2015	2016	2015	2016	2015	2016
14 DPP	59	59	Early pod	Mid pod	Apr 27	May 2	30	29
7 DPP	61	61	Early pod	Mid pod	May 4	May 10	30	29
At planting	64	64	Mid pod	Late pod	May 11	May 16	37	32
7 DAP	64	64	Mid pod	Late pod	May 19	May 24	37	35
14 DAP	64	64	Mid pod	late pod	May 27	May 31	39	39

^aAbbreviations: DPP, days prior to planting; DAP, days after planting.

^bAccording to Zadok's growth staging.

^cDays to POST application is recorded as days after soybean planting until a POST herbicide application was triggered when Palmer amaranth in that termination interval had reached a height of 10 cm.

previous experiences indicated that the damage induced on the cover crop from the no-till planter for the cash crop increases cover crop control. Palmer amaranth emergence and growth was monitored until it reached a height of 10 cm in a particular termination interval, and then treated with a POST herbicide. The 10 cm in height termination timing was selected, as that is the height directed by dicamba labels (Anonymous 2017). The experimental area in each site year had been a weed control research site for >10 previous years and was known to have synonymous Palmer amaranth pressure with adjacent areas. Additionally, in each site year, Palmer amaranth was noted to be emerging in these adjacent areas before the initial 14 DPP treatments were applied. Therefore, delays in Palmer amaranth emergence and growth thereafter could be attributed to the cover crop plus the herbicide applied for termination. All plots for a particular termination interval were treated at the same time. Cover crop termination interval was regressed against the number of days after soybean planting for Palmer amaranth to reach a height of 10 cm (Equation 1). The logistic model was fit using SigmaPlot (ver. 8.02; SPSS, Inc., Chicago, IL) to determine the correlation of termination interval and days until Palmer amaranth triggered a POST application.

$$y = y_0 + \frac{a}{1} \exp\left(-\frac{x - x_0}{b}\right) \quad [1]$$

In this model, y_0 is minimum number of days for Palmer amaranth to reach 10 cm in height, b is the slope around the inflection point, x_0 half the number of days for Palmer amaranth to reach 10 cm, and a is the inflection point, or days before or after planting required to maximize the number of days for Palmer amaranth to reach 10 cm in height. Palmer amaranth control was visually assessed 7, 14, 21, and 28 d after the POST application was made (DAT). An additional evaluation was made when soybeans reached the R6 maturity stage as an end-of-season weed control rating. Days until the POST application were also recorded as DAP (Table 1). Soybeans were harvested in each year of this experiment with a small-plot combine, and yields were adjusted to a moisture content of 13%.

All data were subjected to an ANOVA using PROC GLIMMIX in SAS (ver. 9.4; SAS Institute, Cary, NC). Random effects were years, replications, and replications nested within years (Blouin et al. 2011; Carmer et al. 1989). Cover crop biomass, control, and soybean stands were analyzed using only termination intervals as the fixed effect, as these data were collected prior to the

application of any POST treatments. Palmer amaranth control and soybean yields were analyzed using the fixed effects of termination interval, POST, and the interaction among the fixed effects. The square roots of visual estimates for cover crop and weed control were arcsine transformed, and soybean population, cover crop biomass, and yield data were \log_{10} transformed. The transformations did not improve homogeneity of variance for any data point based on visual inspection of plotted residuals; therefore, nontransformed data were used in analyses. Type III statistics were used to test all fixed effects or interactions between the fixed effects. Least square means were calculated based on $\alpha=0.05$ and utilized for mean separation. The DANDA.sas design and analysis macro collection (Saxton 2013) was utilized to build all PROC GLIMMIX (MMAOV) procedures, examine normality, perform data transformations, and convert mean separation to letter groupings when appropriate.

Results and Discussion

A significant effect of termination interval was detected for cover crop biomass (Tables 2 and 3). Biomass was highest when cover crops were terminated at planting, decreased with the 7 DPP and DAP timings, and again at the 14 DPP and DAP timings. Decreased biomass at the 7- and 14-DAP timings coincided with the authors' previous experience of a no-till planter providing an effect that was similar to, but less efficacious than that of other mechanical termination methods such as a roller crimper, especially in cereal cover crops planted in a mixture with hairy vetch. The cereal + hairy vetch mixtures generally become entangled and are pressed to the ground with the planter. This effect is further explained with the cover crop control ratings. For both wheat and hairy vetch control, there was a significant main effect of termination timing 7 and 14 DAP (Tables 2 and 3). All treatments applied prior to planting provided 99% control of wheat and hairy vetch 7 DAP. Wheat or hairy vetch ranged from 85% to 87% in the 7- and 14-DAP termination timings, immediately prior to the 7-DAP termination treatment being applied. Similarly, 14-DAP control of wheat and hairy vetch was 91% and 90%, respectively, immediately prior to the 14-DAP termination treatment being applied. Data from the 21-DAP rating is not presented, as all treatments provided total control of each cover crop species. Similarly, Curran et al. (2015) reported that dicamba applied at 140 g ai ha⁻¹ provided ≥90% control of hairy vetch whether applied in the fall or spring, despite using a much lower rate of

Table 2. Main effects of termination interval, POST herbicide application, and the interaction of the main effects on wheat + hairy vetch cover crop biomass and control and soybean stand and yield.

Effect ^a	Biomass	Cover crop control ^b				Soybean stand	Yield
		Wheat		Vetch			
		7	14	7	14		
		<i>p</i> value					
Termination interval	<.0001	<.0001	<.0001	<.0001	<.0001	0.1697	0.3745
POST	na ^c	na	na	na	na	na	0.0424
Termination interval*POST	na	na	na	na	na	na	0.8546

^aTermination interval refers to cover crop termination intervals of 14, 7, and 0 d prior to planting, and 7 and 14 d after planting. POST stands for POST herbicide premixes of glyphosate + fomesafen (1,120 and 280 g ae ha⁻¹, respectively) or glyphosate + the diglycolamine salt of dicamba (1,120 and 560 g ae ha⁻¹, respectively) applied when Palmer amaranth for that termination interval reached 10 cm in height.

^bColumn headings of 7 and 14 designate rating intervals of 7 and 14 d after planting soybean.

^cna (not applicable) as no factorial was run.

dicamba than did the current research. Many other studies have documented the efficacy of dicamba on various vetch species at lower rates than used in this study, confirming the viability of this herbicide for controlling vetch species (Curran et al. 2015; McCurdy et al. 2013; Wolfe et al. 2016). Also, the efficacy of glyphosate for controlling wheat is well documented (Ashford and Reeves 2003; Davis 2010; Price et al. 2009; Reddy 2001). The combination of these herbicides can control a wheat + hairy vetch cover crop before or after soybean establishment at rates for use in GDT soybeans.

Significant main effects were not present for crop stand (Tables 2 and 3). However, reported least squared means for crop stand are generally low. A severe early-season infestation of three-cornered alfalfaoppers (*Spissistilus festinus* Say) occurred in 2015. This caused notable stand loss across all treatments. However, stand loss was uniform across all treatments and did not cause an interaction with data for stands between site years ($P=0.8369$, data not shown). Means for crop stands in 2015 and 2016 were 238,200 and 303,200 plants ha⁻¹, respectively. However, because of the aforementioned insect problem in 2015, a blanket application of a pyrethroid insecticide was made in 2016 at planting to alleviate this problem (S. Stewart, personal

communication). Additionally, there was a significant main effect of POST herbicide for soybean yields (Tables 2 and 4). Pooled over all termination timings, treatments receiving a POST application of dicamba yielded 100 kg ha⁻¹ higher than those treated with fomesafen. Although there were differences in yield among POST herbicide treatments, least squared means for each herbicide treatment was still above the 2015 average yield of 3,300 kg ha⁻¹ for Tennessee (Anonymous 2012). Similarly, Reddy et al. (2003) reported no differences in soybean stand or yield when comparing a cereal rye or crimson clover cover crop to a conventional no-till system in Mississippi.

The parameters for the logistic regression of termination interval and days to 10-cm Palmer amaranth produced a model with the parameter estimates $y = 28.3 + 11.3 / (1 + \exp(-(x-0.9)/5.5))$ with an $R^2 = 0.86$ (Figure 1). This would estimate that all termination treatments delayed Palmer amaranth growth to 10 cm in height by at least 28 d. The termination interval for a wheat + vetch cover crop terminated with glyphosate + dicamba to maximize the number of days for Palmer amaranth to reach 10 cm in height is 11 DAP. Similarly, Ryan et al. (2011) reported that increased cereal rye biomass was strongly related to decreasing weed biomass. Although biomass in this study stopped increasing at cash crop planting, the

Table 3. Biomass and control of a wheat + hairy vetch cover crop, and soybean population as affected by termination intervals before and after planting soybean in Tennessee in 2015 and 2016.^{a,b}

Termination interval ^c	Biomass	Cover crop control ^d				Soybean stand
		Wheat		Hairy vetch		
		7	14	7	14	
	kg ha ⁻¹	%				plants ha ⁻¹
14 DPP	7,000c	99a	99a	99a	99a	247,800
DPP	11,300b	99a	99a	99a	99a	275,700
At planting	16,000a	99a	99a	99a	99a	288,000
7 DAP	10,200b	87b	99a	87b	99a	273,200
14 DAP	6,100c	86b	91b	85c	90b	269,100

^aAbbreviations: DPP, days prior to planting; DAP, days after planting.

^bMeans within a column followed by the same letter are not significantly different at $p \leq 0.05$.

^cTermination intervals of -14, -7, 0, 7, and 14 designate the number of days before or after soybean planting that the termination treatment of glyphosate + dicamba (1,120 and 560 g ae ha⁻¹, respectively) was applied.

^dColumn headings of 7 and 14 designate rating intervals of 7 and 14 d after planting soybean.

Table 4. Control of Palmer amaranth and dicamba-resistant soybean yield in a wheat + hairy vetch cover crop mixture in Jackson, TN, in 2015 and 2016^a.

Effect ^b		Palmer amaranth control ^{c,d}					Yield
		7	14	21	28	R6	
		%					kg ha ⁻¹
POST							
	Dicamba	95	98	99a	99a	99	4,300a
	Fomesafen	95	96	97b	97b	98	4,200b
Termination interval							
	14 DPP	96ab	97a	97	98	98	4,200
	7 DPP	92c	98a	97	98	99	4,400
	At planting	97ab	97a	97	97	98	4,200
	7 DAP	97a	99a	99	99	99	4,200
	14 DAP	93bc	93b	99	99	98	4,200
Termination interval*POST							
	Dicamba -14 DPP	97	99	98	99	98	4,300
	Fomesafen --14 DPP	96	96	96	96	97	4,100
	Dicamba -7 DPP	92	99	99	99	99	4,400
	Fomesafen -7 DPP	92	98	96	97	99	4,400
	Dicamba-at planting	98	99	99	99	99	4,300
	Fomesafen-at planting	95	96	95	95	97	4,100
	Dicamba-7 DAP	96	99	99	99	99	4,300
	Fomesafen-7 DAP	98	99	99	99	99	4,100
	Dicamba-14 DAP	91	92	99	99	99	4,300
	Fomesafen-14 DAP	96	95	99	99	98	4,100

^aAbbreviations: DPP, days prior to planting; DAP, days after planting.^bTermination intervals of -14, -7, 0, 7, and 14 designate the number of days before or after soybean planting that the termination treatment of glyphosate + dicamba (1,120 and 560 g ae ha⁻¹, respectively) was applied.^cColumn headings of 7, 14, 21, and 28 designate rating intervals of 7, 14, 21, and 28 d after POST herbicide application. Column heading R6 refers to soybean growth stage of R6 and was taken as an end-of-season rating.^dMeans within a column followed by the same letter are not significantly different at $p \leq 0.05$. Letters are only reflective of means within a main effect. Means followed by no letter are either not significantly different at $p \leq 0.05$, or letter separation is shown in a higher interaction.

effects of delayed termination with glyphosate + dicamba until after planting proved to be an effective way to increase the amount of time necessary for Palmer amaranth to reach 10 cm in height. Palmer amaranth control was significantly affected by termination interval 7 and 14 DAT (Tables 4 and 5). Control was lowest (92% to 93%) 7 DAT in 7-DPP and 14-DAP termination treatments. At this rating, differences in control among the other termination treatments were not significant, and control among these treatments ranged from 96% to 97%. Additionally, 14-DAT control from all treatments except the 14 DAP (93%) termination interval was similar ($\geq 97\%$). However, these differences were transient, as differences among termination intervals were not present 21 or 28 DAT, and, pooled over POST herbicides, all treatments provided $\geq 97\%$ control at these timings. There was a significant main effect of POST herbicide treatment 21 and 28 DAT. Control 21 and 28 DAT ranged from 97% to 99%, with dicamba having greater control at these ratings. Significant main effects or interactions among the

main effects were not detected at the final (R6 soybean stage) rating, and all treatments provided $\geq 97\%$ control of Palmer amaranth.

Although differences in Palmer amaranth control were not apparent at the end of the season, early-season Palmer amaranth control ratings (21 and 28 DAT) followed a trend similar to that of soybean yields (Tables 1 and 4). Van Acker et al. (1993) reported in four of six site years that the critical period of weed removal to prevent 2.5% yield loss in soybeans was ≥ 27 d after emergence. However, in this same study, the critical period of weed removal to prevent a 5% yield loss was ≥ 40 d after emergence in three of six site years. POST treatments for Palmer amaranth control were applied 29 to 39 d after planting, meaning the weed removal in this study fell in a period that could cause 2.5% to 5% yield loss. The differences in yield are thought to be attributed to early-season weed control.

Glyphosate + dicamba can be an effective tool for terminating wheat + hairy vetch cover crops when used 14 DPP to 14 DAP.

Table 5. The main effects of cover crop termination interval, POST herbicide treatment, and the interaction among the main effects on Palmer amaranth control in Jackson, TN, in 2015 and 2016.

Effect ^a	Palmer amaranth ^b				
	7	14	21	28	R6
	<i>p</i> value				
Termination interval	0.0099	0.0001	0.0861	0.1568	0.3389
POST	0.5232	0.2580	0.0026	0.0029	0.0812
Termination interval*POST	0.1560	0.0828	0.1347	0.1568	0.5885

^aTermination interval refers to cover crop termination intervals of 14, 7, and 0 d prior to planting, and 7 and 14 d after planting. POST stands for POST herbicide premixes of glyphosate + fomesafen (1,120 and 280 g ae ha⁻¹, respectively) or glyphosate plus the diglycolamine salt of dicamba (1,120 and 560 g ae ha⁻¹, respectively) applied when Palmer amaranth for that termination interval reached 10 cm in height.


^bColumn headings of 7, 14, 21, and 28 designate rating intervals of 7, 14, 21, and 28 d after POST herbicide application. Column heading R6 refers to soybean growth stage of R6 and was taken as an end-of-season rating.

Additionally, in all treatments, one effective POST herbicide application plus a cover crop was sufficient to provide $\geq 97\%$ control of GR Palmer amaranth while maintaining soybean yields above the state average (Anonymous 2012). The ability to use glyphosate + dicamba shortly before or after planting for cover crop control in GDT soybeans allows producers increased flexibility in managing high-biomass cover crops for control of Palmer amaranth. However, producers should be aware of other possible pests such as insects when delaying cover crop termination until near soybean planting (L. Steckel, unpublished data). Additionally, it is well documented that including at least one other effective MOA will greatly improve the longevity of dicamba as an effective POST for Palmer amaranth (Beckie 2011; Burke et al. 2005; Culpepper 2006; Inman et al. 2016; Miller and Norsworthy 2016; Price et al. 2011; Riar et al. 2013). The results suggest that with a wheat + hairy vetch cover crop terminated 14 DPP with glyphosate + dicamba, delayed Palmer amaranth growth to 10 cm in height by ≥ 28 DAP. Moreover, delaying cover crop termination until 14 DAP can correspondently delay this interval for Palmer amaranth to 38 DAP. When utilizing a wheat + hairy vetch cover crop in GDT soybean, producers should delay cover crop termination until 11 to 14 DPP and make at least one POST application of glyphosate + dicamba + an additional herbicide MOA to maximize Palmer amaranth control and soybean yields.

Acknowledgments. Funding for this research was provided by Monsanto and the Tennessee Soybean Promotion Board and is greatly appreciated by the authors.

References

- Anonymous (2012) Crop Production and Agricultural Chemical Usage in Field Crops. Agricultural Statistics Board, NASS, and USDA. https://www.nass.usda.gov/Publications/Methodology_and_Data_Quality/Agricultural_Chemical_Usage_-_Field_Crops/05_2012/chfcqm12.pdf. Accessed: October 18, 2017
- Anonymous (2017) Xtendimax Supplemental Label for use on dicamba-tolerant cotton and soybeans. <http://www.xtendimaxapplicationrequirements.com>. Accessed: August 23, 2017
- Ashford DL, Reeves DW (2003) Use of a mechanical roller-crimper as an alternative kill method for cover crops. *Am J Altern Ag* 18:37–45
- Beckie HJ (2011) Herbicide-resistant weed management: focus on glyphosate. *Pest Manage Sci* 67:1037–1048
- Blouin DC, Webster EP, Bond JA (2011) On the analysis of combined experiments. *Weed Technol* 25:165–169
- Bond JA, Oliver LR (2006) Comparative growth of Palmer amaranth (*Amaranthus palmeri*) accessions. *Weed Sci* 54:121–126
- Burke IC, Twozler SC, Askew SD, Wilcut JW, Smith WD (2005) Weed management systems in glyphosate-resistant cotton. *Weed Technol* 19:422–429
- Cahoon CW, York AC, Jordan DL, Everman WJ, Seagroves RW, Culpepper AS, Eure PM (2015) Palmer amaranth (*Amaranthus palmeri*) management in dicamba-resistant cotton. *Weed Technol* 29:758–770
- Carmer SG, Nyquist WE, Walker WM (1989) Least significant differences in combined analyses of experiments with two- or three-factor treatment designs. *Agron J* 81:655–672
- Corbett JL, Askew SD, Thomas WE, Wilcut JW (2004) Weed efficacy evaluations for bromoxynil, glufosinate, glyphosate, pyriithiobac, and sulfosate. *Weed Technol* 18:443–453
- Crow WD, Steckel LE, Mueller TC, Hayes RM (2016) Management of large, glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) in corn. *Weed Technol* 30:611–616
- Culpepper AS (2006) Glyphosate-induced weed shifts. *Weed Technol* 20:277–281
- Curran WS, Wallace JM, Mirsky S, Crockett B (2015) Effectiveness of herbicides for control of hairy vetch (*Vicia villosa*) in winter wheat. *Weed Technol* 29:509–518
- Davis AS (2010) Cover-crop roller-crimper contributes to weed management in no-till soybean. *Weed Sci* 58:300–309
- Fisk JW, Hersternan OB, Shrestha A, Kells JJ, Harwood RR, Squire JM, Sheaffer CC (2001) Weed suppression by annual legume cover crops in no-tillage corn. *Agron J* 93:319–325
- Heap I (2016) The International Survey of Herbicide Resistant Weeds. <http://www.weedscience.org>. Accessed: August 30, 2016

**Figure 1.** Termination interval of a wheat + hairy vetch cover crop effects the number of days until Palmer amaranth can reach 10 cm in height.

- Inman MD, Jordan DL, York AC, Jennings KM, Monks DW, Everman WJ, Bollman SL, Fowler JT, Cole RM, Soteres JK (2016) Long-term management of Palmer amaranth (*Amaranthus palmeri*) in dicamba-tolerant cotton. *Weed Sci* 64:161–169
- Isik DE, Ngouajio KM, Mennan H (2009) Weed suppression in organic pepper (*Capiscum annuum* L.) with winter cover crops. *Crop Prot* 28:356–363
- Kenerson D (2014) 2014 Tennessee tillage systems. USDA, National Agricultural Statistics Service. https://www.nass.usda.gov/Statistics_by_State/Tennessee/Publications/Special_Surveys/tillage2014.pdf. Accessed August 15, 2016
- Klingaman TE, Oliver LR (1994) Palmer amaranth (*Amaranthus palmeri*) interference in soybeans (*Glycine max*). *Weed Sci* 42:523–527
- McCurdy JD, McElroy JS, Flessner ML (2013) Differential response to four *Trifolium* species to common broadleaf herbicides: implications for mixed grass–legume swards. *Weed Technol* 27:123–128
- Merchant RM, Sosnoskie LM, Culpepper AS, Steckel LE, York AC, Braxton LB, Ford JC (2013) Weed response to 2,4-D, 2,4-DB, and dicamba applied alone or with glufosinate. *J Cotton Sci* 17:212–218
- Miller MR, Norsworthy JK (2016) Evaluation of herbicide programs for use in a 2,4-D-resistant soybean technology for control of glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*). *Weed Technol* 30:366–376
- Mirsky SB, Curran WS, Mortensen DA, Ryan MR, Shumway DL (2009) Control of cereal rye with a roller/crimper as influenced by cover crop phenology. *Agron J* 101:1589–1596
- Mirsky SB, Ryan MR, Teasdale JR, Curran WS, Reberg-Horton CS, Spargo JT, Wells MS, Keene CL, Moyer JW (2013) Overcoming weed management challenges in cover crop–based organic rotational no-till soybean production in the eastern United States. *Weed Technol* 27:193–203
- Mueller TC, Wright DR, Remund KM (2013) Effect of formulation and application time of day on detecting dicamba in the air under field conditions. *Weed Sci* 61:586–593
- Norsworthy JK, Griffith GM, Scott RC, Smith KL, Oliver LR (2008) Confirmation and control of glyphosate-resistant Palmer Amaranth (*Amaranthus palmeri*) in Arkansas. *Weed Technol* 22:108–113
- Norsworthy JK, McClelland M, Griffith G, Bangarwa SK, Still J (2010) Evaluation of legume cover crops and weed control programs in conservation-tillage, enhanced glyphosate-resistant cotton. *Weed Technol* 24:269–274
- Norsworthy JK, McClelland M, Griffith G, Bangarwa SK, Still J (2011) Evaluation of cereal and brassicaceae cover crops in conservation-tillage enhanced, glyphosate-resistant cotton. *Weed Technol* 25:6–13
- Norsworthy JK, Ward SM, Shaw DR, Llewellyn RS, Nichols RL, Webster TM, Bradely KW, Frisvold G, Powles SB, Burgos NR, Witt WW, Barrett M (2012) Reducing the risks of herbicide resistance: best management practices and recommendations. *Weed Sci* 60(sp1): 31–62
- Price AJ, Arriaga FL, Raper RL, Balkcom KS, Komecki TS, Reeves DW (2009) Comparison of mechanical and chemical winter cereal cover crop termination systems and cotton yield in conservation agriculture. *J Cotton Sci* 13:238–245
- Price AJ, Balkcom KS, Culpepper SA, Kelton JA, Nichols RL, Schomberg H (2011) Glyphosate-resistant Palmer amaranth: a threat to conservation tillage. *J Soil Water Conservation* 66:265–275
- Reddy KN (2001) Effects of cereal and legume cover crop residues on weeds, yield, and net return in soybean (*Glycine max*). *Weed Technol* 15:660–668
- Reddy KN, Zablotowicz RM, Locke MA, Koger CH (2003) Cover crop, tillage, and herbicide effects on weeds, soil properties, microbial populations, and soybean yield. *Weed Sci* 51:987–994
- Riar DS, Norsworthy JK, Steckel LE, Stephenson IV SO, Eubank TW, Bond J, Scott RC (2013) Adoption of best management practices for herbicide-resistant weeds in the midsouthern United States cotton, rice, and soybean. *Weed Technol* 27:788–797
- Ryan MR, Mirsky SB, Mortensen DA, Teasdale JR, Curran WS (2011a) Potential synergistic effects on cereal rye biomass and soybean planting density on weed suppression. *Weed Sci* 59:238–246
- Ryan MR, Curran WS, Grantham AM, Hunsberger LK, Mirsky SB, Mortensen DA, Nord EA, Wilson DO (2011b) Effects of seeding rate and poultry litter on weed suppression from a rolled cereal rye cover crop. *Weed Sci* 59: 438–444
- Samples CA, Dodds DM, Reynolds DZ, Dixon TH, Bond JA, Reynolds DB, Mills A (2013) Evaluation of PRE and POST weed control programs for Palmer amaranth control. Page 173 in *Proceedings of the 2013 Southern Weed Science Society*. Houston, TX: Southern Weed Science Society
- Saxton AM (2013) DandA.sas: Design and analysis macro collection version 2.11. Knoxville, TN: University of Tennessee
- Shaner DL, ed (2014) *Herbicide Handbook*. 10th edn. Lawrence, KS: Weed Science Society of America. Pp 139–141
- [SARE 2007] Sustainable Agriculture Research and Education (2007) *Managing cover crops profitably*, 3rd edn. College Park, MD: SARE. P 244
- [USDA, NASS] United States Department of Agriculture, National Agricultural Statistics Service (2014) <http://www.nass.usda.gov>. Accessed September 29, 2016
- Van Acker RC, Swanton CJ, Weise SF (1993) The critical period of weed control in soybean [*Glycine max* (L.) Merr.]. *Weed Sci* 41:194–200
- VanGessel MJ (2001) Glyphosate-resistant horseweed from Delaware. *Weed Sci* 49:703–705
- Webster TM, Sosnoskie LM (2010) A changing weed spectrum in Georgia cotton. *Weed Sci* 58:73–79
- White RH, Worsham AD (1990) Control of legume cover crops in no-till corn (*Zea mays*) and cotton (*Gossypium hirsutum*). *Weed Technol* 4:57–62
- Wiggins MS, Hayes RM, Steckel LE (2016b) Evaluating cover crops and herbicides for glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) control in cotton. *Weed Technol* 30:415–422
- Wiggins MS, McClure MA, Hayes RM, Steckel LE (2016a) Integrating cover crops and POST herbicides for glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) control in corn. *Weed Technol* 29:412–418
- Wolfe JC, Neal JC, Harlow CD (2016) Selective broadleaf weed control in turfgrass with bioherbicides *Phoma macrostoma* and Thaxtomin A. *Weed Technol* 30:688–700
- Wortman SE, Francis CA, Bernards ML, Drijber RA, Lindquist JL (2012) Optimizing cover crop benefits with diverse mixtures and an alternative termination method. *Agron J* 104:1425–1435
- York AC, Cahoon CW, Oliver GW (2015) Palmer amaranth management with Engenia in BollGard II XtendFlex cotton. Page 174 in *Proceedings of the Southern Weed Science Society*. Savannah, GA: Southern Weed Science Society
- York AC, Culpepper AS, Sosnoskie LM, Bollman S (2012) Palmer amaranth management in dicamba/glufosinate tolerant cotton. Page 98 in *Proceedings of the Southern Weed Science Society*. Charleston, SC: Southern Weed Science Society
- Zadoks JC, Chang TT, Konzak CF (1974) A decimal code for growth stages of cereals. *Weed Res* 14: 415–442
- Zotarelli L, Avila L, Scholberg MS, Alves BJR (2009) Benefits of vetch and rye cover crops to sweet corn under no-tillage. *Agron J* 101:252–260