

# The fastest known globally convergent first-order method for minimizing strongly convex functions

Bryan Van Scoy

University of Wisconsin–Madison

Dec 12, 2017

## Unconstrained optimization:

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & x \in \mathbb{R}^d \end{array}$$

- Need methods which are *fast* and *simple*
- Use *first-order* methods

## Unconstrained optimization:

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & x \in \mathbb{R}^d \end{array}$$

- Need methods which are *fast* and *simple*
- Use *first-order* methods
- In this talk, we will design a first-order method for the case when  $f$  is smooth and strongly convex

## Unconstrained optimization:

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & x \in \mathbb{R}^d \end{array}$$

- Need methods which are *fast* and *simple*
- Use *first-order* methods
- In this talk, we will design a first-order method for the case when  $f$  is smooth and strongly convex

### Main result

Design and analyze a novel method which is both globally convergent and faster than Nesterov's method

**Analysis** Simple convergence proof (time domain)

**Design** Intuition using IQCs (frequency domain)

## Smooth strongly convex

A differentiable function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  is called  $L$ -smooth if

$$\|\nabla f(x) - \nabla f(y)\| \leq L \|x - y\| \quad \text{for all } x, y \in \mathbb{R}^d$$

and  $m$ -strongly convex if

$$f(y) \geq f(x) + \nabla f(x)^\top (y - x) + \frac{m}{2} \|x - y\|^2 \quad \text{for all } x, y \in \mathbb{R}^d.$$


# Smooth strongly convex

A differentiable function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  is called  $L$ -smooth if


$$\|\nabla f(x) - \nabla f(y)\| \leq L \|x - y\| \quad \text{for all } x, y \in \mathbb{R}^d$$

and  $m$ -strongly convex if

$$f(y) \geq f(x) + \nabla f(x)^\top (y - x) + \frac{m}{2} \|x - y\|^2 \quad \text{for all } x, y \in \mathbb{R}^d.$$



$L$ -smooth  $m$ -strongly convex



slope restricted on  $[m, L]$

# Method

gradient method

$$x_{k+1} = x_k - \alpha \nabla f(x_k)$$

heavy ball method

$$x_{k+1} = (1 + \beta)x_k - \beta x_{k-1} - \alpha \nabla f(x_k)$$

fast gradient method

$$x_{k+1} = (1 + \beta)x_k - \beta x_{k-1} - \alpha \nabla f((1 + \beta)x_k - \beta x_{k-1})$$

# Method

gradient method

$$x_{k+1} = x_k - \alpha \nabla f(x_k)$$

heavy ball method

$$x_{k+1} = (1 + \beta)x_k - \beta x_{k-1} - \alpha \nabla f(x_k)$$

fast gradient method

$$x_{k+1} = (1 + \beta)x_k - \beta x_{k-1} - \alpha \nabla f((1 + \beta)x_k - \beta x_{k-1})$$

triple momentum method

$$x_{k+1} = (1 + \beta)x_k - \beta x_{k-1} - \alpha \nabla f((1 + \gamma)x_k - \gamma x_{k-1})$$

Method	Parameters
GM	$(\alpha, 0, 0)$
HBM (Polyak, 1964)	$(\alpha, \beta, 0)$
FGM (Nesterov, 2004)	$(\alpha, \beta, \beta)$
TMM (Van Scoy, Freeman, Lynch, 2017)	$(\alpha, \beta, \gamma)$



# Triple momentum method

$$x_{k+1} = (1 + \beta)x_k - \beta x_{k-1} - \alpha \nabla f((1 + \gamma)x_k - \gamma x_{k-1})$$

**Parameters:**

$$\rho = 1 - \frac{1}{\sqrt{\kappa}}$$

$$\alpha = \frac{1+\rho}{L}$$

$$\beta = \frac{\rho^2}{2-\rho}$$

$$\gamma = \frac{\rho^2}{(1+\rho)(2-\rho)}$$

**Condition ratio**  $\kappa := L/m$

# Triple momentum method

$$x_{k+1} = (1 + \beta)x_k - \beta x_{k-1} - \alpha \nabla f((1 + \gamma)x_k - \gamma x_{k-1})$$

**Parameters:**

$$\rho = 1 - \frac{1}{\sqrt{\kappa}}$$

$$\alpha = \frac{1+\rho}{L}$$

$$\beta = \frac{\rho^2}{2-\rho}$$

$$\gamma = \frac{\rho^2}{(1+\rho)(2-\rho)}$$



**Condition ratio**  $\kappa := L/m$

## Theorem (Van Scoy, Freeman, Lynch, 2017)

Suppose  $f$  is  $L$ -smooth and  $m$ -strongly convex with minimizer  $x_\star \in \mathbb{R}^d$ . Then for any initial conditions  $x_0, x_{-1} \in \mathbb{R}^d$ , there exists a constant  $c > 0$  such that

$$\|x_k - x_\star\| \leq c \rho^k \quad \text{for all } k \geq 1.$$



# $f$ quadratic



Convergence rate:  $\|x_k - x_\star\| \leq c \rho^k$

$$\text{Iterations to converge} \propto -\frac{1}{\log \rho}$$

# $f$ smooth strongly convex



- HBM does **not** converge if  $L/m \geq (2 + \sqrt{5})^2 \approx 17.94$
- For FGM, Nesterov proved the rate  $\sqrt{1 - \sqrt{m/L}}$  which is **loose**!
- TMM converges **faster** than Nesterov's method!

# Simulations

**Objective function:**

$$f(x) = \sum_{i=1}^p g(a_i^T x - b_i) + \frac{m}{2} \|x\|^2, \quad x \in \mathbb{R}^d$$

where

$$g(y) = \begin{cases} \frac{1}{2} y^2 e^{-r/y}, & y > 0 \\ 0, & y \leq 0 \end{cases}$$


with  $A = [a_1, \dots, a_p] \in \mathbb{R}^{d \times p}$ ,  $b \in \mathbb{R}^p$ , and  $\|A\| = \sqrt{L - m}$

$f$  is

- $m$ -smooth
- $L$ -strongly convex
- infinitely differentiable (of class  $C^\infty$ )


# Simulations

**Parameters:**  $m = 1$ ,  $L = 10^4$ ,  $d = 100$ ,  $p = 5$ ,  $r = 10^{-6}$



# Robustness to $m$

**Parameters:**  $m = 1$ ,  $L = 10^4$ ,  $d = 100$ ,  $p = 5$ ,  $r = 10^{-6}$




To prove the bound for **TMM**, use *interpolation*.




To prove the bound for **TMM**, use *interpolation*.

**Interpolation:** The set  $\{y, u, v\}$  is  $\mathcal{F}$ -interpolable if and only if  $u_k = \nabla f(y_k)$  and  $v_k = f(y_k)$  for some  $f \in \mathcal{F}$  and all  $k$ .



To prove the bound for **TMM**, use *interpolation*.

**Interpolation:** The set  $\{y, u, v\}$  is  $\mathcal{F}$ -interpolable if and only if  $u_k = \nabla f(y_k)$  and  $v_k = f(y_k)$  for some  $f \in \mathcal{F}$  and all  $k$ .



### Theorem (Taylor, Hendrickx, Glineur, 2016)

The set  $\{y, u, v\}$  is interpolable by an  $L$ -smooth  $m$ -strongly convex function if and only if  $q_{ij} \geq 0$  for all  $i, j$  where

$$q_{ij} := (L - m)(v_i - v_j) - \frac{1}{2}\|u_i - u_j\|^2 \\ + (mu_i - Lu_j)^\top (y_i - y_j) - \frac{mL}{2}\|y_i - y_j\|^2.$$

# Sketch of proof for TMM

1. Suppose  $f$  is  $L$ -smooth and  $m$ -strongly convex. Then the **interpolation conditions** are satisfied; specifically,  $q_{ij} \geq 0$  for all  $i, j$ .

# Sketch of proof for TMM

1. Suppose  $f$  is  $L$ -smooth and  $m$ -strongly convex. Then the **interpolation conditions** are satisfied; specifically,  $q_{ij} \geq 0$  for all  $i, j$ .
2. Define the **Lyapunov function**

$$V_k := mL \|z_k - x_\star\|^2 + q_{k-1,\star}$$

where  $z_k := (1 + \delta)x_k - \delta x_{k-1}$  and  $\delta := \frac{\rho^2}{1 - \rho^2}$ .

# Sketch of proof for TMM

1. Suppose  $f$  is  $L$ -smooth and  $m$ -strongly convex. Then the **interpolation conditions** are satisfied; specifically,  $q_{ij} \geq 0$  for all  $i, j$ .
2. Define the **Lyapunov function**

$$V_k := mL \|z_k - x_\star\|^2 + q_{k-1,\star}$$

where  $z_k := (1 + \delta)x_k - \delta x_{k-1}$  and  $\delta := \frac{\rho^2}{1 - \rho^2}$ .

3. Using the definition of TMM, it is straightforward to verify that

$$V_{k+1} - \rho^2 V_k = -[(1 - \rho^2)q_{\star,k} + \rho^2 q_{k-1,k}] \leq 0$$

for all  $k \geq 1$ .

# Sketch of proof for TMM

1. Suppose  $f$  is  $L$ -smooth and  $m$ -strongly convex. Then the **interpolation conditions** are satisfied; specifically,  $q_{ij} \geq 0$  for all  $i, j$ .
2. Define the **Lyapunov function**

$$V_k := mL \|z_k - x_\star\|^2 + q_{k-1,\star}$$

where  $z_k := (1 + \delta)x_k - \delta x_{k-1}$  and  $\delta := \frac{\rho^2}{1 - \rho^2}$ .


3. Using the definition of TMM, it is straightforward to verify that

$$V_{k+1} - \rho^2 V_k = -[(1 - \rho^2)q_{\star,k} + \rho^2 q_{k-1,k}] \leq 0$$

for all  $k \geq 1$ .


4. Iterating gives the **bound**  $V_k \leq \rho^{2(k-1)} V_1$  for  $k \geq 1$ .

# Integral Quadratic Constraints (IQCs)



$$G : \quad \begin{aligned} x_{k+1} &= (1 + \beta)x_k - \beta x_{k-1} - \alpha u_k \\ y_k &= (1 + \gamma)x_k - \gamma x_{k-1} \end{aligned}$$

# Integral Quadratic Constraints (IQCs)




$(\Psi, M)$  are chosen such that  $w$  satisfies

$$0 \leq \sum_{j=0}^k \rho^{-2j} (w_j - w_*)^\top M (w_j - w_*)$$

when  $f$  is  $L$ -smooth and  $m$ -strongly convex.



# Integral Quadratic Constraints (IQCs)



$(\Psi, M)$  are chosen such that  $w$  satisfies

$$0 \leq \sum_{j=0}^k \rho^{-2j} (w_j - w_\star)^\top M (w_j - w_\star)$$

when  $f$  is  $L$ -smooth and  $m$ -strongly convex.


## Theorem (Boczar, Lessard, Recht, 2015)

Define  $\Pi(z) := \Psi(z)^* M \Psi(z)$ . If there exists  $\varepsilon > 0$  such that

$$\begin{bmatrix} G(z) \\ I \end{bmatrix}^* \Pi(z) \begin{bmatrix} G(z) \\ I \end{bmatrix} \preceq -\varepsilon I \quad \text{for all } z \in \rho\mathbb{T}$$

then the state of  $G$  converges linearly with rate  $\rho$ .

# Integral Quadratic Constraints (IQCs)



$(\Psi, M)$  are chosen such that  $w$  satisfies

$$0 \leq \sum_{j=0}^k \rho^{-2j} (w_j - w_\star)^\top M (w_j - w_\star)$$

when  $f$  is  $L$ -smooth and  $m$ -strongly convex.

## Theorem (Boczar, Lessard, Recht, 2015)

Define  $\Pi(z) := \Psi(z)^* M \Psi(z)$ . If there exists  $\varepsilon > 0$  such that

$$\begin{bmatrix} G(z) \\ I \end{bmatrix}^* \Pi(z) \begin{bmatrix} G(z) \\ I \end{bmatrix} \preceq -\varepsilon I \quad \text{for all } z \in \rho\mathbb{T}$$

then the state of  $G$  converges linearly with rate  $\rho$ .

The TMM parameters are the unique solution to

$$\begin{bmatrix} G(z) \\ I \end{bmatrix}^* \Pi(z) \begin{bmatrix} G(z) \\ I \end{bmatrix} = 0 \quad \text{for all } z \in \rho\mathbb{T}$$

## Summary

**Triple momentum method:** globally convergent with rate  $1 - \sqrt{m/L}$  when  $f$  is  $L$ -smooth and  $m$ -strongly convex

**Analysis** Simple convergence proof (time domain)

**Design** Intuition using IQCs (frequency domain)

## Summary

**Triple momentum method:** globally convergent with rate  $1 - \sqrt{m/L}$  when  $f$  is  $L$ -smooth and  $m$ -strongly convex

**Analysis** Simple convergence proof (time domain)

**Design** Intuition using IQCs (frequency domain)

## Extension: gradient noise

$$x_{k+1} = (1 + \beta)x_k - \beta x_{k-1} - \alpha u_k$$

$$y_k = (1 + \gamma)x_k - \gamma x_{k-1}$$

**No noise:**  $u = \nabla f(y)$

**Relative gradient noise:**  $\|u - \nabla f(y)\|_2 \leq \delta \|\nabla f(y)\|_2$

---

S. Cyrus, B. Hu, B. Van Scoy, L. Lessard. "A Robust Accelerated Optimization Algorithm for Strongly Convex Functions". In ArXiv e-prints (Oct. 2017). arXiv: 170.04753 [math.OC].

Thanks!

# Gradient noise

What if the measured gradient is *not* the actual gradient?

$$\begin{aligned}x_{k+1} &= (1 + \beta)x_k - \beta x_{k-1} - \alpha u_k \\y_k &= (1 + \gamma)x_k - \gamma x_{k-1}\end{aligned}$$

**No noise:**  $u = \nabla f(y)$

**Relative gradient noise:**  $\|u - \nabla f(y)\|_2 \leq \delta \|\nabla f(y)\|_2$

# Robust momentum method

$$x_{k+1} = (1 + \beta)x_k - \beta x_{k-1} - \alpha \nabla f((1 + \gamma)x_k - \gamma x_{k-1})$$

## Parameters:

$$\rho \in \left[1 - \frac{1}{\sqrt{\kappa}}, 1 - \frac{1}{\kappa}\right]$$

$$\alpha = \frac{\kappa(1-\rho)^2(1+\rho)}{L}$$

$$\beta = \frac{\kappa\rho^3}{\kappa-1}$$

$$\gamma = \frac{\rho^3}{(\kappa-1)(1-\rho)^2(1+\rho)}$$

## Theorem (Cyrus, Hu, Van Scoy, Lessard, 2017)

Suppose  $f$  is  $L$ -smooth and  $m$ -strongly convex with minimizer  $x_\star \in \mathbb{R}^d$ . Then for any initial conditions  $x_0, x_{-1} \in \mathbb{R}^d$ , there exists a constant  $c > 0$  such that

$$\|x_k - x_\star\| \leq c\rho^k \quad \text{for all } k \geq 1.$$

# Robust momentum method

$$x_{k+1} = (1 + \beta)x_k - \beta x_{k-1} - \alpha \nabla f((1 + \gamma)x_k - \gamma x_{k-1})$$


## Parameters:

$$\rho \in \left[1 - \frac{1}{\sqrt{\kappa}}, 1 - \frac{1}{\kappa}\right]$$

$$\alpha = \frac{\kappa(1-\rho)^2(1+\rho)}{L}$$

$$\beta = \frac{\kappa\rho^3}{\kappa-1}$$

$$\gamma = \frac{\rho^3}{(\kappa-1)(1-\rho)^2(1+\rho)}$$



## Theorem (Cyrus, Hu, Van Scoy, Lessard, 2017)

Suppose  $f$  is  $L$ -smooth and  $m$ -strongly convex with minimizer  $x_\star \in \mathbb{R}^d$ . Then for any initial conditions  $x_0, x_{-1} \in \mathbb{R}^d$ , there exists a constant  $c > 0$  such that

$$\|x_k - x_\star\| \leq c \rho^k \quad \text{for all } k \geq 1.$$



# Robust momentum method

$$x_{k+1} = (1 + \beta)x_k - \beta x_{k-1} - \alpha \nabla f((1 + \gamma)x_k - \gamma x_{k-1})$$


## Parameters:

$$\rho \in \left[1 - \frac{1}{\sqrt{\kappa}}, 1 - \frac{1}{\kappa}\right]$$

$$\alpha = \frac{\kappa(1-\rho)^2(1+\rho)}{L}$$

$$\beta = \frac{\kappa\rho^3}{\kappa-1}$$

$$\gamma = \frac{\rho^3}{(\kappa-1)(1-\rho)^2(1+\rho)}$$




## Theorem (Cyrus, Hu, Van Scoy, Lessard, 2017)

Suppose  $f$  is  $L$ -smooth and  $m$ -strongly convex with minimizer  $x_\star \in \mathbb{R}^d$ . Then for any initial conditions  $x_0, x_{-1} \in \mathbb{R}^d$ , there exists a constant  $c > 0$  such that

$$\|x_k - x_\star\| \leq c \rho^k \quad \text{for all } k \geq 1.$$


# Trade-off: Speed vs. Robustness

( $\kappa = 10$ )




# Trade-off: Speed vs. Robustness

( $\kappa = 10$ )




# Trade-off: Speed vs. Robustness

( $\kappa = 10$ )




# Trade-off: Speed vs. Robustness

( $\kappa = 10$ )



# Trade-off: Speed vs. Robustness

( $\kappa = 10$ )



# Conclusion

## Analysis

- **Numerical:** solve SDP to calculate upper bound on convergence rate
- **Closed-form:** have expressions for convergence rate for some methods and functions classes (such as **TMM** on smooth strongly convex functions)

## Design

- **Triple momentum method** - Fastest known convergence rate for first-order methods on smooth strongly convex functions
- **Robust momentum method** - Interpolates **TMM** and **GM** (with  $\alpha = 1/L$ ) to exploit the trade-off between convergence rate and robustness to gradient noise