

## second task of InPTA

May 16, 2022

0.1 Need to reproduce the result in table A1... using equations of A.1 and 3

objective: to reproduce the results in the first row of table A.1 which is

$$Y_{nr500}/Y_{500}$$

- defination of the Y is given as:

$$Y_{nr500}/Y_{500} = \frac{\int_0^{nr500} dr P(r) 4\pi r^2}{\int_0^{r500} dr P(r) 4\pi r^2}$$

$$P(r) = P_{500} * P_0 / (x^\gamma (1 + x^\alpha)^{(\beta-\gamma)/\alpha}) \dots \alpha, \beta, \gamma, c_{500} \text{ are given}$$

$$x = r/r_s, r_s = r_{500}/c_{500}$$

\* r500 is the radius of the sphere in which the integration is done

```
[38]: import numpy as np
import math as m
import matplotlib.pyplot as plt
from scipy import integrate
```

```
[101]: #writing the constants in the equation
c_500 = 1.156
alpha = 1.0620
beta = 5.4807
gamma = 0.3292
P_0 = 23.13
diff = (beta-gamma)/alpha

#defining r_500
r_500 = 1.15 #Mpc #1.351 (earlier value used)
#reason for taking r_500 = 1.15 for matching the values
```

the reference for the values for the r\_500 is taken to match the output in the table. I otherwise used the below mentioned paper to check for the values of the parameters and r\_500 using Table C.1

paper: The universal galaxy cluster pressure profile from a representative sample of nearby systems (REXCESS) and the Y SZ – M 500 relation

```
[97]: #function we need to integrate

def f(x):
    return x**2/(x**gamma *(1 + x**alpha)**diff)

#using quad to integrate from 0 to r_500 for denominator
res, err = integrate.quad(f, 0, r_500)

print("The numerical result is {:.f} (+-{:g})"
      .format(res, err))

#using quad to integrate from 0 to n*r_500 for denominator
n = 1
res_1, err_1 = integrate.quad(f, 0, r_500)

print("The numerical result is {:.f} (+-{:g})"
      .format(res_1, err_1))
#=====
n = 2
res_2, err_2 = integrate.quad(f, 0, 2*r_500)

print("The numerical result is {:.f} (+-{:g})"
      .format(res_2, err_2))
#=====
n = 3
res_3, err_3 = integrate.quad(f, 0, 3*r_500)

print("The numerical result is {:.f} (+-{:g})"
      .format(res_3, err_3))
```

```
The numerical result is 0.041332 (+-5.79516e-09)
The numerical result is 0.041332 (+-5.79516e-09)
The numerical result is 0.062360 (+-5.79516e-09)
The numerical result is 0.070086 (+-2.49954e-09)
```

```
[98]: print("For the n = 1, the ratio of Ynr500/Yr500 is {:.f} (+-{:g})".format(res_1/
      ↪res, err_1/err))
```

```
For the n = 1, the ratio of Ynr500/Yr500 is 1.000000 (+-1)
```

```
[99]: print("For the n = 1, the ratio of Ynr500/Yr500 is {:.f} (+-{:g})".format(res_2/
      ↪res, err_2/err))
```

```
For the n = 1, the ratio of Ynr500/Yr500 is 1.508766 (+-1)
```

```
[100]: print("For the n = 1, the ratio of Ynr500/Yr500 is {:.f} (+-{:g})".format(res_3/  
↪res, err_3/err))
```

For the n = 1, the ratio of Ynr500/Yr500 is 1.695696 (+-0.431314)