This excerpt from

Gateway to Memory.
Mark A. Gluck and Catherine E. Myers.
© 2000 The MIT Press.

is provided in screen-viewable form for personal use only by members of MIT CogNet.

Unauthorized use or dissemination of this information is expressly forbidden.

If you have any questions about this material, please contact cognetadmin@cognet.mit.edu.

gluc_su_ind.qxd 8/29/00 2:29 PM Page 439

Subject Index

Page references followed by *t* and *f* refer to tables and figures, respectively.

Ablati on, 25, 263–264
Acetylcholine (ACh), 253, 305. *See also entries under* Cholinergic
 in memory storage and retrieval, 308–310, 308f, 310f, 313–318, 314f, 316f, 317f, 336, 337t
 as neuromodulator, 306–308, 306f
 regulation of, 332–334, 333f
ACoA. *See* Anterior communicating artery
ACoA syndrome, 338–341, 339f, 340f
Acquired equivalence, 280
Acquisition
 in cortico-hippocampal model, 142–154, 152f, 153f
 definition of, 59
Activation level of node, 47f, 48–49
AD. *See* Alzheimer's disease
Afferents, 44
Aggregate predictions, in S-D model, 168
Agonists, 307
Allocortex, 216
Alzheimer's disease (AD), 294–295, 295f
 atrophy of brain in, 294, 295f, 296–302, 296f, 298f, 300f, 302
 cholinergic regulation in, 341–342
 detection of, 296
 hippocampal region and, 294–297, 295f, 296f
 and memory loss, 17, 294–295, 295f
 treatment of, 295–296
Amnesia
 anterograde, 15, 17–20, 18f, 20f, 123, 339–341, 340f
 connectionist models of, 123f, 124
 contextual learning in, 211–214
 declarative memory in, 23, 24f
 definition of, 20
 episodic memory in, 23, 24f
 event-specific, 19
 and learned irrelevance, 182–184, 183f, 184f
 learning ability preserved in, 21–24, 24f
 procedural memory in, 23
 retrograde, 18f, 19–20, 20f, 123
 training of, 181–184, 183f
 types of, 17–20, 18f
Amnesic, definition of, 20
Amygdala, 11, 11f, 13f, 337
Anatomy of brain, 11–14, 11f, 40
 cerebral cortex, 216–220, 216f, 217f, 219f
 entorhinal cortex, 260–261, 261f, 262f, 266
 hippocampal region, 259–263, 261f, 262f
 hippocampus, 11–14, 11f, 117–119, 118f, 119f
 piriform cortex, 232–234, 233f
Aneurysm, pathology and treatment of, 338–339, 339f
Animal studies, utility of, 347–348
Anoxia, 17
Antagonists, 307
Anterior communicating artery (ACoA), aneurysm in, 338–341, 339f, 340f
Anterograde amnesia, 15, 17–20, 18f, 20f
 in AcoA aneurysm, 339–341, 340f
 hippocampus and, 123
Aricept. *See* Donepezil
Aspiration, lesions by, 25, 263–265, 264f
Associative learning, animal studies, 31–39
Associative *vs.* occasion-setting aspect of context, 196, 198–199
Associative weight, 47f, 48–49. *See also* Internal layer nodes, setting weights in

gluc_su_ind.qxd 8/29/00 2:29 PM Page 440

Associative weight (*continued*)
in autoencoders, 129
cholinergic regulation and, 318–320, 319f
competition between cues for, 65, 67
and Hebbian learning, 114
in learning, 50–51, 50f, 51–61, 57f, 59f
in multi-layered models, 90–91, 91f
Asymptote, on learning curves, 58
Atrophy of brain, in Alzheimer's disease, 294, 295f, 296–297, 296f
behavioral measurement of, 297–301, 298f, 300f
in entorhinal cortex, 302
Attentional processing, in Schmajuk-DiCarlo (S-D) model, 168
Auditory cortex
primary (A1), 218, 219, 219f
training of, 251–253, 252f
Autoassociative networks, 114–117, 115f, 116f, 177f
capacity of, 122–124
cholinergic regulation in, 311–318, 311f, 312f, 314f, 316f, 317f
compression in, 129, 130f, 134–138
consolidation in, 122–124
in hippocampus, 117–118, 119f, 120
hippocampus as, 119–121, 121f, 123, 123f, 124
interference in, 122–124, 122f
pattern completion in, 115–116, 116f, 117
pattern recognition in, 116–117, 117f
runaway excitation in, 311–312, 312f
Autoencoders, 124–127, 125f–127f, 128–129, 128f
description of, 128–129, 128f
compression in, 129, 130f
predictive models, 129–138, 131f–133f, 135f, 138f
Axons, 44, 45f, 216

Backpropagation. *See* Error backpropagation
Basal forebrain, 253, 253f, 308f
cholinergic neurons in, 308–309
Best frequency, in auditory cortex, 251
Blocking effect, 62–65, 62t, 63f, 64f
in cortico-hippocampal model, 170–171, 171t, 172f
in Rescorla-Wagner model, 67–68, 67f, 73, 74f
in Schmajuk-DiCarlo (S-D) model, 170–171, 171t, 172f
stimulus selection and, 178

CA1 region
cholinergic regulation in, 313–315, 314f
regulation of acetylcholine by, 332–334, 333f
CA3 region
and acetylcholine regulation, 332–334, 333f
anatomy of, 117–119, 118f, 119f
as autoassociator, 311
cholinergic regulation in, 313–315, 314f, 316f, 317f. *See also* Cortico-hippocampal model, cholinergic regulation in
role of, 261, 265
Categorization learning, 97–99, 100f
in cortico-hippocampal model, 160
Cerebellum, 11, 11f
in cortico-hippocampal model, 148–150, 149f
as error-correction mechanism, in eyeblink conditioning, 70–74, 71f, 78
Cerebral aneurysm, pathology and treatment of, 338–341, 339f
Cerebral cortex, 11, 11f. *See also* Cortico-hippocampal model; Piriform cortex
anatomy of, 216–220, 216f, 217f, 219f
cholinergic regulation in, 315, 336–338
in memory storage, 123f
plasticity in, 220–223, 221f, 222f, 254. *See also* Competitive networks
Cholinergic agonists
and Alzheimer's disease, 341
modeling of, 321–324, 323f
Cholinergic antagonists, 309
modeling of, 319f, 320–321, 321f
Cholinergic neurons, 305
Cholinergic receptors, types of, 331
Cholinergic regulation. *See also* Acetylcholine
in Alzheimer's disease, 341–342
of cerebral cortex, 336–338
in cortico-hippocampal model, 318–326, 319f–321f, 323f, 325f, 327f, 342–343
disruption of, 305, 307, 309–310, 310f, 347
of hippocampus, 305, 311–318, 311f, 312f, 314f, 316f, 317f, 328–331, 329f
memory disorders and, 338–342, 339f
Chunking, 175

gluc_su_ind.qxd 8/29/00 2:29 PM Page 441

Classical conditioning. *See* Conditioning
Climbing fibers, in error-correction learning, 70–74, 71f
Coarse coding. *See* Distributed representation
Cognex. *See* Tacrine
Cognitive mapping
in animals, 112
hippocampus in, 180
Combinatorial explosion, in configural tasks, 97–102
Competitive networks, 223–232, 224f. *See also* Self-organizing networks
entorhinal cortex as, 267–268, 267f, 283–286, 284f, 285f
piriform cortex as. *See* Piriform cortex, models of
self-organizing feature maps, 225–232, 226f
Component representation, 81–82, 84f, 86
Compression
in autoassociative networks, 129, 130f, 134–138
in cortico-hippocampal model, 157–160, 158f, 159f, 159t, 160f
by hippocampus, 148
in hippocampus atrophy, 301
multimodal *vs.* unimodal, 279
Conditioned inhibition, 169–170, 170f, 205
Conditioned response (CR), 32, 32f
extinction of. *See* Extinction of conditioned response
Conditioned stimulus (CS), 32, 32f
Conditioning. *See also individual paradigms:* Acquisition, Blocking, Conditioned inhibition, Discrimination, Discrimination reversal, Extinction, Feature-negative discrimination, Feature-positive discrimination, Latent inhibition, Learned irrelevance, Negative patterning, Sensory preconditioning, Trace conditioning
cholinergic disruption and, 309–310, 310f
classical, 31–39
Confabulation, 339
Configural learning, 97–99, 98f
combinatorial explosion in, 97–102
definition of, 97
in S-D model *vs.* cortico-hippocampal model, 171–174, 173f, 174f
Configural networks
in humans, 101
power of, 101
Connectionism, 46
models of amnesia in, 123f, 124
Consolidation period, 20
in autoassociative networks, 122–124, 123f
GABA and, 336
Context, 189
associative *vs.* occasion-setting, 196, 198–199
in cortico-hippocampal model, 159
definition of, 177
Context shift effects, 209–210
in contextual processing, 199–201, 199f, 200f, 201f
Contextual processing
in amnesia, 211–214
associative, 193–198, 194f, 195f, 197f, 198f
in cortico-hippocampal model, 196–207, 197f, 199f–201f, 203f, 204f, 206f, 210–214, 211f
entorhinal cortex and, 275–279, 276f, 277f
hippocampus in, 177, 190, 192–193, 196, 201, 202, 207–211
implications for human memory, 211–214
latent inhibition in, 201–205, 203f, 204f
occasion-setting models, 196–207, 197f, 199f–201f, 203f, 204f, 206f
overview of, 189–193, 191f
qualitative theories and, 207–211
Controls, 25
Cortico-hippocampal model, 145–146, 148–151, 148–160, 149f, 151f
acetylcholine regulation in, 332–334
acquisition in, 142–154, 152f, 153f
blocking effect in, 170–171, 171t, 172f
cerebral cortex in, 149–150, 149f
cholinergic regulation in, 318–326, 319f–321f, 323f, 325f, 327f, 342–343
compression in, 157–160, 158f, 159f, 159t, 160f
conditioned inhibition in, 169–170, 170f
configural learning in, 171–174, 173f, 174f
contextual processing in, 196–207, 197f, 199f–201f, 203f, 204f, 206f, 210–214, 211f
differentiation in, 148, 151–157
discrimination in, 154–156, 155f, 156f
with entorhinal cortex, 266–282, 267f, 269f,

gluc_su_ind.qxd 8/29/00 2:29 PM Page 442

Cortico-hippocampal (*continued*)
270f, 272f–274f, 276f, 277f, 280f, 281f, 291–303, 293f
extinction in, 161, 162f, 209–210
latent inhibition in, 159, 202–203, 203f, 204f, 324–325, 325f
learned irrelevance in, 158–159, 159f, 159t, 160f
limitations of, 160–163
mathematical details, 186–187
memory in, 211–214, 254–256, 294–302
negative patterning in, 171–174, 173f, 174f
neurophysiological support for, 163–165, 164f
with piriform cortex, 241–258, 241f, 242f, 244f–247f, 249f
representation in, 145–146, 148–151, 149f, 151f, 169–174, 170f–174f, 186–187
vs. Schmajuk-DiCarlo (S-D) model, 169–174, 170f–174f
CR. *See* Conditioned response
Credit assignment problem, in neural network models, 55–56
Criterion performance, 187
Critical period, in topographical layout of cortex, 220
CS. *See* Conditioned stimulus

Declarative memory
in amnesia, 23, 24f
as complex memory, 30
Deep layers, in entorhinal cortex, 260
Delay conditioning
definition of, 36
hippocampus in, 34–36, 35f
Delayed nonmatch to sample (DNMS) test, 26–27, 26f
Delta rule. *See* Widrow-Hoff Learning Rule
Dendrites, 44, 45f
Dentate gyrus, 13f, 260
as input to hippocampus, 117–118, 119f, 120
cholinergic regulation in, 315
and differentiation, 165, 280–282, 281f
and perforant path, 260–261
Desired output, 53, 54f
Differentiation
in animals, 164–165, 164f
in cortico-hippocampal model, 148, 151–157
dentate gyrus and, 165, 280–282, 281f
hippocampus and, 148, 151–157
in predictive autoencoders, 134, 135f
predictive, in cortico-hippocampal model, 156–157
Discrimination
in cortico-hippocampal model, 154–156, 155f
definition of, 154
feature-positive and feature-negative, 205–207
in Schmajuk-DiCarlo (S-D) model, 168
Discrimination reversal, 155–156, 156f
Distributed representation, 86–90, 88f
error-correction learning and, 91–94, 93f
multilayered models, 90–91, 91f
one-layer models, 84–90, 88f
vs. exemplar models, 101
DNMS. *See* Delayed nonmatch to sample (DNMS) test
Donepezil (Aricept), 307, 341
Dopamine
in Alzheimer's disease, 341
in learning and memory, 334

Easy-hard transfer
in cortico-hippocampal model, 156
definition of, 156
entorhinal cortex and, 273–275, 274f
EC lesion of hippocampus, 268, 269f
EEG. *See* Electroencephalogram
Electroencephalogram (EEG), 335–336, 335f
Entorhinal cortex, 13f, 260
anatomy of, 260–261, 261f, 262f, 266
as input to hippocampus, 117–118, 119f, 120
atrophy of, 302
backprojections from, 286–291, 287f, 288f, 289f
and contextual processing, 275–279, 276f, 277f
cortico-hippocampal model, 266–282, 267f, 269f, 270f, 272f–274f, 276f, 277f, 280f, 281f, 291–303, 293f
and easy-hard transfer, 273–275, 274f
function of, 259, 284
and human memory, 294–302
information flow in, 286–287, 287f

gluc_su_ind.qxd 8/29/00 2:29 PM Page 443

and latent inhibition, 271–272, 272f, 279
and learned irrelevance, 272–273, 273f
qualitative theory and, 291–294, 293f
role of, 260, 261f, 265
Schmajuk-DiCarlo (S-D) model, 282f, 283–286, 285f
and sensory preconditioning, 273, 274f
stimulus competition in, 283–286, 283f, 285f
Episodic memory, 23, 24f
Error backpropagation algorithm, 102–110, 106f–108f, 129
Error-correction learning
anatomical location of, 70, 113
and distributed representation, 91–94, 93f
in eyeblink conditioning, 70–74, 71f, 72f
in models, 54–61, 57f, 59f, 69–78, 71f, 72f, 90–94, 93f
Errorless learning, 213
Etiologies
definition of, 16
in memory loss, 16–17
Event-specific amnesia, 19
Excitatory effect of neurotransmitters, 45
Exclusive-OR tasks
in animals, 94–96, 95f
in neural network models, 95f, 96–97, 96f
Exemplar models, 101
Extinction of conditioned response, 58–61, 60f
contextual sensitivity of, 208–210, 208f
in cortico-hippocampal model, 161, 162f, 209–210
hippocampus and, 209
Eyeblink conditioning, 33–34, 33f. *See also* Conditioning
error-correction learning in, 70–74, 71f, 72f
hippocampus in, 34–36, 35f
Schmajuk-DiCarlo (S-D) model of, 167, 166f–167f

Feature-negative discrimination, 205–207
Feature-positive discrimination, 205–207
Figure completion test, in amnesia, 22–23, 22f
Fimbria, 13, 260
Fornix, 13, 260
lesion effects of, 239–240, 242f, 243, 245–250, 245f–247f, 249f, 347
Fugue, 19

GABA (gamma-aminobutyric acid), 334–336, 335f, 337f
Generalization, 82–84, 83f. *See also* Distributed representation
challenges of, 86
distributed representation in, 89, 90f
gradient, 83, 83f
multilayered models, 90–91, 91f
one-layer models, 84–90, 84f, 85f, 86f
vs. specificity, tension between, 82, 86, 94
Granule cells, 280

Hebbian learning, 114
Hebb's rule, 114
Herpes encephalitis, amnesia in, 17
Hidden layer nodes. *See* Internal layer nodes; Multilayered models
Hierarchical clustering of inputs, in competitive network models, 234–235
Hippocampal formation, 291
Hippocampal region. *See also* Dentate gyrus; Entorhinal cortex; Fornix; Hippocampus; Subiculum
Alzheimer's disease and, 294–297, 295f, 296f
anatomy and physiology of, 259–263, 261f, 262f
cholinergic regulation of, 305
definition of, 13, 259, 260
function of, 13–17, 19–20, 215, 279–280, 280f, 283–284, 283f
information flow in, 260, 261f, 269f, 286–287, 287f
lesions of, 260, 263–265, 264f, 268–271, 269f, 270f, 303
role of, 263, 263f
Hippocampus. *See also* CA1 region; CA3 region; Cortico-hippocampal model; H lesion of hippocampus; HR lesion of hippocampus; Schmajuk-DiCarlo (S-D) model
and amnesia, 123
anatomy of, 11–14, 11f
atrophy of, 297–301, 298f, 300f
as autoassociative network, 119–120, 123, 123f, 124
CA3 anatomy, 117–119, 118f, 119f
and cholinergic regulation, 305, 311–318, 311f, 312f, 314f, 316f, 317f, 328–331, 329f

gluc_su_ind.qxd 8/29/00 2:29 PM Page 444

Hippocampus (*continued*)
in cognitive mapping, 180
in conditioning, 34–39, 35f, 37f
and contextual processing, 177, 190, 192–193, 196, 201, 202, 207–211
damage to, 21–24, 24f
differentiation in, 151–157
and extinction and renewal, 209
as information processor, 345
and latent inhibition, 202
and learning, 21–41, 24f, 29f
lesion of, *vs.* cholinergic disruption, 309–310, 310f, 347
and memory, 19–20, 120–121, 121f, 123–124, 123f, 178–184
models of, 40–41, 78, 181–184, 345–350
and piriform cortex, 234, 237–250, 238f, 239f, 241f, 242f, 244f–247f, 249f
place cells in, 28–29, 28f
representation distortions in, 146–148, 147f
Schmajuk-DiCarlo (S-D) model, 146, 165–174, 166f–167f, 170f–174f
and scopolamine, 328–331, 329f
and sensory preconditioning, 38–39, 38t, 39f, 139
and spatial navigation, 28–31, 29f
in stimulus configuration, 175–176
in stimulus selection, 177–178
in trace and delay conditioning, 34–36, 35f
and unsupervised learning, 112–113
vs. qualitative theories, 175–181
H lesion of hippocampus, 268, 269f
in cortico-hippocampal model, 269–279, 270f, 272f–274f, 276f, 277f, 303
in Schmajuk-DiCarlo (S-D) model, 284–286, 285f
HM (patient), 14–16, 15f, 19, 21–22, 23
Homunculus, 218
HR lesion of hippocampus, 268, 269f, 273, 275, 297
Hypoxia, 17

Ibotenic acid lesions, 263, 264–265, 264f
Inferior olive, in error-correction learning, 70–74, 71f, 72f, 149, 149f
Information flow
in entorhinal cortex, 286–287, 287f
in hippocampal region, 260, 261f, 269f, 286–287, 287f
in neocortex, 286–287, 287f
Information processing, in brain
mechanisms of, 44–47
models of, 40–41. *See also* Models
Information value of stimuli, 65–68. *See also* Competitive networks
Inhibitory effect of neurotransmitters, 45
Input nodes, 47, 47f
Instrumental conditioning, in cortico-hippocampal model, 160
Internal layer nodes, 90, 97–99, 98f. *See also* Multilayered models
and compression, 136–137
setting weights in, 102–109, 106f–108f, 131–133, 132f, 133f, 150
Internal recurrency, in CA3 field, 118–119
Isocortex, 216

Language development in children, 254–256
Language-learning-impaired (LLI) children, and cerebral plasticity, 255–256
Latent inhibition
in contextual processing, 201–205, 203f, 204f
in cortico-hippocampal model, 159, 202–203, 203f, 204f, 324–325, 325f
definition of, 201–202
entorhinal cortex and, 271–272, 272f, 279
and hippocampal damage, 78
in Rescorla-Wagner model, 76–78, 77f, 77t
Latent learning paradigms, 111
autoassociative networks and, 126–127, 126f–127f
Learned irrelevance
amnesia and, 182–184, 183f, 184f
in cortico-hippocampal model, 158–159, 159f, 159t, 160f, 326, 327f
definition of, 158
entorhinal cortex and, 272–273, 273f
Learning. *See also* Categorization learning; Configural learning
associative weight and, 50–51, 50f
contextual, 177, 189–193, 191f
cortical reorganization in, 220–222, 221f, 222f
errorless learning, 213

gluc_su_ind.qxd 8/29/00 2:29 PM Page 445

generalization in. *See* Distributed representation; Generalization in learning
hippocampus and, 21–41, 24f, 29f
and memory, 26–27
models of. *See* Error backpropagation algorithm; Error-correction learning; Widrow-Hoff Learning Rule
perceptual learning, 75–78
supervised, 111–112
unsupervised, 111–114, 113f, 223
Widrow-Hoff Learning Rule and, 58–59, 59f, 60f, 65–67
Learning curve, 58, 59f
Learning rate, in neural network models, 56
Learning Rule. *See* Widrow-Hoff Learning Rule
Least Mean Square Rule. *See* Widrow-Hoff Learning Rule
Lesions, 325, 325f
by aspiration, 25, 263–265, 264f
of hippocampal region, 260, 309–310, 310f, 346, 347
by ibotenic acid, 263, 264–265, 264f
limitations of studies using, 36
neurotoxic, 25, 263, 264–265, 264f, 325, 325f
by saporin, and latent inhibition, 325, 325f
LLI (Language-learning-impaired) children, and cerebral plasticity, 255–256
LMS rule. *See* Widrow-Hoff Learning Rule
Local representation. *See* Component representation
Long-term potentiation (LTP), 69–70, 114

Magnetic resonance imaging (MRI), in Alzheimer's Disease, 296, 296f
Meaning, and plasticity, 251–254, 252f
Medial septum, 305, 309
in neuromodulator regulation, 331–336, 333f
Medial temporal lobes, 12–17, 13f
Metrifonate, 322–323, 323f
Memory. *See also* Storage and retrieval
and acetylcholine, 308–310, 308f, 310f, 313–318, 314f, 316f, 317f, 336, 337t
Alzheimer's disease and, 17, 294–295, 295f
and cholinergic regulation, 338–342, 339f
in cortico-hippocampal model, 211–214, 254–256, 294–302
and entorhinal function, 294–302
episodic, 26–27
flexible, 180–181
in hippocampal damage, 21
in hippocampal lesion, 263, 263f, 309–310, 310f, 347
hippocampus and, 19–20, 120–121, 121f, 123–124, 123f, 178–184
intermediate-term, hippocampus in, 178–179
and learning, in animal studies, 26–27
long-term, in hippocampal damage, 21
loss of, causes, 16–17, 19–20
medial temporal lobes in, 13–17
and neuromodulation, 308–310, 308f, 310f
and parahippocampal region, 179
piriform cortex and, 254–256
and plasticity, 254–256
short-term, in hippocampal damage, 21
working, hippocampus in, 178–179
Memory-enhancing drugs, 322–323
Mere exposure paradigms, 111
Mispairs, 292
Models. *See also* Autoassociative networks; Autoencoders; Competitive networks; Cortico-hippocampal model; Multilayered models; Rescorla-Wagner model; Schmajuk-DiCarlo (S-D) model
basic principles of, 47–49, 68, 348–350
of hippocampus, utility of, 345–350
of learning. *See* Error backpropagation algorithm; Error-correction learning; Widrow-Hoff Learning Rule
of motor-reflex conditioning, 49–51, 50f
testability of, 349–350
utility of, 3–5, 43, 46–47, 175, 348–349
Mossy fibers, in autoassociative learning, 118, 120, 121
Motor-reflex conditioning, network models of, 49–51, 50f
Motor skill learning, and amnesia, 21–22, 24f
Multilayered models, 97–99, 98f
representational weight in, 90–91, 91f
training of, 102–109, 106f–107f, 108f
Muscarinic receptors, 331

gluc_su_ind.qxd 8/29/00 2:29 PM Page 446

Negative patterning
in animals, 94–96, 95f
in models, 95f, 96–97, 96f, 171–174, 173f, 174f
Neocortex, 216, 286–287, 287f
Neural network models. *See* Models
Neuromodulation
basics of, 306–308, 306f
disruptions of, 307
memory and, 308–310, 308f, 310f
Neuromodulators, 120, 305
acetylcholine as, 306–310, 306f, 308f, 310f
in Alzheimer's disease, 341
regulation of, 332–336, 333f
Neurons, 45f
in Alzheimer's disease, 295f
formation and alteration of, 69–70
structure and mechanism of, 44–47
Neurotoxic lesions, 25, 263, 264–265, 264f
Neurotransmitters, 44–45, 305, 306, 306f
Nicotinic receptors, 331
Nodes, 47–49, 47f. *See also* Internal layer nodes
Nonmonotonic development, 156
Nonpairs, 292
Norepinephrine, in Alzheimer's disease, 341
Novelty recognition, 126, 128, 168
Nucleus basalis, 253–254, 253f, 309, 336–338

Occasion setting, 192
and contextual processing, 196–207, 197f, 199f–201f, 203f, 204f, 206f
in phasic cues, 205–207, 206f
vs. association, 196, 198–199
Odor discrimination
simultaneous, 239f, 240, 243–250, 244f–247f, 249f, 258
successive, 239–240, 239f
Olfactory bulb, 233f, 234
Olfactory cortex. *See* Piriform cortex
Outcomes, different, from similar stimuli, 94–97, 95f, 96f
Output nodes, 47f, 48
Overtraining reversal effect, 156
Oxygen deprivation, and memory loss, 17

Paleocortex, 216
Paragraph delayed recall test, 294
Parahippocampal cortex, 13f, 259
Parahippocampal region, 179, 291
Parallel distributed processing, 46–47
Pattern recognition and completion. *See also* Storage and retrieval
in autoassociative networks, 115–117, 116f, 117f
cholinergic regulation and, 313–324, 314f, 316f, 317f, 319f–321f, 323f
Pavlovian conditioning. *See* Conditioning
Perceptrons (Minsky and Papert), 102
Perceptual learning, 75–78
Perforant path, 117–118, 260–261
Periallocortex, 260
Perirhinal cortex, 13f, 259
Phasic cues
in contextual processing, 189–190, 190f
occasion setting and, 168, 205–207, 206f
Phonemes, 228, 231
Phones, 228
Phonetic typewriter, 231–232, 231f
Physostigmine, 321–324, 323f, 341
Piriform cortex
anatomy of, 232–234, 233f
cortico-hippocampal model, 241–258, 241f, 242f, 244f–247f, 249f
hippocampus and, 234, 237–250, 238f, 239f, 241f, 242f, 244f–247f, 249f
and memory, 254–256
models of, 234–237, 235f, 236f
and qualitative theories, 250–254
Place cells, 28–29, 28f
Plaques, in Alzheimer's disease, 294
Plasticity, 70. *See also* Competitive networks; Piriform cortex, models of
in cerebral cortex, 220–223, 221f, 222f
in models, *vs.* stability, 56
and stimulus meaning, 251–254, 252f
Polymodal association cortices, 260
Pons, 149, 149f
Predictive autoencoders, 129–138, 131f, 132f, 133f, 135f
sensory preconditioning in, 138–139, 138f
Predictive differentiation, in cortico-hippocampal model, 156–157
Primary auditory cortex (A1), 218,219, 219f

gluc_su_ind.qxd 8/29/00 2:29 PM Page 447

Primary sensory cortex, topographical layout of, 218–220, 250
Primary visual cortex (V1), 218
Priming, 23
Procedural memory, 23
Psychogenic amnesia, 19

RB (patient), 16–17, 16f
Receptors, 45
Redundancy compression, 148, 157–160, 202, 236–240, 266–268, 271–273, 279–280, 297, 302
Reinforcement modulation theories, 177–178
Representation. *See also* Compression; Differentiation; Distributed representation
 component representation, 81–82, 84f, 86
 cortico-hippocampal model, 145–146, 148–151, 149f, 151f, 169–174, 170f–174f, 186–187
 distortions in, 146–148, 147f
 Schmajuk-DiCarlo (S-D) model, 146, 165–174, 166f–167f, 170f–174f
 in visual coding, 81
Representational weight, in multi-layered models, 90–91, 91f
Rescorla-Wagner model, 65–68
 applicability to animal learning, 69–78, 71f, 72f
 benefits of, 43–44
 hippocampus and, 78
 limitations of, 75–78
 mathematics of, 66
 and multilayered networks, 102
 in neural network models, with distributed representation, 92
 origins of, 61
 power of, 68–69
Retrieval from memory. *See* Storage and retrieval
Retrograde amnesia, 18f, 19–20, 20f
 hippocampus and, 123
Runaway excitation, 311–312
Runaway synaptic modification, 311–312

Saporin lesions, and latent inhibition, 325, 325f
Schmajuk-DiCarlo (S-D) model, 146, 165–169, 166f–167f
 blocking effect in, 170–171, 171t, 172f
 conditioned inhibition in, 169–170, 170f
 configural learning in, 171–174, 173f, 174f
 of entorhinal cortex, 282f, 283–286, 285f, 291–294, 293f
 negative patterning in, 171–174, 173f, 174f
 representation in, 146, 165–174, 166f–167f, 170f–174f
 vs. cortico-hippocampal model, 169–174, 170f–174f
Scopolamine, 309, 317–318
 and Alzheimer's disease, 341
 hippocampus and, 328–331, 329f
 modeling of, 319f, 320–321, 321f, 324, 326, 326f, 327f, 328–331, 329f, 347
S-D model. *See* Schmajuk-DiCarlo (S-D) model
Self-organizing feature maps, 225–232, 226f. *See also* Competitive networks
Self-organizing networks, 223. *See also* Competitive networks; Self-organizing feature maps
Semantic memory, 23, 24f
Sensory areas, in mammalian cortex, 218–220, 219f
Sensory cortex, 11, 11f, 218–220, 250
Sensory modulation theories, 177–178
Sensory preconditioning
 definition of, 38, 157
 entorhinal cortex and, 273, 274f
 hippocampus and, 38–39, 38t, 39f, 78, 139
 in predictive autoencoders, 138–139, 138f
 in Rescorla-Wagner model, 38–39, 38t, 39f, 75–76, 75f, 76f
Septohippocampal projection, 309
Serotonin, in Alzheimer's disease, 341
Sham control, 25
Simulation runs, 105
Simultaneous odor discrimination, 239f, 240
 modeling of, 243–250, 244f–247f, 249f, 258
Source amnesia, 211–212
Spatial navigation
 animal studies, 28–31, 29f
 contextual processing and, 194–196, 195f
Specificity in learning, *vs.* generalization, tension between, 82, 86, 94
Spectograms, 228–229, 228f, 229f
Speech
 phonemic analysis of, 229–230, 231–232
 spectograms of, 228–229, 228f, 229f

gluc_su_ind.qxd 8/29/00 2:29 PM Page 448

Stimulus
configuration of, hippocampus in, 175–176
information value of, 65–68. *See also* Competitive networks
representation of. *See* Representation
selection of, hippocampus in, 177–178
Stop consonants, 229
Storage and retrieval, 286–291, 287f, 288f, 289f. *See also* Memory; Pattern recognition and completion
acetylcholine (ACh) in, 308–310, 308f, 310f, 313–318, 314f, 316f, 317f, 336, 337t
in autoassociative networks, 114–117, 115f, 116f
cerebral cortex in, 123f
hippocampus in, 120–121, 121f, 123f
Stroke, 338
Subiculum, 13f, 260
Successive odor discrimination, 239–240, 239f
Superficial layers, in entorhinal cortex, 260
Supervised learning, 111–112, 223
Synapses, structure and mechanism of, 44–45, 46f
Synaptic plasticity. *See* Plasticity
Systemic administration of drugs, 328

Tacrine (Cognex), 307, 341
Tangles, in Alzheimer's disease, 294
Teaching input, 53, 54f, 69
Thalamus, 233f
Theta rhythm, 335–336, 335f
Thresholds, in multilayered models, 97
Timing effects, in cortico-hippocampal model, 161–163
Tonic cues, in contextual processing, 189–190, 190f, 193–194, 194f, 195f
Topographical layout
in distributed representation, 87
of primary sensory cortex, 218–220, 250
Trace conditioning
in cortico-hippocampal model, 161–162
definition of, 161–162
hippocampus in, 34–36, 35f
in Schmajuk-DiCarlo (S-D) model, 168

Unconditioned response (UR), 32, 32f
Unconditioned stimulus (US), 32, 32f, 111
Unimodal sensory cortices, 260
Unsupervised learning. *See* Learning, unsupervised
UR. *See* Unconditioned response
US. *See* Unconditioned stimulus

Visual cortex, 11, 11f
primary (V1), 218

Water maze, 29, 29f
Weights, internode. *See* Associative weight; Representational weight
Widrow-Hoff Learning Rule, 51–61
applicability to animal learning, 58–59, 59f, 60f, 65–67
applications of, 61
mathematics of, 53
and multilayered networks, 102
in neural network models, with distributed representation, 93f
origins of, 51–52

This excerpt from

Gateway to Memory.
Mark A. Gluck and Catherine E. Myers.
© 2000 The MIT Press.

is provided in screen-viewable form for personal use only by members of MIT CogNet.

Unauthorized use or dissemination of this information is expressly forbidden.

If you have any questions about this material, please contact cognetadmin@cognet.mit.edu.