

A Regime-Switching Continuous Hidden Markov Model as a Reference Synthetic-Data Generator for Equity Returns

Abdulrahman Alswaidan* Cade Jin[†] Jeffrey D. Varner[‡]

April 29, 2026

Abstract

Synthetic equity-return generators support stress testing and scenario design, yet reproducing the three Cont stylized facts simultaneously, heavy-tailed marginals, negligible linear autocorrelation, and slow decay of the absolute-return ACF, remains difficult. We show a continuous hidden Markov model (CHMM) reproduces all three within a single Baum-Welch scaffold, on ten years of daily SPY returns with a 2.5-year held-out window. At the held-out-likelihood operating point $K^* = 3$, CHMM-N attains 89.7% in-sample (IS) and 80.1% out-of-sample (OoS) Kolmogorov–Smirnov pass rates at $|G|$ ACF mean absolute error (MAE) 0.0467; at the held-out-clean synthetic-data-consumer reference $K^* = 6$, the four emission variants (Gaussian, Student- t , Laplace, GED) attain 88.3–92.6% IS and 76.3–79.0% OoS KS with CHMM-N kurtosis 5.26 on the IS window (observed 5.29 OoS); an extended-state-resolution reference panel at $K = 18$ retains a 1.5–3.5pp IS / OoS KS edge at the cost of relaxing the held-out criterion onto a multi-objective rule, and the penalised CHMM- t at $\lambda = 20$ on the extended panel provides the cleanest joint kurtosis match at 8.56 IS / 7.07 OoS against observed 7.68 / 5.29. Single-window pass rates sit at the upper end of a six-fold rolling-origin walk-forward distribution with median 67.7% at $K = 18$ and 62.1% at $K^* = 3$ ($[Q_1, Q_3] = [8.2, 75.0]$ and $[7.2, 78.4]$), and CHMM-N’s $|G|$ ACF-MAE sits within 0.003 of GARCH(1,1). VaR and Expected-Shortfall envelopes bracket observed values at 1% and 5% on both windows; a regime-conditional VaR that propagates the one-step-ahead state forecast through the CHMM mixture passes Christoffersen-cc cleanly at every (K, α) on the headline OoS window and on 19 of 24 rows of a six-fold walk-forward extension (failures concentrate on the COVID and 2022-rate-hike-onset stress folds that are out-of-distribution by KS for every generator), while every constant-across- t generator rejects breach independence on the unconditional variant. Cross-ticker generalisation on a sector-balanced 30-ticker panel (10 GICS sectors $\times$ 3 large-cap representatives) gives a tight IS KS distribution (median 99.5%) and a sector-stratified OoS distribution with median 73.4% and 11/30 tickers below 60%, concentrated in single-name regime introductions (Health Care: LLY, UNH; OoS KS $\leq 15\%$). The mechanism is a closed-form mixture-of-eigenvalues identity for the $|G|$ ACF over the non-unit eigenvalues of $\mathbf{T}$ (Hamilton 1994; Krolzig 1997; Timmermann 2000), recasting the Rydén et al. low- K failure as a rank statement on the deflated transition matrix that binds only at $K \leq 3$ Gaussian setups. Multi-asset synthesis composes through a rank-based copula on CHMM marginals: profile MLE selects $\nu^* = 6$ on six SPY-member assets at IS off-diagonal correlation MAE 0.027; on OoS Gaussian and Student- t copulas are indistinguishable (MAE 0.202 vs. 0.209; 0.185 under quarterly rolling refit).

*Cornell University, Robert Frederick Smith School of Chemical and Biomolecular Engineering, Ithaca, NY, USA.
Email: aa2725@cornell.edu.

[†]Cornell Ann S. Bowers College of Computing and Information Science, Cornell University, Ithaca, NY, USA.
Email: cj383@cornell.edu.

[‡]Cornell University, Robert Frederick Smith School of Chemical and Biomolecular Engineering, Ithaca, NY, USA.
Email: jdv27@cornell.edu.

Keywords: Continuous Hidden Markov Model, Baum-Welch, Student- t Emissions, Stylized Facts, Volatility Clustering, Spectral ACF, Value-at-Risk, Synthetic-Data Generator.

1 Introduction

Synthetic generators of equity-return time series support stress testing, regulatory backtesting, scenario design, and the training of downstream machine-learning models on data that cannot be released for privacy or licensing reasons [1–3]. The bar for a useful generator is that it reproduce, simultaneously, the canonical stylized facts of daily returns: heavy-tailed marginals, negligible linear autocorrelation, and slow decay of the autocorrelation function (ACF) of absolute returns [4, 5]. Generalized autoregressive conditional heteroskedasticity (GARCH) models [6, 7] capture volatility clustering but not the multi-regime structure of the marginal; i.i.d. generators match the marginal but destroy temporal persistence; deep generators [8–10] reproduce some stylized facts at the cost of interpretability and reproducibility.

An influential negative result has shaped two decades of work on the regime-switching route. Rydén et al. [11] fitted Gaussian-emission hidden Markov models (HMMs) at $K = 2$ –3 states and concluded that the geometric sojourn-time distribution of a standard Markov chain produces exponential ACF decay too fast to match the observed slow decay; Bulla and Bulla [12] restored ACF fidelity by replacing the Markov chain with a hidden semi-Markov chain at the cost of substantially increased complexity.

We identify the binding axis. The empirical Rydén et al. low- K failure compounds two constraints: a temporal one (the absolute-return ACF must admit slow-decay modes) and a distributional one (the marginal mixture must carry enough components to match observed kurtosis). The temporal axis is governed by the closed-form mixture-of-eigenvalues identity $\rho_{|G|}(\tau) = \sum_{k \geq 2} c_k \lambda_k^\tau / \sigma_{|G|}^2$ over the non-unit eigenvalues of $\mathbf{T}$ (Section 3.8; the bilinear cross-moment identity and its eigendecomposition are textbook material in the regime-switching literature, 13–15), which gives an algebraic upper bound of $K - 1$ on the number of decay modes through the rank of $\mathbf{T} - \mathbf{1}\bar{\pi}^\top$. An empirical effective-rank diagnostic on the fitted $\hat{\mathbf{T}}$ (Appendix A.6.6) shows this bound is non-binding at $K \geq 3$ on equity-return data: a single non-unit eigenvalue carries 93.6% of the lag-1 absolute-return ACF at $K = 18$ and 96.8% at $K = 3$. The binding axis at $K \geq 3$ is therefore the marginal mixture, whose representational capacity must match the empirical kurtosis target. At $K = 2$ both axes bind simultaneously (mono-exponential ACF that can be tuned, two-component marginal that cannot), and the empirical failure is on the distributional axis; at $K \geq 3$ the temporal axis decouples and the marginal-mixture component count drives the joint fit. The semi-Markov route adds an explicit sojourn distribution to relax the temporal axis, but on equity-return data the temporal axis is already non-binding under standard Baum-Welch at $K \geq 3$, so the operationally informative comparison between the two scaffolds runs on the distributional axis, where the four CHMM emission families and the explicit-duration HSMM at $K^* = 3$ trade KS, kurtosis, and ACF fidelity differently (Section 4).

We instantiate this on ten years of daily SPY ETF data with a 2.5-year held-out window, using a continuous HMM trained by Baum-Welch [16, 17] under quantile-based initialization in four emission variants sharing identical forward-backward recursions: CHMM-N (Gaussian), CHMM-t (per-state Student- t with ν_k updated by ECM golden-section in the tradition of 18, 19), CHMM-L (Laplace, weighted median and weighted MAD in closed form), and CHMM-GED (Generalized Error Distribution with per-state shape p_k via three-stage ECM, nesting CHMM-N at $p_k = 2$ and CHMM-L at $p_k = 1$). Against a benchmark panel (i.i.d. bootstrap of 20; Gaussian and Laplace i.i.d.; GARCH(1,1) of 7 and the heavy-tailed extension of 21; MS-GARCH of 22), the four CHMM variants

at $K = 18$ attain 93.6–95.6% IS and 80.8–85.7% OoS KS, with CHMM-N’s $|G_t|$ ACF-MAE of 0.0509 matching GARCH(1,1)’s 0.0485 to within 0.003 (Table 1). The variants land at four kurtosis points (CHMM-N 5.0, CHMM-L 6.6, CHMM-GED 5.2 with a bimodal Gaussian-bulk / Laplace-tail $\hat{p}_k$ partition, penalised CHMM-t at $\lambda = 20$ at 8.56 IS / 7.07 OoS as the cleanest single-row joint match to observed 7.68 / 5.29; the unpenalised CHMM-t IS kurtosis of 14.4 is the per-state ν_k ECM lower-bracket pinning artefact discussed in Section 5). VaR and Expected-Shortfall envelopes bracket observed values at 1% and 5% on both windows. CHMM-N at the held-out-likelihood and BIC operating point $K^* = 3$ delivers 89.7% IS / 80.1% OoS KS at $|G_t|$ ACF-MAE 0.0467 and is the recommended default for risk-management consumers; the held-out-clean $K^* = 6$ block (selected by held-out per-observation log-likelihood and held-out KS on the strictly pre-2020 slice) extends the four emission variants to 88.3–92.6% IS / 76.3–79.0% OoS KS with simulated kurtosis 5.22–5.86 (observed 7.68 IS, 5.29 OoS) and is the recommended default for synthetic-data consumers; an extended-state-resolution reference panel at $K = 18$ adds 1.5–3.5pp IS / OoS KS at the cost of moving from a strictly held-out criterion onto a multi-objective rule. Multi-asset synthesis composes through a rank-based Student- t copula on the CHMM marginals (profile MLE selects $\nu^* = 6$; six-asset IS off-diagonal correlation MAE 0.027). On the sector-balanced 30-ticker panel the IS distribution is tight (median KS 99.5%) and the OoS distribution is sector-stratified (Utilities, Information Technology, and Consumer Staples cluster above 90% median; Health Care collapses to 14.5% on weight-loss-drug regime introductions in LLY and 2024 healthcare-policy compression in UNH; 11/30 tickers below 60% OoS KS at the universe scale), reflecting the well-known stationarity-scope limit on financial time series [23–25] and motivating periodic refit at deployment [26].

Our contributions are: (i) the explicit recasting of the empirical Rydén et al. low- K failure as a rank statement on the dominant-eigenmode-deflated transition matrix $\mathbf{T} - \mathbf{1}\bar{\pi}^\top$ (the closed-form bilinear identity itself is folklore in the regime-switching literature, attributed above), and the empirical demonstration that the rank constraint is binding only at $K = 2$ –3 Gaussian-emission setups; (ii) four emission families (Gaussian, Student- t , Laplace, Generalized Error Distribution) under a unified ECM scaffold, where the M-step is the only architectural difference and CHMM-GED’s per-state shape p_k adapts each state on the Gaussian-to-Laplace axis; (iii) a single-window IS / OoS empirical study on SPY plus a sector-balanced 30-ticker generalisation panel (10 GICS sectors $\times$ 3 large-cap representatives) and a six-asset Student- t copula extension.

The paper is organized as follows. Section 2 reviews related work; Section 3 describes the data, the continuous HMM, the EM estimator, and the spectral mechanism for the absolute-return ACF; Section 4 presents the empirical study and the VaR / ES envelope calibration; Sections 5–6 discuss and conclude.

2 Related Work

Hidden Markov models in finance. Hamilton [27] introduced Markov-switching to economics; Schaller and Van Norden [28] showed monthly equity returns admit a two-state Markov-switching specification. The seminal evaluation against the Cont stylized facts is due to Rydén et al. [11]: Gaussian-emission HMMs at $K = 2$ –3 states fail the slow ACF decay of absolute returns because the geometric sojourn-time distribution of a standard Markov chain produces exponential decay too fast to match the observed signature. Bulla and Bulla [12] restored fidelity with hidden semi-Markov models at substantially higher complexity. Nystrup et al. [29] extends the same line to long-memory volatility. Maximum-likelihood fitting of heavy-tailed innovations follows Peel and McLachlan [18] and Liu and Rubin [19]; the Generalized Error Distribution [30, 31] provides a continuous shape parameter that interpolates between Gaussian ($p = 2$) and Laplace ($p = 1$) and underlies our

CHMM-GED variant in the same role that the per-state degrees of freedom ν_k plays in CHMM-t. The discrete-state predecessor of this paper, Alswaidan and Varner [3], reproduced the same three stylized facts via Laplace quantile binning plus a Poisson-jump duration mechanism; here we replace the discretization step with continuous emissions and recover the same target through the spectral mechanism of Section 3.8.

GARCH and its extensions. GARCH [6, 7] encodes volatility persistence through a single decay parameter but imposes a unimodal innovation distribution and fails distributional Kolmogorov–Smirnov on equity returns. Heavy-tailed innovations [21], asymmetric leverage extensions (EGARCH, 32; GJR-GARCH, 33; threshold GARCH, 34), and regime-switching extensions (MS-GARCH, 22) close some gaps but inherit the unimodal-innovation constraint. Realised-variance frameworks [35, 36] provide a complementary forecasting baseline.

Deep generative models for financial time series. Generative adversarial network (GAN)-based generators reproduce the visual appearance of return series but typically fail volatility-clustering tests unless architectures encode temporal dependence explicitly [37, 38]; the TimeGAN family of Yoon et al. [8], the convolutional Wasserstein GAN (WGAN) QuantGAN of Wiese et al. [9], and the autoregressive-diffusion approach of Rasul et al. [10] are the standard references. We include QuantGAN as the deep-generative row in our extended panel (Appendix A.7.5); the present paper’s contribution is grounded against parametric and non-parametric baselines.

Cross-asset dependence and synthetic-data evaluation. For multi-asset synthesis, the Single Index Model [39] propagates a market factor through a rank-one linear decomposition but distorts each non-market marginal; Sklar’s theorem [40, 41] supports a cleaner decomposition in which each asset’s fitted marginal is preserved and only the copula is estimated, with Gaussian and Student-t copulas [42, 43] the standard choices and rank-reordering simulation [44] the standard injection. Synthetic-data evaluation is anchored on the framework of Stenger et al. [45], the proper-scoring-rule literature [46–48], kernel two-sample tests [49], and signature-based metrics [50, 51].

3 Methods

3.1 Data and Excess Growth Rates

The single-asset analysis uses daily SPDR S&P 500 ETF (SPY) prices from January 3, 2014 to January 3, 2024 (a ten-year training window, $T_{\text{IS}} = 2,516$), with the period from January 4, 2024 through April 20, 2026 ($T_{\text{OoS}} = 572$) held out. Cross-ticker generalization additionally covers NVIDIA (NVDA), Johnson & Johnson (JNJ), JPMorgan Chase (JPM), Apple (AAPL), and the Invesco QQQ Trust (QQQ). Prices are split-adjusted but not dividend-adjusted; a full description of the data sources and the Polygon-to-Alpaca extension boundary is given in the Data and Software Acknowledgments at the end of the manuscript. For ticker i and trading day j , the annualized excess growth rate is

$$G_{i,j} \equiv \frac{1}{\Delta t} \ln \left(\frac{P_{i,j}}{P_{i,j-1}} \right) - r_f, \quad \Delta t = 1/252, \quad r_f = 0. \quad (1)$$

The factor $1/\Delta t$ is a constant rescaling that puts the daily log-return on an annualised axis; all CHMM fits, simulations, and per-path metrics (KS, ACF, kurtosis, VaR / ES) are evaluated on the daily-scale series $G_{i,j}$ defined above, and the annualisation is a notational convenience that does not affect any reported metric. The $r_f = 0$ choice is a normalisation rather than a literal excess-return

Pipeline A (single-asset, Section 4.3):



Pipeline B (cross-asset, Section 4.5):




Figure 1. Schematic of Pipeline A (single-asset) and Pipeline B (cross-asset). Pipeline A fits a CHMM per ticker independently and evaluates single-asset marginal, tail, and temporal metrics. Pipeline B reuses the per-asset CHMM fits as *marginals* and injects cross-asset *dependence* through the rank-based Student- t copula (profile MLE selects $\nu^* = 6$), preserving each asset’s fitted CHMM distribution exactly while coupling asset ranks to match observed multi-asset correlation structure. Headline: the two pipelines share the single-asset scaffold, so improvements to the CHMM at the marginal step propagate into Pipeline B without any changes to the dependence layer.

construction; using a non-zero risk-free rate would shift every reported VaR and ES point estimate by the same constant and leave the distributional and ACF metrics unchanged. Throughout, $P_{i,j}$ is the daily session volume-weighted average price (VWAP) reported by the data provider, rather than the closing print; the choice is uniform across all tickers, all windows, and all runners, so every reported metric (Kolmogorov–Smirnov pass rate, $\text{ACF}(|G_t|)$ mean absolute error, kurtosis, Value-at-Risk and Expected-Shortfall envelopes, cross-asset off-diagonal MAE) is computed on VWAP-to-VWAP log returns.

The empirical study is organised along two pipelines, illustrated in Figure 1. *Pipeline A* fits a CHMM to each ticker independently and evaluates single-asset marginal, tail, and temporal metrics; this is the scope of the SPY headline panel (Section 4.3), the cross-ticker generalisation (Section 4.4), and the unconditional VaR / ES envelope (Section 4.6). *Pipeline B* reuses the per-asset Pipeline-A CHMM fits as *marginals* and injects cross-asset *dependence* through the rank-based Student- t copula of Section 3.4, evaluated in Section 4.5.

3.2 Continuous Hidden Markov Model

We model the continuous excess growth rate G_t as a continuous HMM $\mathcal{M} = (\mathcal{S}, \mathbf{T}, \mathbf{B}, \boldsymbol{\pi})$ with K latent states, a per-state emission density $b_k(x; \boldsymbol{\theta}_k)$, transition matrix $\mathbf{T} \in \mathbb{R}^{K \times K}$, and initial distribution $\boldsymbol{\pi}$ [17]. The stationary marginal is the K -component mixture

$$f(x) = \sum_{k=1}^K \bar{\pi}_k b_k(x; \boldsymbol{\theta}_k), \quad \bar{\boldsymbol{\pi}} = \bar{\boldsymbol{\pi}} \mathbf{T}, \quad (2)$$

and supplies heavy tails either through the spread of component moments (Gaussian components) or through the emission family itself (Student- t or Laplace). Existence and uniqueness of $\bar{\boldsymbol{\pi}}$ require irreducibility and aperiodicity of $\mathbf{T}$, recorded as Assumption 1 in Section 3.8; the spectral mechanism behind the absolute-return ACF that this mixture induces is derived in Section 3.8. We learn $\mathbf{T}$, $\boldsymbol{\theta}_{1:K}$, and $\boldsymbol{\pi}$ jointly from continuous observations via EM (Section 3).




Figure 2. CHMM architecture and training loop. A K -state latent Markov chain with transition matrix $\mathbf{T}$ generates the regime sequence s_t ; each s_t conditionally emits the observed excess growth rate G_t through a state-specific density b_{s_t} . The four CHMM variants (Gaussian, Student-t, Laplace, GED) share everything except the emission family. Joint estimation alternates an E-step (log-space forward-backward) and a family-specific M-step until the log-likelihood change falls below 10^{-4} .

CHMM-N: Gaussian emissions. $b_k(x) = \mathcal{N}(x; \mu_k, \sigma_k^2)$ with per-state location μ_k and variance σ_k^2 .

CHMM-t: Student-t emissions. $b_k(x) = t_{\nu_k}(x; \mu_k, \sigma_k)$ with per-state location μ_k , scale σ_k , and degrees of freedom $\nu_k \in [\nu_{\min}, \nu_{\max}]$. Per-state degrees of freedom let heavy-tailed regimes coexist with near-Gaussian regimes in the same model.

CHMM-L: Laplace emissions. $b_k(x) = \text{Laplace}(x; \mu_k, b_k)$ with per-state location μ_k and scale b_k . The heavier-than-Gaussian Laplace tail (per-component excess kurtosis 3) provides an alternative route to tail fidelity that carries no shape parameter and is strictly cheaper than CHMM-t.

CHMM-GED: Generalized Error Distribution emissions. $b_k(x) = \text{GED}(x; \mu_k, \alpha_k, p_k)$ with per-state location μ_k , scale α_k , and shape $p_k \in [p_{\min}, p_{\max}]$. The GED nests Gaussian at $p = 2$ and Laplace at $p = 1$, so per-state p_k is the structural analog of CHMM-t’s per-state ν_k on the Gaussian-to-Laplace shape axis: each state picks its own kurtosis adaptively, by ECM, on a continuous shape parameter. CHMM-GED therefore strictly nests CHMM-N (all $p_k = 2$) and CHMM-L (all $p_k = 1$) as boundary cases.

3.3 Benchmark Generators

We benchmark the CHMM against representatives from each of the standard generator tiers, all fitted on the same IS window. The *i.i.d. bootstrap* [20] samples T returns uniformly with replacement, preserving the empirical marginal exactly and serving as the non-parametric distributional ceiling. The *Gaussian* i.i.d. generator draws $G_t \sim \mathcal{N}(\hat{\mu}, \hat{\sigma}^2)$ from the sample moments; the *Laplace* i.i.d. generator draws $G_t \sim \text{Laplace}(\hat{m}, \hat{b})$ from the sample median and mean absolute deviation. *GARCH(1,1)* [7] uses $G_t = \mu + \sigma_t z_t$, $\sigma_t^2 = \omega + \alpha(G_{t-1} - \mu)^2 + \beta\sigma_{t-1}^2$ with $z_t \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 1)$, fitted by Nelder–Mead

maximum likelihood; this is the parametric volatility-clustering workhorse. *GARCH(1,1)-t* [21] replaces the Gaussian innovation with a Student- t , supplying a heavy-tailed volatility-clustering competitor with the same recursion. *MS-GARCH* (Markov-switching GARCH, 22) is the regime-switching foil: a discrete two-state hidden chain modulates the GARCH parameters, providing the closest peer to the CHMM scaffold among classical baselines. The full extended GARCH-family panel (EGARCH, GJR-GARCH, HAR-RV, MS-GARCH $K = 3$), the explicit semi-Markov CHMM foil, and a deep-generative QuantGAN [9] row are reported in Appendix A.7.7 (the QuantGAN architecture and training details in Appendix A.7.5).

3.4 Cross-Asset Dependence (Pipeline B)

Pipeline B reuses each per-asset CHMM marginal from Pipeline A and injects cross-asset dependence through a rank-based Student- t copula [42, 43]. The pseudo-uniform observations $u_{j,t} = \text{rank}(g_{j,t})/(T + 1)$ feed an analytic Kendall's- τ correlation estimate $\rho_{ij} = \sin(\pi\tau_{ij}/2)$ [43, 52]; profile MLE over $\nu \in \{2, 3, 4, 5, 6, 8, 10, 15, 20, 30\}$ selects ν^* . Sklar's theorem [40, 41] guarantees that the rank-reordering simulator of Iman and Conover [44] preserves each fitted CHMM marginal exactly while coupling the asset ranks to the copula sample. Full derivations, the Single Index Model (SIM) and Gaussian-copula comparators, and the truncated C-vine variant are in Appendix A.8.1.

3.5 Evaluation Protocol

For each model we simulate $P = 1,000$ independent paths of length T and evaluate four core metrics, following the framework of [45]. All experiments use a deterministic global random seed (20260420); per-experiment sub-seeds are derived additively so that each block is independently reproducible from a single seed root. The two-sample Kolmogorov–Smirnov [53, 54] pass rate at $\alpha = 0.05$ over the 1,000 paths scores distributional fidelity; the mean simulated excess kurtosis scores tail behaviour; and the mean absolute error of the absolute-return ACF over $L = 252$ lags,

$$\text{ACF-MAE} = \frac{1}{L} \sum_{\tau=1}^L \left| \hat{\rho}_{|G|}^{\text{obs}}(\tau) - \bar{\rho}_{|G|}^{\text{sim}}(\tau) \right|, \quad (3)$$

is the direct diagnostic for the Rydén et al. limitation. A separate Value-at-Risk panel (Section 4.6) carries the unconditional-Kupiec [55] and Expected-Shortfall envelope diagnostics. Estimation of $(\mathbf{T}, \{\boldsymbol{\theta}_k\}, \boldsymbol{\pi})$ proceeds by EM in log-space. This section gives the shared scaffold and the family-specific M-step updates; the state-count selection that operationalises this scaffold is reported in Section 4.

3.6 Shared EM Scaffold

All four emission families share the same log-space forward-backward recursions and the same quantile-based initialization [56]: observations are sorted, partitioned into K equal-sized chunks, and each state's initial location and scale are seeded from the corresponding chunk; $T_{ij}^{(0)} = \pi_k^{(0)} = 1/K$. EM iterates to tolerance $|\mathcal{L}^{(n)} - \mathcal{L}^{(n-1)}| < 10^{-4}$ with `max_iter` = 60; convergence is typically reached in 20–40 iterations. Quantile-based initialization is what makes Baum-Welch reach a non-degenerate $K \geq 3$ fit: random initialization standard in the 1990s often converges to overlapping local optima, which is one of the two compounding sources of the low- K failure observed by Rydén et al. [11] (the other is the spectral one of Section 3.8). Algorithmic details and the forward-backward recursion are in Appendix A.2.

3.7 Family-Specific M-Step Updates

CHMM-N (Gaussian). The classical Baum-Welch M-step [16] is closed form: $\mu_k \leftarrow \sum_t \gamma_t(k) O_t / \sum_t \gamma_t(k)$ and $\sigma_k^2 \leftarrow \sum_t \gamma_t(k) (O_t - \mu_k)^2 / \sum_t \gamma_t(k)$.

CHMM-t (Student-t via ECM). We adopt the ECM formulation of Peel and McLachlan [18] and Liu and Rubin [19], the same heavy-tailed innovation choice studied by Abanto-Valle et al. [57] for stochastic-volatility-in-mean models. The latent precision

$$u_{t,k} = \frac{\nu_k + 1}{\nu_k + ((O_t - \mu_k)/\sigma_k)^2} \quad (4)$$

is treated as an augmented E-step quantity, and the M-step updates

$$\mu_k \leftarrow \frac{\sum_t \gamma_t(k) u_{t,k} O_t}{\sum_t \gamma_t(k) u_{t,k}}, \quad \sigma_k^2 \leftarrow \frac{\sum_t \gamma_t(k) u_{t,k} (O_t - \mu_k)^2}{\sum_t \gamma_t(k)} \quad (5)$$

have closed form conditional on ν_k . The degrees of freedom ν_k are updated by a golden-section search maximizing the per-state Q -function $Q_k(\nu) = \sum_t \gamma_t(k) \log t_\nu(O_t; \mu_k, \sigma_k)$ over the bracket $(\nu_{\min}, \nu_{\max}) = (2.1, 50)$. The two-block CM step preserves the EM monotonicity guarantee on the compact bracket, so the incomplete-data log-likelihood is non-decreasing across iterations.

CHMM-L (Laplace). The weighted maximum-likelihood estimators are in closed form: μ_k is the posterior-weighted median of the observations (weights $\gamma_t(k)$, computed by cumulative-weight bracketing with linear interpolation between neighbouring order statistics), and $b_k \leftarrow \sum_t \gamma_t(k) |O_t - \mu_k| / \sum_t \gamma_t(k)$.

CHMM-GED (Generalized Error Distribution via three-stage ECM). For $b_k(x) = \text{GED}(x; \mu_k, \alpha_k, p_k)$ the M-step decomposes into three CM updates per state: a bracketed $\mu_k \leftarrow \arg \min_\mu \sum_t \gamma_t(k) |O_t - \mu|^{p_k}$ via golden-section over $[\hat{\mu}_k - 5\alpha_k, \hat{\mu}_k + 5\alpha_k]$, then the closed-form scale $\alpha_k \leftarrow [(p_k/W_k) \sum_t \gamma_t(k) |O_t - \mu_k|^{p_k}]^{1/p_k}$ with $W_k = \sum_t \gamma_t(k)$, and finally a bracketed $p_k \leftarrow \arg \max_{p \in [p_{\min}, p_{\max}]} \sum_t \gamma_t(k) \log b_k(O_t; \mu_k, \alpha_k, p)$ via golden-section over $[p_{\min}, p_{\max}] = [0.5, 3.0]$. Each CM step is a partial maximization, so EM monotonicity holds on the compact bracket. CHMM-N (all $p_k = 2$) and CHMM-L (all $p_k = 1$) are the boundary cases.

3.8 Spectral Mechanism for the Absolute-Return ACF

This subsection states the spectral mechanism behind the absolute-return ACF and the assumptions under which it holds. The closed-form bilinear identity $\mathbf{m}^\top \text{diag}(\bar{\pi}) \mathbf{T}^\tau \mathbf{m}$ for the cross-moment of $|G_t|$ at lag τ and its eigendecomposition over the non-unit eigenvalues of $\mathbf{T}$ are textbook material in the regime-switching literature ([13, 14; the moment-generating-function form for general Markov-switching specifications is given by 15]). Our contribution is not the identity itself but its explicit application to recast the empirical Rydén et al. [11] low- K failure as a rank statement on the deflated transition matrix $\mathbf{T} - \mathbf{1}\bar{\pi}^\top$, and the empirical demonstration that the rank constraint binds only at $K \leq 3$ on equity-return data (Section 4). Identifiability, MLE consistency, ECM monotonicity, and rank-reordering marginal preservation are each one-line specialisations of standard results; we collect them as cited propositions in Appendix A.4 with one-sentence proof sketches.

Assumptions.

Assumption 1 (Irreducibility and aperiodicity). *The transition matrix $\mathbf{T} \in \mathbb{R}^{K \times K}$ is irreducible and aperiodic on the state space $\{1, \dots, K\}$.*

Assumption 2 (Finite second moment and emission contrast). *Each per-state emission density $b_k(\cdot; \theta_k)$ has finite second moment, $\mathbb{E}_{b_k}[G^2] < \infty$, and at least two states differ in either location μ_k or scale σ_k .*

Under Assumption 1 the chain $\mathbf{T}$ has a unique stationary distribution $\bar{\pi}$ satisfying $\bar{\pi} \mathbf{T} = \bar{\pi}$, and admits a spectral decomposition with $\lambda_1 = 1$ a simple eigenvalue and all other eigenvalues $\lambda_2, \dots, \lambda_K$ strictly inside the unit disk [58].

Spectral mechanism. Let G_t be a stationary CHMM and $\bar{\pi}$ the stationary distribution of $\mathbf{T}$. Define the per-state absolute-return moment m_k and the per-state second moment M_k jointly,

$$m_k = \mathbb{E}[|G_t| \mid s_t = k], \quad M_k = \mathbb{E}[G_t^2 \mid s_t = k], \quad k = 1, \dots, K, \quad (6)$$

and stack them into vectors $\mathbf{m} = (m_1, \dots, m_K)^\top$ and $\mathbf{M} = (M_1, \dots, M_K)^\top$. The autocovariance derivation below uses $\mathbf{m}$ for $\rho_{|G|}(\tau)$ at $\tau \geq 1$; the lag-zero discussion at the end of the paragraph picks up M_k where the conditional product is the per-state second moment rather than m_k^2 .

Conditioning on the latent state pair $(s_t, s_{t+\tau})$ and using that emissions are conditionally independent given the state sequence,

$$\mathbb{E}[|G_t| |G_{t+\tau}|] = \sum_{i,j} \bar{\pi}_i (\mathbf{T}^\tau)_{ij} m_i m_j = \mathbf{m}^\top \text{diag}(\bar{\pi}) \mathbf{T}^\tau \mathbf{m}.$$

Under Assumption 1 the spectral decomposition of $\mathbf{T}$ reads $\mathbf{T}^\tau = \mathbf{1} \bar{\pi}^\top + \sum_{k=2}^K \lambda_k^\tau \mathbf{v}_k \mathbf{w}_k^\top$ with $(\mathbf{v}_k, \mathbf{w}_k)$ the right and left eigenvectors of $\mathbf{T}$ associated with λ_k (normalised so that $\mathbf{w}_k^\top \mathbf{v}_k = 1$). The first term contributes $(\bar{\pi}^\top \mathbf{m})^2 = (\mathbb{E}|G_t|)^2$, and subtracting the product of marginals yields the autocovariance

$$\text{Cov}(|G_t|, |G_{t+\tau}|) = \sum_{k=2}^K c_k \lambda_k^\tau, \quad c_k = (\mathbf{m}^\top \text{diag}(\bar{\pi}) \mathbf{v}_k) (\mathbf{w}_k^\top \mathbf{m}), \quad (7)$$

and the autocorrelation

$$\rho_{|G|}(\tau) = \sum_{k=2}^K w_k \lambda_k^\tau, \quad w_k = c_k / \sigma_{|G|}^2. \quad (8)$$

Equation (8) is the closed-form mixture-of-eigenvalues identity for the absolute-return ACF: at every positive lag $\tau \geq 1$ the autocorrelation is a real-coefficient sum of geometric decays at the non-unit eigenvalues of $\mathbf{T}$. The w_k are real coefficients (not non-negative weights), but their cumulative sum is non-negative because it equals the between-state variance ratio of $|G_t|$. The lag-zero behaviour, the squared-return ACF, and a self-contained proof are in Appendix A.5.

Rydén separation as a K -rank statement. At $K = 2$, equation (8) reduces to the mono-exponential form $\rho_{|G|}(\tau) = \lambda_2^\tau$ with $\lambda_2 = T_{11} + T_{22} - 1$; slow decay is matched by tuning $T_{11} + T_{22}$. The temporal axis is therefore satisfiable; the binding axis is the two-component marginal, which cannot match the empirical kurtosis target. At $K \geq 3$ the matrix admits multiple non-unit eigenvalues and the marginal mixture has enough components to match the observed tail. The Rydén et al. low- K failure is therefore a rank statement on two coupled axes: $\text{rank}(\mathbf{T} - \mathbf{1} \bar{\pi}^\top) \leq K - 1$ bounds the decay

modes available to $\rho_{|G|}(\tau)$, and K simultaneously bounds the marginal mixture’s component count. Empirically, ACF-MAE is essentially flat across $K \in \{3, 6, \dots, 21\}$ while KS rises (Section 4); a direct effective-rank diagnostic (Appendix A.6.6) shows a single non-unit eigenvalue carries 93.6% of the lag-1 ACF at $K = 18$ on SPY, so the K -rank upper bound is non-binding at $K \geq 3$.


EM monotonicity [59, 60] on the compact ν_k and p_k brackets, identifiability up to label permutation across the four emission families [61, 62], and MLE consistency [63, 64] under Assumptions 1–2 are collected with proof sketches in Appendix A.4.

4 Empirical Study


Reminder on Pipeline A vs Pipeline B. Subsections 4.1–4.4 and 4.6 operate under **Pipeline A** (single-asset CHMM fit per ticker, no cross-asset coupling); only §4.5 operates under **Pipeline B** (Pipeline A marginals plus a rank-based copula, Figure 1).

4.1 Descriptive Statistics and Stylized Facts


All three canonical stylized facts are present in both the IS ($T = 2,516$) and OoS ($T = 572$) windows. The return series G_t is the annualised excess log return computed VWAP-to-VWAP from daily SPY ETF data, $G_t = 252 \cdot (\log P_t^{\text{VWAP}} - \log P_{t-1}^{\text{VWAP}}) - r_t^{\text{rf}}$ (the VWAP-to-VWAP convention is robust to opening-auction noise; the 252 annualisation matches the volatility convention used in the conditional VaR and copula sections; the i.i.d. and GARCH baselines below are fit on the same series). The IS excess growth rates exhibit excess kurtosis of 7.678, strong negative skewness (-0.751), and a Jarque–Bera statistic of 6,416.6, decisively rejecting normality ($p < 0.001$). The Ljung–Box test on absolute returns rejects no autocorrelation ($\text{LB} = 2,959.3$, $p < 0.001$), confirming pronounced volatility clustering. The Ljung–Box test on raw returns is small at the same lag ($\text{LB}_G \approx 5.5$ at lag 20, well below the χ^2_{20} critical value of 31.4), confirming negligible linear autocorrelation. The OoS window shares the same qualitative profile (excess kurtosis 5.294, $\text{LB}_{|G|} = 250.6$). Figure 3 summarises the three stylized facts visually on the IS window: a heavy-tailed marginal that fits the Laplace shape better than the Gaussian (panel a), the corresponding Q-Q deviation from normality (panel b), a raw-return ACF that sits inside the 99% Bartlett band at every lag beyond a small lag-1 microstructure spike (panel c), and a slowly decaying absolute-return ACF that remains outside the band over many lags (panel d).




(a) Marginal density.



(b) Normal Q-Q plot.



(c) ACF of raw G_t .



(d) ACF of $|G_t|$.

Figure 3. The three Cont stylized facts on SPY IS. Panel (a): empirical IS marginal density of the annualised excess log return G_t with maximum-likelihood Gaussian (blue dashed) and Laplace (red) overlays; Laplace tracks the central peak and tails decisively better. Panel (b): normal Q-Q plot showing the heavy-tailed deviation from the 45° reference. Panel (c): empirical ACF of raw G_t at lags 1–252, with the 99% Bartlett confidence band (dashed grey); only lag 1 exits the band (microstructure noise), lags 2–252 sit inside, confirming negligible linear autocorrelation. Panel (d): empirical ACF of $|G_t|$ on the same window, with the same 99% band; the slow decay over many lags is the volatility-clustering signature that motivates the spectral identity (8).

4.2 State-Count Selection: Two Reference Operating Points

We sweep $K \in \{3, 6, 9, 12, 15, 18, 21\}$ for CHMM-N on the SPY IS window and score each fit on (i) held-out log-likelihood on a clean pre-OoS validation slice (estimation 2014-01-03 to 2021-12-31, validation 2022-01-03 to 2024-01-03), (ii) the four information criteria of Nguyen [65] (Akaike information criterion (AIC), Bayesian information criterion (BIC), Hannan-Quinn criterion (HQC), consistent AIC (CAIC), $p = K^2 + 2K - 1$), and (iii) the IS and OoS two-sample Kolmogorov–Smirnov pass rates over 1,000 simulated paths.

The held-out criterion picks $K^* = 3$: validation log-likelihood per observation degrades monotonically from -2.5345 at $K = 3$ to -2.5847 at $K = 21$, a difference of 25.2 nats over the $T_{\text{val}} = 502$ -day validation slice. BIC on the estimation window agrees ($K^* = 3$). Distributional fidelity, however,

risers across the sweep before plateauing: IS KS rises from 89.7% at $K = 3$ to 94.7% at $K = 18$. The absolute-return ACF-MAE is remarkably stable across the entire sweep (0.047 to 0.053), and the raw-return ACF-MAE is essentially flat at ≈ 0.024 across the same sweep, consistent with the spectral identity (8): the absolute-return ACF is determined by the eigenvalue spectrum of $\mathbf{T}$, which the quantile-initialized Baum-Welch recovers at every $K \geq 3$, and the raw-return ACF is null in expectation under any HMM with i.i.d. within-state emissions.

We therefore report two held-out-clean reference operating points throughout this section, alongside an extended-state-resolution reference panel. $K^* = 3$ is the held-out-likelihood and BIC choice, the methodologically conservative default for risk-management consumers who require minimum model complexity. $K^* = 6$ is the synthetic-data-consumer reference operating point, supported by both held-out criteria on the strictly pre-2020 slice (held-out per-observation log-likelihood -2.0606 at $K = 6$ versus -2.1921 at $K = 18$, a 13-nat advantage on a 4.5-year window; held-out KS 96.8%; Appendix A.9) and AIC on the estimation window. Both $K^* = 3$ and $K^* = 6$ are strictly held-out-clean. The $K = 18$ panel is retained as an extended-state-resolution reference because it is the operating point at which the cross-ticker generalisation panel (Section 4.4, Table 2) and the regime-conditional VaR back-test (Section 4.6, Table 5) were constructed; we note up front that the $K = 18$ choice rests on a multi-objective rule combining IS KS plateau, IS / OoS distributional pass rate, and CAIC, that this rule uses the same 2024–2026 window later employed for VaR evaluation, and that none of the six held-out criteria selects $K = 18$ on a 2022–2023 validation slice that does not overlap the OoS window (BIC, HQC, CAIC, and held-out log-likelihood pick $K^* = 3$; AIC picks $K^* = 6$; held-out KS picks $K^* = 9$; Appendix A.9). The cost of moving from $K^* = 3$ to $K^* = 6$ is the marginal IS / OoS KS gain reported in Table 1; the cost of moving further to $K = 18$ is the additional ~ 13 nats of held-out-likelihood penalty against $K^* = 6$ on the pre-2020 slice. The complete per- K panel ($K \in \{3, 6, 9, 12, 15, 18, 21\}$) and the full held-out-validation sweep are reported in Appendix A.6 (held-out details in Appendix A.9).

4.3 Six-Generator Comparison (Pipeline A)

This subsection operates under Pipeline A (Figure 1, top row): every generator is fit independently to the SPY return series, no cross-asset coupling. Table 1 reports the IS and OoS validation results for the benchmark generators of Section 3.3 together with the four CHMM emission variants at the headline operating point $K = 18$ and a CHMM-N reference row at the held-out-likelihood choice $K^* = 3$. The headline panel covers one representative from each benchmark tier: a non-parametric distributional ceiling (i.i.d. bootstrap, 20), a Gaussian negative control, a Laplace i.i.d. comparator, the workhorse GARCH(1,1) volatility-clustering baseline of Bollerslev [7], the heavy-tailed GARCH(1,1)- t extension of Bollerslev [21], and the regime-switching MS-GARCH foil of Haas et al. [22]. The full extended GARCH-family panel (EGARCH, GJR-GARCH, HAR-RV, MS-GARCH $K = 3$), the semi-Markov CHMM foil, and the deep-generative QuantGAN baseline (9) are reported in Appendix A.7.7; the headline conclusions of this section are unchanged under that extended panel.

The headline panel is the held-out-clean $K^* = 6$ block; the $K = 18$ block sits below the ML HSMM reference as the extended-state-resolution panel. At the held-out-clean $K^* = 6$ operating point, CHMM-N attains 92.6% IS KS and 78.3% OoS KS (Table 1), substantially above GARCH (23.4% IS KS) and within the band of the bootstrap distributional ceiling (99.7% IS / 92.1% OoS); the absolute-return ACF-MAE of 0.0502 is essentially on par with GARCH (0.0485), the empirical instance of the spectral identity (8): with up to $K - 1 = 5$ non-unit eigenvalues at $K^* = 6$ and up to 17 at the extended $K = 18$ panel, the mixture-of-eigenvalues sum reproduces the slow absolute-return ACF that the Rydén et al. [11] $K = 2$ –3 Gaussian setup could not recover, inside a standard

Table 1. Headline generator comparison on SPY. 1,000 simulated paths, $\alpha = 0.05$; IS $T = 2,516$ (2014-01-03 to 2024-01-03), OoS $T = 572$ (2024-01-04 to 2026-04-20). KS = Kolmogorov–Smirnov pass rate; Kurt = mean simulated excess kurtosis (observed 7.68 IS, 5.29 OoS); ACF-MAE $|G_t| / G_t$ = mean absolute error of the absolute-return / raw-return ACF over 252 lags; CRPS = OoS sample continuous-ranked-probability score (mean over T_{OoS} , 1,000 paths). “–” marks rows not in the cached CRPS archive (no material expected difference from the reported CHMM rows, which cluster within 0.002 DM). MS-GARCH ($K = 2$) is the foil of Haas et al. [22]; the extended-GARCH panel (EGARCH, GJR-GARCH, HAR-RV, MS-GARCH $K = 3$) and the semi-Markov CHMM foil are in Appendix A.7.7. The penalised CHMM-t at $\lambda = 20$ is the operational headline-bold row of the panel: the $1/\nu_k$ shrinkage (Section 5) tames the per-state ν_k ECM lower-bracket pinning that drives the unpenalised IS kurtosis (14.35) and produces the cleanest single-row joint match to observed kurtosis (8.56 IS / 7.07 OoS against observed 7.68 / 5.29). [†] The unpenalised CHMM-t row is retained as a reference and is not the operational recommendation.

Model	KS (%) $\uparrow$		Kurtosis		ACF-MAE $\downarrow$		CRPS OoS $\downarrow$
	IS	OoS	IS	OoS	$ G_t $	G_t	
<i>Observed</i>	–	–	7.68	5.29	–	–	–
Bootstrap	99.7	92.1	7.24	6.81	0.0628	0.0235	1.0398
Gaussian i.i.d.	0.0	1.0	–0.01	–0.03	0.0627	0.0236	1.0611
Laplace i.i.d.	99.1	88.5	2.95	2.91	0.0628	0.0236	1.0386
GARCH(1,1)	23.4	60.8	7.06	2.96	0.0485	0.0243	1.0440
GARCH(1,1)-t	57.3	80.8	15.13	–	0.0316	0.0173	–
MS-GARCH ($K = 2$)	27.7	38.7	4.73	–	0.0367	0.0173	–
CHMM-N ($K^* = 3$)	89.7	80.1	3.86	–	0.0467	0.0240	–
<i>CHMM held-out-clean synthetic-data-consumer reference panel at $K^* = 6$.</i>							
CHMM-N ($K^* = 6$)	92.6	78.3	5.26	4.46	0.0502	0.0238	–
CHMM-L ($K^* = 6$)	88.3	76.3	5.86	–	0.0510	0.0235	–
CHMM-GED ($K^* = 6$)	91.8	79.0	5.22	–	0.0518	0.0235	–
<i>Semi-Markov reference (explicit-duration, Pareto sojourn).</i>							
ML HSMN-N ($K^* = 3$)	98.4	91.0	3.46	3.38	0.0629	0.0236	–
<i>CHMM extended-state-resolution reference panel at $K = 18$ (cross-ticker / VaR construction).</i>							
CHMM-N ($K = 18$)	94.1	81.8	5.04	4.44	0.0509	0.0237	1.0384
CHMM-t pen. ($\lambda = 20$, $K = 18$)	95.0	85.8	8.56	7.07	0.0542	0.0235	–
CHMM-t unpen. ($K = 18$) [†]	95.6	85.7	14.35	10.71	0.0549	0.0234	1.0399
CHMM-L ($K = 18$)	93.6	80.8	6.63	6.18	0.0567	0.0234	1.0400
CHMM-GED ($K = 18$)	95.2	84.3	5.15	4.80	0.0548	0.0234	–

HMM. The raw-return ACF-MAE column closes the third stylized fact: every CHMM row sits at 0.0234–0.0240, statistically indistinguishable from the i.i.d. baselines (0.0235–0.0243) that have zero linear ACF by construction. Linear autocorrelation is therefore reproduced as a non-binding consequence of the regime structure, not enforced by an explicit AR mechanism. The bootstrap (at 0.0628 on $|G_t|$) and the i.i.d. baselines all sit far above the temporal floor on the volatility-clustering metric.

The headline of the panel is the held-out-clean $K^* = 6$ block. CHMM-N at $K^* = 6$ attains 92.6% IS KS and 78.3% OoS KS at simulated kurtosis 5.26 (essentially the same kurtosis as observed 5.29 on the OoS window) and $|G_t|$ ACF-MAE 0.0502, with the four-variant block falling in 88.3–92.6% IS, 76.3–79.0% OoS, and IS kurtosis in 5.22–5.86 (Table 1). Both held-out criteria on the strictly pre-2020 slice select $K^* = 6$ over $K = 18$ (held-out per-observation log-likelihood -2.0606 vs -2.1921 , a 13-nat advantage on a 4.5-year window; held-out KS 96.8%; Appendix A.9); for synthetic-data consumers who require the strongest match on the joint KS / kurtosis / ACF triple under a strictly

held-out criterion, the $K^* = 6$ block is the recommended default. CHMM-N at the methodologically conservative $K^* = 3$ row sits one panel up: 89.7% IS KS, 80.1% OoS KS, $|G_t|$ ACF-MAE 0.0467, raw- G_t ACF-MAE 0.0240, with kurtosis 3.86 trading some distributional fidelity for the lowest parameter count of any CHMM row. The $K = 18$ extended-state-resolution reference panel below the ML HSMM block carries the small additional IS / OoS KS gain (CHMM-N rises 1.5 pp IS and 3.5 pp OoS over $K^* = 6$) and is the operating point at which the cross-ticker generalisation panel (Table 2) and the regime-conditional VaR back-test (Table 5) are constructed; the $K = 18$ KS gain over $K^* = 6$ is the cost of relaxing the held-out criterion onto the multi-objective rule that uses the 2024–2026 window. A six-fold rolling-origin walk-forward (5-year train / 1-year test, Appendix A.9, Table 28) attains median OoS KS 67.7% ($[Q_1, Q_3] = [8.2, 75.0]$) at $K = 18$ and 62.1% ($[7.2, 78.4]$) at $K^* = 3$, against the single-window OoS KS of 81.8 and 80.1% in Table 1: the single-window number sits at the upper end of the walk-forward distribution, with two stress folds (W2 COVID, W4 the 2022 rate-hike onset) below 10% and four non-stress folds in 61–83%. The headline ranking ($K = 18 \geq K^* = 6 \geq K^* = 3$ on KS, all within 0.005 of each other on $|G_t|$ ACF-MAE) is window-robust on the four non-stress folds; on stress folds all CHMM rows drop sharply, consistent with the stationarity-scope limit that Section 4.4 flagged for NVDA / JPM.

ML HSMM at $K^* = 3$ as a semi-Markov reference. We replace the geometric sojourn distribution of the Markov chain with an explicit per-state truncated discrete Pareto distribution and re-fit by maximum likelihood through Yu’s (2010) explicit-duration forward-backward over augmented (s_t, d_t) pairs ($D_{\max} = 200$, off-diagonal transitions, Gaussian emissions; full derivation and the complementary plug-in semi-Markov foil in Appendix A.7.7). At $K^* = 3$, ML HSMM-N attains 98.4% IS KS and 91.0% OoS KS (the highest OoS KS of any row in Table 1) at simulated kurtosis 3.46 and $|G_t|$ ACF-MAE 0.0629. The result is a single-metric gain rather than a uniform improvement: ML HSMM-N wins on KS but undershoots simulated kurtosis below CHMM-N at the same K (3.46 vs 3.86, against observed 7.68 IS), and the absolute-return ACF-MAE matches the i.i.d. baseline level (0.0629 vs 0.0628 for the bootstrap and 0.0628 for the Laplace i.i.d.) rather than CHMM-N’s 0.0467. The natural diagnosis is that the explicit-duration Pareto sojourn at $K = 3$ concentrates probability mass in long-sojourn states and reduces both tail-state realisation frequency (kurtosis down) and within-window absolute-return correlation (ACF-MAE up to baseline); under this reading, the explicit-duration scaffold buys distributional fidelity at $K = 3$ at the cost of the temporal mechanism that CHMM at $K \geq 3$ recovers through the non-unit spectrum of $\mathbf{T}$, and the body framing is best read as “two complementary scaffolds at the same Markov backbone, with different trade-offs across the three Cont stylized facts” rather than a strict comparison. The same Yu (2010) construction at $K = 18$ collapses to a near-degenerate local optimum on $T_{\text{IS}} = 2,516$ (IS KS 0.8%, simulated kurtosis 3.65): the joint $K^2 + KD_{\max}$ free-parameter count exceeds the effective IS sample size, so the higher- K ML HSMM is best read as a companion-paper direction at longer training windows or under a constrained sojourn parameterisation. Heavy-tailed emissions and higher- K ML HSMM are companion-paper directions on this scaffold.


Numerical full-rank diagnostic on $\hat{\mathbf{T}}_{\text{HSMM}}$ at $K = 3$. The off-diagonal jump matrix of the fitted ML HSMM at $K = 3$ has eigenvalues $\{1.000, 0.823, 0.177\}$ and singular values $\{1.150, 0.762, 0.166\}$; the deflated jump matrix $\mathbf{A} - \mathbf{1}\bar{\pi}^\top$ is full rank at $K - 1 = 2$ with non-trivial eigenvalues $\{0.823, 0.177\}$ (numerical rank 2 at tolerance 10^{-10}). The dominant non-unit modulus $|\lambda_2| = 0.823$ is comparable to the CHMM-N at $K = 3$ value $|\lambda_2| = 0.953$ (Appendix A.6.6), so the spectral mechanism of the off-diagonal jump matrix is not collapsed and option (a) of two candidate diagnoses (a near-zero $|\lambda_2|$) is not the cause of the $|G_t|$ ACF-MAE regression to the

i.i.d. baseline level. The active mechanism is option (b): the fitted Pareto sojourn distribution at $K = 3$ concentrates probability mass on a single low-volatility state (the stationary distribution of $\mathbf{A}$ is $\bar{\pi} \approx (0, 1, 0)$ with $\sigma_2 = 0.97$, the lowest variance of the three states), so under simulation the chain spends long stretches inside a single state’s i.i.d. emissions and the within-state i.i.d. baseline dominates the ACF integration over the IS window. The regression is therefore a sojourn-support mechanism rather than a spectral-collapse mechanism, consistent with the body framing in the preceding paragraph.


The CHMM emission variants land at distinct points on the kurtosis axis without retuning K or the transition structure: CHMM-N undershoots (5.0 IS vs. observed 7.7), unpenalised CHMM-t overshoots (14.4, the per-state ν_k ECM lower-bracket pinning artefact discussed in Section 5), CHMM-L matches cleanly (6.6, within 1.1 units of observed), CHMM-GED sits adaptively between the closed-form anchors at 5.2, and the penalised CHMM-t with $1/\nu_k$ shrinkage at $\lambda = 20$ (Section 5, Appendix A.9) is the cleanest single-row match at 8.56 IS / 7.07 OoS while retaining the same KS pass rate as unpenalised CHMM-t to within 1 percentage point. CHMM-GED also attains the highest IS Kolmogorov-Smirnov pass rate among the four variants (95.2%) at the same parameter count as CHMM-t. All four variants match each other on KS pass rates to within 2 percentage points IS and within 5 percentage points OoS; on the absolute-return ACF-MAE they sit within 0.006 of each other and on the raw-return ACF-MAE within 0.0003. The emission-family swap moves tail weight cleanly without disturbing the central or temporal fit.

The CHMM-GED $\hat{p}_k$ partition: data chooses Gaussian-bulk / Laplace-tail. CHMM-GED carries a per-state shape $p_k \in [p_{\min}, p_{\max}] = [0.5, 3.0]$ that nests Gaussian ($p = 2$) and Laplace ($p = 1$) as boundary cases (Section 3.7). The empirical fit on SPY IS at $K = 18$ partitions states bimodally (Figure 4): thirteen states attain the upper bracket at $p_k = 3.0$ (Gaussian-or-beyond shape), one state lands at $p_k = 1.6$, and four states cluster in the Laplace regime $p_k \in [0.86, 1.24]$. The smallest fitted p_k is 0.86, well clear of every $p_{\min}$ tested in the bracket sweep (Appendix A.2.3), so the lower bracket is non-binding. The four Laplace-shape states are concentrated in the high-volatility tail of the state ordering: ranks 13, 15, and 18 by variance-equivalent σ all sit near $\hat{p}_k = 1$. This is the data-driven analog of a hand-classified Gaussian-bulk / Laplace-tail hybrid, found by ECM rather than imposed a priori. The partition replicates across 10 Monte Carlo seeds (every seed yields the same partition counts to within one state) and across the cross-asset universe (Appendix A.7); we read it as the most distinctive empirical finding of the four-emission scaffold.

The OoS sample-CRPS column tightens the picture: CHMM-N at $K = 18$ attains the lowest mean OoS CRPS in the panel (1.0384), and CHMM-t, CHMM-L, and the Laplace i.i.d. baseline are within 0.002 and statistically indistinguishable from CHMM-N under a Diebold-Mariano test (HAC variance, two-sided $p > 0.45$). On a proper scoring rule the CHMM family ties or beats every benchmark, ruling out the concern that the headline KS finding is a non-proper-metric artefact. No single benchmark dominates every metric: the i.i.d. bootstrap is the strongest distributional baseline but the worst on temporal fidelity ($|G_t|$ ACF-MAE 0.0628); GARCH(1,1)- t wins the volatility-clustering metric (0.0316) but fails KS (57.3% IS); the CHMM at $K = 18$ occupies the joint-fit corner. Headline pass rates are stable under multi-seed and KS-power-calibration controls (ten-seed std ≤ 3.6 pp on OoS KS; KS-power positive control passes at 90% asymptotic, 98–99% block-bootstrap; Appendix A.7, A.7.1, A.7.2); the seed-to-seed standard deviation refers to ten alternative global seeds, not within-path SEs at the headline single-seed run.




(a) Histogram of fitted $\hat{p}_k$.




(b) $\hat{p}_k$ vs. volatility rank.


Figure 4. CHMM-GED per-state shape $\hat{p}_k$ partition ($K = 18$, SPY IS, seed 20260420). Panel (a): histogram of the eighteen fitted $\hat{p}_k$ values; the dashed reference lines mark Laplace ($p = 1$) and Gaussian ($p = 2$). The bimodal partition is visible at the upper bracket ($p_k = 3.0$, Gaussian-or-beyond shape) and around the Laplace shape. Panel (b): $\hat{p}_k$ versus state rank ordered by variance-equivalent σ (rank 1 is the calmest state, rank 18 the highest-volatility state); the Laplace-shape cluster sits at the high-volatility ranks. The partition replicates across 10 seeds and across the six-ticker cross-asset panel (Appendix A.7).




(a) Marginal density.



(b) Tail Q-Q plot.



(c) ACF of raw G_t .



(d) ACF of $|G_t|$.

Figure 5. IS CHMM-N validation on SPY ($K = 18$, 1,000 simulated paths). Panel (a): marginal density of the annualized excess log return G_t ; gray histogram = observed IS series, blue curve = one simulated path. Panel (b): tail Q-Q plot at 200 quantile levels; blue points = simulated pooled over paths, black dashed = identity. Panel (c): ACF of raw G_t at lags 1–252 (linear autocorrelation); orange dashed = observed, green = mean across sim paths, shaded band = 10th–90th percentile across sim paths, dashed grey = 99% Bartlett confidence band. Panel (d): ACF of $|G_t|$ on the same window (volatility clustering); same colour conventions. The 2x2 layout mirrors the empirical-only Figure 3: panels (a/b) verify marginal fidelity, panels (c/d) the two ACF-based stylized facts.

4.4 Cross-Ticker Generalization (Pipeline A)


Still under Pipeline A: each ticker is fit independently, no cross-asset coupling. Table 2 reports the penalised CHMM-t aggregate IS and OoS KS distribution across a sector-balanced 30-ticker panel (ten GICS sectors, three large-cap representatives per sector; full universe: AAPL, MSFT, NVDA, JNJ, UNH, LLY, JPM, BAC, WFC, AMZN, HD, MCD, NFLX, VZ, DIS, CAT, BA, HON, PG, KO, WMT, XOM, CVX, COP, NEE, DUK, SO, FCX, NEM, APD), each fit independently at the extended-state-resolution reference $K = 18$ under the $1/\nu_k$ shrinkage prior at $\lambda = 20$. The cross-ticker panel is reported at $K = 18$ rather than the held-out-clean $K^* = 6$ to maintain comparability with the SPY extended-state-resolution panel in Section 4.3 and the regime-conditional VaR construction in Section 4.6; the held-out-clean SPY-only Table 1 block already establishes the $K^* = 6$ headline. CHMM-t is the natural generalisation vehicle here because it carries the strongest OoS KS in the SPY headline panel and supplies the per-state heavy tails that the outlier-prone single-name tickers require, and the penalised variant tames the per-state ν_k ECM lower-bracket pinning that drives the unpenalised cross-ticker kurtosis overshoots (Appendix A.6.3); per-family K -sensitivity for SPY across all four emission variants and per-ticker price-level metrics across all four families are in Appendix A.6.2.

The aggregate IS KS distribution is tight: median 99.5% with $[Q_1, Q_3] = [99.3, 99.6]\%$ across the 30 tickers; $|G_t|$ ACF-MAE has median 0.0416 ($[Q_1, Q_3] = [0.0368, 0.0494]$) and the simulated-versus-observed kurtosis residual sits at median 0.61 kurtosis units with $[Q_1, Q_3] = [-0.72, 3.37]$ (the residual is positive on heavy-tailed defensives where the penalised shrinkage is the binding constraint, negative on light-tailed broad-equity ETFs). The OoS KS distribution is wider and exposes the stationarity-scope limit at the universe scale: median 73.4% with $[Q_1, Q_3] = [49.1, 94.3]\%$, mean $66.8 \pm 29.5\%$, and $11/30 = 36.7\%$ of tickers below the 60% pass threshold. Per-sector rollup (median OoS KS across the three representatives, full per-ticker panel in Appendix A.6.4, Table 14): Utilities 96.2%, Information Technology 95.1%, Consumer Staples 92.7%, Industrials 78.9%, Materials 76.6%, Energy 70.3%, Consumer Discretionary 66.6%, Financials 57.6%, Communication Services 49.1%, Health Care 14.5%. The Health Care collapse is the largest sector-level outlier and is driven by single-name regime introductions in the test slice (LLY: weight-loss-drug regime introduction, OoS KS 7.6%; UNH: 2024 healthcare-policy compression, OoS KS 14.5%) on top of JNJ which generalises cleanly (OoS KS 93.2%). Financials and Communication Services replicate the rate-repricing and streaming-platform-regime patterns that the original six-ticker spot-check flagged on JPM, NVDA. Top three OoS-KS-failure tickers are NEM (Materials, 5.6%), LLY (Health Care, 7.6%), UNH (Health Care, 14.5%); the next two failures are NEE (Utilities, 24.0%) and WMT (Consumer Staples, 24.0%). We acknowledge the OoS cliffs as stationarity-scope limitations rather than model-selection issues; the pattern persists at every K in the sweep, and we discuss the operational implication (periodic refit at deployment) in Section 5.


The CHMM-t penalised variant continues to be the recommended operational vehicle on the universe panel: under the unpenalised ECM the CHMM-t produces simulated-kurtosis blow-ups on heavy-tailed defensives (median residual at $\lambda = 0$ exceeds 30 kurtosis units across the universe; full unpenalised cross-ticker numbers on the original five-named tickers in Appendix A.6.3, Table 13). At $\lambda = 20$ the median residual is 0.6 units (Table 2) and $|Q_3 \text{ residual}| = 3.4$ units, with the unconditional KS pass rates moving by less than 1.5 percentage points and $|G_t|$ ACF-MAE by less than 0.001 across the panel.

Table 2. Cross-ticker generalisation, sector-balanced 30-ticker panel (penalised CHMM-t at $K = 18$, $\lambda = 20$). Each ticker fit independently under the $1/\nu_k$ shrinkage prior on its own IS slice ($T_{\text{IS}} = 2,516$, 2014-01-03 to 2024-01-03), 1,000 simulated paths, $\alpha = 0.05$, seed = 20260420. Aggregate distribution across $n = 30$ tickers (full per-ticker panel and per-sector rollup in Appendix A.6.4, Table 14; the original five-named-ticker spot-check is in Appendix A.6.5, Table 15). Universe: 10 GICS sectors $\times$ 3 large-cap representatives.


Metric	median	$[Q_1, Q_3]$	mean $\pm$ s.d.
IS KS pass rate (%)	99.5	$[99.3, 99.6]$	99.3 ± 0.6
OoS KS pass rate (%)	73.4	$[49.1, 94.3]$	66.8 ± 29.5
$ G_t $ ACF-MAE	0.0416	$[0.0368, 0.0494]$	—
Kurtosis residual (sim – obs)	0.61	$[-0.72, 3.37]$	—
<i>Per-sector OoS KS median (3 representatives per sector):</i>			
Utilities	96.2% (NEE 24.0, DUK 96.9, SO 96.2)		
Information Technology	95.1% (AAPL 95.1, MSFT 96.9, NVDA 55.7)		
Consumer Staples	92.7% (PG 94.3, KO 92.7, WMT 24.0)		
Industrials	78.9% (CAT 78.9, BA 62.2, HON 96.0)		
Materials	76.6% (FCX 76.6, NEM 5.6, APD 90.2)		
Energy	70.3% (XOM 70.3, CVX 62.3, COP 83.6)		
Consumer Discretionary	66.6% (AMZN 96.9, HD 41.4, MCD 66.6)		
Financials	57.6% (JPM 50.5, BAC 84.8, WFC 57.6)		
Communication Services	49.1% (NFLX 45.1, VZ 49.1, DIS 96.4)		
Health Care	14.5% (JNJ 93.2, UNH 14.5, LLY 7.6)		
<i>Tickers with OoS KS < 60% (failure rate at universe scale):</i>			
Count	11/30 = 36.7%		




(a) OoS density fan.



(b) KS p -value histogram.



(c) OoS ACF of raw G_t .



(d) OoS ACF of $|G_t|$.

Figure 6. OoS CHMM-N validation on SPY ($K = 18$, 1,000 paths, $T_{\text{OoS}} = 572$). Panel (a): OoS return-density fan; blue thin curves = simulated densities, orange thick curve = observed OoS density. Panel (b): histogram of the 1,000 OoS KS p -values against the observed OoS series with the $\alpha = 0.05$ threshold (orange dashed). Panel (c): ACF of raw G_t on the OoS window (linear autocorrelation); orange dashed = observed, green = mean across sim paths, shaded band = 10th–90th percentile, dashed grey = 99% Bartlett band (wider than IS because T_{OoS} is shorter). Panel (d): ACF of $|G_t|$ on the OoS window (volatility clustering); same colour conventions. The 2x2 layout parallels the IS comparison Figure 5.

4.5 Cross-Asset Dependence (Pipeline B): Student- t Copula

This is the only Pipeline B subsection in the main text (Figure 1, bottom row): we reuse the per-asset CHMM-N marginals from §4.4 and add a rank-based Student- t dependence layer (Section 3.4, full derivation in Appendix A.8.1). Profile MLE on the six-asset universe (SPY, NVDA, JNJ, JPM, AAPL, QQQ; $T_{\text{IS}} = 2,516$, 200 simulated paths) selects $\nu^* = 6$ on the unit-spaced grid $\nu \in [4, 12]$ with a Wilks 95% profile-LL CI of $[6, 7]$ (cutoff -1.92 profile-LL units below the optimum; Appendix A.8), consistent with values typically reported for equity indices [42, 43].

Rank reordering preserves each fitted CHMM marginal exactly, so the per-asset KS column is a sanity check after dependence injection rather than a substitute for Table 2. The headline dependence-layer result is the off-diagonal correlation MAE: 0.027 on IS for the Student- t copula, against 0.030 Gaussian, 0.068 truncated C-vine, and 0.076 SIM factor (Appendix A.8.1); the copula advantage is intrinsic to rank reordering rather than specific to the marginal model. On OoS the off-diagonal MAE rises to 0.209 and the Gaussian and Student- t copulas are statistically indistinguishable. The degradation localises to a single-name regime shift in the JNJ-equity-factor relationship: a per-pair breakdown (Appendix A.8.1) shows three JNJ pairs with $|\Delta| > 0.48$ on OoS, while extending the universe with GLD (a non-equity commodity) reduces the OoS off-diagonal MAE to 0.179 (Appendix A.8.6); a quarterly rolling-window refit reduces it further to 0.185 on the original six-asset universe (Appendix A.8.5).

Table 3. Cross-asset Student-t copula on CHMM-N marginals (Pipeline B; $\nu^* = 6$ by profile MLE; 200 paths). Per-asset KS pass rate at $\alpha = 0.05$ (rank reordering preserves marginals exactly, so the per-asset KS column is a sanity check after dependence injection rather than a substitute for Table 2). The lower block reports correlation reproduction averaged over paths: Frobenius norm of $\hat{\Sigma}_{\text{sim}} - \hat{\Sigma}_{\text{obs}}$ and off-diagonal mean absolute error. Comparison against SIM, the Gaussian copula, and a truncated C-vine on the same marginals is in Appendix A.8; the Student-t copula is the best of the four on the headline IS off-diagonal MAE (0.027 vs. 0.030 Gaussian, 0.068 truncated C-vine, 0.076 SIM).

Ticker	KS (%) $\uparrow$	
	IS	OoS
SPY	96.0	79.5
NVDA	98.5	53.5
JNJ	99.0	97.5
JPM	99.5	48.0
AAPL	99.0	93.5
QQQ	95.5	92.0
Mean	97.9	77.3
Median	98.8	85.8
<i>Correlation reproduction (averaged over 200 paths)</i>		
Frobenius IS		0.183
Off-diag MAE IS		0.027
Frobenius OoS		1.507
Off-diag MAE OoS		0.209
<i>Top-three OoS off-diagonal Δ (full ten-pair ranking in App. A.8.5):</i>		
JNJ-QQQ	IS $ \Delta = 0.027$, OoS $ \Delta = 0.544$	
SPY-JNJ	IS $ \Delta = 0.037$, OoS $ \Delta = 0.509$	
NVDA-JNJ	IS $ \Delta = 0.017$, OoS $ \Delta = 0.489$	

4.6 Value-at-Risk and Expected-Shortfall Back-Test (Pipeline A)

Still under Pipeline A (Figure 1): VaR and Expected-Shortfall are evaluated on the per-ticker SPY simulations, no cross-asset coupling. The headline of this subsection is the regime-conditional Christoffersen-cc result: a VaR threshold built from the CHMM one-step-ahead state forecast passes the Christoffersen joint conditional-coverage test cleanly at every (K, α) on OoS, where every constant-across- t generator (CHMM unconditional, GARCH, bootstrap) rejects breach independence. The remainder of the subsection arranges the diagnostics around this result in three layers: (iii) the regime-conditional Christoffersen-cc test (the headline), (ii) the unconditional Kupiec coverage rate (a sanity check at $T_{\text{OoS}} = 572$, integer-breach grid behaviour), and (i) the simulated envelope bracketing of observed VaR / ES (a baseline sanity check that does not require any breach-independence claim).

(i) Envelope bracketing. We simulate 1,000 independent paths from each of the six most relevant generators (Bootstrap, GARCH(1,1), CHMM-N/-t/-L/-GED) and report median and [5%, 95%] envelope across paths in Table 4. All four CHMM variants bracket observed historical VaR and ES at both 1% and 5% levels within the [5%, 95%] envelopes on both windows; the IS CHMM-N median $\text{VaR}_{0.01}$ at -6.64 (observed -6.38) and $\text{ES}_{0.01}$ at -8.80 (observed -9.00) are the closest point estimates across the six generators. The CHMM envelopes are roughly $1.7\times$ tighter than GARCH’s. CHMM-t’s IS kurtosis overshoot does *not* translate into a systematically more conservative VaR/ES point estimate; CHMM-L is mildly more conservative than CHMM-N, and CHMM-GED sits between

Table 4. VaR and Expected-Shortfall envelope for SPY ($N_{\text{paths}} = 1,000$). Each entry is the median and [5%, 95%] envelope across paths of the left-tail VaR / ES at $\alpha \in \{0.01, 0.05\}$; observed historical values are in the first row.

Model	IS VaR _{0.01} median [5%, 95%]	IS ES _{0.01} median [5%, 95%]	IS VaR _{0.05} median [5%, 95%]	IS ES _{0.05} median [5%, 95%]
<i>Observed</i>	−6.38	−9.00	−3.43	−5.50
Bootstrap	−6.38 [−7.04, −5.99]	−8.86 [−10.37, −7.78]	−3.43 [−3.68, −3.20]	−5.46 [−5.99, −5.05]
GARCH	−5.80 [−8.62, −4.52]	−7.86 [−13.54, −5.77]	−3.19 [−3.92, −2.71]	−4.89 [−7.11, −3.91]
CHMM-N	−6.64 [−8.01, −5.47]	−8.80 [−10.16, −7.34]	−3.45 [−4.05, −2.95]	−5.45 [−6.33, −4.61]
CHMM-t	−6.58 [−7.51, −5.83]	−9.06 [−10.70, −7.79]	−3.40 [−3.90, −2.90]	−5.51 [−6.18, −4.85]
CHMM-L	−6.89 [−7.85, −6.01]	−9.15 [−10.42, −7.97]	−3.30 [−3.77, −2.88]	−5.53 [−6.17, −4.87]
CHMM-GED	−6.85 [−7.70, −6.06]	−8.79 [−9.83, −7.82]	−3.43 [−3.98, −2.93]	−5.55 [−6.23, −4.91]
	OoS VaR _{0.01}	OoS ES _{0.01}	OoS VaR _{0.05}	OoS ES _{0.05}
<i>Observed</i>	−5.51	−7.94	−2.77	−4.67
Bootstrap	−6.33 [−7.60, −5.28]	−8.43 [−11.76, −6.59]	−3.39 [−3.91, −2.88]	−5.34 [−6.38, −4.50]
GARCH	−5.35 [−9.98, −3.61]	−6.72 [−13.51, −4.38]	−3.14 [−4.94, −2.36]	−4.59 [−7.92, −3.23]
CHMM-N	−6.30 [−9.02, −4.30]	−8.44 [−11.36, −5.42]	−3.39 [−4.74, −2.47]	−5.27 [−7.23, −3.74]
CHMM-t	−6.32 [−8.13, −4.80]	−8.60 [−12.20, −6.29]	−3.29 [−4.27, −2.45]	−5.34 [−6.72, −4.08]
CHMM-L	−6.69 [−8.59, −4.91]	−8.88 [−11.48, −6.73]	−3.27 [−4.30, −2.51]	−5.47 [−6.82, −4.20]
CHMM-GED	−6.60 [−8.30, −4.82]	−8.51 [−10.68, −6.47]	−3.32 [−4.38, −2.46]	−5.41 [−6.66, −4.06]

the two, consistent with their kurtosis residuals.

(ii) **Unconditional Kupiec coverage.** The four CHMM rows pass the unconditional Kupiec test cleanly on both windows: at $\alpha = 0.05$ on OoS, CHMM-N $\text{LR}_{\text{uc}} = 3.83$ against $\chi_1^2(0.05) = 3.841$, CHMM-t 3.83, CHMM-L 3.03, CHMM-GED 3.83. The repeated 3.83 across rows is a feature of $T_{\text{OoS}} = 572$: the Kupiec statistic is a function of the integer breach count k , and at $\alpha = 0.05$ the values $k \in \{14, 19, 20\}$ produce $\text{LR}_{\text{uc}} \in \{3.03, 3.83, 3.83\}$ that pile against the χ_1^2 critical value rather than spreading continuously. We treat the unconditional Kupiec as a sanity check rather than the primary diagnostic at this T_{OoS} ; the substantive risk-management diagnostic is the conditional construction below.

(iii) **Regime-conditional Christoffersen-cc (headline).** The Christoffersen [66] independence ($\text{LR}_{\text{ind}} \sim \chi_1^2$) and joint conditional-coverage ($\text{LR}_{\text{cc}} = \text{LR}_{\text{uc}} + \text{LR}_{\text{ind}} \sim \chi_2^2$) tests check whether breaches cluster in time. Under the unconditional pooled-archive variant (constant-across- t VaR threshold from the IS-only simulated envelope), every panel generator rejects independence on the OoS window at $\alpha = 0.05$ for both VaR levels (CHMM-N at 5%: $\text{LR}_{\text{ind}} = 5.26$, $p = 0.022$; $\text{LR}_{\text{cc}} = 9.08$, $p = 0.011$; full panel in Appendix A.3.2, Table 9). The rejection is structurally driven by volatility-clustering in R_{oos} that any unconditional generator cannot track.

The regime-conditional variant addresses this directly. At each OoS day t , run the forward filter through $\mathcal{F}_{t-1} = R_{\text{IS}} \cup R_{\text{oos}}[1:t-1]$ to get the one-step-ahead state forecast

$$\mathbb{P}(s_t = k \mid \mathcal{F}_{t-1}) = \sum_i \mathbb{P}(s_{t-1} = i \mid \mathcal{F}_{t-1}) T_{ik}, \quad (9)$$

and define $\widehat{\text{VaR}}_t(\alpha) = F_t^{-1}(\alpha)$ where $F_t(\cdot) = \sum_k \mathbb{P}(s_t = k \mid \mathcal{F}_{t-1}) \Phi(\cdot; \mu_k, \sigma_k)$ is the K -component Gaussian-mixture predictive CDF under IS-fixed $(\mathbf{T}, \{\mu_k, \sigma_k\})$. Table 5 reports the back-test for CHMM-N at $K^* = 3$ and $K = 18$ on OoS.

Table 5. Regime-conditional VaR back-test, CHMM-N and penalised CHMM-t ($\lambda = 20$) on OoS ($T_{\text{oos}} = 572$, seed 20260420). Conditional VaR is the α -quantile of the one-step-ahead family-appropriate predictive mixture density under IS-fixed parameters (Gaussian for CHMM-N, per-state Student- t for CHMM-t). Critical values: $\chi_1^2(0.05) = 3.841$, $\chi_2^2(0.05) = 5.991$. Every $(K, \alpha, \text{family})$ row passes Kupiec, Christoffersen-ind, and Christoffersen-cc cleanly at $\alpha = 0.05$. The full four-family extension (CHMM-N, CHMM-t at $\lambda = 20$, CHMM-L, CHMM-GED) is in Appendix A.3.2, Table 29.

Family	K	α	breaches	br rate	med VaR	LR _{uc}	LR _{ind}	LR _{cc}	p_{cc}
CHMM-N	3	0.01	9	1.57%	−4.56	1.62	2.36	3.98	0.14
CHMM-N	3	0.05	35	6.12%	−2.87	1.41	0.01	1.42	0.49
CHMM-N	18	0.01	9	1.57%	−5.20	1.62	2.36	3.98	0.14
CHMM-N	18	0.05	26	4.55%	−3.02	0.26	0.52	0.78	0.68
CHMM-t ($\lambda=20$)	3	0.01	8	1.40%	−5.59	0.82	2.81	3.63	0.16
CHMM-t ($\lambda=20$)	3	0.05	32	5.59%	−2.73	0.41	0.77	1.19	0.55
CHMM-t ($\lambda=20$)	18	0.01	10	1.75%	−6.25	2.65	1.98	4.62	0.10
CHMM-t ($\lambda=20$)	18	0.05	29	5.07%	−3.15	0.01	1.39	1.40	0.50

The conditional construction passes all three tests at every (K, α) , with breach rates 1.57% and 4.55–6.12% against nominal 1% and 5%. The independence-test improvement from the unconditional variant (LR_{ind} = 5.26 at $K = 18, \alpha = 0.05$) to the conditional variant (LR_{ind} = 0.52) is the regime-switching value proposition that the unconditional Kupiec headline did not exercise: the latent-state forecast tightens the threshold in clustered-volatility windows so that breaches do not pile up at sub-runs of high volatility. The conditional Christoffersen-cc is the substantive risk-management contribution of the paper; the unconditional Kupiec serves as a sanity check at this T_{OoS} . The construction reuses IS-fixed CHMM-N parameters, requires no retuning of K on a stress slice, and runs the forward filter on the OoS series under frozen parameters. The conditional pass extends to every emission family with the family-appropriate predictive density (CHMM-t at $\lambda = 20$, CHMM-L, CHMM-GED): all sixteen $(K, \alpha, \text{family})$ rows pass Kupiec, Christoffersen-ind, and Christoffersen-cc at $\alpha = 0.05$ with $p_{cc} \geq 0.089$ throughout (Appendix A.3.2, Table 29). The regime-switching value proposition is therefore intrinsic to the latent-state forecast and not specific to the Gaussian emission family. The full unconditional Christoffersen panel across the six benchmark rows is in Appendix A.3.2.

Walk-forward stability of the conditional construction. The conditional pass reported above is for the headline OoS window 2024–2026. Extending the construction to the six rolling-origin folds of Table 28, with CHMM-N refit per fold from scratch on the train slice and forward-filtered through the test slice under fold-IS-fixed parameters, gives a 24-row panel (Appendix A.9, Table 30). The conditional Christoffersen-cc passes at $\alpha = 0.05$ on 10 of 12 fold- K combinations: W1, W3, W4, W5, W6 at both $K \in \{3, 18\}$ all clear the $\chi_2^2(0.05) = 5.991$ critical value, with $p_{cc} \in [0.11, 0.92]$. The two failures concentrate on W2 (COVID, $p_{cc} = 0.011$ at $K = 3$ and 0.002 at $K = 18$); at $\alpha = 0.01$ a third failure appears on W4 at $K = 18$ ($p_{cc} = 0.022$ on a LR_{ind} tail-event clustering). The pass-rate is therefore 19/24 aggregate at the 5% level, and 22/24 if one excludes the two stress folds that the univariate walk-forward already flagged as out-of-distribution by KS in Table 28. The reading is that the conditional construction is window-conditional under regime shifts of the W2 / W4 magnitude (a regime introduction the IS distribution does not span; Section 5, stationarity-scope paragraph) but window-robust on the four non-stress folds plus W4 at $K = 3$ at $\alpha = 0.05$; the body claim that the conditional construction is the regime-switching value proposition is supported beyond the headline single-window result. Periodic refit at quarterly cadence is the standard mitigation for the

stress folds (26); the four-family extension on the headline window in Table 29 establishes that the construction is not Gaussian-emission-specific.

5 Discussion

A continuous HMM trained by Baum-Welch reproduces the three Cont stylized facts simultaneously on SPY at two operating points: at the held-out-likelihood and BIC choice $K^* = 3$, CHMM-N attains 89.7% IS KS and 80.1% OoS KS at $|G_t|$ ACF-MAE 0.0467 (Table 1); at the tail-fidelity panel $K = 18$, the four emission variants attain 93.6–95.6% IS KS and 80.8–85.7% OoS KS, with CHMM-N $|G_t|$ ACF-MAE within 0.003 of GARCH(1,1). Four emission variants share the same scaffold and land at four distinct points on the kurtosis axis without retuning K or the transition structure: CHMM-N undershoots, CHMM-L matches observed kurtosis cleanly without a shape parameter, CHMM-t overshoots while supplying per-state heavy tails for tail-conditional consumers, and CHMM-GED nests Gaussian and Laplace as boundary cases ($p_k = 2$ and $p_k = 1$, respectively) and lets the data choose each state’s kurtosis adaptively on the continuous shape axis. The mechanism behind the simultaneous KS / ACF fit is the spectral identity of equation (8): at $K \geq 3$ with quantile-initialised Baum-Welch, the absolute-return ACF is a population-level real-coefficient sum of geometric decays at the non-unit eigenvalues of $\mathbf{T}$.

Replicating the slow-ACF behaviour at $K \geq 3$. Two issues compound in the Rydén et al. [11] low- K setup: a single self-transition rate produces a single exponential mode that cannot be simultaneously fast (good marginal) and slow (good ACF), and the random initialisations standard in the 1990s often converge to degenerate local optima with overlapping states. Our $K = 2$ replication on SPY (Appendix A.6.11) clarifies the mechanism: ACF-MAE is essentially constant at ≈ 0.050 across quantile and five random-seed initialisations, matching the $K = 18$ CHMM-N. The binding low- K constraint is therefore distributional rather than temporal (KS at $K = 2$ drops to 67–72%); with modern initialisation the ACF is reproduced at any $K \geq 2$, while distributional fidelity is what actually drives the choice of K . The effective-rank diagnostic in Appendix A.6.6 (Table 16) makes this microscopically explicit: at the fitted $\hat{\mathbf{T}}$ for $K = 18$, a single non-unit eigenvalue carries 93.6% of the lag-1 absolute-return ACF and the next two modes bring the cumulative share to 95.5%; at $K = 3$ the dominant non-unit mode alone carries 96.8%. The K -rank statement on $\mathbf{T} - \mathbf{1}\bar{\pi}^\top$ is the algebraic upper bound on the ACF identity (8); the empirical effective rank on equity-return data is one to three, whether $K = 3$ or $K = 18$, which is the spectral expression of the temporal axis decoupling from K at the operating points the paper considers.

OoS KS pass rates against the test-power ceiling. At $T_{\text{OoS}} = 572$ the two-sample KS test does not reach 100% on truly-correct generators. A length-matched i.i.d. resample of the IS distribution itself passes at 90.0%, while the Gaussian i.i.d. negative control is rejected at 0%, so the test retains ample power to refuse clearly-wrong generators. The CHMM OoS KS pass rates of 79–86% sit just below the 90% ceiling (a genuine fidelity claim, not a low-power artefact); GARCH’s 58% is correctly rejected. The asymptotic two-sample KS test assumes i.i.d. samples, but the simulated paths from CHMM and GARCH carry volatility clustering, so the asymptotic null is mildly lenient; under stationary block-bootstrap recalibration (Appendix A.7.2, Table 19) the CHMM family retains 88–93% pass rate at block length $L = 10$ while GARCH(1,1) drops sharply from 26.8% to 13.6%, consistent with the i.i.d.-baseline null being mildly favourable to volatility-clustering generators with the wrong marginal. The block-aware reading reinforces the asymptotic conclusion: CHMM

passes the temporal-aware distributional benchmark at the same rate it passes the asymptotic one; GARCH’s failure widens.

Closing the kurtosis gap with heavy-tailed emissions. CHMM-N produces excess kurtosis 5.02 against observed 7.68: a Gaussian mixture’s kurtosis is bounded by the variance of component means relative to their variances, and pushing tail states out degrades the central fit. CHMM-t addresses this through per-state Student- t emissions (ν_k updated by ECM one-dimensional search), at the cost of an IS overshoot to 14.57. The overshoot is a single-state artefact rather than a lower-bracket pinning: a sweep $\nu_{\min} \in \{2.1, 2.5, 3.0, 4.0\}$ leaves the median ν_k at 50 with only 2/18 states near the lower edge, and an exponential $1/\nu_k$ shrinkage at rate $\lambda = 20$ brings simulated kurtosis to ≈ 8 at a 1pp KS cost (Appendix A.9). CHMM-L sits between CHMM-N and CHMM-t at 6.75 without any shape parameter, and CHMM-GED’s bimodal $\hat{p}_k$ partition (body Figure 4) is the data-driven analog of a hand-classified Gaussian-bulk / Laplace-tail hybrid. The four-way decision is summarised in Table 6; all four variants’ 5–95% envelopes bracket observed VaR/ES on both windows.

Table 6. Variant decision guide. Recommended CHMM emission family by use-case priority. All four variants share the forward-backward scaffold and the quantile-based initialisation; the M-step is the only architectural difference.

Use-case priority	Recommended variant
Simplest closed-form fit; kurtosis fidelity secondary	CHMM-N
Cleanest IS kurtosis match; no shape parameter	CHMM-L
Per-state heavy tails for tail-conditional consumers	CHMM-t with $1/\nu_k$ shrinkage ($\lambda = 20$)
Adaptive per-state shape; data-chosen Gaussian-bulk / Laplace-tail mixture	CHMM-GED

Per-ticker $\hat{\lambda}^*$ for the penalised CHMM-t. The body Table 2 reports the operational variant at a uniform $\lambda = 20$. A per-ticker sweep $\lambda \in \{0, 5, 10, 20, 50, 100\}$ on each of the six body tickers (Appendix A.7) defines $\hat{\lambda}_i^* = \arg \min_{\lambda} |\text{kurt}_{\text{sim}, \lambda} - \text{kurt}_{\text{obs}, i}|$ subject to IS KS degradation ≤ 1.5 percentage points from $\lambda = 0$. The per-ticker optima distribute as: $\hat{\lambda}_{\text{SPY}}^* = 10$, $\hat{\lambda}_{\text{NVDA}}^* = 20$, $\hat{\lambda}_{\text{JNJ}}^* = 20$, $\hat{\lambda}_{\text{JPM}}^* = 10$, $\hat{\lambda}_{\text{AAPL}}^* = 10$, $\hat{\lambda}_{\text{QQQ}}^* = 0$. The body uniform choice $\lambda = 20$ is the right rate on the heavy-tailed defensives (NVDA, JNJ) where unpenalised kurtosis blows up by an order of magnitude, but $\lambda = 10$ suffices on the moderate-tail tickers and $\lambda = 0$ is appropriate for QQQ whose unpenalised fit already matches observed kurtosis to within 0.05 kurtosis units. We commit to the following operational rule: *set $\hat{\lambda}_i^* = 20$ if the residual kurtosis of CHMM-t at $\lambda = 0$ exceeds the observed kurtosis on ticker i by more than one unit; set $\hat{\lambda}_i^* = 10$ if the gap is between 0.1 and 1 unit; set $\hat{\lambda}_i^* = 0$ if the gap is below 0.1 units.* Under this rule the per-ticker assignment is $\hat{\lambda}_{\text{SPY}}^* = 10$, $\hat{\lambda}_{\text{NVDA}}^* = 20$, $\hat{\lambda}_{\text{JNJ}}^* = 20$, $\hat{\lambda}_{\text{JPM}}^* = 10$, $\hat{\lambda}_{\text{AAPL}}^* = 10$, $\hat{\lambda}_{\text{QQQ}}^* = 0$, which matches the per-ticker sweep optima above. The body uniform $\lambda = 20$ is therefore a conservative default that is correct on the two defensives where the unpenalised pinning artefact is severe, slightly heavier than the rule on three moderate-tail tickers (with the 1.5pp KS / 0.001 ACF-MAE penalty quantified above), and not the rule’s choice on QQQ.

Stationarity scope and operational deployment. The CHMM is trained on the assumption that returns are locally stationary across the 10-year IS window. Single-name equity returns are well known to carry sector-specific structural breaks (23; 24; surveyed by 25), and Cross-ticker generalization (Section 4.4) reflects this directly: at $K = 18$, OoS distributional fidelity is preserved

on SPY, JNJ, AAPL, and QQQ but degrades on NVDA (57.2%) and JPM (53.0%), where the IS-fixed transitions and emissions cannot capture the AI-driven NVDA regime or the rate-repricing cycle for financials. The pattern persists at every K in the sweep, so it is not a model-selection issue. Operational deployment of the CHMM should follow standard practice for time-series generators on financial data, with periodic refit at quarterly or finer cadence (26; an online-EM formulation is given by 67). The static IS / OoS panel reported here is the reference comparison, not the operational protocol.

Why the W4 stress fold fails where the headline OoS holds. The walk-forward distribution (Appendix A.9, Table 28) shows two stress folds with KS pass rates below 10%: W2 (2020 COVID) and W4 (2022 rate-hike onset, $KS \leq 1\%$). The headline OoS window is the rate-cycle *reversal* of 2024–2026 ($KS \sim 82\%$). The asymmetry between the W4 onset (catastrophic) and the headline OoS reversal (clean) is consistent with two compounding mechanisms. First, the IS window 2014–2019 saw only one mild rate-hike cycle (2015–2018 in 25-bp steps), so the IS-fitted volatility-regime distribution does not include the magnitude of the 2022 hike (a 525-bp move over 18 months) and the IS-fixed CHMM has no state with the right σ to absorb it. The 2024–2026 window, by contrast, covers the *aftermath* of that hike, which the IS-period regime distribution can partially span (the 2018 December selloff and the 2020 COVID volatility cluster jointly cover the high- σ tail of the 2024–2026 distribution). Second, the 2022 hike was structurally a regime *introduction* (a new high-rate, high-real-yield regime that did not exist in the IS window), whereas the 2024–2026 reversal is a regime *attenuation* (a return to a regime closer to the IS volatility level). The IS-fixed transitions and emissions can track an attenuation but not an introduction, which is the operational reading of the asymmetry: walk-forward stress folds at regime introductions are out of scope for the IS-fixed protocol regardless of K , while regime attenuations and same-regime-with-different-amplitude OoS slices remain in scope.

Limitations. The cross-ticker study covers six tickers; a truncated C-vine variant on the same universe is reported in Appendix A.8.4 and does not improve over the elliptical copulas at this scale, so scaling to larger d via untruncated regular vines or factor copulas is the natural next step. State selection is via information-criterion plus distributional pass-rate rather than held-out log-likelihood; a clean re-selection on a non-OoS validation slice picks $K^* = 3$ by held-out log-lik (Appendix A.9), so the $K = 18$ operating point is best read as a multi-metric choice that weights tail fidelity alongside likelihood. Skew-heavy-tailed emissions (skew- t or skew-Laplace) and explicit-duration semi-Markov sojourns are deferred as companion-paper directions; regime-conditional VaR via one-step-ahead latent-state forecast propagation is delivered in Section 4.6 (Table 5) and passes all three Kupiec / Christoffersen-ind / Christoffersen-cc tests at every (K, α) on OoS.

6 Conclusion

A continuous HMM trained by Baum-Welch with quantile-based initialization reproduces the three Cont stylized facts simultaneously on SPY at two held-out-clean operating points and an extended-state-resolution reference: the held-out-likelihood and BIC choice $K^* = 3$ for risk-management consumers (CHMM-N: 89.7% IS / 80.1% OoS KS, $|G_t|$ ACF-MAE 0.0467); the held-out-likelihood / held-out-KS reference $K^* = 6$ on the strictly pre-2020 slice for synthetic-data consumers (four-variant block at 88.3–92.6% IS, 76.3–79.0% OoS KS; CHMM-N kurtosis 5.26 matches observed 5.29 OoS within 0.03); and an extended-state-resolution reference panel at $K = 18$ at the cost of relaxing the held-out criterion onto a multi-objective rule, with the four emission variants attaining 93.6–95.6% IS,

80.8–85.7% OoS KS and the penalised CHMM-t at $\lambda = 20$ providing the cleanest IS / OoS kurtosis match at 8.56 / 7.07 against observed 7.68 / 5.29 (Table 1). The mechanism is the closed-form mixture-of-eigenvalues identity for the absolute-return ACF (13–15); we apply it to recast the Rydén et al. low- K failure as a rank statement on $\mathbf{T} - \mathbf{1}\bar{\pi}^\top$ and show the rank binds only at $K \leq 3$ Gaussian setups. The four emission variants share the same scaffold and land at four kurtosis points; CHMM-GED’s per-state shape p_k nests Gaussian and Laplace as boundary cases and produces a bulk/tail partition that replicates across seeds and tickers. VaR and ES envelopes bracket observed values at 1% and 5%; a regime-conditional VaR that propagates the one-step-ahead state forecast through the CHMM mixture passes Kupiec, Christoffersen-ind, and Christoffersen-cc cleanly at every (K, α) on OoS (Table 5), and the conditional construction extends to a six-fold rolling-origin walk-forward with a 19/24 aggregate pass-rate at the 5% level (Table 30); the two failures concentrate on the COVID and 2022-rate-hike-onset stress folds that the univariate walk-forward already flags as out-of-distribution by KS, where every constant-across- t generator rejects breach independence in any case. Cross-ticker generalisation is exercised on a sector-balanced 30-ticker panel (10 GICS sectors $\times$ 3 large-cap representatives; Section 4.4, Table 2): IS KS distribution is tight (median 99.5%), and OoS KS distribution is sector-stratified with $11/30 = 36.7\%$ of tickers below 60%, concentrated in single-name regime introductions (Health Care: LLY weight-loss-drug, UNH 2024 policy compression) and the rate-cycle / streaming-platform patterns previously identified on the original five-named-ticker spot-check. Multi-asset synthesis composes through a rank-based Student- t copula on the CHMM marginals at $\nu^* = 6$ (six-asset IS off-diagonal correlation MAE 0.027); on OoS the Gaussian and Student- t copulas are statistically indistinguishable, with a quarterly rolling refit reducing the off-diagonal MAE to 0.185 (Appendix A.8.5).

Skew-heavy-tailed emissions and explicit-duration semi-Markov sojourns are companion-paper directions; scaling to larger d via untruncated regular vines or factor copulas is the natural multi-asset follow-up.

Data and code availability. Reproducer at <https://github.com/altashly1/CHMM-paper>; companion Julia package CHMM-Model.jl at <https://github.com/altashly1/CHMM-Model> (public API in Appendix A.1, Julia ≥ 1.12). Daily prices: Polygon.io via Massive (2014-01-03 to 2025-11-18) and Alpaca Markets / IEX (held-out extension through 2026-04-20); we thank both providers. The authors declare no competing interests. **Author contributions:** J.V. directed the study; A.A. developed the continuous model, implemented Baum-Welch, conducted the empirical analysis, and generated all figures and tables; C.J. contributed to methodology review and validation; all authors edited and reviewed the final manuscript.

References

- [1] James Jordon, Lukasz Szpruch, Florimond Houssiau, Mirko Bottarelli, Giovanni Cherubin, Carsten Maple, Samuel N. Cohen, and Adrian Weller. Synthetic data: what, why and how? *arXiv preprint arXiv:2205.03257*, 2022.
- [2] Samuel A. Assefa, Danial Dervovic, Mahmoud Mahfouz, Robert E. Tillman, Prashant Reddy, and Manuela Veloso. Generating synthetic data in finance: opportunities, challenges and pitfalls. In *Proceedings of the First ACM International Conference on AI in Finance*, pages 1–8, 2020.
- [3] Abdulrahman Alswaidan and Jeffrey D. Varner. Hybrid hidden markov model for modeling equity excess growth rate dynamics: A discrete-state approach with jump-diffusion. *arXiv preprint arXiv:2603.10202*, 2026.

- [4] Benoit Mandelbrot. The variation of certain speculative prices. *The Journal of Business*, 36(4): 394–419, 1963.
- [5] R. Cont. Empirical properties of asset returns: stylized facts and statistical issues. *Quantitative Finance*, 1(2):223–236, 2001.
- [6] Robert F. Engle. Autoregressive conditional heteroscedasticity with estimates of the variance of united kingdom inflation. *Econometrica*, 50(4):987–1007, 1982.
- [7] Tim Bollerslev. Generalized autoregressive conditional heteroscedasticity. *Journal of Econometrics*, 31(3):307–327, 1986.
- [8] Jinsung Yoon, Daniel Jarrett, and Mihaela van der Schaar. Time-series generative adversarial networks. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 32, 2019.
- [9] Magnus Wiese, Robert Knobloch, Ralf Korn, and Peter Kretschmer. Quant GANs: Deep generation of financial time series. *Quantitative Finance*, 20(9):1419–1440, 2020.
- [10] Kashif Rasul, Calvin Seward, Ingmar Schuster, and Roland Vollgraf. Autoregressive denoising diffusion models for multivariate probabilistic time series forecasting. In *Proceedings of the 38th International Conference on Machine Learning*, pages 8857–8868, 2021.
- [11] Tobias Rydén, Timo Teräsvirta, and Stefan Åsbrink. Stylized facts of daily return series and the hidden Markov model. *Journal of Applied Econometrics*, 13(3):217–244, 1998.
- [12] Jan Bulla and Ingo Bulla. Stylized facts of financial time series and hidden semi-Markov models. *Computational Statistics & Data Analysis*, 51(4):2192–2209, 2006.
- [13] James D. Hamilton. *Time Series Analysis*. Princeton University Press, Princeton, NJ, 1994.
- [14] Hans-Martin Krolzig. *Markov-Switching Vector Autoregressions: Modelling, Statistical Inference, and Application to Business Cycle Analysis*, volume 454 of *Lecture Notes in Economics and Mathematical Systems*. Springer, Berlin, 1997.
- [15] Allan Timmermann. Moments of Markov switching models. *Journal of Econometrics*, 96(1): 75–111, 2000.
- [16] Leonard E. Baum, Ted Petrie, George Soules, and Norman Weiss. A maximization technique occurring in the statistical analysis of probabilistic functions of markov chains. *The Annals of Mathematical Statistics*, 41(1):164–171, 1970.
- [17] L. Rabiner and B. Juang. An introduction to hidden Markov models. *IEEE ASSP Magazine*, 3(1):4–16, 1986.
- [18] David Peel and Geoffrey J. McLachlan. Robust mixture modelling using the t distribution. *Statistics and Computing*, 10(4):339–348, 2000.
- [19] Chuanhai Liu and Donald B. Rubin. ML estimation of the t distribution using EM and its extensions, ECM and ECME. *Statistica Sinica*, 5(1):19–39, 1995.
- [20] Dimitris N. Politis and Joseph P. Romano. The stationary bootstrap. *Journal of the American Statistical Association*, 89(428):1303–1313, 1994.

- [21] Tim Bollerslev. A conditionally heteroskedastic time series model for speculative prices and rates of return. *Review of Economics and Statistics*, 69(3):542–547, 1987.
- [22] Markus Haas, Stefan Mittnik, and Marc S. Paoletta. A new approach to Markov-switching GARCH models. *Journal of Financial Econometrics*, 2(4):493–530, 2004.
- [23] Ľuboš Pástor and Robert F. Stambaugh. The equity premium and structural breaks. *Journal of Finance*, 56(4):1207–1239, 2001.
- [24] Elena Andreou and Eric Ghysels. Detecting multiple breaks in financial market volatility dynamics. *Journal of Applied Econometrics*, 17(5):579–600, 2002.
- [25] Andrew Ang and Allan Timmermann. Regime changes and financial markets. *Annual Review of Financial Economics*, 4(1):313–337, 2012.
- [26] M. Hashem Pesaran and Allan Timmermann. Selection of estimation window in the presence of breaks. *Journal of Econometrics*, 137(1):134–161, 2007.
- [27] James D. Hamilton. A new approach to the economic analysis of nonstationary time series and the business cycle. *Econometrica*, 57(2):357–384, 1989.
- [28] Huntley Schaller and Simon Van Norden. Regime switching in stock market returns. *Applied Financial Economics*, 7(2):177–191, 1997.
- [29] Peter Nystrup, Henrik Madsen, and Erik Lindström. Long memory of financial time series and hidden markov models with time-varying parameters. *Journal of Forecasting*, 36(8):989–1002, 2017.
- [30] M. T. Subbotin. On the law of frequency of error. *Matematicheskii Sbornik*, 31(2):296–301, 1923.
- [31] George E. P. Box and George C. Tiao. *Bayesian Inference in Statistical Analysis*. Addison-Wesley, Reading, MA, 1973.
- [32] Daniel B. Nelson. Conditional heteroskedasticity in asset returns: A new approach. *Econometrica*, 59(2):347–370, 1991.
- [33] Lawrence R. Glosten, Ravi Jagannathan, and David E. Runkle. On the relation between the expected value and the volatility of the nominal excess return on stocks. *The Journal of Finance*, 48(5):1779–1801, 1993.
- [34] Jean-Michel Zakoian. Threshold heteroskedastic models. *Journal of Economic Dynamics and Control*, 18(5):931–955, 1994.
- [35] Torben G. Andersen and Tim Bollerslev. Heterogeneous information arrivals and return volatility dynamics: Uncovering the long-run in high frequency returns. *The Journal of Finance*, 52(3): 975–1005, 1997.
- [36] Fulvio Corsi. A simple approximate long-memory model of realized volatility. *Journal of Financial Econometrics*, 7(2):174–196, 2009.
- [37] Shuntaro Takahashi, Yu Chen, and Kumiko Tanaka-Ishii. Modeling financial time-series with generative adversarial networks. *Physica A: Statistical Mechanics and its Applications*, 527: 121261, 2019.

- [38] Sohyeon Kwon and Yongjae Lee. Can GANs learn the stylized facts of financial time series? In *Proceedings of the 5th ACM International Conference on AI in Finance*, pages 126–133, 2024.
- [39] William F. Sharpe. A simplified model for portfolio analysis. *Management Science*, 9(2):277–293, 1963.
- [40] A. Sklar. Fonctions de répartition à n dimensions et leurs marges. *Publications de l’Institut de Statistique de l’Université de Paris*, 8:229–231, 1959.
- [41] Roger B. Nelsen. *An Introduction to Copulas*. Springer, 2 edition, 2006.
- [42] Stefano Demarta and Alexander J. McNeil. The t copula and related copulas. *International Statistical Review*, 73(1):111–129, 2005.
- [43] Alexander J. McNeil, Rüdiger Frey, and Paul Embrechts. *Quantitative Risk Management: Concepts, Techniques and Tools*. Princeton University Press, revised edition, 2015.
- [44] Ronald L. Iman and W. J. Conover. A distribution-free approach to inducing rank correlation among input variables. *Communications in Statistics–Simulation and Computation*, 11(3):311–334, 1982.
- [45] Michael Stenger, André Bauer, Thomas Prantl, Robert Leppich, Nathaniel Hudson, Kyle Chard, Ian Foster, and Samuel Kounev. Thinking in categories: A survey on assessing the quality for time series synthesis. *Journal of Data and Information Quality*, 16(2):1–32, 2024.
- [46] Tilmann Gneiting and Adrian E. Raftery. Strictly proper scoring rules, prediction, and estimation. *Journal of the American Statistical Association*, 102(477):359–378, 2007.
- [47] Tilmann Gneiting and Matthias Katzfuss. Probabilistic forecasting. *Annual Review of Statistics and Its Application*, 1:125–151, 2014.
- [48] Francis X. Diebold and Roberto S. Mariano. Comparing predictive accuracy. *Journal of Business & Economic Statistics*, 13(3):253–263, 1995.
- [49] Arthur Gretton, Karsten M. Borgwardt, Malte J. Rasch, Bernhard Schölkopf, and Alexander Smola. A kernel two-sample test. *Journal of Machine Learning Research*, 13(1):723–773, 2012.
- [50] Ilya Chevyrev and Andrey Kormilitzin. A primer on the signature method in machine learning. *arXiv preprint arXiv:1603.03788*, 2016.
- [51] Hao Ni, Lukasz Szpruch, Magnus Wiese, Shujian Liao, and Baoren Xiao. Conditional Sig-Wasserstein GANs for time series generation. *arXiv preprint arXiv:2006.05421*, 2020.
- [52] Paul Embrechts, Alexander McNeil, and Daniel Straumann. Correlation and dependence in risk management: Properties and pitfalls. *Risk Management: Value at Risk and Beyond*, pages 176–223, 2002.
- [53] A. N. Kolmogorov. Sulla determinazione empirica di una legge di distribuzione. *Giornale dell’Istituto Italiano degli Attuari*, 4:83–91, 1933.
- [54] N. Smirnov. Table for estimating the goodness of fit of empirical distributions. *The Annals of Mathematical Statistics*, 19(2):279–281, 1948.

- [55] Paul H. Kupiec. Techniques for verifying the accuracy of risk measurement models. *Journal of Derivatives*, 3(2):73–84, 1995.
- [56] Jeff A. Bilmes. A gentle tutorial of the em algorithm and its application to parameter estimation for gaussian mixture and hidden markov models. *Technical Report TR-97-021, International Computer Science Institute*, 1998.
- [57] Carlos A. Abanto-Valle, Roland Langrock, Ming-Hui Chen, and Marcos V. Cardoso. Maximum likelihood estimation for stochastic volatility in mean models with heavy-tailed distributions. *Applied Stochastic Models in Business and Industry*, 33(4):394–408, 2017.
- [58] David A. Levin and Yuval Peres. *Markov Chains and Mixing Times*. American Mathematical Society, 2 edition, 2017.
- [59] C. F. Jeff Wu. On the convergence properties of the EM algorithm. *The Annals of Statistics*, 11(1):95–103, 1983.
- [60] Xiao-Li Meng and Donald B. Rubin. Maximum likelihood estimation via the ECM algorithm: A general framework. *Biometrika*, 80(2):267–278, 1993.
- [61] Elizabeth S. Allman, Catherine Matias, and John A. Rhodes. Identifiability of parameters in latent structure models with many observed variables. *The Annals of Statistics*, 37(6A):3099–3132, 2009.
- [62] Sidney J. Yakowitz and John D. Spragins. On the identifiability of finite mixtures. *The Annals of Mathematical Statistics*, 39(1):209–214, 1968.
- [63] Peter J. Bickel, Ya’acov Ritov, and Tobias Rydén. Asymptotic normality of the maximum-likelihood estimator for general hidden Markov models. *The Annals of Statistics*, 26(4):1614–1635, 1998.
- [64] Randal Douc, Eric Moulines, and Tobias Rydén. Asymptotic properties of the maximum likelihood estimator in autoregressive models with Markov regime. *The Annals of Statistics*, 32(5):2254–2304, 2004.
- [65] Nguyet Nguyen. Hidden Markov model for stock trading. *International Journal of Financial Studies*, 6(2):36, 2018.
- [66] Peter F. Christoffersen. Evaluating interval forecasts. *International Economic Review*, 39(4):841–862, 1998.
- [67] Olivier Cappé. Online EM algorithm for hidden Markov models. *Journal of Computational and Graphical Statistics*, 20(3):728–749, 2011.
- [68] Paul Glasserman. *Monte Carlo Methods in Financial Engineering*. Springer, 2003.
- [69] Bruce G. Lindsay. *Mixture Models: Theory, Geometry and Applications*, volume 5 of *NSF-CBMS Regional Conference Series in Probability and Statistics*. Institute of Mathematical Statistics, Hayward, CA, 1995.
- [70] Martin Arjovsky, Soumith Chintala, and Léon Bottou. Wasserstein generative adversarial networks. In *Proceedings of the 34th International Conference on Machine Learning*, pages 214–223, 2017.

- [71] Mike Innes. Flux: Elegant machine learning with Julia. *Journal of Open Source Software*, 3 (25):602, 2018.
- [72] Abdulrahman Alswaidan, Cade Jin, and Jeffrey D. Varner. Long memory from markov mixing: A semi-markov regime-switching generator for the cboe volatility index. 2026. Manuscript in preparation.
- [73] Shun-Zheng Yu. Hidden semi-Markov models. *Artificial Intelligence*, 174(2):215–243, 2010.

A Supplementary Material

This appendix collects supplementary material referenced in the main body: the public API of the companion package, the EM forward-backward recursions and algorithm pseudocode, the validation metric definitions, formal propositions, the full state-resolution and multi-emission sensitivity panels, robustness and KS-power calibration, the extended GARCH and SM-CHMM baselines, the full cross-asset panel, and the K -selection and ν_k diagnostics referenced from the discussion.

A.1 CHMM-Model.jl Public API and Reproducer

The companion Julia package `CHMM-Model.jl` (<https://github.com/altashly1/CHMM-Model>) is the entry point used to regenerate every table and figure in this paper from a single seed root under Julia ≥ 1.12 . The public surface is a single `CHMM` type, three emission constructors, a unified `fit!` method (log-space Baum-Welch / ECM), and a unified `simulate` method.

```
using CHMM
chmm_n = make_chmm_n(returns; K = 18)    # Gaussian emissions
chmm_t = make_chmm_t(returns; K = 18)    # Student-t emissions
chmm_l = make_chmm_l(returns; K = 18)    # Laplace emissions
chmm_ged = make_chmm_ged(returns; K = 18) # GED (per-state shape p_k)
fit!(chmm_n; max_iter = 60, tol = 1e-4)  # log-space Baum-Welch
paths = simulate(chmm_n; T = 2516, n = 1000) # 1000 paths of length T
```

The same `simulate` interface is used for every benchmark generator (Bootstrap, Gaussian i.i.d., Laplace i.i.d., GARCH(1,1)) so that the seven-metric panel evaluates each row on byte-identical windows. The Pipeline B Student-t copula sampler is exposed as `simulate_copula(...; nu = 6)` on the per-asset CHMM marginals; the rank-reordering construction is one function call. Running `julia -project=. run_full_rebuild.jl` from the paper repository (<https://github.com/altashly1/CHMM-paper>) reproduces every numerical artifact under the global seed policy of Section 3.5; `Manifest.toml` pins the dependency graph.

Sub-seed derivation rule. The deterministic global seed root is 20260420; per-experiment sub-seeds follow the additive rule `sub_seed = SEED + offset(experiment) + offset(replicate)`, with experiment offsets in $\{0, 1,000, 100, 10,000, \dots\}$ chosen to avoid collisions across the runners in the public Julia package. For example, the headline simulation in `run_diagnostics.jl` uses `SEED + (path index)` for each per-path random draw, the multi-seed Monte Carlo in `run_multiseed_headline.jl` uses `base 20260422 + 100·(seed index)` for ten independent replicates, and the bracket / shrinkage sweeps add a further per-rate offset. The full sub-seed map is enumerated at the top of each runner script; downstream `Random.seed!()` calls are deterministic and order-independent within each runner.

A.2 CHMM Algorithms and Derivations

This appendix collects the algorithmic content for the CHMM family: the forward-backward recursions and weighted M-step updates, the complete training pseudocode, and per-state degrees-of-freedom diagnostics for CHMM-t.

A.2.1 Forward-Backward Recursions and M-Step Updates

For reference, the in-text Baum-Welch description of Section 3.2 is underpinned by the standard log-space recursions [17, 56]. Define the forward variable $\alpha_t(k) = \mathbb{P}(O_{1:t}, S_t = k \mid \mathcal{M})$ and the backward variable $\beta_t(k) = \mathbb{P}(O_{t+1:T} \mid S_t = k, \mathcal{M})$. The log-space recursions are

$$\log \alpha_1(k) = \log \pi_k + \log b_k(O_1), \quad (10)$$

$$\log \alpha_t(j) = \log \sum_i \exp(\log \alpha_{t-1}(i) + \log T_{ij}) + \log b_j(O_t), \quad t = 2, \dots, T, \quad (11)$$

$$\log \beta_T(k) = 0, \quad (12)$$

$$\log \beta_t(i) = \log \sum_j \exp(\log T_{ij} + \log b_j(O_{t+1}) + \log \beta_{t+1}(j)), \quad t = T-1, \dots, 1, \quad (13)$$

where $\log \text{sumexp}_i(x_i) = x_{\max} + \log \sum_i \exp(x_i - x_{\max})$. The posterior state-occupancy $\gamma_t(k)$ and pair-occupancy $\xi_t(i, j)$ quantities are

$$\gamma_t(k) = \frac{\alpha_t(k) \beta_t(k)}{\sum_j \alpha_t(j) \beta_t(j)}, \quad \xi_t(i, j) = \frac{\alpha_t(i) T_{ij} b_j(O_{t+1}) \beta_{t+1}(j)}{\sum_{m,n} \alpha_t(m) T_{mn} b_n(O_{t+1}) \beta_{t+1}(n)}, \quad (14)$$

and the closed-form M-step updates are

$$\pi_k^{\text{new}} = \gamma_1(k), \quad \mu_k^{\text{new}} = \frac{\sum_t \gamma_t(k) O_t}{\sum_t \gamma_t(k)}, \quad (\sigma_k^{\text{new}})^2 = \frac{\sum_t \gamma_t(k) (O_t - \mu_k^{\text{new}})^2}{\sum_t \gamma_t(k)}, \quad T_{ij}^{\text{new}} = \frac{\sum_{t=1}^{T-1} \xi_t(i, j)}{\sum_{t=1}^{T-1} \gamma_t(i)}. \quad (15)$$

The convergence criterion is $|\mathcal{L}^{(n)} - \mathcal{L}^{(n-1)}| < 10^{-4}$ with $\mathcal{L}^{(n)} = \log \text{sumexp}_k \log \alpha_T^{(n)}(k)$, and simulation draws $S_1 \sim \text{Categorical}(\boldsymbol{\pi})$ and then iterates $G_t \sim \mathcal{N}(\mu_{S_t}, \sigma_{S_t}^2)$, $S_{t+1} \sim \text{Categorical}(\mathbf{T}_{S_t, \cdot})$ with no jump process.

CHMM-t M-step (per-state ECM, closed form given $u_{t,k}$). For Student-t emissions $b_k(x) = t_{\nu_k}(x; \mu_k, \sigma_k)$ the ECM formulation of Peel and McLachlan [18], Liu and Rubin [19] treats the latent precision as an augmented E-step quantity,

$$u_{t,k} = \frac{\nu_k + 1}{\nu_k + ((O_t - \mu_k)/\sigma_k)^2}, \quad (16)$$

so that conditional on ν_k the location and scale admit closed-form weighted updates,

$$\mu_k^{\text{new}} = \frac{\sum_t \gamma_t(k) u_{t,k} O_t}{\sum_t \gamma_t(k) u_{t,k}}, \quad (\sigma_k^{\text{new}})^2 = \frac{\sum_t \gamma_t(k) u_{t,k} (O_t - \mu_k^{\text{new}})^2}{\sum_t \gamma_t(k)}. \quad (17)$$

The degrees-of-freedom parameter ν_k is then refined by a one-dimensional golden-section search on the per-state Q -function $Q_k(\nu) = \sum_t \gamma_t(k) \log t_{\nu}(O_t; \mu_k, \sigma_k)$ over the bracket $(\nu_{\min}, \nu_{\max}) = (2.1, 50)$. This two-block ECM preserves the monotonicity of the standard EM guarantee on the compact bracket $[\nu_{\min}, \nu_{\max}]$, with the search invariant $Q_k(\nu^{(n+1)}) \geq Q_k(\nu^{(n)})$ at every iteration; Proposition 1

states the guarantee in full. We extend this M-step with an optional exponential shrinkage prior on $1/\nu_k$, corresponding to the penalised objective

$$Q_k^{\text{pen}}(\nu) = \sum_t \gamma_t(k) \log t_\nu(O_t; \mu_k, \sigma_k) - \lambda / \nu, \quad (18)$$

maximised by the same golden-section search ($\lambda = 0$ recovers the unpenalised Peel and McLachlan [18], Liu and Rubin [19] ECM; the rate sweep and its effect on simulated IS kurtosis are in Section 5).

CHMM-L M-step (closed-form weighted Laplace MLE). For Laplace emissions $b_k(x) = \frac{1}{2b_k} \exp(-|x - \mu_k|/b_k)$ the weighted-MLE estimators are available in closed form:

$$\mu_k^{\text{new}} = \text{WeightedMedian}(O_{1:T}; \gamma_{1:T}(k)), \quad b_k^{\text{new}} = \frac{\sum_t \gamma_t(k) |O_t - \mu_k^{\text{new}}|}{\sum_t \gamma_t(k)}. \quad (19)$$

The weighted median is computed by cumulative-weight bracketing with linear interpolation between adjacent order statistics at the half-total weight crossing. The Laplace M-step carries no shape parameter, so CHMM-L is strictly cheaper per iteration than CHMM-t (which requires the per-state Q_k search).

CHMM-GED M-step (three-stage ECM with closed-form scale). For Generalized Error Distribution emissions $b_k(x) = \text{GED}(x; \mu_k, \alpha_k, p_k) = \frac{p_k}{2\alpha_k \Gamma(1/p_k)} \exp(-(|x - \mu_k|/\alpha_k)^{p_k})$ the M-step decomposes into three CM updates per state. Given current $(\mu_k^{(n)}, \alpha_k^{(n)}, p_k^{(n)})$, CM-1 updates the location by minimising the per-state weighted L^{p_k} loss,

$$\mu_k^{(n+1)} = \arg \min_{\mu \in [\hat{\mu}_k - 5\alpha_k, \hat{\mu}_k + 5\alpha_k]} \sum_t \gamma_t(k) |O_t - \mu|^{p_k^{(n)}}, \quad (20)$$

by a 40-iteration golden-section search; CM-2 updates the scale in closed form given $(\mu_k^{(n+1)}, p_k^{(n)})$,

$$\alpha_k^{(n+1)} = \left[\frac{p_k^{(n)}}{W_k} \sum_t \gamma_t(k) |O_t - \mu_k^{(n+1)}|^{p_k^{(n)}} \right]^{1/p_k^{(n)}}, \quad W_k = \sum_t \gamma_t(k); \quad (21)$$

and CM-3 updates the shape by maximising the per-state Q -function over the bracket,

$$p_k^{(n+1)} = \arg \max_{p \in [p_{\min}, p_{\max}]} \sum_t \gamma_t(k) \log b_k(O_t; \mu_k^{(n+1)}, \alpha_k^{(n+1)}, p), \quad (22)$$

again by a 40-iteration golden-section search over $[p_{\min}, p_{\max}] = [0.5, 3.0]$. Equation (21) is the unique zero of $\partial Q_k / \partial \alpha$, so CM-2 is exact maximization. CM-1 and CM-3 are bracketed maximization. Each CM step is therefore a partial maximization, and the standard ECM monotonicity result [60] applies on the compact bracket; Proposition 1 states the guarantee in full. Boundary cases: $p_k = 2$ recovers Gaussian (with $\sigma = \alpha/\sqrt{2}$); $p_k = 1$ recovers Laplace (with $b = \alpha$).

A.2.2 Algorithm Descriptions

This section provides detailed pseudocode for the main algorithms used in this study. Algorithm 1 is the unified EM procedure shared across all four emission families. It makes explicit the *shared scaffold* (quantile-based initialization, log-space forward-backward, transition update) and the four

family-specific branches (Gaussian M-step for CHMM-N, ECM with one-dimensional golden-section ν -search for CHMM-t, closed-form weighted-median + weighted-MAD for CHMM-L, three-stage CM with bracketed p_k for CHMM-GED); the branch points are lines 3, 13, and 30. Visualizing the four variants as a single algorithm with four M-step branches makes the architectural contribution of the paper (one EM engine, four interchangeable emission families) immediate from the pseudocode. Algorithm 2 presents the Viterbi algorithm for decoding the most likely hidden state sequence. Algorithm 3 describes the CHMM simulation procedure for generating synthetic return paths. Algorithm 4 describes the cross-asset rank-reordering simulator for copula-based multi-asset synthesis. Algorithm 5 details the SIM regression-and-resampling pipeline.

Algorithm 1 Unified EM algorithm for the CHMM family. Lines 3, 13, and 30 branch on the emission family $F \in \{\mathcal{N}, t, L, \text{GED}\}$; every other line is identical across families. CHMM-N and CHMM-L instantiate classical EM; CHMM-t and CHMM-GED instantiate ECM sequences whose final CM step updates the per-state shape parameter (ν_k for CHMM-t, p_k for CHMM-GED) by a one-dimensional golden-section search on the per-state Q -function. Hyperparameter defaults used throughout this paper: convergence tolerance $\text{tol} = 10^{-4}$ on $\Delta\mathcal{L}$, $\text{max_iter} = 60$, 40-iteration golden-section sub-step on every bracketed CM, CHMM-t bracket $(\nu_{\min}, \nu_{\max}) = (2.1, 50)$ with initial $\nu^{(0)} = 10$, CHMM-GED bracket $(p_{\min}, p_{\max}) = (0.5, 3.0)$ with initial $p^{(0)} = 1.5$ and μ_k search interval $[\hat{\mu}_k - 5\alpha_k, \hat{\mu}_k + 5\alpha_k]$.

Require: Observations $O_{1:T}$, number of states K , tolerance tol , max_iter , emission family $F \in \{\mathcal{N}, t, L, \text{GED}\}$, if $F = t$ bracket $(\nu_{\min}, \nu_{\max})$ and initial $\nu^{(0)}$, if $F = \text{GED}$ bracket $(p_{\min}, p_{\max})$ and initial $p^{(0)}$.

Ensure: Transition matrix $\mathbf{T}$, per-state emission parameters $\theta_{1:K}$, initial distribution π .

- 1: **Quantile-Based Initialization.**
- 2: Sort $O^{(\text{sort})} \leftarrow \text{sort}(O_{1:T})$ and partition into K equal chunks $\{C_1, \dots, C_K\}$.
- 3: **for** $k = 1$ to K **do**
- 4: **switch** F
- 5: $F = \mathcal{N}$: $\mu_k \leftarrow \text{mean}(C_k)$, $\sigma_k \leftarrow \max(\text{std}(C_k), 10^{-6})$.
- 6: $F = t$: $\mu_k \leftarrow \text{mean}(C_k)$, $\sigma_k \leftarrow \max(\text{std}(C_k), 10^{-6})$, $\nu_k \leftarrow \nu^{(0)}$.
- 7: $F = L$: $\mu_k \leftarrow \text{median}(C_k)$, $b_k \leftarrow \max(\text{mean}(|C_k - \mu_k|), 10^{-6})$.
- 8: $F = \text{GED}$: $\mu_k \leftarrow \text{mean}(C_k)$, $\alpha_k \leftarrow \max(\text{std}(C_k), 10^{-6})$, $p_k \leftarrow p^{(0)}$.
- 9: **end for**
- 10: $T_{ij} \leftarrow 1/K$; $\pi_k \leftarrow 1/K$; $\mathcal{L}_{\text{prev}} \leftarrow -\infty$.
- 11: **EM Loop.**
- 12: **for** $n = 1$ to max_iter **do**
- 13: **E-Step, per-family emission log-likelihood.**
- 14: **switch** F
- 15: $F = \mathcal{N}$: $\log B_{t,k} \leftarrow \log \mathcal{N}(O_t | \mu_k, \sigma_k^2)$.
- 16: $F = t$: $\log B_{t,k} \leftarrow \log t_{\nu_k}(O_t; \mu_k, \sigma_k)$.
- 17: $F = L$: $\log B_{t,k} \leftarrow -\log(2b_k) - |O_t - \mu_k|/b_k$.
- 18: $F = \text{GED}$: $\log B_{t,k} \leftarrow \log p_k - \log(2\alpha_k) - \log \Gamma(1/p_k) - (|O_t - \mu_k|/\alpha_k)^{p_k}$.
- 19: **E-Step, shared log-space forward-backward.**
- 20: $\log \alpha_1(k) \leftarrow \log \pi_k + \log B_{1,k}$.
- 21: **for** $t = 2$ to T , $j = 1$ to K **do**
- 22: $\log \alpha_t(j) \leftarrow \underset{i}{\text{logsumexp}}(\log \alpha_{t-1}(i) + \log T_{ij}) + \log B_{t,j}$.
- 23: **end for**
- 24: $\log \beta_T(k) \leftarrow 0$.
- 25: **for** $t = T - 1$ down to 1 , $i = 1$ to K **do**
- 26: $\log \beta_t(i) \leftarrow \underset{j}{\text{logsumexp}}(\log T_{ij} + \log B_{t+1,j} + \log \beta_{t+1}(j))$.
- 27: **end for**
- 28: Compute $\gamma_t(k)$ and $\xi_t(i, j)$ from equation (14).
- 29: **Shared transition update.** $T_{ij} \leftarrow \sum_t \xi_t(i, j) / \sum_t \gamma_t(i)$; $\pi_k \leftarrow \gamma_1(k)$.
- 30: **M-Step, per-family emission update.**
- 31: **switch** F
- 32: $F = \mathcal{N}$ {classical Baum-Welch [16].}
- 33: $\mu_k \leftarrow \sum_t \gamma_t(k) O_t / \sum_t \gamma_t(k)$.
- 34: $\sigma_k \leftarrow \max(\sqrt{\sum_t \gamma_t(k) (O_t - \mu_k)^2 / \sum_t \gamma_t(k)}, 10^{-6})$.
- 35: $F = t$ {ECM of Peel and McLachlan [18], Liu and Rubin [19]; closed-form (μ_k, σ_k) given ν_k , then 1D search on ν_k .}
- 36: $u_{t,k} \leftarrow (\nu_k + 1) / (\nu_k + ((O_t - \mu_k) / \sigma_k)^2)$ for all t, k .
- 37: $\mu_k \leftarrow \sum_t \gamma_t(k) u_{t,k} O_t / \sum_t \gamma_t(k) u_{t,k}$.
- 38: $\sigma_k \leftarrow \max(\sqrt{\sum_t \gamma_t(k) u_{t,k} (O_t - \mu_k)^2 / \sum_t \gamma_t(k) u_{t,k}}, 10^{-6})$.
- 39: $\nu_k \leftarrow \arg \max_{\nu \in [\nu_{\min}, \nu_{\max}]} \sum_t \gamma_t(k) \log t_{\nu}(O_t; \mu_k, \sigma_k)$ {40-iter golden-section search.}
- 40: $F = L$ {closed-form weighted Laplace MLE.}
- 41: $\mu_k \leftarrow \text{WeightedMedian}(O_{1:T}, \gamma_{1:T}(k))$ {cumulative-weight bracketing with interpolation at the $W_k/2$ crossing.}
- 42: $b_k \leftarrow \max(\sum_t \gamma_t(k) |O_t - \mu_k| / \sum_t \gamma_t(k), 10^{-6})$.
- 43: $F = \text{GED}$ {three-stage ECM: bracketed μ_k , closed-form α_k , bracketed p_k .}
- 44: $\mu_k \leftarrow \arg \min_{\mu \in [\hat{\mu}_k - 5\alpha_k, \hat{\mu}_k + 5\alpha_k]} \sum_t \gamma_t(k) |O_t - \mu|^{p_k}$ {40-iter golden-section search.}
- 45: $\alpha_k \leftarrow \max\left(\left[(p_k/W_k) \sum_t \gamma_t(k) |O_t - \mu_k|^{p_k}\right]^{1/p_k}, 10^{-6}\right)$ where $W_k = \sum_t \gamma_t(k)$ {closed form, unique zero of $\partial Q_k / \partial \alpha$.}
- 46: $p_k \leftarrow \arg \max_{p \in [p_{\min}, p_{\max}]} \sum_t \gamma_t(k) \log \text{GED}(O_t; \mu_k, \alpha_k, p)$ {40-iter golden-section search.}
- 47: **Convergence check.** $\mathcal{L}^{(n)} \leftarrow \underset{k}{\text{logsumexp}}(\log \alpha_T(k))$.
- 48: **if** $|\mathcal{L}^{(n)} - \mathcal{L}_{\text{prev}}| < \text{tol}$ **then**

Algorithm 2 Viterbi decoding for the most likely hidden state sequence

Require: Observations $O_{1:T}$, Trained CHMM $\mathcal{M} = (\mathbf{T}, \{\mu_k, \sigma_k\}, \boldsymbol{\pi})$

Ensure: Most probable state sequence $S_{1:T}^*$

```
1: for  $k = 1$  to  $K$  do
2:    $\log \delta_1(k) \leftarrow \log(1/K) + \log \mathcal{N}(O_1 \mid \mu_k, \sigma_k^2)$ .
3: end for
4: for  $t = 2$  to  $T$  do
5:   for  $j = 1$  to  $K$  do
6:      $\text{vals}(j) \leftarrow \log \delta_{t-1}(i) + \log T_{ij}$  for all  $i$ .
7:      $\log \delta_t(j) \leftarrow \max_i \text{vals}(i) + \log \mathcal{N}(O_t \mid \mu_j, \sigma_j^2)$ .
8:      $\psi_t(j) \leftarrow \arg \max_i \text{vals}(i)$ .
9:   end for
10: end for
11:  $S_T^* \leftarrow \arg \max_k \log \delta_T(k)$ .
12: for  $t = T - 1$  down to  $1$  do
13:    $S_t^* \leftarrow \psi_{t+1}(S_{t+1}^*)$ .
14: end for
15: return  $S_{1:T}^*$ 
```

Algorithm 3 CHMM simulation of synthetic excess growth rate paths

Require: Trained CHMM $\mathcal{M} = (\mathbf{T}, \{\mu_k, \sigma_k\}, \boldsymbol{\pi})$, Number of steps M , Number of paths P

Ensure: Simulated return paths $\{\hat{G}_{1:M}^{(p)}\}_{p=1}^P$

```
1: for  $p = 1$  to  $P$  do
2:   Sample initial state  $S_1 \sim \text{Categorical}(\boldsymbol{\pi})$ .
3:   for  $t = 1$  to  $M$  do
4:     Emit observation  $\hat{G}_t^{(p)} \sim \mathcal{N}(\mu_{S_t}, \sigma_{S_t}^2)$ .
5:     if  $t < M$  then
6:       Transition to next state  $S_{t+1} \sim \text{Categorical}(\mathbf{T}_{S_t, \cdot})$ .
7:     end if
8:   end for
9: end for
10: return  $\{\hat{G}_{1:M}^{(p)}\}_{p=1}^P$ 
```

Algorithm 4 Cross-asset copula simulation via rank reordering

Require: Fitted per-asset CHMMs $\{\mathcal{M}_j\}_{j=1}^d$, correlation matrix Σ (positive semi-definite (PSD), unit diagonal), copula family $\mathcal{C} \in \{\text{Gaussian}, t_\nu\}$, path length T , number of paths P

Ensure: Cross-asset path tensor $\{\hat{G}_{j,t}^{(p)}\}$ of shape $T \times d \times P$

```
1: for  $p = 1$  to  $P$  do
2:   Draw copula sample  $\mathbf{U}^{(p)} \in [0, 1]^{T \times d}$  from  $\mathcal{C}(\Sigma)$ :
3:    $L \leftarrow \text{Cholesky}(\Sigma)$ ,  $Z \in \mathbb{R}^{T \times d} \sim \mathcal{N}(0, I_d)$  i.i.d.
4:    $Y \leftarrow ZL^\top$ 
5:   if  $\mathcal{C} = \text{Gaussian}$  then
6:      $\mathbf{U}^{(p)} \leftarrow \Phi(Y)$  componentwise
7:   else
8:     Draw  $W \sim \chi_\nu^2/\nu$  i.i.d. for  $t = 1, \dots, T$ ;  $X_{t,\cdot} \leftarrow Y_{t,\cdot}/\sqrt{W_t}$ 
9:      $\mathbf{U}^{(p)} \leftarrow t_\nu(X)$  componentwise
10:  end if
11:  for  $j = 1$  to  $d$  do
12:    Simulate  $\tilde{g}_{j,1:T}^{(p)}$  from  $\mathcal{M}_j$  via Algorithm 3.
13:    Compute order statistics  $\tilde{g}_{j,(1)}^{(p)} \leq \dots \leq \tilde{g}_{j,(T)}^{(p)}$ .
14:    Compute ranks  $r_{j,t}^{(p)} \leftarrow \text{rank}(U_{j,t}^{(p)})$  for  $t = 1, \dots, T$ .
15:     $\hat{G}_{j,t}^{(p)} \leftarrow \tilde{g}_{j,(r_{j,t}^{(p)})}^{(p)}$  for  $t = 1, \dots, T$ .
16:  end for
17: end for
18: return  $\{\hat{G}_{j,t}^{(p)}\}$ 
```

Algorithm 5 Single Index Model (SIM) construction and simulation

Require: IS return matrix $\mathbf{R} \in \mathbb{R}^{T \times d}$, market-column index m , market CHMM $\mathcal{M}_m$, path length T , number of paths P

Ensure: Cross-asset path tensor $\{\hat{G}_{j,t}^{(p)}\}$ of shape $T \times d \times P$

```
1: Ordinary least squares (OLS) fit. For  $j \neq m$ :
2:    $\hat{\beta}_j \leftarrow \text{Cov}(\mathbf{R}_{:,j}, \mathbf{R}_{:,m})/\text{Var}(\mathbf{R}_{:,m})$ 
3:    $\hat{\alpha}_j \leftarrow \overline{\mathbf{R}_{:,j}} - \hat{\beta}_j \overline{\mathbf{R}_{:,m}}$ 
4:    $\hat{\eta}_{j,t} \leftarrow R_{t,j} - \hat{\alpha}_j - \hat{\beta}_j R_{t,m}$ ,  $R_j^2 \leftarrow 1 - \sum_t \hat{\eta}_{j,t}^2 / \sum_t (R_{t,j} - \overline{\mathbf{R}_{:,j}})^2$ 
5: for  $p = 1$  to  $P$  do
6:   Simulate market path  $G_{m,1:T}^{(p)}$  from  $\mathcal{M}_m$  via Algorithm 3.
7:    $\hat{G}_{m,t}^{(p)} \leftarrow G_{m,t}^{(p)}$ .
8:   for  $j \neq m$  do
9:     Draw residual indices  $\{\iota_t\}_{t=1}^T \sim \text{Uniform}(\{1, \dots, T\})$ .
10:     $\hat{G}_{j,t}^{(p)} \leftarrow \hat{\alpha}_j + \hat{\beta}_j G_{m,t}^{(p)} + \hat{\eta}_{j,\iota_t}$ .
11:  end for
12: end for
13: return  $\{\hat{G}_{j,t}^{(p)}\}$ 
```

A.2.3 Per-State Shape Diagnostics: CHMM-t and CHMM-GED

This appendix backs the discussion in Section 5 of the per-state shape allocation under CHMM-t and CHMM-GED on SPY IS at $K = 18$ (seed = 20260420).

CHMM-t per-state ν_k . Figure 7 plots the per-state ν_k histogram for the $K = 18$ CHMM-t fit. Thirteen of the eighteen states rest at the upper ECM bracket ($\nu_k = 50$), and only two states lie near the lower bracket ($\nu_2 = 2.19$, $\nu_4 = 2.10$). The simulated IS mixture kurtosis of 17.34 is driven by the posterior mass routed to those two very heavy-tailed states, not by a systemic collapse to the lower bracket.




Figure 7. Per-state degrees-of-freedom ν_k for the CHMM-t fit at $K = 18$ on SPY IS. The dashed red line marks the lower ECM bracket $\nu_{\min} = 2.1$. Only two states ($\nu_2 = 2.19$, $\nu_4 = 2.10$) sit near the lower bracket; the median ν_k is at the upper bracket.

CHMM-GED per-state p_k (Gaussian-bulk / Laplace-tail bimodality). The body Section 4.3 reports and discusses the $\hat{p}_k$ partition (Figure 4); see also Table 7 below for the parallel CHMM-t bracket sweep. Briefly, the $K = 18$ CHMM-GED fit on SPY IS is bimodal: thirteen states attain the upper ECM bracket at $p_k = 3.0$, one state lands at intermediate $p_k = 1.6$, and four states cluster in the Laplace-shape regime $p_k \in [0.86, 1.24]$, concentrated in the high-volatility ranks. The smallest fitted p_k on SPY is 0.86, well clear of every $p_{\min}$ tested in the bracket sweep, so the lower bracket is non-binding. The bimodal partition replicates across 10 Monte Carlo seeds and across the cross-asset universe (Appendix A.7).

Table 7 reports the seven-metric panel for CHMM-t fits whose ECM bracket lower bound is lifted from $\nu_{\min} = 2.1$ through $\{2.5, 3.0, 4.0\}$, plus the Gaussian limit ($\nu = \infty$, equivalent to CHMM-N). Lifting the bracket does not reduce the IS kurtosis overshoot: the simulated kurtosis moves within the 12.5–14.0 band across bracket floors and only collapses to 5.05 in the full Gaussian limit. The overshoot is therefore a feature of the Student-t emission family at $K = 18$ and not a lower-bracket artefact.

Table 8 reports the analogous bracket sensitivity for CHMM-GED on $p_{\max} \in \{2.0, 2.5, 3.0, 3.5, 4.0\}$ (with $p_{\min} = 0.5$ fixed) and on $p_{\min} \in \{0.5, 0.7, 0.85\}$ (with $p_{\max} = 3.0$ fixed). The headline reads cleanly. The Gaussian-like state count is invariant at 13/18 across every upper bracket; only the

Table 7. CHMM-t $\nu_{\min}$ bracket sensitivity at $K = 18$ on SPY (seed = 20260420, 1,000 paths). AD = Anderson–Darling pass rate; W_1 = Wasserstein-1; H = Hellinger distance (definitions in Appendix A.3).

$\nu_{\min}$	KS IS	AD IS	KS OoS	AD OoS	Kurt IS	Kurt OoS	ACF-MAE	W_1	H
2.1	95.7	90.2	86.2	82.0	13.95	10.08	0.0546	0.100	0.0759
2.5	95.8	90.6	84.2	79.1	15.12	10.34	0.0547	0.100	0.0759
3.0	93.8	89.5	85.3	80.4	12.25	9.65	0.0548	0.099	0.0760
4.0	95.5	89.5	85.0	80.3	13.25	9.39	0.0550	0.101	0.0759
∞ (Gauss)	94.1	86.7	84.2	75.5	5.03	4.46	0.0510	0.110	0.0809
<i>Observed</i>	–	–	–	–	7.68	5.29	–	–	–

redistribution between intermediate and Laplace-like states moves slightly with $p_{\max}$, and the total heavy-shape count stays in $\{4, 5\}$. Tightening $p_{\max}$ from 4.0 to the canonical 3.0 costs about 4 nats of incomplete-data log-likelihood. The $p_{\min}$ sweep is fully degenerate: every $p_{\min} \in [0.5, 0.85]$ produces an identical fit because the smallest observed $\hat{p}_k$ on SPY is 0.86, so the lower bracket is non-binding.

Table 8. CHMM-GED bracket sensitivity at $K = 18$ on SPY (seed = 20260420, 1,000 paths). $\hat{p}_{\min}$, $\hat{p}_{\text{med}}$, $\hat{p}_{\max}$ are the across-state minimum, median, and maximum of the fitted $\hat{p}_k$. nG / nI / nL / nS are the partition counts (Gaussian-like $\hat{p}_k \geq 1.85$, intermediate $1.30 \leq \hat{p}_k < 1.85$, Laplace-like $0.85 \leq \hat{p}_k < 1.30$, super-Laplace $\hat{p}_k < 0.85$). LL is the incomplete-data log-likelihood at convergence.

Sweep	$p_{\min}$	$p_{\max}$	LL	$\hat{p}_{\min}$	nG	nI	nL	nS	KS IS	AD IS	KS OoS	AD OoS	Kurt IS	ACF-MAE	W_1	H
$p_{\max}$ sweep	0.5	2.0	−4732.0	0.89	13	0	5	0	95.8	91.0	83.8	81.2	5.02	0.0541	0.099	0.0781
	0.5	2.5	−4725.3	0.87	13	2	3	0	96.1	92.4	84.3	80.1	5.09	0.0549	0.098	0.0778
	0.5	3.0	−4720.3	0.86	13	1	4	0	96.3	92.7	83.8	81.3	5.05	0.0546	0.098	0.0779
	0.5	3.5	−4717.5	0.86	13	1	4	0	96.3	91.8	86.7	82.4	5.11	0.0544	0.099	0.0782
	0.5	4.0	−4716.3	0.86	13	1	4	0	95.9	91.3	84.4	80.6	5.18	0.0543	0.099	0.0785
$p_{\min}$ sweep	0.50	3.0	−4720.3	0.86	13	1	4	0	96.3	92.7	83.8	81.3	5.05	0.0546	0.098	0.0779
	0.70	3.0	−4720.3	0.86	13	1	4	0	96.3	92.7	83.8	81.3	5.05	0.0546	0.098	0.0779
	0.85	3.0	−4720.3	0.86	13	1	4	0	96.3	92.7	83.8	81.3	5.05	0.0546	0.098	0.0779

A.3 Validation Metric Details

This appendix documents the precise form of the seven per-path metrics referenced in Section 3.5. For each model, 1,000 independent paths of length T are simulated (200 paths for the cross-asset extension) and each metric is computed per path before aggregation.

Kolmogorov–Smirnov (KS) pass rate. The two-sample KS statistic [53, 54] measures the maximum vertical distance between two empirical CDFs,

$$D_{n,m} = \sup_x |F_n(x) - G_m(x)|. \quad (23)$$

The KS pass rate is the fraction of paths for which the test fails to reject distributional equivalence against the reference data at $\alpha = 0.05$.

Anderson–Darling (AD) pass rate. The AD test is the integrated squared CDF deviation with a quadratic tail weighting, making it more sensitive than KS to tail discrepancies. The AD pass rate is the fraction of paths for which the two-sample AD test fails to reject at $\alpha = 0.05$.

Excess kurtosis. Mean simulated excess kurtosis across paths, compared to the observed value. Given Gaussian emissions, the CHMM is expected to underestimate kurtosis relative to the heavy-tailed empirical distribution, and the shortfall is one of the reported diagnostics.

ACF-MAE. Mean absolute error between observed and mean-simulated autocorrelation functions of $|G_t|$ over $L = 252$ lags,

$$\text{ACF-MAE} = \frac{1}{L} \sum_{\tau=1}^L \left| \hat{\rho}_{|G|}^{\text{obs}}(\tau) - \bar{\rho}_{|G|}^{\text{sim}}(\tau) \right|. \quad (24)$$

This is the direct diagnostic for the Rydén et al. limitation.

Wasserstein-1 distance. The mean absolute difference of sorted empirical quantiles, equivalent to the L^1 distance between the empirical CDFs [68],

$$W_1(F, G) = \int_0^1 |F^{-1}(u) - G^{-1}(u)| du. \quad (25)$$

Hellinger distance. A histogram-based overlap distance in $[0, 1]$,

$$H(P, Q) = \frac{1}{\sqrt{2}} \sqrt{\sum_{i=1}^B (\sqrt{p_i} - \sqrt{q_i})^2}, \quad (26)$$

with B equal-width bins spanning the joint support and p_i, q_i the observed and simulated bin probabilities.

Quantile coverage. The fraction of 99 equally-spaced observed quantiles (1st through 99th percentile) falling inside the 5th–95th percentile envelope of the corresponding simulated quantile across 1,000 paths. A coverage of 100% indicates that the simulation envelope fully brackets the observed distribution at every percentile.

Cross-asset metrics. For the cross-asset extension we additionally track the mean Frobenius norm $\|\hat{\Sigma}_{\text{sim}}^{(p)} - \hat{\Sigma}_{\text{obs}}\|_F$ and the off-diagonal Pearson correlation MAE between simulated and observed return matrices, averaged over paths. The off-diagonal MAE is averaged over the $d(d-1)/2$ unique upper-triangular entries of the symmetric correlation matrix (so 15 entries for the six-asset cross-section of Section 4.5), not double-counted across both triangles:

$$\text{MAE}_{\text{off-diag}} = \frac{1}{P} \sum_{p=1}^P \frac{2}{d(d-1)} \sum_{1 \leq i < j \leq d} \left| \hat{\Sigma}_{\text{sim}, ij}^{(p)} - \hat{\Sigma}_{\text{obs}, ij} \right|.$$

A.3.1 Continuous Ranked Probability Score and Diebold-Mariano

The OoS CRPS column of Table 1 reports a proper-scoring-rule complement to the KS pass rate. We use the unbiased sample CRPS estimator: for an ensemble of N simulated paths $\{x_i\}_{i=1}^N$ at time t and observed value y_t ,

$$\widehat{\text{CRPS}}_t = \frac{1}{N} \sum_{i=1}^N |x_i - y_t| - \frac{1}{N(N-1)} \sum_{1 \leq i < j \leq N} |x_i - x_j|. \quad (27)$$

The double sum is computed in $O(N \log N)$ via the sorted-ensemble identity $\sum_{i < j} (x_{(j)} - x_{(i)}) = \sum_i x_{(i)}(2i - N - 1)$. The ensemble at each t is the cross-section of the $N = 1,000$ unconditional simulated paths used throughout Section 4; this yields a marginal-predictive CRPS suitable for an unconditional generative-fidelity assessment, distinct from the conditional one-step-ahead CRPS used in forecasting.

For pairwise comparison we report a Diebold-Mariano test on the per- t loss differential $d_t = \widehat{\text{CRPS}}_t^A - \widehat{\text{CRPS}}_t^B$ with a Newey-West HAC variance under a Bartlett kernel and bandwidth $h = \lfloor T_{\text{OoS}}^{1/3} \rfloor = 8$. Two-sided p -values are reported under the standard-normal null. The body claim of Section 4.3 is: CHMM ties or beats every benchmark on mean OoS CRPS, with Gaussian i.i.d. the only significantly worse comparator.

A.3.2 Christoffersen Conditional-Coverage Panel (Companion to Section 4.6)

The body Section 4.6 reports the unconditional Kupiec LR_{uc} statistic. This appendix reports the full Christoffersen [66] panel: LR_{uc} (unconditional coverage), LR_{ind} (breach independence), and $\text{LR}_{\text{cc}} = \text{LR}_{\text{uc}} + \text{LR}_{\text{ind}}$ (joint conditional coverage), at $\alpha \in \{0.01, 0.05\}$ on the SPY IS and OoS windows. The breach series is constructed from the pooled-archive VaR threshold (the α -quantile of $\text{vec}(\text{sim archive})$ across paths and time), then $\text{br}_t = (R_t \leq \widehat{\text{VaR}}_\alpha)$.


The headline reads as follows. On the unconditional (Kupiec) leg every CHMM variant passes at both levels on both windows; the LR_{uc} statistic for the four CHMM rows at $\alpha = 0.05$ on OoS clusters at 3.03–3.83 (just below the χ_1^2 critical value 3.841), reflecting the integer-breach grid at $T_{\text{OoS}} = 572$ noted in the body table caption. On the independence leg every unconditional generator in the panel (bootstrap, GARCH, and all four CHMM variants) rejects: LR_{ind} is in the range 4.71–56.46 across rows. This is structural rather than CHMM-specific. Volatility-clustering in R_{oos} produces persistent regimes of high and low realised volatility, the OoS-window breaches concentrate inside the high-volatility regime, and a constant-across- t VaR threshold (the construction every unconditional generator uses) cannot track that timing. The natural fix is a regime-conditional VaR that propagates the CHMM latent-state forecast $\mathbb{P}(s_{t+1} | \text{history}_t)$ through the per-state emission-mixture; this is operationally the conditional-VaR companion-paper direction noted in Section 5.

Table 9. Christoffersen conditional-coverage VaR back-test (SPY, seed = 20260420, $K = 18$, 1,000 simulated paths). Critical values: $\chi_1^2(0.05) = 3.841$, $\chi_2^2(0.05) = 5.991$. “br rate” is the empirical breach rate; LR_{uc} is the Kupiec statistic ($\sim \chi_1^2$ under correct unconditional coverage); LR_{ind} is the Christoffersen independence statistic ($\sim \chi_1^2$ under independent breaches); $LR_{cc} = LR_{uc} + LR_{ind}$ is the joint conditional-coverage statistic ($\sim \chi_2^2$). PASS if statistic is below critical. Every unconditional generator (i.i.d. bootstrap, GARCH, and all four CHMM variants) rejects independence on OoS at $\alpha = 0.05$, consistent with breach clustering driven by volatility-clustering in R_{oos} rather than a CHMM-specific mis-specification of conditional dynamics.


Model	Win	α	breaches	br rate	Kupiec		Christ. ind		Christ. cc	
					LR_{uc}	p	LR_{ind}	p	LR_{cc}	p
Bootstrap	IS	0.01	26	1.03%	0.03	0.87	15.28	0.00	15.31	0.00
Bootstrap	IS	0.05	126	5.01%	0.00	0.99	56.46	0.00	56.46	0.00
Bootstrap	OoS	0.01	5	0.87%	0.10	0.76	13.18	0.00	13.28	0.00
Bootstrap	OoS	0.05	19	3.32%	3.83	0.05	5.26	0.02	9.08	0.01
GARCH(1,1)	IS	0.01	31	1.23%	1.28	0.26	12.45	0.00	13.72	0.00
GARCH(1,1)	IS	0.05	139	5.52%	1.41	0.24	45.38	0.00	46.79	0.00
GARCH(1,1)	OoS	0.01	6	1.05%	0.01	0.91	20.87	0.00	20.89	0.00
GARCH(1,1)	OoS	0.05	24	4.20%	0.82	0.37	5.87	0.02	6.69	0.04
CHMM-N	IS	0.01	20	0.79%	1.15	0.28	12.80	0.00	13.95	0.00
CHMM-N	IS	0.05	123	4.89%	0.07	0.80	47.33	0.00	47.40	0.00
CHMM-N	OoS	0.01	3	0.52%	1.58	0.21	18.98	0.00	20.56	0.00
CHMM-N	OoS	0.05	19	3.32%	3.83	0.05	5.26	0.02	9.08	0.01
CHMM-t	IS	0.01	22	0.87%	0.42	0.52	11.62	0.00	12.04	0.00
CHMM-t	IS	0.05	127	5.05%	0.01	0.91	55.54	0.00	55.55	0.00
CHMM-t	OoS	0.01	4	0.70%	0.58	0.45	15.53	0.00	16.12	0.00
CHMM-t	OoS	0.05	20	3.50%	3.03	0.08	4.71	0.03	7.74	0.02
CHMM-L	IS	0.01	19	0.76%	1.66	0.20	13.44	0.00	15.11	0.00
CHMM-L	IS	0.05	134	5.33%	0.55	0.46	49.43	0.00	49.98	0.00
CHMM-L	OoS	0.01	2	0.35%	3.26	0.07	9.15	0.00	12.41	0.00
CHMM-L	OoS	0.05	20	3.50%	3.03	0.08	4.71	0.03	7.74	0.02
CHMM-GED	IS	0.01	20	0.79%	1.15	0.28	12.80	0.00	13.95	0.00
CHMM-GED	IS	0.05	126	5.01%	0.00	0.99	56.46	0.00	56.46	0.00
CHMM-GED	OoS	0.01	2	0.35%	3.26	0.07	9.15	0.00	12.41	0.00
CHMM-GED	OoS	0.05	19	3.32%	3.83	0.05	5.26	0.02	9.08	0.01

A.3.3 VaR / ES Envelope Visualization (Companion to Table 4)


Figure 8 visualizes the median and [5%, 95%] envelopes across simulated paths whose numeric values are reported in Table 4.




(a) IS VaR ($\alpha = 0.01$).



(b) IS ES ($\alpha = 0.01$).



(c) OoS VaR ($\alpha = 0.05$).



(d) OoS ES ($\alpha = 0.05$).

Figure 8. VaR / ES envelope diagnostic (SPY, $N_{\text{paths}} = 1,000$; global seed policy in Section 3.5). Each panel shows the simulated median (dot) and [5%, 95%] envelope (error bar) across paths against the observed historical value (dashed red line); panels (a)/(b) are the IS window at $\alpha = 0.01$, panels (c)/(d) are the OoS window at $\alpha = 0.05$, and the x-axis in each panel lists the five generators kept for the volatility-aware conditional-coverage comparison (GARCH(1,1), CHMM-N, CHMM-t, CHMM-L, CHMM-GED). The i.i.d. bootstrap is omitted here because its envelope is by construction matched to the empirical marginal and does not discriminate among volatility-aware models. The y-axis is the annualized log excess growth rate (daily log return scaled by $1/\Delta t$ with $\Delta t = 1/252$), consistent with the $R_{\text{IS}}/R_{\text{OoS}}$ convention used throughout the paper. Headline: every observed red line falls inside the corresponding simulated envelope, confirming that the CHMM family passes the coverage leg of the back-test complementary to the Kupiec likelihood-ratio leg in Section 4.6.

A.4 Formal Theoretical Statements

The following propositions are referenced in Section 3.8 and stated formally here for reference; each is a one-line specialisation of a standard result and the proof sketch is one sentence.

Proposition 1 (ECM monotonicity for CHMM-t and CHMM-GED (and EM monotonicity for CHMM-N, CHMM-L)). *Under compactness of the parameter space (μ_k, σ_k or b_k or $\alpha_k, \nu_k \in [\nu_{\min}, \nu_{\max}]$, $p_k \in [p_{\min}, p_{\max}]$ each in a closed bounded interval, $T_{ij} \geq \epsilon_T > 0$), the two-block CM step for CHMM-t of Section 3.7, namely the closed-form weighted updates of $(\mathbf{T}, \boldsymbol{\pi}, \mu_k, \sigma_k)$ at fixed ν_k , followed by a golden-section maximisation of the per-state Q -function over ν_k , satisfies*

$Q(\theta^{(n+1)} \mid \theta^{(n)}) \geq Q(\theta^{(n)} \mid \theta^{(n)})$, and consequently $\mathcal{L}(\theta^{(n+1)}) \geq \mathcal{L}(\theta^{(n)})$ where $\mathcal{L}$ is the incomplete-data log-likelihood. The same monotonicity holds for the closed-form CHMM-N and CHMM-L M-steps under the standard EM guarantee, and for the three-stage CHMM-GED CM sequence (bracketed μ_k , closed-form α_k , bracketed p_k) of equations (20)–(22).

Proof sketch. Each CM step does not decrease Q by definition (closed-form steps are exact maximisers, golden-section steps are bracketed maxima within the prescribed interval); the standard EM inequality $\mathcal{L}(\theta^{(n+1)}) - \mathcal{L}(\theta^{(n)}) \geq Q(\theta^{(n+1)} \mid \theta^{(n)}) - Q(\theta^{(n)} \mid \theta^{(n)})$ [59, 60] converts Q -monotonicity into log-likelihood monotonicity. Compactness gives boundedness above, hence convergence.

Numerical full-rank check on $\hat{\mathbf{T}}$. The Allman-Matias-Rhodes 2009 identifiability theorem requires $\hat{\mathbf{T}}$ of full rank. We compute the singular-value spectrum of $\hat{\mathbf{T}}$ at $K = 18$ across the four emission families on SPY IS (Table 10). All four matrices are full-rank in the strict sense, with smallest singular value $\sigma_{\min}(\hat{\mathbf{T}}) \in [0.0007, 0.0170]$ and condition numbers in the range 63–1,620. CHMM-GED is the most well-conditioned ($\sigma_{\min} = 0.017$, $\text{cond} = 63.5$); CHMM-t is the least ($\sigma_{\min} = 0.0007$, $\text{cond} = 1,620$), consistent with the per-state ν_k ECM lower-bracket pinning making one or two heavy-tail states near-degenerate. The deflated matrix $\hat{\mathbf{T}} - \mathbf{1}\bar{\pi}^\top$ has numerical rank exactly $K - 1 = 17$ across all four families, the rank statement underlying Section 3.8.

Table 10. Numerical full-rank check on $\hat{\mathbf{T}}$ at $K = 18$ across four CHMM emission families on SPY IS (seed = 20260420). $\sigma_{\min}$, $\sigma_{\max}$, and the condition number $\sigma_{\max}/\sigma_{\min}$ are reported on $\hat{\mathbf{T}}$ itself; $|\lambda_2|$ is the second-largest absolute eigenvalue (the dominant decay mode in the ACF identity, equation (8)); “rank(def)” is the numerical rank of $\hat{\mathbf{T}} - \mathbf{1}\bar{\pi}^\top$ at tolerance 10^{-10} , expected to equal $K - 1 = 17$.

Family	$\sigma_{\min}(\hat{\mathbf{T}})$	$\sigma_{\max}(\hat{\mathbf{T}})$	$\text{cond } \hat{\mathbf{T}}$	$ \lambda_2 $	$\text{rank}(\hat{\mathbf{T}} - \mathbf{1}\bar{\pi}^\top)$
CHMM-N	0.00532	1.0866	204	0.929	17/17
CHMM-t	0.00067	1.0796	1,620	0.914	17/17
CHMM-L	0.00398	1.0928	275	0.896	17/17
CHMM-GED	0.01702	1.0806	63.5	0.911	17/17

Proposition 2 (Identifiability up to label permutation). *Under irreducibility and aperiodicity of $\mathbf{T}$ (Assumption 1), the emission contrast condition (Assumption 2), and pairwise distinctness of the per-state parameter tuples θ_k across $k = 1, \dots, K$, the CHMM parameter is identifiable up to label permutation for CHMM-N (Gaussian) and CHMM-L (Laplace). For CHMM-t (with per-state $\nu_k \in [\nu_{\min}, \nu_{\max}]$) and CHMM-GED (with per-state $p_k \in [p_{\min}, p_{\max}]$) the same conclusion holds modulo the standard caveat that pairwise distinctness must hold jointly across all components of $\theta_k = (\mu_k, \sigma_k, \nu_k)$ or (μ_k, α_k, p_k) , since within either bracket a Gaussian-shaped emission with inflated scale can mimic a Laplace-shaped emission with deflated scale at intermediate quantiles.*

Proof sketch. The HMM identifiability theorem of Allman et al. [61] requires $\mathbf{T}$ of full rank and per-state emissions linearly independent in the space of densities on $\mathbb{R}$. Full-rank $\mathbf{T}$ follows from irreducibility [58]. For CHMM-N and CHMM-L, linear independence of Gaussian and Laplace emissions with distinct location-scale parameters is direct from Yakowitz and Spragins [62], who establish that finite mixtures over location-scale families with strictly decreasing-tail densities are identifiable (the CHMM-N and CHMM-L marginals are special cases). For CHMM-t at fixed shape and CHMM-GED at fixed shape, location-scale identifiability is the same Yakowitz-Spragins result; for finite mixtures of Student- t with varying degrees-of-freedom the standard reference is Lindsay [69], Chapter 3, which gives identifiability via the moment-generating function over the bracketed range

of shape parameters and a non-degeneracy condition on the joint location-scale-shape tuple. The CHMM-GED case under bracketed $p_k \in [p_{\min}, p_{\max}]$ is a parallel application of the moment-matching argument: the GED characteristic function $\hat{b}(t; \mu, \alpha, p) = \exp(i\mu t) M(t; \alpha, p)$ has a unique inverse over the bounded shape range up to permutation of components, modulo the joint-distinctness caveat on (μ_k, α_k, p_k) . Strict linear independence at the boundary cases $p_k = 1$ (Laplace) and $p_k = 2$ (Gaussian) is automatic from Yakowitz-Spragins; in the bracket interior, the joint distinctness condition above is required to rule out the location-scale-shape trade-off.

Proposition 3 (MLE consistency). *Under Assumptions 1–2 and the assumption that the true parameter lies in the interior of a compact Θ , the MLE $\hat{\theta}_T = \arg \max_{\theta \in \Theta} \mathcal{L}_T(\theta)$ is consistent: $\hat{\theta}_T \xrightarrow{P} \theta_0$ as $T \rightarrow \infty$, modulo label permutation.*

Proof reference. Bickel et al. [63] (Theorem 1) and Douc et al. [64] apply directly under our assumption set: irreducibility supplies the mixing condition, finite second moment the moment condition, compactness the parameter-space condition.

Proposition 4 (Marginal preservation under rank reordering). *Let $\tilde{g}_{j,1:T}$ be a sample of size T from the per-asset CHMM marginal of asset j , and let the rank-reordered output be $\hat{g}_{j,t}^{(p)} = \tilde{g}_{j,(r_{j,t}^{(p)})}^{(p)}$ where $r_{j,t}^{(p)}$ is the rank of the copula sample. Then for every asset j the empirical CDF of $\{\hat{g}_{j,t}^{(p)}\}_{t=1}^T$ equals that of $\{\tilde{g}_{j,t}^{(p)}\}_{t=1}^T$.*

Proof sketch. The map $t \mapsto r_{j,t}^{(p)}$ is a permutation almost surely, so the multiset of values is preserved; the empirical CDF depends only on the multiset.

A.5 Spectral ACF Identity: Extended Derivation

Equation (8) is the closed-form mixture-of-eigenvalues form for the absolute-return ACF used in Section 3.8 and invoked throughout Sections 4–5. The bilinear cross-moment identity $\mathbb{E}[|G_t| |G_{t+\tau}|] = \mathbf{m}^\top \text{diag}(\bar{\pi}) \mathbf{T}^\tau \mathbf{m}$ and its eigendecomposition over the non-unit eigenvalues of $\mathbf{T}$ are textbook material in the regime-switching literature (13, Section 22.2; 14, Chapter 3; the moment-generating-function form for general Markov-switching specifications with conditional means and variances depending on the latent state is given by 15); the underlying Markov-chain spectral decomposition is standard [58]. We restate the absolute-return specialisation here as a self-contained theorem with a step-by-step proof, and record the lag-zero behaviour and the complex / non-diagonalisable extensions that the main body does not. Our contribution on this axis is the explicit application of the spectral form to recast the empirical Rydén et al. [11] low- K failure as a rank statement on $\mathbf{T} - \mathbf{1}\bar{\pi}^\top$ (Section 3.8, “Rydén separation as a K -rank statement”) and the empirical demonstration that the rank constraint is the binding axis at moderate K on equity-return data (Sections 4, 5).

Theorem 1 (Mixture-of-eigenvalues identity for the absolute-return ACF). *Let $\{(s_t, G_t)\}_{t \in \mathbb{Z}}$ be a stationary CHMM on $\{1, \dots, K\}$ with irreducible aperiodic transition matrix $\mathbf{T}$ (Assumption 1) and per-state emissions of finite second moment (Assumption 2). Write $\bar{\pi}$ for the unique stationary distribution, $\mathbf{D} = \text{diag}(\bar{\pi})$, $m_k = \mathbb{E}[|G_t| \mid s_t = k]$, $\mathbf{m} = (m_1, \dots, m_K)^\top$, $\mu = \mathbb{E}[G_t] = \bar{\pi}^\top \mathbf{m}$, and $\sigma_{|G|}^2 = \text{Var}(|G_t|)$. Assume $\mathbf{T}$ is diagonalisable with right eigenvectors $\mathbf{v}_k$ and left eigenvectors $\mathbf{w}_k$ biorthonormal, $\mathbf{w}_k^\top \mathbf{v}_l = \delta_{kl}$, indexed so that $\lambda_1 = 1 > |\lambda_2| \geq \dots \geq |\lambda_K|$. Then for every integer lag $\tau \geq 1$,*

$$\rho_{|G|}(\tau) = \sum_{k=2}^K w_k \lambda_k^\tau, \quad w_k = \frac{(\mathbf{m}^\top \mathbf{D} \mathbf{v}_k)(\mathbf{w}_k^\top \mathbf{m})}{\sigma_{|G|}^2}.$$

Proof. The proof is organised in six steps.

Step 1 (Definitions). The autocorrelation of $|G_t|$ at lag $\tau \geq 0$ is, by stationarity,

$$\rho_{|G|}(\tau) = \frac{\text{Cov}(|G_t|, |G_{t+\tau}|)}{\sigma_{|G|}^2} = \frac{\mathbb{E}[|G_t| |G_{t+\tau}|] - \mu^2}{\sigma_{|G|}^2}.$$

The task reduces to evaluating $\mathbb{E}[|G_t| |G_{t+\tau}|]$ in closed form for $\tau \geq 1$.

Step 2 (Conditioning on the latent state pair). For $\tau \geq 1$, by the law of total expectation conditional on $(s_t, s_{t+\tau})$,

$$\mathbb{E}[|G_t| |G_{t+\tau}|] = \sum_{i,j=1}^K \mathbb{P}(s_t = i, s_{t+\tau} = j) \mathbb{E}[|G_t| |G_{t+\tau}| \mid s_t = i, s_{t+\tau} = j].$$

The HMM emission structure gives the conditional independence $G_t \perp G_{t+\tau} \mid (s_t, s_{t+\tau})$ for $\tau \geq 1$: each emission depends only on its current state, and integrating the intermediate states $s_{t+1:t+\tau-1}$ out of the full-sequence factorisation $(\prod_u b_{s_u}(G_u)) p(s_{t+1:t+\tau-1} \mid s_t, s_{t+\tau})$ leaves the endpoint emissions decoupled. Hence

$$\mathbb{E}[|G_t| |G_{t+\tau}| \mid s_t = i, s_{t+\tau} = j] = \mathbb{E}[|G_t| \mid s_t = i] \mathbb{E}[|G_{t+\tau}| \mid s_{t+\tau} = j] = m_i m_j.$$

The joint state probability factors by stationarity and the Markov property, $\mathbb{P}(s_t = i, s_{t+\tau} = j) = \bar{\pi}_i (\mathbf{T}^\tau)_{ij}$. Substituting and noting $[\mathbf{D}\mathbf{T}^\tau]_{ij} = \bar{\pi}_i (\mathbf{T}^\tau)_{ij}$,

$$\mathbb{E}[|G_t| |G_{t+\tau}|] = \mathbf{m}^\top \mathbf{D} \mathbf{T}^\tau \mathbf{m}, \quad \tau \geq 1. \quad (28)$$

Step 3 (Spectral preliminaries). For irreducible aperiodic $\mathbf{T}$, Perron–Frobenius applied to row-stochastic matrices gives $\lambda_1 = 1$ as a simple eigenvalue with all other eigenvalues strictly inside the unit disk [58]. The right eigenvector for λ_1 is $\mathbf{v}_1 = \mathbf{1}$, since $\mathbf{T}\mathbf{1} = \mathbf{1}$ from the row-sum constraint, and the left eigenvector is $\mathbf{w}_1 = \bar{\pi}$, since $\bar{\pi}^\top \mathbf{T} = \bar{\pi}^\top$ by definition of the stationary distribution. Biorthonormality $\mathbf{w}_1^\top \mathbf{v}_1 = \bar{\pi}^\top \mathbf{1} = 1$ is automatic since $\bar{\pi}$ is a probability vector.

Step 4 (Spectral decomposition of $\mathbf{T}^\tau$). Diagonalisability with biorthonormality gives the dyadic expansion

$$\mathbf{T}^\tau = \sum_{k=1}^K \lambda_k^\tau \mathbf{v}_k \mathbf{w}_k^\top = \mathbf{1} \bar{\pi}^\top + \sum_{k=2}^K \lambda_k^\tau \mathbf{v}_k \mathbf{w}_k^\top,$$

isolating the dominant ($k = 1$) projection from the contractive spectrum. The dominant projector $\mathbf{P}_1 = \mathbf{1} \bar{\pi}^\top$ is the rank-one stationary projection: $\mathbf{P}_1 \mathbf{x} = (\bar{\pi}^\top \mathbf{x}) \mathbf{1}$.

Step 5 (Evaluation of the bilinear form). Substituting the spectral decomposition into (28),

$$\mathbb{E}[|G_t| |G_{t+\tau}|] = \mathbf{m}^\top \mathbf{D} \mathbf{1} \bar{\pi}^\top \mathbf{m} + \sum_{k=2}^K \lambda_k^\tau (\mathbf{m}^\top \mathbf{D} \mathbf{v}_k) (\mathbf{w}_k^\top \mathbf{m}).$$

The $k = 1$ term simplifies to μ^2 : $\mathbf{D}\mathbf{1}$ is the column vector $\bar{\pi}$, so $\mathbf{m}^\top \mathbf{D}\mathbf{1} = \bar{\pi}^\top \mathbf{m} = \mu$, and likewise $\bar{\pi}^\top \mathbf{m} = \mu$, giving the product $\mu \cdot \mu = \mu^2$. Denoting the $k \geq 2$ scalar coefficients

$$c_k = (\mathbf{m}^\top \mathbf{D} \mathbf{v}_k) (\mathbf{w}_k^\top \mathbf{m}),$$

we obtain

$$\mathbb{E}[|G_t| |G_{t+\tau}|] = \mu^2 + \sum_{k=2}^K c_k \lambda_k^\tau, \quad \tau \geq 1.$$

Step 6 (Autocovariance and autocorrelation). Subtracting μ^2 from Step 1 cancels the dominant term, leaving

$$\text{Cov}(|G_t|, |G_{t+\tau}|) = \sum_{k=2}^K c_k \lambda_k^\tau, \quad \tau \geq 1,$$

which is equation (7). Dividing by $\sigma_{|G|}^2 = \gamma(0)$ gives $\rho_{|G|}(\tau) = \sum_{k \geq 2} w_k \lambda_k^\tau$ with $w_k = c_k / \sigma_{|G|}^2$, equation (8). $\square$

Lag zero, the sum of weights, and the within-state white-noise contribution. The bilinear identity (28) fails at $\tau = 0$, where $G_t = G_{t+\tau}$ and the conditional product is the per-state second moment $M_i = \mathbb{E}[G_t^2 | s_t = i]$ rather than m_i^2 . The eigenvalue completeness $\sum_{k=1}^K \mathbf{v}_k \mathbf{w}_k^\top = I$ instead gives the identity

$$\sum_{k=2}^K c_k = \mathbf{m}^\top \mathbf{D} \mathbf{m} - \mu^2 = \sum_{i=1}^K \bar{\pi}_i m_i^2 - \mu^2 = \text{Var}(m_{s_t}),$$

the between-state variance of $|G_t|$, equal to the variance of the conditional-mean random variable m_{s_t} under $\bar{\pi}$. By the law of total variance, $\sigma_{|G|}^2 = \text{Var}(m_{s_t}) + \mathbb{E}[\text{Var}(|G_t| | s_t)]$, so

$$\sum_{k=2}^K w_k = \frac{\text{Var}(m_{s_t})}{\sigma_{|G|}^2} \in [0, 1],$$

with equality to 1 if and only if every per-state distribution is degenerate at m_i . The residual $1 - \sum_{k \geq 2} w_k = \mathbb{E}[\text{Var}(|G_t| | s_t)] / \sigma_{|G|}^2$ is the within-state variance ratio: it contributes to $\sigma_{|G|}^2$ but not to $\text{Cov}(|G_t|, |G_{t+\tau}|)$ at any positive lag, and is the source of the discontinuity from $\rho_{|G|}(0) = 1$ to $\rho_{|G|}(1) = \sum_{k \geq 2} w_k \lambda_k$ at the first lag. The substantive consequence for moderate- K CHMM is unchanged: at every $\tau \geq 1$ the autocorrelation lies in the span of the geometric decays $\{\lambda_k^\tau\}_{k \geq 2}$, and the slow-decay envelope is governed by the largest non-unit eigenvalue.

Squared-return analogue. Replacing $|G_t|$ with G_t^2 and $\mathbf{m}$ with $\mathbf{M} = (M_1, \dots, M_K)^\top$ in Steps 1–6 above is termwise valid for $\tau \geq 1$ (the conditional independence and stationarity arguments are unaffected), giving the squared-return analogue

$$\rho_{G^2}(\tau) = \sum_{k=2}^K \tilde{w}_k \lambda_k^\tau, \quad \tilde{w}_k = \frac{(\mathbf{M}^\top \mathbf{D} \mathbf{v}_k)(\mathbf{w}_k^\top \mathbf{M})}{\sigma_{G^2}^2}.$$

The non-unit spectrum of $\mathbf{T}$ that determines $\rho_{|G|}(\tau)$ also determines $\rho_{G^2}(\tau)$; only the linear projection of the per-state moment vector onto the eigenmodes changes.

Remark (complex eigenvalues). For real $\mathbf{T}$ without self-adjoint structure, eigenvalues outside $\mathbb{R}$ occur in complex conjugate pairs with conjugate eigenvectors. A pair $(\lambda_k, \bar{\lambda}_k)$ at $\lambda_k = r e^{i\theta}$ contributes a real damped-oscillation term of the form $A r^\tau \cos(\theta\tau) + B r^\tau \sin(\theta\tau)$ to $\gamma(\tau)$, with (A, B) determined from $(\text{Re } c_k, \text{Im } c_k)$. Theorem 1 as stated remains valid; only the interpretation of the individual modes shifts from monotone geometric to damped sinusoidal. For $\mathbf{T}$ estimated on equity returns at the operating point of Section 4, the slow-decay mode at $\lambda_2 \approx \exp(-1/\tau^*)$ is real and positive, and the mono-exponential interpretation of the dominant timescale used in Section 3.8 carries through.

Remark (non-diagonalisable $\mathbf{T}$). Diagonalisability is generic: for $\mathbf{T}$ estimated by Baum–Welch in the interior of a non-degenerate parameter region, the eigenvalues are almost surely distinct, and the sample paths used in this paper realise this generic case at every K in the sweep. Should $\mathbf{T}$ admit a non-trivial Jordan block of size J , the dyadic expansion of Step 4 acquires polynomial-in- τ prefactors $\binom{\tau}{j} \lambda_k^{\tau-j}$ for $j = 0, \dots, J-1$ on the off-diagonal blocks, and $\rho_{|G|}(\tau)$ becomes a finite sum of polynomial-times-geometric terms with the same exponential rate $\max_{k \geq 2} |\lambda_k|$. The substantive content of Theorem 1, that the absolute-return ACF is determined entirely by the non-unit spectrum of $\mathbf{T}$, is unchanged.

A.6 State-Resolution and Multi-Emission Sensitivity

This appendix reports the full K -sweep underlying the headline operating point of Section 4, the multi-emission family extension confirming that $K = 18$ is stable under Student-t and Laplace emissions, the EM convergence curves at representative K , the IS and OoS visual comparison panels at non-selected $K \in \{3, 6, 12, 21\}$, the multi-emission figure panels at $K = 18$, and the Rydén $K = 2$ replication.

A.6.1 Full State-Resolution Sensitivity Table

Table 11 reports the full K -sweep referenced in Section 4.2 of the main text. Observed excess kurtosis is 7.68 IS and 5.29 OoS at all K .

Table 11. State resolution sensitivity for SPY CHMM-N (1,000 paths, $\alpha = 0.05$). Every metric is computed on the identical simulation pipeline used for Table 1 in the main text. IS coverage was 100% at every K ; OoS coverage is reported separately. ACF-MAE columns: $|G_t|$ measures the volatility-clustering reproduction; G_t measures linear-autocorrelation reproduction (parallel to $|G_t|$, both averaged over lags 1–252).

K	KS (%)		AD (%)		Kurtosis		ACF-MAE		W_1	H	Cov OoS (%)
	IS	OoS	IS	OoS	Obs	Sim	$ G_t $	G_t			
3	89.0	77.2	78.9	68.6	7.68	3.82	0.0463	0.0240	0.131	0.0839	94.9
6	92.9	78.7	84.3	74.3	7.68	5.38	0.0496	0.0238	0.116	0.0829	90.9
9	93.0	81.4	84.4	74.4	7.68	4.63	0.0492	0.0238	0.118	0.0810	91.9
12	94.9	81.6	85.8	75.0	7.68	4.57	0.0500	0.0238	0.113	0.0801	91.9
15	93.9	81.0	84.8	73.5	7.68	4.89	0.0508	0.0238	0.114	0.0804	88.9
18	94.8	85.5	88.2	79.7	7.68	5.04	0.0515	0.0237	0.110	0.0807	91.9
21	95.6	88.0	89.1	78.8	7.68	3.78	0.0534	0.0236	0.112	0.0801	99.0

A.6.2 Multi-Emission State-Resolution Sensitivity

Table 12 extends Table 11 across the four emission families, confirming that the qualitative operating point $K = 18$ is stable when the Gaussian component is replaced with Student-t, Laplace, or GED emissions. For CHMM-t the per-state degrees of freedom ν_k are refit by ECM-plus-golden-section at every K ; for CHMM-L the weighted-median location and weighted MAD scale are closed form and impose no additional cost; for CHMM-GED the three-stage CM update of Section 3.7 refits (μ_k, α_k, p_k) at every K with p_k bracketed in $[0.5, 3.0]$. Three patterns extend from Table 11 to the new families. First, the distributional pass rates plateau in the $K \in \{12, 15, 18, 21\}$ band for all four families: CHMM-t reaches its highest IS KS at $K = 18$ (94.6%), CHMM-L reaches its highest IS KS at $K = 18$ (94.2%) and its highest IS AD at $K = 18$ (91.4%), and CHMM-GED reaches its highest IS KS at $K = 12$ (97.6%) and the highest entry across the entire panel. Once a family

enters the plateau the operating point is flat with respect to K , so we report the main-text results at the same $K = 18$ used for CHMM-N. Second, the absolute-return ACF-MAE remains within the narrow 0.046–0.058 band across the four families, and the raw-return ACF-MAE remains essentially constant at ≈ 0.024 across all 28 (family, K) combinations. Both temporal axes are properties of the regime structure rather than the emission family: $|G_t|$ ACF reproduction is governed by the eigenvalue spectrum of $\mathbf{T}$ (Section 3.8), and raw- G_t ACF is null in expectation under any HMM with i.i.d. within-state emissions. Third, simulated kurtosis behaves as expected from the component families: the Gaussian sweep peaks around 5.0 at the high- K end, the Laplace sweep lands squarely in the 5.2–6.7 range (closest to the observed 7.68), the Student-t sweep climbs sharply above 13 across the entire sweep because the ECM search assigns small ν_k to a subset of tail states, and the GED sweep mirrors the Gaussian curve at 4.7–5.5 because the bulk of $\hat{p}_k$ values pin near the upper bracket and the Laplace-like tail states contribute too little stationary mass to lift the aggregate moment.

Table 12. Multi-emission state resolution sensitivity for SPY (1,000 paths, $\alpha = 0.05$). CHMM-N: Gaussian; CHMM-t: Student-t with per-state ν_k refit by ECM-plus-golden-section; CHMM-L: Laplace with weighted-median location and weighted-MAD scale; CHMM-GED: Generalized Error Distribution with per-state shape $p_k \in [0.5, 3.0]$ via three-stage ECM (Section 3.7). Every metric is computed on the identical simulation pipeline used for Table 1. Observed excess kurtosis is 7.68 IS and 5.29 OoS. ACF-MAE columns parallel the main-text Table 1: $|G_t|$ for volatility clustering, G_t (raw) for linear autocorrelation. **Sub-seed convention:** this multi-emission sweep uses a different sub-seed offset from the body-table headline run, so per-row metrics may drift by ≤ 1 percentage point from the corresponding entries in Table 1 (e.g. CHMM-GED at $K = 18$: this table 96.3% / 5.05 / 0.0546 versus body 95.2% / 5.15 / 0.0548).

Family	K	KS (%)		AD (%)		Kurt (IS sim)	ACF-MAE		W_1 IS	H IS
		IS	OoS	IS	OoS		$ G_t $	G_t		
CHMM-N	3	89.7	80.1	79.2	70.7	3.86	0.0467	0.0240	0.128	0.0833
	6	93.8	79.1	87.2	72.8	5.29	0.0504	0.0238	0.114	0.0825
	9	94.9	83.0	84.3	74.8	4.58	0.0490	0.0238	0.115	0.0806
	12	93.7	82.7	85.5	75.0	4.56	0.0501	0.0237	0.115	0.0804
	15	93.6	83.0	87.0	75.1	4.89	0.0509	0.0238	0.114	0.0803
	18	94.7	83.4	88.5	76.6	4.97	0.0511	0.0237	0.108	0.0805
	21	95.1	88.0	88.2	79.7	3.80	0.0532	0.0236	0.110	0.0796
CHMM-t	3	92.2	82.9	88.0	79.1	33.68	0.0538	0.0235	0.108	0.0759
	6	91.8	80.4	86.7	74.8	22.74	0.0527	0.0235	0.108	0.0772
	9	95.4	84.0	89.9	78.0	16.60	0.0535	0.0234	0.102	0.0752
	12	95.2	85.0	89.8	78.4	14.98	0.0543	0.0234	0.100	0.0757
	15	94.7	83.7	88.8	79.5	16.36	0.0539	0.0234	0.101	0.0758
	18	96.0	84.5	91.7	80.8	13.19	0.0551	0.0234	0.099	0.0759
	21	94.5	83.3	88.5	78.1	14.66	0.0539	0.0234	0.102	0.0758
CHMM-L	3	80.5	62.7	81.2	66.5	5.24	0.0531	0.0235	0.113	0.0855
	6	88.3	76.3	83.1	71.8	5.86	0.0510	0.0235	0.112	0.0821
	9	90.7	77.2	84.2	72.2	5.92	0.0519	0.0235	0.112	0.0823
	12	93.8	79.3	88.7	76.7	6.17	0.0541	0.0234	0.105	0.0793
	15	92.5	78.3	88.2	75.5	5.98	0.0539	0.0234	0.107	0.0794
	18	94.2	80.5	91.4	80.6	6.74	0.0566	0.0234	0.104	0.0795
	21	91.5	79.6	87.5	77.7	5.95	0.0539	0.0234	0.107	0.0807
CHMM-GED	3	90.3	78.4	85.5	73.6	5.45	0.0531	0.0236	0.109	0.0818
	6	91.8	79.0	84.9	73.5	5.22	0.0518	0.0235	0.108	0.0805
	9	94.6	81.6	89.1	75.9	5.03	0.0535	0.0234	0.101	0.0787
	12	97.6	84.2	94.0	81.2	4.67	0.0563	0.0234	0.094	0.0775
	15	94.2	82.0	87.4	76.5	5.14	0.0535	0.0235	0.104	0.0784
	18	96.3	83.8	92.7	81.3	5.05	0.0546	0.0235	0.098	0.0779
	21	94.1	85.3	88.7	81.2	4.82	0.0542	0.0235	0.102	0.0781

A.6.3 Cross-Ticker Per-Family Kurtosis Pattern

The body Table 2 reports the penalised CHMM-t cross-ticker performance ($1/\nu_k$ shrinkage at $\lambda = 20$). This appendix documents the unpenalised cross-ticker behaviour that motivates the penalty. The CHMM-t simulated kurtosis blow-ups under the unpenalised ECM (NVDA 40.8, JNJ 104.1, JPM 17.6 at $K = 18$) are specific to the Student-t emission family, and Table 13 extends the kurtosis column across all four CHMM emission variants on the same six-ticker universe to make this explicit. The pattern is consistent: unpenalised CHMM-t produces single-ticker overshoots of an order of magnitude on the heavy-tailed defensives (JNJ, NVDA), CHMM-L sits closest to observed across

every ticker (median absolute residual 1.4 kurtosis units), CHMM-N undershoots uniformly, and CHMM-GED interpolates between them. The CHMM-t blow-up mechanism is the same per-state ν_k ECM lower-bracket pinning documented for SPY in Section A.2.3: on tickers whose tail regime is concentrated in a small subset of states (NVDA’s pre-AI-rally drawdowns, JNJ’s outlier-driven defensive-stock dynamics), one or two states pin at $\nu_k \approx 2.1$ and dominate the simulated-kurtosis sample moment. Under the $1/\nu_k$ penalty at $\lambda = 20$ (the body operating point in Table 2; full rate sweep in Section 5, Appendix A.9) the simulated kurtosis on every ticker drops into the same band as CHMM-L: SPY $14.7 \rightarrow 8.0$, NVDA $40.8 \rightarrow 5.2$, JNJ $104.1 \rightarrow 12.3$, JPM $17.6 \rightarrow 7.7$, AAPL $6.3 \rightarrow 2.8$, QQQ $3.3 \rightarrow 2.9$, while IS / OoS KS pass rates and $|G_t|$ ACF-MAE move by less than 1.5 percentage points and 0.001 respectively.

Table 13. Per-ticker simulated excess kurtosis across the four CHMM emission families ($K = 18$, 1,000 paths, seed = 20260420). Columns report mean simulated IS excess kurtosis under each variant; the “Obs” column is the IS sample value. CHMM-t is the only family that produces order-of-magnitude blow-ups (NVDA, JNJ, JPM); the remaining three sit within 4 kurtosis units of observed across every ticker. Numbers are sourced from the canonical multi-emission cross-ticker run used for Table 12; minor drift from the body Table 2 CHMM-t column reflects a single re-run with a refreshed sub-seed.

Ticker	Obs	CHMM-N	CHMM-t	CHMM-L	CHMM-GED
SPY	7.7	5.0	15.0	6.8	5.1
NVDA	5.5	3.7	40.8	3.2	5.0
JNJ	9.0	5.5	104.1	6.1	9.3
JPM	7.6	7.2	17.6	6.0	14.8
AAPL	3.2	2.3	6.3	3.9	3.2
QQQ	3.4	2.0	3.3	4.1	2.7

A.6.4 Sector-Balanced 30-Ticker Panel: Full Per-Ticker Rollup

The body Table 2 reports the aggregate distribution and per-sector medians for the sector-balanced 30-ticker panel. Table 14 below gives the full per-ticker rollup. Each ticker is fit independently under the penalised CHMM-t scaffold ($K = 18$, $\lambda = 20$, 1,000 simulated paths, seed 20260420) on its own IS slice covering 2014-01-03 to 2024-01-03 and validated on the same 2024-01-04 to 2026-04-20 OoS window used for SPY in Table 1. The per-ticker ν_k medians sit at the upper bracket (50) on every ticker, indicating that the $1/\nu_k$ shrinkage is in the operative regime across the panel; minimum and maximum ν_k values are in the per-ticker CSV in `results/sector_panel/sector_panel_summary.csv`.

Table 14. Sector-balanced 30-ticker panel, per-ticker rollup (penalised CHMM-t at $K = 18$, $\lambda = 20$). Sectors ordered by GICS classification; tickers within sector ordered by the panel construction (top three large-cap representatives at IS-window-median market cap). Kurt resid is simulated minus observed IS excess kurtosis. Aggregate distribution and per-sector medians are summarised in body Table 2.

Sector	Ticker	IS KS%	OoS KS%	Kurt obs	Kurt sim	Kurt resid	$ G_t $ ACF-MAE
Information Technology	AAPL	99.1	95.1	3.19	2.84	-0.35	0.0418
Information Technology	MSFT	99.7	96.9	4.20	3.48	-0.72	0.0389
Information Technology	NVDA	99.6	55.7	5.53	6.23	0.70	0.0474
Health Care	JNJ	99.3	93.2	8.97	11.65	2.68	0.0335
Health Care	UNH	99.3	14.5	8.85	10.31	1.46	0.0382
Health Care	LLY	99.7	7.6	13.60	66.51	52.91	0.0295
Financials	JPM	99.5	50.5	7.62	8.14	0.52	0.0458
Financials	BAC	99.6	84.8	5.76	5.47	-0.29	0.0394
Financials	WFC	98.2	57.6	7.53	10.90	3.37	0.0791
Consumer Discretionary	AMZN	99.4	96.9	4.31	3.68	-0.63	0.0413
Consumer Discretionary	HD	99.2	41.4	12.92	17.03	4.11	0.0368
Consumer Discretionary	MCD	99.5	66.6	18.38	26.93	8.55	0.0394
Communication Services	NFLX	99.3	45.1	12.60	4.67	-7.93	0.0320
Communication Services	VZ	99.3	49.1	5.14	5.15	0.01	0.0270
Communication Services	DIS	99.5	96.4	10.77	13.94	3.17	0.0644
Industrials	CAT	99.7	78.9	4.34	3.48	-0.86	0.0321
Industrials	BA	97.6	62.2	20.59	52.00	31.41	0.0793
Industrials	HON	99.3	96.0	21.93	17.09	-4.84	0.0571
Consumer Staples	PG	99.7	94.3	8.15	9.51	1.36	0.0347
Consumer Staples	KO	99.5	92.7	10.57	12.84	2.27	0.0457
Consumer Staples	WMT	99.7	24.0	13.84	32.63	18.79	0.0279
Energy	XOM	96.9	70.3	5.61	10.99	5.38	0.0988
Energy	CVX	99.6	62.3	12.22	14.78	2.56	0.0498
Energy	COP	99.4	83.6	17.11	8.28	-8.83	0.0494
Utilities	NEE	98.6	24.0	10.71	9.87	-0.84	0.0455
Utilities	DUK	99.7	96.9	9.75	8.11	-1.64	0.0466
Utilities	SO	99.6	96.2	14.52	12.73	-1.79	0.0481
Materials	FCX	99.8	76.6	4.83	4.11	-0.72	0.0573
Materials	NEM	99.6	5.6	3.74	3.88	0.14	0.0376
Materials	APD	99.2	90.2	8.40	16.74	8.34	0.0381

A.6.5 Five-Named-Ticker Spot-Check (Original Six-Ticker Panel)

The original peer-review-cycle six-ticker spot-check (SPY, NVDA, JNJ, JPM, AAPL, QQQ; broad-equity ETF and five named single-name representatives) is retained in Table 15 as a hand-curated reference for downstream consumers who need to compare against the cross-asset Pipeline-B copula construction in Section 4.5 (whose six-asset universe is the same five named tickers plus QQQ). The aggregate-distribution, sector-rollup, and universe-scale failure-rate framing is moved to the body Table 2 on the 30-ticker panel.

Table 15. Original five-named-ticker spot-check (penalised CHMM-t at $K = 18$, $\lambda = 20$). Hand-curated panel covering broad-equity (SPY, QQQ), AI / semiconductors (NVDA), healthcare (JNJ), financials (JPM), and consumer-tech (AAPL); reproduced from `results/chmm_t_penalised/Cross_Ticker_CHMM_t_Pen.txt`. The expanded sector-balanced 30-ticker panel is in Table 14.

Ticker	KS (%) $\uparrow$		Kurtosis		ACF-MAE $\downarrow$
	IS	OoS	Obs	Sim	
SPY	95.8	85.0	7.7	8.01	0.0541
NVDA	99.5	53.8	5.5	5.23	0.0473
JNJ	99.8	92.3	9.0	12.28	0.0334
JPM	97.8	53.8	7.6	7.70	0.0457
AAPL	99.1	95.1	3.2	2.84	0.0418
QQQ	96.1	91.8	3.4	2.88	0.0569

A.6.6 Effective Spectral Rank of the Absolute-Return ACF

The body Section 3.8 states the algebraic upper bound: the deflated transition matrix $\mathbf{T} - \mathbf{1}\bar{\pi}^\top$ has rank at most $K - 1$, so the absolute-return ACF identity (8) contains at most $K - 1$ non-unit decay modes. The empirical question is how many of those modes carry non-trivial contribution to $\rho_{|G|}(\tau)$ at the relevant lags. Table 16 reports the per-mode contribution $w_k \lambda_k^\tau$ at $\tau \in \{1, 5, 20, 50\}$ for the fitted CHMM-N at $K = 18$ and at $K = 3$ on SPY IS, with modes ranked by lag-1 contribution magnitude $|w_k \lambda_k|$. Per-state moments $m_k = \mathbb{E}[|G_t| \mid s_t = k]$ are estimated by 200,000 draws from each emission; eigenvectors are normalised so $\mathbf{w}_k^\top \mathbf{v}_k = 1$.

Table 16. Per-mode contribution to the absolute-return ACF identity, CHMM-N at $K = 18$ (top panel; top six modes plus tail aggregate) and at $K = 3$ (bottom panel; full). Modes ordered by $|w_k \lambda_k|$ descending. “cum” is cumulative share of $\sum_k |w_k \lambda_k|$. At $K = 18$ the top mode ($|\lambda| = 0.929$) carries 93.6% of the lag-1 ACF, three modes reach 95%, and at lag 20 only the dominant mode contributes non-negligibly. At $K = 3$ the dominant mode ($|\lambda| = 0.953$) alone carries 96.8% of lag-1 ACF; the temporal axis is therefore as well-satisfied at $K = 3$ as at $K = 18$, consistent with the body finding that the binding constraint at low K is the marginal mixture, not the eigenvalue spectrum.

rank	$ \lambda_k $	$ w_k $	$ w_k \lambda_k $	$w_k \lambda_k^5$	$w_k \lambda_k^{20}$	cum
$K = 18, \rho_{ G }(1) = 0.301, \sigma_{ G }^2 = 2.555$						
1	0.929	0.324	0.301	0.224	0.0744	0.936
2	0.335	0.012	0.004	0.0001	0.0000	0.948
3	0.854	0.003	0.002	-0.001	-0.0001	0.955
4	0.231	0.008	0.002	≈ 0	≈ 0	0.961
5	0.231	0.008	0.002	≈ 0	≈ 0	0.967
6	0.203	0.008	0.002	≈ 0	≈ 0	0.971
7-17	various	various	≤ 0.001 each	≈ 0	≈ 0	1.000
$K = 3, \rho_{ G }(1) = 0.287, \sigma_{ G }^2 = 2.517$						
1	0.953	0.292	0.278	0.230	0.112	0.968
2	0.866	0.011	0.009	0.005	0.0006	1.000

The empirical effective rank confirms the body framing of Section 3.8: the ACF-MAE flatness across $K \in \{3, 6, 9, 12, 15, 18, 21\}$ documented in Appendix A.6.1, Table 11, is the macroscopic signature of the temporal axis being driven by a single dominant decay mode at every operating point in the sweep. The K -rank statement on $\mathbf{T} - \mathbf{1}\bar{\pi}^\top$ is correctly interpreted as a non-binding

upper bound at $K \geq 3$; at $K = 2$ the bound is tight (only one non-unit eigenvalue available), but the failure mode is then the marginal mixture, which two Gaussian components cannot tile against the empirical equity-return target. The bilinear identity (8) therefore plays a pedagogical role in this paper rather than serving as a direct K -selection criterion: it makes the rank constraint explicit, but the operational K choice is driven by distributional fidelity (KS, AD, kurtosis), not by the count of non-unit eigenvalues in the spectral expansion.

A.6.7 Baum-Welch Convergence

Figure 9 shows the Baum-Welch convergence curves for representative state resolutions. All models converged within the 60-iteration budget, with the log-likelihood monotonically increasing (as guaranteed by the EM algorithm). Lower K values typically converged faster (15–25 iterations), while higher K values required 25–40 iterations to reach the convergence tolerance of $|\Delta\mathcal{L}| < 10^{-4}$. The monotonic increase confirms that the quantile-based initialization provided a valid starting point from which the EM algorithm could improve without encountering degenerate solutions.






Figure 9. Baum-Welch convergence for the SPY CHMM-N fit (Pipeline A, IS) at selected state resolutions. Log-likelihood is monotonically increasing (EM guarantee) and plateaus well before `max_iter` = 60.

A.6.8 IS Comparison at Non-Selected State Resolutions


Figures 10–13 show the IS model comparison panels for $K \in \{3, 6, 12, 21\}$, complementing the $K = 18$ comparison in Figure 5 of the main text. Across all resolutions, the CHMM’s simulated density closely overlays the observed density, and the ACF of absolute returns tracks the slow decay pattern. The visible improvement from $K = 3$ to $K = 18$ is concentrated in the tails of the Q-Q panel and the steady-state level of the ACF panel. $K = 9$ and $K = 15$ panels are omitted because they interpolate smoothly between the shown resolutions; their numerical metrics are in Table 11.




(a) Marginal density.



(b) Tail Q-Q plot.




(c) ACF of raw G_t .




(d) ACF of $|G_t|$.


Figure 10. IS SPY CHMM-N comparison at $K = 3$ (Pipeline A, 1,000 simulated paths; panels as in Figure 5). The $K = 18$ counterpart is Figure 5 of the main text.




(a) Marginal density.



(b) Tail Q-Q plot.




(c) ACF of raw G_t .




(d) ACF of $|G_t|$.


Figure 11. IS SPY CHMM-N comparison at $K = 6$ (Pipeline A, 1,000 simulated paths; panels as in Figure 5).




(a) Marginal density.



(b) Tail Q-Q plot.




(c) ACF of raw G_t .




(d) ACF of $|G_t|$.


Figure 12. IS SPY CHMM-N comparison at $K = 12$ (Pipeline A, 1,000 simulated paths; panels as in Figure 5).




(a) Marginal density.



(b) Tail Q-Q plot.



(c) ACF of raw G_t .




(d) ACF of $|G_t|$.


Figure 13. IS SPY CHMM-N comparison at $K = 21$ (Pipeline A, 1,000 simulated paths; panels as in Figure 5).

A.6.9 OoS Validation at Non-Selected State Resolutions


Figures 14–17 show the OoS validation panels for $K \in \{3, 6, 12, 21\}$. KS pass rates remain above 91% at all resolutions; the $K = 18$ panel is reproduced in Figure 6 of the main text. As with the IS panels, $K = 9$ and $K = 15$ are omitted; numerical OoS KS pass rates are in Table 11.




(a) OoS density fan.



(b) KS p -value histogram.




(c) OoS ACF of raw G_t .




(d) OoS ACF of $|G_t|$.


Figure 14. OoS SPY CHMM-N validation at $K = 3$ (Pipeline A, 1,000 simulated paths; panels as in Figure 6). The $K = 18$ counterpart is Figure 6 of the main text.




(a) OoS density fan.



(b) KS p -value histogram.




(c) OoS ACF of raw G_t .




(d) OoS ACF of $|G_t|$.


Figure 15. OoS SPY CHMM-N validation at $K = 6$ (Pipeline A, 1,000 simulated paths; panels as in Figure 6).




(a) OoS density fan.



(b) KS p -value histogram.




(c) OoS ACF of raw G_t .




(d) OoS ACF of $|G_t|$.


Figure 16. OoS SPY CHMM-N validation at $K = 12$ (Pipeline A, 1,000 simulated paths; panels as in Figure 6).




(a) OoS density fan.



(b) KS p -value histogram.



(c) OoS ACF of raw G_t .



(d) OoS ACF of $|G_t|$.


Figure 17. OoS SPY CHMM-N validation at $K = 21$ (Pipeline A, 1,000 simulated paths; panels as in Figure 6).

A.6.10 Multi-Emission Figures at $K = 18$


This appendix reproduces the standard diagnostic suite (emission PDFs, residence times, IS comparison, OoS validation, and convergence) for the two heavy-tailed emission families (CHMM-t and CHMM-L) at the selected state resolution $K = 18$; transition matrices and stationary distributions are discussed in prose only below, since they are visually indistinguishable across families at $K = 18$. The Gaussian counterparts are already shown as Figures 5 and 6 in the main text and as Figure 9 above in this appendix. The figures confirm that the shared forward–backward scaffold produces internally-similar regime structure across the four families: the near-uniform interior of the transition matrix, the K -component dense emission cover, the longer central-state residence times, and the monotone log-likelihood convergence are features of the CHMM framework and not of the Gaussian component specifically.

Emission PDFs. Figure 18 compares the learned $K = 18$ emissions for the four families. The CHMM-t panel’s per-state ν_k allocation is visible in the narrower modes of certain tail states; the CHMM-L panel’s Laplace components have sharper peaks and slower exponential tail decay than the Gaussians; the CHMM-GED panel mixes both, with bulk states near the Gaussian shape ($\hat{p}_k$


near the upper bracket) and a small set of tail states near the Laplace shape ($\hat{p}_k$ near 1).




(a) CHMM-N (Gaussian)



(b) CHMM-t (Student-t)




(c) CHMM-L (Laplace)




(d) CHMM-GED (per-state p_k)

Figure 18. Learned emission distributions at $K = 18$ across the four emission families (CHMM-N Gaussian, CHMM-t Student-t with per-state ν_k , CHMM-L Laplace, CHMM-GED with per-state p_k), overlaid on the observed SPY IS return histogram (Pipeline A). Heavier-tailed emission families (t, L) give individual components more tail mass, which is what closes the kurtosis gap of the Gaussian CHMM; CHMM-GED interpolates between Gaussian and Laplace per state.


Transition matrices and stationary distributions. Figure 19 shows the $\log_{10}$ transition-matrix heatmaps at $K = 18$ across the four emission families. The four panels are visually indistinguishable: the same near-uniform interior, the same characteristic boundary states (rows 1 and K) with a small handful of low-mass entries, and self-transition probabilities of comparable magnitude. The eigenvalue spectrum of $\mathbf{T}$ that drives the absolute-return ACF identity (8) is therefore set by the K -rank regime structure rather than by the emission family. This is consistent with the ACF-MAE being stable across the family axis (Table 12). The per-family residence times $\tau_k = 1/(1 - T_{kk})$, which magnify small differences in the self-transition probabilities each family recovers, are compared in Figure 20.




(a) CHMM-N (Gaussian).



(b) CHMM-t (Student-t).



(c) CHMM-L (Laplace).



(d) CHMM-GED (per-state p_k).

Figure 19. $\log_{10}$ transition-matrix heatmaps at $K = 18$ across the four emission families on SPY IS (Pipeline A). Color scale is $\log_{10} T_{ij}$ with a 10^{-10} floor for the near-zero entries. The cross-family similarity supports the rank/spectral interpretation of Section 3.8: the regime structure (and thus the eigenvalue spectrum that drives the absolute-return ACF) is largely a property of K and the data, not of the emission family.






Figure 20. Natural residence times $\tau_k = 1/(1 - T_{kk})$ at $K = 18$ for the SPY IS fit (Pipeline A) across the four emission families (CHMM-N Gaussian, CHMM-t Student-t, CHMM-L Laplace, CHMM-GED).

Sample simulated trajectories. Figure 21 overlays one simulated SPY return path against the observed series under CHMM-N at $K = 18$, separately for the IS and OoS windows. Both panels show the simulated path tracking the volatility-clustering envelope of the observed series at the same overall amplitude. CHMM-N is shown as the headline reference; per-family marginal differences are governed by the emission family rather than by the trajectory representation, and are visualised more directly in Figure 18.




(a) In-sample window, full $T_{IS} = 2,516$ trading days.




(b) Out-of-sample window, full $T_{OoS} = 572$ trading days.

Figure 21. Sample simulated return trajectory at $K = 18$ for CHMM-N on SPY (Pipeline A). Orange is the observed series; blue is one simulated path drawn from the fit. Panel (a): full in-sample window. Panel (b): full out-of-sample window. Both panels share the same y-axis convention (annualized excess log return G_t).


IS comparison. Figures 22–24 reproduce the four-panel IS comparison (marginal density, tail Q-Q, raw-return ACF, $|G_t|$ ACF) at $K = 18$ for the three non-Gaussian emission families; the CHMM-N panel is Figure 5 in the main text.




(a) Marginal density.



(b) Tail Q-Q plot.




(c) ACF of raw G_t .




(d) ACF of $|G_t|$.


Figure 22. IS SPY model comparison at $K = 18$ for CHMM-t Student-t emissions (Pipeline A, 1,000 simulated paths). Panels show (a) marginal density with KS pass rate, (b) tail Q-Q plot, (c) ACF of raw G_t (linear autocorrelation, with the 99% Bartlett band), and (d) ACF of $|G_t|$ (volatility clustering). The CHMM-N counterpart is Figure 5; the CHMM-L and CHMM-GED counterparts are Figures 23 and 24.




(a) Marginal density.



(b) Tail Q-Q plot.




(c) ACF of raw G_t .




(d) ACF of $|G_t|$.


Figure 23. IS SPY model comparison at $K = 18$ for CHMM-L Laplace emissions (Pipeline A, 1,000 simulated paths). Panels as in Figure 22.




(a) Marginal density.



(b) Tail Q-Q plot.




(c) ACF of raw G_t .




(d) ACF of $|G_t|$.

Figure 24. IS SPY model comparison at $K = 18$ for CHMM-GED emissions with per-state shape $p_k \in [0.5, 3.0]$ (Pipeline A, 1,000 simulated paths). Panels as in Figure 22. The fitted $\hat{p}_k$ partition is reported in Figure 4 and Appendix A.2.3.


OoS validation. Figures 25–27 show the OoS validation panels for CHMM-t, CHMM-L, and CHMM-GED at $K = 18$. The CHMM-t panel demonstrates the OoS kurtosis alignment noted in the main text, the CHMM-L panel shows the tighter density fan produced by the heavier Laplace tails relative to the Gaussian, and the CHMM-GED panel shows the density fan tightening that follows from the data-chosen p_k partition.




(a) OoS density fan.



(b) KS p -value histogram.




(c) OoS ACF of raw G_t .




(d) OoS ACF of $|G_t|$.


Figure 25. OoS SPY validation at $K = 18$ for CHMM-t Student-t emissions (Pipeline A, 1,000 simulated paths). Panels show (a) OoS density fan, (b) KS p -value histogram with the $\alpha = 0.05$ threshold, (c) OoS ACF of raw G_t (linear autocorrelation, with the 99% Bartlett band), and (d) OoS ACF of $|G_t|$ (volatility clustering). The CHMM-N counterpart is Figure 6; the CHMM-L and CHMM-GED counterparts are Figures 26 and 27.




(a) OoS density fan.



(b) KS p -value histogram.




(c) OoS ACF of raw G_t .




(d) OoS ACF of $|G_t|$.


Figure 26. OoS SPY validation at $K = 18$ for CHMM-L Laplace emissions (Pipeline A, 1,000 simulated paths). Panels as in Figure 25.




(a) OoS density fan.



(b) KS p -value histogram.



(c) OoS ACF of raw G_t .



(d) OoS ACF of $|G_t|$.

Figure 27. OoS SPY validation at $K = 18$ for CHMM-GED emissions with per-state shape $p_k \in [0.5, 3.0]$ (Pipeline A, 1,000 simulated paths). Panels as in Figure 25.

Convergence. Figure 28 shows the Baum-Welch / ECM / weighted-EM log-likelihood histories at $K = 18$ for all four families. The CHMM-t, CHMM-L, and CHMM-GED curves inherit the monotone increase of the classical EM guarantee; the CHMM-t and CHMM-GED curves have slightly longer pre-plateau phases because each ECM sub-iteration refines a shape parameter (per-state ν_k for CHMM-t, per-state p_k for CHMM-GED) with a golden-section search.




Figure 28. Log-likelihood histories for the SPY IS fit (Pipeline A) at $K = 18$ across the four emission families (CHMM-N Gaussian classical EM; CHMM-t Student-t ECM with per-state ν_k golden-section sub-step; CHMM-L Laplace weighted-EM; CHMM-GED with per-state p_k three-stage ECM). All four curves are monotonically non-decreasing and plateau well before the `max_iter` = 60 budget.

A.6.11 Rydén $K = 2$ Replication

Table 17 reports the direct replication of Rydén et al.’s 1998 $K = 2$ setting on SPY IS under both initialization strategies, as discussed in Section 5. The quantile-based initialization used in the main body is row 1; five independent random-seed initializations are rows 2–6. Random init here draws per-state (μ_k, σ_k) from a moderate perturbation of the sample moments, which is representative of the random-init practice common in the 1990s HMM literature.

Table 17. Rydén replication: $K = 2$ CHMM-N on SPY IS under quantile vs random initialization (seed sub-index varies the random init).

Init	KS IS (%)	AD IS (%)	ACF-MAE	Kurt IS	KS OoS (%)
Quantile (main body)	79.0	71.5	0.0501	2.57	72.4
Random seed = 1	78.7	70.7	0.0502	2.57	73.2
Random seed = 7	75.3	69.6	0.0498	2.56	74.0
Random seed = 42	77.6	70.0	0.0498	2.59	71.8
Random seed = 100	76.2	69.4	0.0500	2.59	72.3
Random seed = 2024	76.4	69.8	0.0499	2.54	72.9

The key observation is that ACF-MAE is essentially constant at ≈ 0.050 across all six rows, matching the $K = 18$ CHMM-N of the main body. IS KS pass rate remains below the $K = 18$ operating point (75.3–79.0%) because two Gaussian components cannot tile the full multi-regime return distribution. The combination is consistent with the reading that the binding low- K constraint in the Rydén et al. [11] setup is distributional rather than temporal: the ACF is already well reproduced at $K = 2$, and it is distributional fidelity that drives the preference for $K = 18$.

A.7 Robustness, Test Power, Multi-Seed, and Auxiliary Baselines

This appendix collects the robustness diagnostics and auxiliary baseline panel referenced from Section 4: the KS test-power calibration, block-bootstrap KS recalibration, bandwidth/bin-count and multi-seed sensitivity, the auxiliary block-bootstrap baseline, the QuantGAN deep-generative baseline, the per-ticker OoS price-simulation figures, and the expanded baseline panel that adds the GARCH family and the SM-CHMM foil to the headline table.

A.7.1 OoS KS Test-Power Calibration

Table 18 reports the two-sample KS test-power calibration used to anchor the OoS KS pass rates of Section 4.3. For each of 1,000 Monte Carlo replications (seed = 20260420), a simulated series of length $T_{\text{OoS}} = 572$ is drawn from each of three reference generators and tested against the observed OoS series using the same $\alpha = 0.05$ threshold as the main-body pass-rate metric.

Table 18. OoS KS pass-rate calibration (1,000 replications, seed = 20260420).

Reference generator	Pass rate (%)
I.i.d. resamples of R_{OoS} at length $T_{\text{OoS}} = 572$ (known-correct ceiling)	100.0
I.i.d. resamples of R_{IS} at length $T_{\text{OoS}} = 572$ (nearly correct)	90.0
Gaussian($\hat{\mu}_{\text{IS}}, \hat{\sigma}_{\text{IS}}$) at length T_{IS} (negative control)	0.0

The 90.0% "nearly correct" rate is the practical ceiling for the CHMM OoS KS pass rates reported in Section 4.3 (80–84%); the 0% rate on the Gaussian misspecified generator confirms that the KS test retains ample power at IS length to reject clearly wrong generators.

A.7.2 Block-Bootstrap KS Recalibration and Mean p-value Distribution

Two complementary concerns can be raised about the headline IS KS pass-rate metric: (i) the asymptotic two-sample KS test assumes i.i.d. samples, but the simulated paths from CHMM and GARCH all carry volatility clustering, so the asymptotic null distribution may not apply; and (ii) the binary "pass / fail at $\alpha = 0.05$ " reduces a continuous p-value to a one-bit summary. Table 19 reports two recalibrations addressing each concern.

For (i), we generate the empirical null distribution of the two-sample KS statistic under stationary block bootstrap (Politis-Romano 1994) of the SPY IS series at mean block lengths $L \in \{5, 10, 20\}$, $B = 1,000$ replications. The resulting 95% block-bootstrap critical values are 0.0306, 0.0338, and 0.0350 respectively, all *smaller* than the asymptotic critical value $1.36\sqrt{2/n_{\text{IS}}} = 0.0383$; the block-bootstrap test is therefore stricter than the asymptotic one because it accounts for the within-window dependence in the IS series itself. For each generator we then compute the per-path KS statistic and report the fraction of paths whose KS statistic falls below the block-bootstrap critical value.

For (ii), we report the mean two-sample KS p-value across simulated paths plus its 5/25/50/75/95 quantiles, giving a continuous picture of how distinguishable each generator's paths are from the observed IS series.

Two patterns emerge that the asymptotic pass-rate metric of the main panel does not surface. First, the cross-generator ordering on the mean p-value scale (bootstrap 0.82, CHMM-t 0.57, CHMM-GED 0.55, CHMM-N 0.55, CHMM-L 0.50, GARCH 0.05) is much more compressed for CHMM-vs-bootstrap than the 94.8–99.6% asymp pass rates suggest: the CHMM family lands at mean p-value ~ 0.5 , the natural value for paths that are statistically indistinguishable from the observed series. Second, GARCH(1,1)'s drop from 26.8% asymp $\rightarrow$ 6.8–17.4% block-aware confirms concern (i): the asymptotic test is mildly lenient on volatility-clustering generators with the wrong marginal, and the block-aware test catches GARCH's misspecification more sharply. The CHMM family, by contrast, sees only a small block-bootstrap penalty (95% $\rightarrow$ 88–93%); CHMM-GED is the strongest entry on the block-aware metric across all three block lengths (83.2%, 90.2%, 93.2% at $L = 5, 10, 20$).

Table 19. Block-bootstrap KS recalibration and p-value distribution. All numbers under seed 20260422 with $N_{\text{paths}} = 500$. “asyp pass” is the original Section 4.3 metric (two-sample KS p-value ≥ 0.05). “mean pv” is the mean two-sample KS p-value across paths, with q_5 and q_{95} as the lower / upper p-value quantiles. “blkL%” is the fraction of paths whose KS statistic falls below the stationary-block-bootstrap 95% critical value at mean block length L . Block-bootstrap critical values: 0.0306 ($L = 5$), 0.0338 ($L = 10$), 0.0350 ($L = 20$); asymptotic critical value 0.0383.

Generator	asyp pass% $\uparrow$	mean pv $\uparrow$	pv q_{05}	pv q_{95}	blk5% $\uparrow$	blk10% $\uparrow$	blk20% $\uparrow$
i.i.d. bootstrap	99.6	0.820	0.352	0.999	97.6	99.4	99.6
GARCH(1,1)	26.8	0.048	0.000	0.227	6.8	13.6	17.4
CHMM-N	94.8	0.546	0.047	0.976	81.2	89.4	91.6
CHMM-t	95.8	0.571	0.060	0.987	82.4	89.2	91.6
CHMM-L	94.8	0.498	0.048	0.948	81.0	88.2	90.0
CHMM-GED	95.6	0.547	0.064	0.968	83.2	90.2	93.2

A.7.3 Bandwidth, Bin-Count, and Multi-Seed Robustness

Multi-seed Monte Carlo on headline metrics. The paper reports headline statistics under a single global seed. We re-run the CHMM-N / -t / -L / -GED pipeline at $K = 18$ across ten alternative global seeds (base 20260422, increment 100) with $N_{\text{paths}} = 250$ per seed and report mean $\pm$ std for the headline metrics in Table 20. Seed-to-seed variability is small: IS KS std is 1.2–1.8%; OoS KS std is 1.8–3.6%; simulated IS kurtosis std is 0.06 for CHMM-N, 0.05 for CHMM-L, 0.09 for CHMM-GED, and 1.95 for CHMM-t (reflecting the well-known instability of the per-state ν_k ECM at moderate K , already discussed in Section 5). Kupiec $\text{LR}_{\text{uc}}^{1\%}$ has std up to 1.25 for CHMM-N because the breach count moves on a coarse 0–10 grid at $T_{\text{OoS}} = 572$. The single-seed headline numbers in Table 1 sit comfortably inside the multi-seed bands. CHMM-GED has the highest mean IS KS pass rate among the four families (95.9%), consistent with its single-seed headline value.

Table 20. Multi-seed Monte Carlo on headline metrics. Ten global seeds; $N_{\text{paths}} = 250$ per seed (reduced from the headline panel’s 1,000 for tractability across 40 fits, so the seed-to-seed std reported here is an upper bound on the std the headline panel would show at $N_{\text{paths}} = 1,000$). Both ACF-MAE columns parallel Table 1: $|G_t|$ for volatility clustering, G_t for linear autocorrelation.

Model	IS KS %	OoS KS %	sim Kurt	ACF-MAE $ G_t $	ACF-MAE G_t	$\text{LR}_{\text{uc}}^{1\%}$	$\text{LR}_{\text{uc}}^{5\%}$
CHMM-N	94.7 ± 1.2	82.6 ± 1.8	5.03 ± 0.06	0.0467 ± 0.0006	0.0173 ± 0.0001	2.12 ± 1.25	3.83 ± 0.00
CHMM-t	94.6 ± 1.2	85.6 ± 1.9	13.95 ± 1.95	0.0518 ± 0.0003	0.0170 ± 0.0001	0.54 ± 0.15	3.19 ± 0.34
CHMM-L	94.4 ± 1.8	80.8 ± 3.6	6.71 ± 0.05	0.0537 ± 0.0003	0.0171 ± 0.0001	3.26 ± 0.00	2.71 ± 0.59
CHMM-GED	95.9 ± 1.2	83.2 ± 2.7	5.15 ± 0.09	0.0513 ± 0.0003	0.0171 ± 0.0001	2.93 ± 0.71	3.75 ± 0.25

A.7.4 Block-Bootstrap Baseline

Table 21 reports the full seven-metric panel for the stationary block bootstrap at block length $b = 10$ trading days, the temporally-aware non-parametric baseline added to Table 1.

Table 21. Stationary block bootstrap (block length $b = 10$ trading days) on SPY (1,000 paths, seed = 20260420).

Window	KS	AD	Kurt	ACF-MAE	W_1	H	Cov
IS	99.2	96.4	7.07	0.0568	0.092	0.0534	100.0
OoS	87.5	86.4	5.34	0.0512	0.225	0.1570	86.9

A.7.5 QuantGAN Deep-Generative Baseline

Architecture and training details for the QuantGAN deep-generative baseline used in Section 4.3 are summarised here. The construction is a repo-native approximation to the convolutional WGAN of Wiese et al. [9]: a one-dimensional convolutional generator and critic, Wasserstein loss with critic weight clipping, trained on rolling windows of standardised SPY returns and synthesising long paths by stitching generated windows.

- **Generator.** Three 1D convolutional layers ($G_{\text{ch}} = 32$, kernel size 5, ReLU, same-padding), followed by a 1×1 projection to a single output channel. Input is i.i.d. Gaussian noise of latent dimension $L = 8$ at every position of the rolling window.
- **Critic.** Three 1D convolutional layers ($D_{\text{ch}} = 32$, kernel size 5, leaky-ReLU slope 0.2, same-padding), global mean pooling along the time axis, and a single linear scalar output.
- **Loss.** Wasserstein loss with critic weight clipping at ± 0.01 [70]. The critic minimises $\mathbb{E}[D(\tilde{G})] - \mathbb{E}[D(G)]$ on real / fake mini-batches; the generator maximises $\mathbb{E}[D(\tilde{G})]$.
- **Optimiser.** Adam, learning rate 1.0×10^{-4} for both networks, batch size 64.
- **Training schedule.** 15 epochs, 120 outer steps per epoch, 4 critic updates per generator update, training on rolling windows of length $W = 64$ standardised by the IS mean and standard deviation.
- **Synthesis.** Independent draws of length W are stitched end-to-end to produce paths of length T_{IS} or T_{OoS} ; outputs are then re-scaled back to the natural return scale by the IS mean and standard deviation.
- **Implementation.** Julia with Flux.jl [71] and Optimisers.Adam state. Random seed for weight initialisation, training shuffling, and per-path sampling: 20260422, with deterministic per-path sub-seeds.

The trained model is callable through the same `simulate` interface as the CHMM, so the seven-metric panel evaluates QuantGAN on byte-identical windows to the other generators. The QuantGAN row is reported alongside the extended-GARCH and SM-CHMM rows in Table 23; the deep-generative row reaches 0% IS and OoS KS pass rates and undershoots IS kurtosis decisively (2.1 vs. observed 7.7), consistent with the documented difficulty of GAN-based generators on financial-time-series volatility-clustering benchmarks [37, 38].

A.7.6 Per-Ticker OoS Price-Simulation Figures and Full Path-Level Metrics (Unconditional CHMM)

This appendix collects (i) the full per-family path-level probabilistic-forecast metrics for an auxiliary OoS price-simulation diagnostic on the six SPY-member tickers and (ii) the per-ticker price-fan and

terminal-distribution figures for NVDA, JNJ, JPM, AAPL, and QQQ. Plotting conventions: blue shaded bands are the 5–95% and 25–75% quantile envelopes of the 500 simulated paths, blue solid line is the simulated median, red solid line is the observed OoS price track. Terminal-histogram panels show the cross-path distribution of $\tilde{P}_T$ at T_{OoS} in gray, with the observed terminal P_T^{obs} as a red vertical line and the simulated median as a blue dashed line. For brevity, the figures show CHMM-N only (the representative family; differences across the four emission families in the price-level metrics are small, see Table 22).

Table 22. Full path-level probabilistic-forecast metrics for OoS equity-price simulation, all (ticker, family) cells ($K = 18$, 500 paths, $T_{\text{OoS}} = 572$; global seed policy in Section 3.5). $\overline{C}_{0.90}$: fraction of OoS steps at which the observed price falls inside the simulated 5–95% band (calibration); $\text{MAPE}_{0.50}$: mean absolute percentage error between observed path and simulated median (sharpness); $\overline{\text{CRPS}}$: horizon-average bias-corrected sample CRPS of Gneiting and Raftery [46]; $\overline{\text{CRPS}}/S_0$: CRPS scaled by S_0 to remove price-level effects. Lower MAPE and CRPS, higher $\overline{C}_{0.90}$, are better. **Headline:** no CHMM emission family dominates on all four metrics across all six tickers; differences between CHMM-N, CHMM-t, CHMM-L, and CHMM-GED live mostly in the return tails and propagate only weakly into aggregated price-level scores. CHMM-GED is the strongest entry on JNJ (lowest CRPS, lowest MAPE) and effectively ties CHMM-t on QQQ; the other four tickers show no clear ordering.

Ticker	Family	$\overline{C}_{0.90}$	$\text{MAPE}_{0.50}$	$\overline{\text{CRPS}}$	$\overline{\text{CRPS}}/S_0$
SPY	CHMM-N	1.000	0.100	39.25	0.0835
	CHMM-t	1.000	0.112	42.66	0.0908
	CHMM-L	1.000	0.102	38.68	0.0823
	CHMM-GED	1.000	0.116	44.54	0.0948
NVDA	CHMM-N	0.698	0.339	30.62	0.6424
	CHMM-t	0.636	0.383	34.77	0.7293
	CHMM-L	0.666	0.411	36.58	0.7674
	CHMM-GED	0.682	0.375	33.26	0.6977
JNJ	CHMM-N	1.000	0.101	12.55	0.0780
	CHMM-t	1.000	0.098	12.62	0.0784
	CHMM-L	1.000	0.101	12.65	0.0786
	CHMM-GED	1.000	0.095	12.21	0.0759
JPM	CHMM-N	1.000	0.195	32.23	0.1881
	CHMM-t	1.000	0.241	40.38	0.2357
	CHMM-L	1.000	0.211	35.25	0.2057
	CHMM-GED	0.993	0.253	43.73	0.2552
AAPL	CHMM-N	1.000	0.108	21.30	0.1156
	CHMM-t	1.000	0.075	18.90	0.1025
	CHMM-L	1.000	0.077	18.82	0.1021
	CHMM-GED	1.000	0.089	19.28	0.1046
QQQ	CHMM-N	0.998	0.073	30.48	0.0763
	CHMM-t	0.995	0.079	31.63	0.0792
	CHMM-L	1.000	0.118	41.12	0.1029
	CHMM-GED	0.997	0.080	31.76	0.0795




Figure 29. NVDA OoS price fan, CHMM-N ($S_0 = 47.67 \rightarrow P_T^{\text{obs}} = 198.07$, a $4.15\times$ rally). The observed path exits the 25–75% interquartile band by mid-horizon but remains inside the 5–95% envelope; horizon-wide coverage under CHMM-N is 0.70 (CHMM-t: 0.64, CHMM-L: 0.67; Table 22), the lone calibration outlier in the panel.






Figure 30. OoS terminal-price distributions under CHMM-N for SPY and NVDA. The right-skew is characteristic of log-returns aggregated to prices; the SPY terminal sits close to the simulated median, while the NVDA terminal sits in the upper-right of the simulated distribution (the realised rally is in the upper tail of the CHMM-N terminal density).




(a) Price fan.




(b) Terminal distribution.

Figure 31. JNJ OoS price fan and terminal distribution, CHMM-N (Pipeline A, $K = 18$, 500 simulated paths, $T_{\text{OoS}} = 572$; plotting conventions as above).




(a) Price fan.




(b) Terminal distribution.

Figure 32. JPM OoS price fan and terminal distribution, CHMM-N (Pipeline A, $K = 18$, 500 simulated paths, $T_{\text{OoS}} = 572$; plotting conventions as above).




(a) Price fan.




(b) Terminal distribution.

Figure 33. AAPL OoS price fan and terminal distribution, CHMM-N (Pipeline A, $K = 18$, 500 simulated paths, $T_{\text{OoS}} = 572$; plotting conventions as above).



(a) Price fan.



(b) Terminal distribution.

Figure 34. QQQ OoS price fan and terminal distribution, CHMM-N (Pipeline A, $K = 18$, 500 simulated paths, $T_{\text{OoS}} = 572$; plotting conventions as above).

A.7.7 Expanded Baseline Panel: GARCH Family, Semi-Markov CHMM, and QuantGAN

We add five conditional-volatility baselines, the three semi-Markov CHMM variants, and the QuantGAN deep-generative baseline (Appendix A.7.5) to the single GARCH(1,1) Gaussian row of Table 1. The conditional-volatility models are fit by grid-initialised Nelder-Mead maximum likelihood on the SPY IS window ($T_{\text{IS}} = 2,516$); simulation follows the same recursion used in estimation. Model specifications for the conditional-volatility rows are below; the SM-CHMM and QuantGAN constructions are described in their own paragraphs further down.

EGARCH(1,1) Gaussian. The exponential GARCH of Nelson [32] evolves $\log \sigma_t^2$ in levels:

$$\log \sigma_t^2 = \omega + \beta \log \sigma_{t-1}^2 + \alpha \left(|z_{t-1}| - \sqrt{2/\pi} \right) + \gamma z_{t-1}, \quad z_t = (r_t - \mu)/\sigma_t, \quad z_t \sim \mathcal{N}(0, 1). \quad (29)$$

The log-variance formulation removes the GARCH positivity constraints, and the leverage term γz_{t-1} captures the asymmetric response to negative returns.

GJR-GARCH(1,1) Gaussian. The threshold GARCH of Glosten et al. [33] / Zakoian [34] augments GARCH(1,1) with an indicator on negative innovations:

$$\sigma_t^2 = \omega + \alpha \varepsilon_{t-1}^2 + \gamma \varepsilon_{t-1}^2 \mathbf{1}\{\varepsilon_{t-1} < 0\} + \beta \sigma_{t-1}^2, \quad (30)$$

with $\varepsilon_t = r_t - \mu$. Stationarity requires $\alpha + \beta + \gamma/2 < 1$; positive γ indicates leverage.

GARCH(1,1) Student-t innovations. A direct tail-aware competitor to CHMM-t: the GARCH(1,1) variance recursion is unchanged, but innovations follow a standardised Student-t with $\nu > 2$ degrees of freedom rescaled to unit variance. The log-likelihood uses the Student-t density:

$$r_t - \mu = \sigma_t \cdot z_t, \quad z_t = \eta_t \sqrt{(\nu - 2)/\nu}, \quad \eta_t \sim t_\nu. \quad (31)$$

The fitted $\hat{\nu} = 6.89$ on SPY IS sits inside the moderate-tail regime the equity-risk literature typically reports.

HAR-RV on daily squared returns. The heterogeneous-autoregressive model of Corsi [36] models realised variance as a mixture of daily, weekly, and monthly components:

$$\log \text{RV}_t = \beta_0 + \beta_d \log \text{RV}_{t-1} + \beta_w \log \overline{\text{RV}}_{t-5:t-1} + \beta_m \log \overline{\text{RV}}_{t-22:t-1} + \eta_t. \quad (32)$$

In the absence of intraday bars we use daily squared demeaned returns as the RV proxy; simulation samples $\eta_t \sim \mathcal{N}(0, \hat{\sigma}_\eta^2)$ and rolls forward the recursion, then draws $r_t - \mu = \sqrt{\text{RV}_t} z_t$ with $z_t \sim \mathcal{N}(0, 1)$.

MS-GARCH $K = 3$. A three-regime extension of the Markov-switching GARCH(1,1) ($G_t = \mu + \sigma_t z_t$ with regime-specific recursion $\sigma_t^2(k) = \omega_k + \alpha_k (G_{t-1} - \mu)^2 + \beta_k \sigma_{t-1}^2(k)$). Parameter layout is three regime triples $(\omega_k, \alpha_k, \beta_k)$ plus six transition-matrix logits (softmax-normalised per row, enforcing a non-trivial diagonal); fit by grid-initialised Nelder-Mead. States are canonicalised by ascending unconditional variance, producing the calm / medium / stress decomposition $\sigma = [0.10, 1.77, 10.04]$ on SPY IS.

MS-GARCH $K = 2$ (Haas-Mittnik-Paoletta 2004). The path-independent two-regime variant of Haas et al. [22]. Regime stickiness is high: $p_{11} = 0.915$ (calm) and $p_{22} = 0.547$ (stress), with unconditional standard deviations 0.97 (calm) and 12.67 (stress); the calm regime is near-GARCH behaviour and the stress regime is a rare, transient high-variance mode. Common $\mu = 0.094$, log-likelihood $-4,965$.

MS-GARCH $K = 4$ and $K = 6$. A natural concern with the $K = 2$ baseline is that it is unfairly low against a $K = 18$ CHMM. We refit the same Haas-Mittnik-Paoletta specification at $K = 4$ (24 parameters: $3K = 12$ GARCH triples plus $K(K - 1) = 12$ transition logits) and $K = 6$ (48 parameters), via Nelder-Mead with diagonal-dominant $\mathbf{T}$ initialisation, geometric variance-ratio spread across regimes, and ascending-variance regime canonicalisation. Higher- K MS-GARCH does not close the gap to the CHMM family on the headline distributional metric: IS KS rises only marginally from 27.7% at $K = 2$ to 37.6% at $K = 4$ and falls back to 34.5% at $K = 6$, against 94.1% for CHMM-N at $K = 18$ (a 56-percentage-point gap). The OoS KS exhibits the same plateau pattern (38.7% at $K = 2$, 38.2% at $K = 4$, 33.4% at $K = 6$). Log-likelihoods on the $T_{IS} = 2,516$ window saturate at the same value within Monte Carlo noise ($-4,965, -4,965, -4,952, -4,956$ for $K = 2, 3, 4, 6$). The MS-GARCH ceiling on this universe is therefore not a low- K issue: the unimodal-innovation constraint (each regime’s emission is a Gaussian rescaled by the regime’s conditional variance) is the binding constraint, and adding regimes does not relax it. The CHMM family decouples this constraint by giving each state its own emission-mixture component, which is the structural advantage at moderate K .

Semi-Markov CHMM (SM-CHMM, the explicit hidden semi-Markov model (HSMM) foil). The conceptual foil of the present paper, drawing on the explicit-duration HSMM construction of the companion volatility paper [72]: a hidden semi-Markov extension of the CHMM in which the geometric sojourn distribution of the Markov chain is replaced by a per-state explicit-duration family chosen by AIC from {Pareto, negative binomial, geometric}. Plug-in estimator: Viterbi state path on the IS series under the converged CHMM fit, then per-state AR(1) emissions and a sojourn family chosen by AIC. On SPY IS at $K = 18$ the AIC-selected sojourn family is Pareto for 17 of the 18 states (negative binomial for one) across all three emission families, indicating heavy-tailed sojourns that the geometric CHMM cannot capture by construction.

Caveat on the SM-CHMM estimator. The plug-in construction above (single-shot Viterbi decoding under the converged CHMM fit, then AR(1)-residual + AIC-sojourn refit) is not maximum-likelihood for the underlying HSMM: it does not iterate the joint forward-backward over (state, duration) pairs in the spirit of Yu [73]. A full ML HSMM at $K = 18$ would refit the duration-augmented log-likelihood jointly. The empirical reading we draw from the rows below is therefore strictly: *under the plug-in estimator*, the SM-CHMM weakens the headline KS / ACF / kurtosis pattern relative to the standard CHMM at the same K . We do not claim that an ML-estimated HSMM would do the same; an ML HSMM at $K = 18$ on this universe is left as a companion-paper extension. The Bulla-Bulla 2006 ML HSMM result at low K on monthly returns is the closest published anchor, but its $K \in \{2, 3\}$ regime resolution and monthly aggregation do not directly transfer.

Table 23 reports the IS / OoS distributional fidelity, kurtosis, ACF-MAE, and Kupiec breach diagnostics for the extended GARCH family, the SM-CHMM rows, and the QuantGAN deep-generative row, alongside the headline CHMM panel. Three structural observations.

First, none of the extended GARCH variants challenges the CHMM family on the joint (KS, ACF) corner. GARCH(1,1)- t is the strongest extended row on KS (57.3% IS, 80.8% OoS) but its IS excess kurtosis of 15.1 overshoots the observed 7.7 by a factor of two and its ACF-MAE of 0.0316 is

Table 23. Extended baseline panel. SPY IS / OoS, $T_{\text{IS}} = 2,516$, $T_{\text{OoS}} = 572$, $N_{\text{paths}} = 1,000$, $\alpha = 0.05$ for KS. Columns: IS / OoS KS pass rate (%); IS / OoS AD pass rate (%); simulated IS excess kurtosis; ACF-MAE on $|G_t|$ (volatility clustering) and on raw G_t (linear autocorrelation, parallel column); pooled-archive Kupiec breach rate (%) and unconditional likelihood-ratio statistic at the 1% and 5% tails. Raw-ACF-MAE values for SM-CHMM and QuantGAN are reported as “–” here pending a full re-run of those baseline scripts; the qualitative reading carries through under the methodology of the other rows (every regime-switching or i.i.d.-emission generator produces a near-zero raw ACF in expectation).

Model	KS (%) $\uparrow$		AD (%) $\uparrow$		Kurt	ACF-MAE		1% tail		5% tail	
	IS	OoS	IS	OoS		$ G_t $	G_t	br%	LR _{uc}	br%	LR _{uc}
GARCH(1,1) Gaussian	28.0	59.4	12.5	59.5	7.0	0.0309	0.0173	0.9	0.10	4.0	1.23
EGARCH(1,1)	6.3	34.9	4.1	41.7	2.8	0.0430	0.0173	1.0	0.01	3.5	3.03
GJR-GARCH(1,1)	40.1	57.7	28.7	61.4	7.6	0.0330	0.0173	1.2	0.27	4.7	0.10
GARCH(1,1)- t	57.3	80.8	36.6	65.3	15.1	0.0316	0.0173	1.0	0.01	4.9	0.01
HAR-RV	0.0	0.0	0.0	0.0	189	0.0566	0.0173	0.2	5.99	3.5	3.03
MS-GARCH $K = 2$	27.7	38.7	24.0	44.4	4.7	0.0367	0.0173	1.0	0.01	4.2	0.82
MS-GARCH $K = 3$	36.1	33.1	28.8	38.0	4.1	0.0284	0.0173	1.0	0.01	4.2	0.82
MS-GARCH $K = 4$	37.6	38.2	–	–	3.8	0.0437	–	–	–	–	–
MS-GARCH $K = 6$	34.5	33.4	–	–	4.4	0.0429	–	–	–	–	–
CHMM-N ($K = 18$)	94.1	81.8	36.6	73.0	5.0	0.0509	0.0237	0.7	1.58	4.0	3.83
CHMM-t ($K = 18$)	95.6	85.7	33.4	69.8	14.4	0.0549	0.0234	1.2	0.58	5.0	3.83
CHMM-L ($K = 18$)	93.6	80.8	30.1	65.4	6.6	0.0567	0.0234	1.4	3.26	5.6	3.03
CHMM-GED ($K = 18$)	95.2	84.3	34.1	71.0	5.2	0.0548	0.0234	0.9	0.58	4.4	3.83
SM-CHMM-N ($K = 18$)	79.2	66.7	19.5	57.1	4.7	0.0598	–	1.1	0.01	5.4	0.82
SM-CHMM-t ($K = 18$)	77.1	63.4	17.2	52.3	13.9	0.0625	–	1.4	0.10	5.7	1.74
SM-CHMM-L ($K = 18$)	80.4	69.1	21.0	58.9	6.4	0.0581	–	1.4	3.26	5.5	3.03
QuantGAN	0.0	0.0	0.0	0.0	2.1	0.0591	–	1.0	0.01	3.3	3.83

below the CHMM’s only because the GARCH variance recursion is hard-wired to track conditional volatility; CHMM at $K = 18$ matches GARCH on ACF-MAE within 0.02 while gaining 36 to 36.5 percentage points on IS KS. EGARCH undershoots kurtosis decisively, GJR-GARCH improves on threshold asymmetry but not on the headline gap, and HAR-RV is degenerate on the daily-RV proxy used here (see methods note above).

Second, the SM-CHMM rows are weaker than the headline CHMM rows on every metric in the panel: IS KS 77–80% vs CHMM 93–94%, OoS KS 63–69% vs CHMM 79–86%. *Under the Viterbi-AR(1) plug-in estimator*, the geometric-sojourn restriction of the standard CHMM is therefore not the bottleneck for the headline KS / ACF / kurtosis story; the spectral mechanism of Section 3.8 carries the comparison. We caveat strongly that this conclusion is conditional on the plug-in estimator (paragraph above): a maximum-likelihood explicit-duration HSMM under a Pareto sojourn family at the same K may close part of the gap, and we do not claim otherwise. An ML HSMM benchmark at $K = 18$ on this universe is left as a companion-paper extension.

Third, the QuantGAN deep-generative row (Appendix A.7.5) is the weakest entry in the panel on every distributional metric. IS and OoS KS pass rates collapse to 0% at $\alpha = 0.05$ across all 1,000 simulated paths, IS and OoS AD pass rates are also 0%, and simulated IS excess kurtosis of 2.1 undershoots the observed 7.7 by more than a factor of three. ACF-MAE of 0.0591 is comparable to the i.i.d. baselines of Table 1 (0.0628), so the GAN does not recover absolute-return autocorrelation either. The convolutional WGAN reproduces some surface-level visual properties of the return series but the stitched-window simulation scheme has no mechanism for modelling either the heavy-tailed marginal or the persistent absolute-return autocorrelation; this matches the documented failure

mode of GAN-based generators on financial-time-series volatility-clustering benchmarks [37, 38]. The runner is a repo-native convolutional WGAN with critic weight clipping (Appendix A.7.5), materially smaller than the original temporal convolutional network of Wiese et al. [9] (a seven-block TCN with 127-day receptive field, hidden width 80, plus Lambert-W input pre-processing that the original authors credit with most of the tail-fidelity); the row therefore reads as the panel’s deep-generative negative control rather than a verdict on the WGAN literature as a whole. The QuantGAN row complements the i.i.d. Gaussian negative control of the headline Table 1.

A.8 Cross-Asset Dependence: Full Pipeline B Panel

This appendix collects the full Pipeline B detail: cross-asset dependence model derivations and diagnostics, the Student-t copula profile log-likelihood, the cross-asset correlation heat maps, and the non-US asset-class extension with the per-pair OoS off-diagonal breakdown.

A.8.1 Cross-Asset Dependence Models: Derivations and Diagnostics (Pipeline B)

This appendix provides the technical background for the SIM and copula constructions used in Section 4.5 of the main text. These constructions constitute *Pipeline B*: per-asset marginals come from the independent CHMMs of Pipeline A (Section 4.4) and joint dependence is overlaid on top. Pipeline A stays purely univariate; Pipeline B is the sole multi-asset pipeline in the paper.

Sklar’s theorem and the rank-reordering simulator. Sklar’s theorem [40, 41] states that any d -dimensional joint CDF $H(g_1, \dots, g_d)$ with continuous marginals $F_1, \dots, F_d$ can be written as $H = C(F_1, \dots, F_d)$ for a unique copula C . The rank-reordering simulator exploits this uniqueness. If $\tilde{\mathbf{G}}$ is a matrix whose j -th column is an i.i.d. sample from F_j , and $\mathbf{U}$ is an independent sample from C , then reordering each column of $\tilde{\mathbf{G}}$ so that its ranks match $\mathbf{U}$ ’s produces a sample from H , up to discrete approximation error in the empirical ranks. Crucially, reordering never creates new values: each column of the output is a permutation of the corresponding column of $\tilde{\mathbf{G}}$, so marginal distributions are preserved exactly [44].

Kendall’s τ inversion. Given IS returns $\mathbf{R} \in \mathbb{R}^{T \times d}$, we estimate the pairwise Kendall’s τ_{ij} from concordant/discordant pair counts and invert to the correlation scale via the analytic relationship $\rho_{ij} = \sin(\pi\tau_{ij}/2)$, which holds exactly for the Gaussian copula family and is commonly used as a robust correlation estimator for the Student-t copula [43, 52]. The resulting matrix $\hat{\Sigma}$ is symmetric but not guaranteed positive semi-definite; we project to the nearest PSD matrix by clipping negative eigenvalues to 10^{-8} and rescaling the diagonal to unity.

Profile MLE for ν . The Student-t copula log-density at a single pseudo-uniform observation $\mathbf{u} \in [0, 1]^d$, with $x_j = t_\nu^{-1}(u_j)$ and $\mathbf{x} = (x_1, \dots, x_d)^\top$, is

$$\begin{aligned} \log c_t(\mathbf{u}; \Sigma, \nu) &= \log \Gamma\left(\frac{\nu+d}{2}\right) + (d-1) \log \Gamma\left(\frac{\nu}{2}\right) - d \log \Gamma\left(\frac{\nu+1}{2}\right) - \frac{1}{2} \log |\Sigma| \\ &\quad - \frac{\nu+d}{2} \log \left(1 + \frac{\mathbf{x}^\top \Sigma^{-1} \mathbf{x}}{\nu}\right) + \sum_{j=1}^d \frac{\nu+1}{2} \log \left(1 + \frac{x_j^2}{\nu}\right). \end{aligned} \quad (33)$$

Summing over $t = 1, \dots, T$ pseudo-observations yields the profile log-likelihood as a function of ν with Σ held fixed at the Kendall’s- τ estimate. We evaluate equation (33) on the discrete grid $\nu \in \{2, 3, 4, 5, 6, 8, 10, 15, 20, 30\}$ and select $\hat{\nu}$ as the grid point with the largest total log-likelihood. In Section 4.5 the selected value was $\hat{\nu} = 6$ on the six-asset universe.

Sampling the copula. To sample T rows from the d -dimensional Gaussian copula with correlation Σ , we (i) compute the Cholesky factor $\Sigma = LL^\top$, (ii) draw $Z \sim \mathcal{N}(0, I_d)$ i.i.d. and form $Y = LZ$, and (iii) return $U = \Phi(Y)$ componentwise. For the Student-t copula with degrees of freedom ν , we draw $W \sim \chi_\nu^2/\nu$ independent of Y and form the multivariate- t variate $X = Y/\sqrt{W}$, then return $U = t_\nu(X)$ componentwise.

SIM regression summary. Table 24 reports the fitted SIM regression coefficients and goodness-of-fit statistics for the non-market assets in Section 4.5. QQQ’s high $R^2 = 0.840$ reflects that the Nasdaq-100 ETF is essentially a linear combination of large-cap technology names that also drive the S&P 500 index, while JNJ’s lower $R^2 = 0.260$ reflects the limited systematic exposure of low-beta healthcare stocks. These factor loadings and residual magnitudes are what the SIM simulation re-injects at simulation time, and they explain the uneven per-asset KS pass rates reported in Table 32.

Table 24. SIM regression $\hat{G}_{j,t} = \hat{\alpha}_j + \hat{\beta}_j G_{\text{SPY},t} + \hat{\eta}_{j,t}$ for the five non-market assets, fitted on the 2014-01-03 through 2024-01-03 IS window ($T = 2,516$).

Ticker	$\hat{\alpha}$	$\hat{\beta}$	R^2
NVDA	0.321	1.694	0.371
JNJ	0.002	0.576	0.260
JPM	−0.008	1.223	0.507
AAPL	0.111	1.205	0.492
QQQ	0.047	1.122	0.840

A.8.2 Student-t Copula Profile Log-Likelihood (Pipeline B)

Figure 35 plots the profile log-likelihood of the Student-t copula used inside Pipeline B, computed on the six-asset SPY cross-section over the grid $\nu \in \{2, 3, 4, 5, 6, 8, 10, 15, 20, 30\}$ used in Section 4.5. The curve is smooth and single-peaked, with $\nu^* = 6$ attaining the maximum profile log-likelihood (6157.47), and the neighbouring grid points $\nu = 5$ (6143.37) and $\nu = 8$ (6148.37) each within 15 profile-log-likelihood units of the peak, indicating a shallow-but-clean optimum that agrees with the 4–10 range typically reported for equity-portfolio Student-t copulas in the risk-management literature. The AIC penalty for the additional ν parameter is 2 (one degree of freedom relative to the Gaussian copula limit $\nu \rightarrow \infty$), and the AIC differential between the Student-t at $\nu^* = 6$ and the Gaussian copula limit on these data is $-2 \cdot (6157.47 - 6143.37) - 2 = -30.2$, decisively in favour of the Student-t copula on IS; the same $\nu^* = 6$ is selected by AIC as by profile MLE on this grid.




Figure 35. Profile log-likelihood of the Student-t copula used inside Pipeline B on the six-asset SPY cross-section (SPY, NVDA, JNJ, JPM, AAPL, QQQ; IS window 2014-01-03 through 2024-01-03). The vertical axis is the profile log-likelihood evaluated at each degrees-of-freedom value ν on the grid $\{2, 3, 4, 5, 6, 8, 10, 15, 20, 30\}$; the dashed red line marks the profile MLE $\nu^* = 6$ (peak value 6157.47). The CHMM-N marginals are held fixed at the Pipeline A per-asset fits while ν is varied.

A.8.3 Cross-Asset Correlation Heat Maps (Pipeline B, Companion to Table 32)

Figure 36 visualises the observed and path-averaged simulated correlation matrices for the dependence models of Section 4.5. The panel ordering is SIM, Gaussian copula, Student-t copula, and a truncated C-vine variant with SPY as root, each reported quantitatively in Table 32 and (for the C-vine) Section A.8.4. All five panels share the six-asset SPY cross-section and the same CHMM-N marginals at $K = 18$, so any visible differences between the simulated panels reflect only the dependence construction.




Figure 36. Cross-asset IS correlation reproduction across dependence models (Pipeline B, companion to Table 32). Panel (a): *observed* correlation matrix on 2014-01-03 through 2024-01-03 daily excess log returns for SPY, NVDA, JNJ, JPM, AAPL, QQQ (IS window, $T_{IS} = 2,516$). Panel (b): path-averaged correlation under the Single Index Model with SPY as market factor. Panel (c): path-averaged correlation under the Gaussian copula on CHMM marginals. Panel (d): path-averaged correlation under the Student- t copula on CHMM marginals, $\nu^* = 6$. Panel (e): path-averaged correlation under a truncated C-vine copula with SPY as root. Each simulated panel averages over 200 paths of length T_{IS} . All marginals are the per-asset CHMM-N fits at $K = 18$ from Table 3; color scale in $[-1, 1]$, red-white-blue (RdBu) diverging. Closer visual match to panel (a) is better.

A.8.4 Truncated C-vine Copula on CHMM Marginals (Pipeline B)

The truncated C-vine variant in panel (e) of Figure 36 uses SPY as the root and selects each pair-copula family edge-wise from $\{\text{Gaussian, Student-}t, \text{Clayton, Gumbel}\}$ by Akaike Information Criterion (AIC); on the six-asset universe every selected pair is Student- t , with $\hat{\rho}$ and $\hat{\nu}$ as in Table 25. On the same CHMM-N marginals at $K = 18$ and 200 paths under seed 20260422, the truncated C-vine attains an off-diagonal MAE of 0.068 on IS and 0.235 on OoS (Table 32), strictly above both elliptical copulas on the IS column and on OoS off-diag MAE, and only modestly below the SIM baseline. The reason is the level-1 truncation: a root-centred C-vine models only the SPY-vs-asset bivariate copulas and treats the remaining pairs as conditionally independent given SPY, which is a stronger restriction than either elliptical copula and only a small relaxation of the SIM factor structure. Untruncated regular vines or factor copulas are the natural way to recover non-root cross-pair structure at larger d , but at $d = 6$ with $T_{IS} = 2,516$ the additional flexibility does not justify the extra estimation cost.

Table 25. Truncated C-vine edge summary, Pipeline B with SPY as root on the six-asset universe. Pair family selected edge-wise by AIC over {Gaussian, Student-t, Clayton, Gumbel}; on this universe every selected pair is Student-t. $\hat{\rho}$ is the Kendall’s- τ -inverted correlation parameter and $\hat{\nu}$ is the per-edge degrees-of-freedom estimate.

Child	Family	$\hat{\rho}$	$\hat{\nu}$
NVDA	Student-t	0.637	5
JNJ	Student-t	0.483	4
JPM	Student-t	0.686	4
AAPL	Student-t	0.698	4
QQQ	Student-t	0.911	3

A.8.5 Quarterly Rolling-Window Copula Refit on the OoS Window

The body Section 4.5 reports a static-IS-fit OoS off-diagonal MAE of 0.209, an order of magnitude larger than the IS value of 0.027. Section 5 attributes this to the stationarity-scope limit and recommends periodic refit. Reviewer Question Q4 asked for a concrete rolling-window-refit number on the same six-asset universe.

We refit the Pipeline-B Student-t copula on a 252-day rolling window (one trading year) sliding by 63 days (one trading quarter) across the OoS span, holding the per-asset CHMM-N marginals fixed at the IS fits used for Table 3. Each refit yields a Kendall’s- τ correlation matrix and a profile-MLE-selected ν^* on the discrete grid $\{2, 3, 4, 5, 6, 8, 10, 15, 20, 30\}$; we then simulate forward 63 days under the new copula and compare the path-averaged simulated correlation matrix to the realised next-quarter correlation. Table 26 reports the per-quarter detail.

Table 26. Quarterly rolling-window refit of the Pipeline-B Student-t copula on the OoS window (six-asset universe; CHMM-N marginals at $K = 18$ held fixed at IS fits; 200 paths per refit; seed = 20260422). Per-quarter window slice and refitted ν^* , plus the off-diagonal MAE between the path-averaged simulated correlation matrix under the refitted copula and the realised next-quarter correlation matrix. Headline: rolling-quarterly refit reduces mean OoS off-diagonal MAE from 0.207 (static-IS-fit baseline) to 0.185, a $\sim 10\%$ relative improvement; the gain concentrates in the later OoS quarters where the rolling window incorporates more recent OoS data and the refitted ν^* shifts from the IS-fit value of 6 down to 5 or up to 10–15 as the empirical tail-coupling rolls.

Quarter	Window slice	Forecast horizon	Refit ν^*	Off-diag MAE
1	[2265, 2516]	[2517, 2579]	15	0.198
2	[2328, 2579]	[2580, 2642]	15	0.205
3	[2391, 2642]	[2643, 2705]	10	0.241
4	[2454, 2705]	[2706, 2768]	6	0.133
5	[2517, 2768]	[2769, 2831]	6	0.255
6	[2580, 2831]	[2832, 2894]	6	0.232
7	[2643, 2894]	[2895, 2957]	5	0.147
8	[2706, 2957]	[2958, 3020]	5	0.119
9	[2769, 3020]	[3021, 3083]	6	0.138
<i>mean</i>	—	—	—	0.185
<i>median</i>	—	—	—	0.198
<i>Static IS-fit</i>	[1, 2516]	[2517, 3083]	6	0.207

The quarterly refit closes part of the OoS gap but not most of it: mean off-diagonal MAE moves from 0.207 static to 0.185 rolling ($\sim 10\%$ relative improvement). The improvement is

concentrated in the later OoS quarters (Q4, Q7, Q8, Q9 all below 0.15) where the rolling window incorporates more recent OoS data, while early-OoS quarters (Q1, Q3, Q5, Q6) remain at the static-IS level because the rolling window is still dominated by IS observations. The refitted ν^* sequence (15, 15, 10, 6, 6, 6, 5, 5, 6) tracks the gradual shift from the equity-cluster IS regime toward the OoS regime, supporting the operational recommendation in Section 5 of quarterly refit at deployment. The residual gap (mean MAE ≈ 0.185 versus IS 0.027) is consistent with the per-pair JNJ-equity-factor regime shift documented in Table 27: even a perfectly-refitted copula on the historical 252 days cannot anticipate the regime shift at the next quarter’s start, only track it once it has unfolded.

A.8.6 Non-US Asset Class Extension and Per-Pair OoS Off-Diagonal Breakdown

The body universe (SPY, NVDA, JNJ, JPM, AAPL, QQQ) is six US-listed equity tickers from one country-index family. To probe whether the cross-asset construction generalises off the equity cluster we extend the universe with GLD (SPDR Gold Trust), a US-listed ETF whose underlying is physically-backed gold, i.e. a non-equity commodity asset class.

GLD Pipeline-A univariate fidelity. Fitting CHMM-N, CHMM-t, and CHMM-L on the GLD return series at $K = 18$ produces 100% IS KS pass rate for all three families and observed IS excess kurtosis 3.6 matched closely by CHMM-L (2.2); on OoS, however, the KS pass rate collapses to 0% for all three families against an observed OoS excess kurtosis of 6.5. Inspection of the GLD price path on the 2024–2026 OoS window shows the asset re-priced from ~ 200 to ~ 300 on a persistent demand shock not represented in the 2014–2024 IS regime library. This is the same stationarity-scope artefact as the NVDA / JPM single-name OoS cliffs reported in Section 4.4, on a non-equity asset; the limitation is not equity-specific.

Pipeline-B: 7-ticker Student-t copula off-diagonal MAE. Refitting the Student-t copula on the augmented seven-ticker universe at $\nu^* = 6$ gives IS off-diagonal MAE of 0.029 (six-ticker baseline: 0.027) and OoS off-diagonal MAE of 0.179 (six-ticker baseline: 0.209). Adding the low-equity-correlation gold ETF *reduces* the OoS off-diagonal MAE, because the equity-cluster correlation degradation is the dominant component; pairs involving GLD have moderate IS-to-OoS shifts ($|\Delta| \in [0.04, 0.19]$) while the largest equity-cluster pairs (JNJ-QQQ, SPY-JNJ, NVDA-JNJ) have $|\Delta| > 0.48$ on OoS.

Per-pair OoS gap localisation. Table 27 reports the per-pair off-diagonal absolute deviation between observed and path-averaged simulated correlation matrices on IS and OoS, sorted by OoS $|\Delta|$. The OoS gap concentrates in pairs involving JNJ (the defensive single-name proxy): JNJ’s IS correlation with the broad equity factor (QQQ, SPY) collapses on OoS, in some pairs flipping sign, and the IS-fixed copula has no mechanism for tracking that shift. The gold-ETF pairs (SPY-GLD, JPM-GLD, JNJ-GLD, QQQ-GLD) sit in the middle of the ranking with $|\Delta|$ between 0.10 and 0.19 on OoS. The dominant cause of the OoS off-diagonal gap is therefore neither equity-cluster size nor copula-degree mis-specification but a single-name regime shift in the JNJ-equity-factor relationship that the IS-fitted dependence layer cannot anticipate.

Table 27. Per-pair off-diagonal correlation MAE on the seven-ticker universe (Pipeline B, Student-t copula on CHMM-N marginals at $K = 18$, 200 paths, $\nu^* = 6$). For each ticker pair (i, j) the table reports $|\Sigma_{\text{sim,avg}}[i, j] - \Sigma_{\text{obs}}[i, j]|$ on IS and OoS, ranked by OoS $|\Delta|$. Only the ten largest-OoS-gap pairs are shown for compactness.

Pair	IS $ \Delta $	OoS $ \Delta $	OoS – IS
JNJ-QQQ	0.027	0.544	0.517
SPY-JNJ	0.037	0.509	0.473
NVDA-JNJ	0.017	0.489	0.473
NVDA-AAPL	0.011	0.278	0.268
JNJ-JPM	0.046	0.261	0.216
JNJ-AAPL	0.042	0.244	0.202
AAPL-QQQ	0.003	0.227	0.224
JPM-GLD	0.027	0.187	0.160
SPY-GLD	0.038	0.127	0.089
JNJ-GLD	0.015	0.117	0.101

A.9 Pre-OoS Validation K -Selection and ν_k Diagnostics

Pre-OoS validation K -selection. The $K = 18$ operating point in the main panel was chosen via a multi-objective criterion combining information-criterion (IC) rank with IS and OoS distributional pass rates, which uses the same 2024–2026 window later used for VaR. As a clean re-selection that decouples the two, we fit CHMM-N on 2014-01-03 through 2021-12-31 ($n = 2,013$) and score every K in the sweep by the forward-algorithm held-out log-likelihood on 2022-01-03 through 2024-01-03 ($n = 502$); the 2024–2026 window is not touched. Held-out log-lik, BIC, HQC, and CAIC all select $K^* = 3$ (per-observation log-lik -2.5345 at $K = 3$ vs. -2.5694 at $K = 18$); AIC selects $K^* = 6$; held-out two-sample KS selects $K^* = 9$. None of the six criteria that avoid the OoS window select $K = 18$. The $K = 18$ choice is best read as a multi-metric one that weights tail fidelity (kurtosis, AD) and downstream VaR behaviour alongside likelihood; a parallel main panel at $K = 3$ would re-rank some rows in CHMM’s favour on ACF-MAE and against it on tail fidelity. The 2022–2023 validation slice is itself partly a structural-break period (the rate-hike cycle), so any model with substantial IS-specific structure generalises worse on it.

Pre-2020 held-out K -selection (no rate-hike confound). Reviewer R1’s Question 1 asked whether a strictly pre-2020 validation slice (avoiding the rate-hike confound that contaminates the 2022-2023 slice above) selects the same K^* . We re-ran the held-out procedure on a 2014-01-03 through 2018-06-29 estimation window ($n_{\text{est}} = 1,130$) with a 2018-07-02 through 2019-12-31 validation slice ($n_{\text{val}} = 377$), both fully pre-COVID and pre-2022-rate-hike (the Q4 2018 drawdown and 2019 recovery sit inside the validation window as a moderate-stress non-regime-shift event). Per-observation held-out log-likelihood: -2.0681 at $K = 3$, -2.0606 at $K = 6$, -2.0763 at $K = 9$, -2.1287 at $K = 12$, -2.1921 at $K = 18$, -2.2609 at $K = 21$. **The held-out log-likelihood selects $K^* = 6$, and the held-out two-sample KS pass rate also selects $K^* = 6$ (96.8% at $K = 6$ versus 84.6% at $K = 3$).** AIC, BIC, HQC, and CAIC continue to select $K^* = 3$ on this slice, reflecting the parameter-count penalty on a 4.5-year estimation window where the per-state sample is small. The qualitative reading is robust to the slice choice: held-out distributional fidelity rises from $K = 3$ to a peak in the $K = 6$ to $K = 9$ band and then degrades, with neither slice selecting $K = 18$ by any held-out criterion, but the pre-2020 result favours a moderately larger K (6 vs 3) than the rate-hike-contaminated 2022-2023 slice.

Penalised-ECM rate sweep. The bracket sensitivity is reported with its table in Section A.2.3; here we report the complementary $1/\nu_k$ shrinkage rate sweep referenced in Section 5. An exponential $1/\nu_k$ prior at rate $\lambda \in \{0, 5, 20, 50, 100, 200\}$ reduces simulated excess kurtosis monotonically: 14.30, 11.19, 8.43, 5.41, 3.96, 3.67. The recommended rate $\lambda = 20$ brings simulated kurtosis to ≈ 8 (close to the observed 7.69) at a 1pp IS KS cost. Only the $\nu_{\min}$ state parameter actually moves ($2.1 \rightarrow 4.89$); upper bracket and median remain at 50. The overshoot is therefore a single-state artefact of the two tail regimes hitting the lower bracket under unpenalised ECM.

Walk-forward / rolling-origin OoS for CHMM-N. We define six rolling-origin folds, each train 5 years and test 1 year, refit CHMM-N at $K \in \{3, 18\}$ from scratch on each fold’s train slice, and report KS pass rate, simulated kurtosis, and $|G_t|$ ACF-MAE on the fold’s test slice ($N_{\text{paths}} = 500$). Across the six folds (Table 28), CHMM-N at $K = 18$ attains median (IQR) KS 67.7% [8.2, 75.0] and $|G_t|$ ACF-MAE 0.0542 [0.0508, 0.0571]; CHMM-N at $K = 3$ attains 62.1% [7.2, 78.4] and 0.0563 [0.0552, 0.0592]. Two folds carry sharply lower KS pass rates: W2 (COVID test slice, 7–8%) and W4 (rate-hike test slice, 0–1%). The four non-stress folds attain KS 61–83%, consistent with the headline OoS pass rate. The headline ranking (CHMM-N at $K = 18 \geq$ CHMM-N at $K^* = 3$ on KS, both within 0.005 of each other on $|G_t|$ ACF-MAE) is window-robust on the four non-stress folds; on stress folds (COVID, rate hike) all CHMM rows drop sharply, reflecting the stationarity-scope limit rather than a model-selection issue. The body framing ("CHMM is the joint-fit row in the panel under stationary OoS conditions; periodic refit is recommended for production deployment") is unchanged under the walk-forward stress test.

Table 28. Walk-forward / rolling-origin OoS for CHMM-N. Six folds, each train 5 years and test 1 year; $N_{\text{paths}} = 500$ per fold. KS pass rate at $\alpha = 0.05$. W2 covers the COVID drawdown; W4 covers the start of the 2022 rate-hike cycle. The bracketed pairs in the bottom row report $[Q_1, Q_3]$ across the six folds (the empirical first and third quartiles, not a scalar IQR width).

Fold (test window)	KS (%)		Kurt sim		ACF-MAE $ G_t $	
	$K = 3$	$K = 18$	$K = 3$	$K = 18$	$K = 3$	$K = 18$
W1 (2019)	63.2	82.6	2.51	2.66	0.0572	0.0525
W2 (2020 COVID)	7.2	8.2	2.42	2.50	0.0552	0.0584
W3 (2021)	82.4	74.0	5.73	6.76	0.0602	0.0571
W4 (2022 rate hike)	0.8	0.0	4.72	4.35	0.0592	0.0558
W5 (2023)	78.4	75.0	3.47	2.35	0.0511	0.0508
W6 (2024)	61.0	61.4	2.81	2.86	0.0555	0.0493
median	62.1	67.7	3.14	2.76	0.0563	0.0542
IQR	[7.2, 78.4]	[8.2, 75.0]	[2.51, 4.72]	[2.50, 4.35]	[0.0552, 0.0592]	[0.0508, 0.0571]

Four-family conditional VaR back-test. Extending the regime-conditional VaR construction of §4.6 to all four emission families (CHMM-N, CHMM-t at $\lambda = 20$, CHMM-L, CHMM-GED) at $K \in \{3, 18\}$ and $\alpha \in \{0.01, 0.05\}$ yields sixteen (K, α, family) rows on the OoS window (Table 29). Every row passes Kupiec, Christoffersen-ind, and Christoffersen-cc cleanly: $p_{\text{cc}} \geq 0.089$ throughout, with all LR_{cc} statistics below the $\chi^2_2(0.05) = 5.991$ critical value. The regime-switching value proposition is therefore intrinsic to the latent-state forecast and not specific to the Gaussian emission family.

Table 29. Regime-conditional VaR back-test across four CHMM emission families on SPY OoS ($T_{\text{OoS}} = 572$, seed 20260420). Critical values: $\chi_1^2(0.05) = 3.841$, $\chi_2^2(0.05) = 5.991$. Every $(K, \alpha, \text{family})$ row passes Kupiec, Christoffersen-ind, and Christoffersen-cc at $\alpha = 0.05$.

Family	K	α	breaches	br rate	med VaR	LR _{uc}	LR _{ind}	LR _{cc}
CHMM-N	3	0.01	9	1.57%	-4.56	1.62	2.36	3.98
CHMM-N	3	0.05	35	6.12%	-2.87	1.41	0.01	1.42
CHMM-N	18	0.01	9	1.57%	-5.20	1.62	2.36	3.98
CHMM-N	18	0.05	26	4.55%	-3.02	0.26	0.52	0.78
CHMM-t ($\lambda = 20$)	3	0.01	8	1.40%	-5.59	0.82	2.81	3.63
CHMM-t ($\lambda = 20$)	3	0.05	32	5.59%	-2.73	0.41	0.77	1.19
CHMM-t ($\lambda = 20$)	18	0.01	10	1.75%	-6.25	2.65	1.98	4.62
CHMM-t ($\lambda = 20$)	18	0.05	29	5.07%	-3.15	0.01	1.39	1.40
CHMM-L	3	0.01	5	0.87%	-5.45	0.10	4.74	4.84
CHMM-L	3	0.05	26	4.55%	-2.63	0.26	2.23	2.48
CHMM-L	18	0.01	9	1.57%	-6.18	1.62	0.29	1.91
CHMM-L	18	0.05	30	5.24%	-2.80	0.07	1.16	1.23
CHMM-GED	3	0.01	7	1.22%	-5.73	0.27	3.34	3.61
CHMM-GED	3	0.05	31	5.42%	-2.54	0.21	0.96	1.16
CHMM-GED	18	0.01	9	1.57%	-6.55	1.62	2.36	3.98
CHMM-GED	18	0.05	25	4.37%	-3.10	0.50	2.56	3.06

Walk-forward conditional-VaR back-test. Extending the regime-conditional VaR construction to the six rolling-origin folds of Table 28, with CHMM-N refit per fold from scratch on the train slice and forward-filtered through the test slice under fold-IS-fixed parameters, gives twenty-four (fold, K, α) rows in Table 30. The conditional Christoffersen-cc passes at $\alpha = 0.05$ on ten of twelve fold- K combinations (W1, W3, W4, W5, W6 at both $K \in \{3, 18\}$); the two failures concentrate on W2 (COVID, $p_{\text{cc}} \in \{0.011, 0.002\}$ at $K \in \{3, 18\}$). At $\alpha = 0.01$ the construction passes on nine of twelve combinations: W2 fails at both K ($p_{\text{cc}} < 10^{-3}$) and W4 at $K = 18$ fails at $p_{\text{cc}} = 0.022$ on a LR_{ind} tail-event clustering rather than a LR_{uc} coverage miss (LR_{ind} = 7.49 at LR_{uc} = 0.11). The pass-rate is therefore 19/24 aggregate at the 5% level and 22/24 if we exclude the two stress folds that the univariate walk-forward already flagged as out-of-distribution by KS in Table 28. The reading is that the conditional construction is window-conditional: under regime shifts of the W2 / W4 magnitude (a regime introduction the IS distribution does not span; Section 5, stationarity-scope paragraph), the conditional VaR inherits the same out-of-scope behaviour as the unconditional generators. On the four non-stress folds (W1, W3, W5, W6) plus W4 at $K = 3$ at the 5% level the conditional construction is window-robust, supporting the body framing of §4.6 extending beyond the headline single-window result.

Per-ticker $\hat{\lambda}^*$ sweep. A per-ticker sweep $\lambda \in \{0, 5, 10, 20, 50, 100\}$ on each of the six body tickers under the constraint that IS KS at λ exceeds the $\lambda = 0$ baseline by no more than 1.5 percentage points (Table 31) shows that the body uniform $\lambda = 20$ is the right rate on the heavy-tailed defensives (NVDA, JNJ) but is over-shrunk on the moderate-tail tickers. The per-ticker optima are $\lambda_{\text{SPY}}^* = 10$, $\lambda_{\text{NVDA}}^* = 20$, $\lambda_{\text{JNJ}}^* = 20$, $\lambda_{\text{JPM}}^* = 10$, $\lambda_{\text{AAPL}}^* = 10$, $\lambda_{\text{QQQ}}^* = 0$. The honest operational reading is that $\lambda = 20$ is a reasonable uniform default for production deployment but per-ticker tuning is recommended whenever a ticker’s residual kurtosis at $\lambda = 0$ is within ~ 1 unit of observed.

Cross-asset summary. The full per-construction Pipeline B panel is in Section A.8; for reference, the headline ordering is summarised in Table 32.

Table 30. Walk-forward regime-conditional VaR back-test for CHMM-N on six rolling-origin folds (train 5y / test 1y; seed 20260420). At each test day t , conditional $\widehat{\text{VaR}}_t(\alpha)$ is the α -quantile of the K -component Gaussian mixture predictive density under fold-IS-fixed parameters, evaluated on the full-timeline filter $\Pr(s_{t+1} | \mathcal{F}_t = \text{train} \cup \text{test}[1:t-1])$. Critical values: $\chi_1^2(0.05) = 3.841$, $\chi_2^2(0.05) = 5.991$. Bold p_{cc} rows fail at the 5% level.

Fold	K	α	breaches	br rate	med VaR	LR _{uc}	LR _{ind}	LR _{cc}	p_{cc}
W1	3	0.01	2	0.80%	−4.01	0.11	0.03	0.15	0.93
W1	3	0.05	15	5.98%	−2.43	0.48	1.18	1.65	0.44
W1	18	0.01	0	0.00%	−5.03	5.05	0.00	5.05	0.08
W1	18	0.05	12	4.78%	−2.66	0.03	0.38	0.40	0.82
W2	3	0.01	16	6.35%	−6.54	32.93	0.87	33.80	0.00
W2	3	0.05	23	9.13%	−4.59	7.34	1.71	9.05	0.01
W2	18	0.01	14	5.56%	−6.67	25.59	1.56	27.15	0.00
W2	18	0.05	25	9.92%	−4.54	10.11	2.56	12.68	0.00
W3	3	0.01	6	2.39%	−4.48	3.53	0.30	3.82	0.15
W3	3	0.05	20	7.97%	−2.47	3.98	0.30	4.28	0.12
W3	18	0.01	2	0.80%	−5.57	0.11	0.03	0.15	0.93
W3	18	0.05	15	5.98%	−3.18	0.48	1.18	1.65	0.44
W4	3	0.01	5	2.00%	−7.98	1.96	0.21	2.16	0.34
W4	3	0.05	18	7.20%	−4.16	2.26	1.99	4.24	0.12
W4	18	0.01	2	0.80%	−9.62	0.11	7.49	7.60	0.02
W4	18	0.05	10	4.00%	−6.63	0.56	3.80	4.36	0.11
W5	3	0.01	0	0.00%	−6.06	5.01	0.00	5.01	0.08
W5	3	0.05	8	3.21%	−3.90	1.91	0.53	2.44	0.30
W5	18	0.01	1	0.40%	−7.10	1.16	0.01	1.17	0.56
W5	18	0.05	7	2.81%	−3.85	2.96	0.41	3.37	0.19
W6	3	0.01	2	0.80%	−4.68	0.11	0.03	0.15	0.93
W6	3	0.05	13	5.18%	−2.85	0.02	0.15	0.17	0.92
W6	18	0.01	1	0.40%	−5.83	1.19	0.01	1.20	0.55
W6	18	0.05	7	2.79%	−2.98	3.06	0.40	3.46	0.18

Table 31. Per-ticker $\hat{\lambda}^*$ for the penalised CHMM-t at $K = 18$ ($N_{\text{paths}} = 500$, seed 20260420, KS-degradation tolerance $\leq 1.5\text{pp}$ from $\lambda = 0$). $\hat{\lambda}^*$ is the value of λ in $\{0, 5, 10, 20, 50, 100\}$ that minimises $|\text{kurt}_{\text{sim}} - \text{kurt}_{\text{obs}}|$ subject to the KS-degradation constraint.

Ticker	kurt_{obs}	$\hat{\lambda}^*$	$\text{kurt}_{\text{sim}}(\lambda^*)$	IS KS(λ^*)
SPY	7.69	10	11.82	96.4%
NVDA	5.54	20	5.68	99.4%
JNJ	8.98	20	10.74	99.8%
JPM	7.63	10	8.62	98.6%
AAPL	3.20	10	3.24	99.2%
QQQ	3.40	0	3.35	95.2%

Table 32. Cross-asset dependence summary (Pipeline B; CHMM-N marginals at $K = 18$; 200 paths; seed 20260422). Median per-asset KS pass rate (%) and off-diagonal MAE of the simulated correlation matrix vs. observed. Bold marks the column winner. The Student-t copula at $\nu^* = 6$ is the strongest dependence layer on the two IS columns; on OoS the Gaussian copula attains marginally lower off-diagonal MAE (0.202 vs. 0.209) while the truncated C-vine sits between the elliptical copulas and the SIM baseline. Full per-asset detail is in Section A.8; the C-vine edge summary is in Table 25.

	Median IS KS	Median OoS KS	Off-diag MAE IS	Off-diag MAE OoS
Single Index Model	77.0	87.0	0.076	0.252
Gaussian copula	97.8	87.5	0.030	0.202
Student-t copula ($\nu^* = 6$)	98.8	85.8	0.027	0.209
Truncated C-vine (root SPY)	97.2	86.0	0.068	0.235