

Mechanistic Resource Competition Modeling of a Cell-Free Gluconate Biosensor

Abhinav Adhikari, Abhishek Murti, Anirudh M. Narayanan, Ha Eun Lim,
and Jeffrey D. Varner *

Robert Frederick Smith School of Chemical and Biomolecular Engineering
College of Engineering, Cornell University, Ithaca, NY 14853

Running Title: Mechanistic Resource Model for Cell-Free Biosensors

*Corresponding author:

Jeffrey D. Varner,

Professor, School of Chemical and Biomolecular Engineering,

244 Olin Hall, Cornell University, Ithaca NY, 14853

Email: jdv27@cornell.edu

Phone: (607) 255 - 4258

Fax: (607) 255 - 9166

Abstract

Cell-free synthetic biology systems offer a promising platform for rapid prototyping of biosensor circuits because they eliminate the confounding effects of cellular growth and membrane transport. However, the finite transcriptional and translational resources in cell-free reactions impose time-dependent capacity constraints that are difficult to capture with empirical decay functions. In this study, we developed and validated a mechanistic resource competition model for a D-gluconate biosensor circuit operating in the reconstituted cell-free system PURExpress. The circuit employed the transcription factor GntR to repress expression of the fluorescent reporter Venus, with D-gluconate relieving repression in a dose-dependent manner. We formulated a two-layer resource model in which the first layer described the allocation of RNA polymerase and ribosome between competing genes using Michaelis-Menten-like expressions, and the second layer captured the irreversible depletion of consumable resources (nucleotides, energy cofactors, and amino acids) through ordinary differential equations driven by cumulative transcription and translation flux. This mechanistic formulation replaced the ad hoc logistic and exponential decay functions used previously and reduced the number of estimated parameters from 28 to 25 while providing a physical interpretation of resource exhaustion. Model parameters were estimated using a multi-objective and single-objective cycling strategy implemented in ParetoEnsembles.jl, yielding an ensemble of 5464 parameter sets that captured the transcription and translation kinetics of the circuit. The model predicted that 81% of transcriptional resources and 26% of translational resources were consumed by 12 hours, consistent with the observed saturation of mRNA and protein trajectories. Without re-estimation of parameters, the model accurately predicted the dose-response relationship of Venus protein as a function of D-gluconate concentration over four orders of magnitude, yielding a dissociation constant of $K_{\text{gluconate}} = 0.25 \pm 0.20$ mM and a Hill coefficient of $n_{\text{gluconate}} = 1.92 \pm 0.86$. Taken together, this study demonstrated that a mechanistic

41 resource competition framework improved the predictive capability and interpretability of
42 cell-free biosensor models and provided quantitative design rules for the allocation of
43 transcriptional and translational capacity in synthetic gene circuits.

44 **Keywords:** cell-free biosensors, mathematical modeling, synthetic biology, resource com-
45 petition, parameter estimation, transcription-translation dynamics

46 Introduction

47 Synthetic biology seeks to program biological systems to perform user-specified func-
48 tions. Traditionally, synthetic biology involved engineering the functionality of living cells
49 using genetic tools; however, recent applications have increasingly focused on cell-free
50 platforms, which offer several advantages over their in vivo counterparts [1–7]. The ab-
51 sence of cellular growth and maintenance requirements, together with the removal of the
52 cell wall barrier, facilitates the interrogation and manipulation of reaction conditions and
53 directs the allocation of carbon and energy resources toward a product of interest. Fur-
54 thermore, cell-free platforms enable the use of linear DNA fragments instead of plasmids,
55 which significantly reduces the time required for design-build-test cycles. Fundamentally,
56 a cell-free system includes only the components required for transcription and translation
57 (TXTL) processes, such as RNA polymerase (RNAP), ribosomes, transcription and trans-
58 lation elongation factors, energy compounds, salts, amino acids, and ribonucleotides.
59 There are two classes of cell-free systems: reconstituted and extract-based. Extract-
60 based systems are prepared by lysing whole cells and separating the unwanted debris,
61 while reconstituted systems are well-defined and prepared using only the factors essential
62 for protein synthesis, including purified enzymes, tRNAs, ribosomes, elongation factors,
63 amino acids, and energy molecules. The well-defined composition of reconstituted sys-
64 tems makes them an attractive platform for studying biomolecular interactions and for
65 quantitative modeling without the confounding effects of unknown components present in
66 extract-based systems [8–10].

67 One promising application area for cell-free systems is biosensing. Traditionally, biosen-
68 sors based on live cells have faced a significant challenge: transport limitations caused
69 by the cell membrane. Cell-free biosensors overcome this barrier by removing the con-
70 straint of the cell membrane and allowing direct access to the transcription and translation
71 machinery, which leads to increased sensitivity and faster response times. Transcription-

factor-based biosensors, which utilize a transcription factor that is allosterically modified upon binding to a small molecule to control the expression of a reporter gene, are predominantly used to detect small molecules and ions. Such sensing systems have recently been used to detect water contaminants, biomarkers, and antibiotics [11–18]. Because cell-free systems can be freeze-dried and later reconstituted with water, they can be deployed for on-demand, point-of-care applications [14]. Moreover, the detection range of existing transcription factors can be expanded through metabolic engineering; for example, Voyvodic and coworkers used the transcription factor BenR to detect benzoic acid and then employed metabolic enzymes to convert hippuric acid and cocaine into benzoic acid, thereby broadening the spectrum of detectable molecules [19].

Despite the experimental progress in cell-free biosensors, quantitative mathematical models that capture the full dynamics of these systems remain underdeveloped. A central challenge is the finite and non-renewable nature of transcriptional and translational resources in cell-free reactions. As nucleotides, amino acids, and energy cofactors are consumed, the rates of mRNA and protein production decline in ways that are difficult to predict from first principles. Several groups have developed resource-aware models for gene expression systems. Gyorgy and Murray developed a framework for resource competition in synthetic genetic circuits where shared cellular machinery couples otherwise independent modules [20]. Weisse et al. proposed a whole-cell model in which host and circuit compete for a shared ribosome pool [21]. In the cell-free context, Stogbauer et al. characterized the time-dependent loss of translational activity in extract-based systems due to ribosome inactivation [22], Karzbrun et al. developed a coarse-grained model of TX/TL dynamics [10], and Vilkhovoy et al. used constraint-based modeling to evaluate the performance limits of *E. coli* cell-free protein synthesis [8]. More recently, Piorino et al. experimentally characterized plasmid crosstalk in cell-free expression systems [23], and Han et al. developed a mathematical model that explicitly captured resource com-

petition between co-expressed genes [24]. Yurchenko et al. reviewed both mechanism-based and data-driven modeling approaches for cell-free synthetic biology [25], while Ganesh and Maerkl explored the performance limits of the PURE reconstituted system [26]. Our earlier work on the gluconate sensor circuit [27] addressed resource depletion using empirical functions: a logistic decay for transcriptional capacity and an exponential decay for translational capacity. While these ad hoc functions could fit the observed time courses, they lacked a mechanistic basis, did not couple resource consumption to the actual biosynthetic load, and provided limited insight into the underlying resource dynamics.

In this study, we addressed these limitations by developing a mechanistic two-layer resource competition model for a D-gluconate biosensor circuit operating in the reconstituted cell-free system PURExpress. D-gluconate was chosen as the target analyte because its cognate transcription factor, GntR, is well-characterized in *Bacillus subtilis*, *Escherichia coli*, and *Pseudomonas aeruginosa* [28–32], and because gluconate detection is relevant to food safety and industrial fermentation monitoring, where D-gluconate serves as an indicator of microbial activity and product quality [33]. The first layer of the resource model described the allocation of RNA polymerase and ribosome between competing transcription units using Michaelis-Menten-like saturation expressions, while the second layer captured the irreversible depletion of consumable resources through differential equations driven by the cumulative transcription and translation flux. This formulation replaced the previous empirical decay functions with a physically interpretable framework and simultaneously reduced the number of estimated parameters. We estimated model parameters using a multi-objective optimization approach implemented in ParetoEnsembles.jl, which cycled between multi-objective Pareto front exploration and single-objective refinement. The resulting model accurately captured the time-resolved mRNA and protein dynamics under training conditions and, without re-estimation of parameters, predicted the dose-response relationship of Venus protein as a function of D-

¹²⁴ gluconate concentration over four orders of magnitude.

Materials and Methods

Construction of the gluconate-responsive transcription units. Design of the transcription unit was adapted from the P70 series developed by Noireaux and coworkers [34]. A 16 bp *E. coli* GntR operator (ATGTTACCCGTATCAT) was added between the –35 and –10 region of the P70 promoter. The –10 binding site of the RpoD holoenzyme was altered as a result from *GATAAT* in the original promoter to *TATCAT* in our modified P70 promoter (mP70). The gene sequence of the reporter protein (Venus) was added downstream of this modified promoter. The 5-prime untranslated region and the T500 terminator were unchanged from ref [34]. The second transcription unit had the GntR repressor gene downstream of the original (P70) promoter (without the GntR operator). The full constructs were ordered as linear DNA fragments, with 150–200 bp flanker sequences on both ends, from Twist Bioscience. The sequences of the GntR operator and the repressor were obtained from the literature [30, 35].

Cell-free protein synthesis reactions. The cell-free protein synthesis (CFPS) reactions were carried out using the PURExpress In Vitro Protein Synthesis Kit (New England Biolabs Inc) in 384-well plates (Thermofisher NUNC, flat-bottom) in a Varioskan Lux plate reader at 37°C. The working volume of all reactions was 25 μ L, composed of the PURE solutions A (10 μ L) and B (7.5 μ L), RNase inhibitor, Murine (0.5 μ L), NEB *E. coli* RNA Polymerase holoenzyme (2 μ L), and the linear DNA: Venus (7 nM), GntR (10 nM). For the control reactions without GntR, an equal volume of nuclease-free water was used in its stead. For the de-repression reactions, D-gluconate was added while maintaining the same total volume. The *E. coli* RNA Polymerase, Holoenzyme is the core enzyme saturated with sigma factor 70 and initiates RNA synthesis from sigma 70 specific promoters. The gene constructs were ordered as linear DNA fragments, with 150–200 bp flanker sequences on both ends, from Twist Bioscience. Prior to the cell-free protein synthesis reactions, the DNA fragments were PCR amplified using Q5[®] Hot Start High-Fidelity 2X

Master Mix (New England Biolabs Inc) following manufacturer instructions and the PCR reactions were purified using PureLink™ PCR Purification Kit (Invitrogen).

mRNA quantification. Following each CFPS run, the total RNA was extracted from 5 μ L of the reaction mixture using PureLink RNA Mini Kit (Thermo Fisher Scientific) and stored immediately at -80°C . The quantitative RT-PCR reactions were done using Applied Biosystems™ TaqMan™ RNA-to-CT™ 1-Step Kit. An mRNA standard curve was used to determine absolute mRNA concentrations for each of the samples. The mRNA standards were prepared as follows: separate CFPS reactions for 5 nM of plasmids (Venus and GntR) were carried out for 2 hours. Total RNA was extracted using the full reaction volume. This was followed by the removal of 16S and 23S rRNA using the MICROBExpress™ Bacterial mRNA Enrichment Kit (Life Technologies Corporation). Lastly, the MEGAclear™ Kit (Life Technologies Corporation) was used to further purify the mRNA. The mRNA concentrations were determined using the Qubit™ RNA assay kit (ThermoFisher Scientific).

Protein quantification. Venus fluorescence was measured using the Varioskan Lux plate reader at 513 nm (excitation) and 531 nm (emission). The fluorescence was measured with 25 μ L for each of the mixtures. For all measurements, at least three biological replicates were performed. A protein standard curve was used to determine the absolute protein concentrations for each of the samples. The standard curve was prepared by using Venus standards of known concentration. To prepare the standards, histidine-tagged Venus was expressed using CFPS and purified using a Ni spin column (New England Biolabs). The concentration was measured using SDS-PAGE on Mini-PROTEAN Tetra Cell (Bio-Rad Laboratories) using Any kD™ Mini-PROTEAN® TGX Stain-Free™ Protein Gels (Bio-Rad Laboratories). This was followed by staining the gel with Bio-Safe™ Coomassie Stain (Bio-Rad Laboratories). The stained gel was visualized using the ChemiDoc MP and then the concentrations were measured using Image Lab software.

Formulation and solution of model equations. The model formulation built upon the previous study of Adhikari et al. [27], with significant modifications to the resource limitation framework. The concentration of mRNA m_j and protein p_j for each gene j were described using the balance equations:

$$\dot{m}_j = r_{X,j} \bar{u}_j \epsilon_X - \theta_{m,j} m_j \quad (1)$$

$$\dot{p}_j = r_{L,j} \bar{w}_j \epsilon_L - \theta_{p,j} p_j \quad (2)$$

where $r_{X,j} \bar{u}_j$ and $r_{L,j} \bar{w}_j$ denote the transcription and translation rates, respectively (units: concentration/time), and $\theta_{m,j}$ and $\theta_{p,j}$ denote the respective degradation rate constants (units: 1/time). The $\bar{u}_j(\dots) \in [0, 1]$ and $\bar{w}_j(\dots) \in [0, 1]$ terms describe the control input governing the transcription of gene j and the translation of mRNA j , respectively. The $\epsilon_X(\dots) \in [0, 1]$ and $\epsilon_L(\dots) \in [0, 1]$ terms describe the availability of consumable transcriptional and translational resources, respectively, and are governed by differential equations described below.

Transcription and translation kinetics. The rate of transcription was modeled as the product of a kinetic limit $r_{X,j}$, a control term $\bar{u}_j(\dots) \in [0, 1]$, and the consumable resource term $\epsilon_X(\dots) \in [0, 1]$. The kinetic limit of transcription was derived from elementary reactions (initiation, elongation, and abortive initiation) leading to the formation of mRNA m_j , as described previously [27]:

$$r_{X,j} = V_{X,j}^{\max} \left(\frac{\mathcal{G}_j}{\tau_{X,j} K_{X,j} + (\tau_{X,j} + 1) \mathcal{G}_j} \right) \quad (3)$$

193 where the initiation time constant $\tau_{X,j}$ and saturation constant $K_{X,j}$ were experimentally
 194 measured by McClure [36]. The maximum transcription rate $V_{X,j}^{\max}$ was formulated as:

$$V_{X,j}^{\max} = R_{X,\text{free}} \left(\frac{\dot{v}_X}{l_{G,j}} \right) \quad (4)$$

195 where $R_{X,\text{free}}$ denotes the free RNA polymerase concentration, $\dot{v}_X$ denotes the RNA poly-
 196 merase elongation rate (nt/h), and $l_{G,j}$ denotes the length of gene j in nucleotides. By
 197 analogy, the kinetic limit of translation was formulated as:

$$r_{L,j} = V_{L,j}^{\max} \left(\frac{m_j}{\tau_{L,j} K_{L,j} + (\tau_{L,j} + 1) m_j} \right) \quad (5)$$

198 where m_j denotes the mRNA concentration and $V_{L,j}^{\max}$ denotes the maximum translation
 199 rate:

$$V_{L,j}^{\max} = K_P R_{L,\text{free}} \left(\frac{\dot{v}_L}{l_{P,j}} \right) \quad (6)$$

200 where $R_{L,\text{free}}$ denotes the free ribosome concentration, $\dot{v}_L$ denotes the translation elonga-
 201 tion rate (aa/s), $l_{P,j}$ denotes the length of protein j in amino acids, and K_P is a polysome
 202 amplification constant.

203 *Transcription and translation control functions.* The formulation of the control function
 204 $\bar{u}(\dots)$ was similar to the previous study of Adhikari et al. [27] and the earlier work of
 205 Moon et al. [37], with pseudo-energy bounds informed by the analysis of Bintu et al. [38].
 206 Suppose the promoter P can exist in one of $s = 1, 2, \dots, S$ possible discrete independent
 207 microstates, where each microstate s has some pseudo energy ϵ_s . Some microstates will
 208 lead to transcription, while others will not. For each microstate s , let us assign a pseudo
 209 energy ϵ_s , where $\epsilon_1 = 0$; we assumed the base state of a promoter P , i.e., the state in
 210 which nothing is bound to the promoter-DNA, has zero pseudo energy. Further, suppose

211 the probability that promoter P is in microstate s follows a Boltzmann distribution:

$$p_i = \frac{1}{Z} \times f_i \exp(-\beta \epsilon_i) \quad i = 1, 2, \dots, \mathcal{S} \quad (7)$$

212 The quantity p_i denotes the probability that promoter P is in microstate i , f_i denotes
 213 a state-specific factor $f_i \in [0, 1]$ (typically a Hill-type expression describing the fraction
 214 of bound effector), β denotes the thermodynamic beta, and Z denotes a normalization
 215 factor. Using the summation law of discrete probability, $\sum_s p_s = 1$, the partition function
 216 is:

$$Z = \sum_{s=1}^{\mathcal{S}} f_s \exp(-\beta \epsilon_s) \quad (8)$$

217 The control function $\bar{u}(\dots)$ was then computed as the probability that the promoter is in
 218 a transcription-competent microstate, given by the sum of probabilities over the subset
 219 $\mathcal{A} \subseteq \Omega$ in which transcription occurs:

$$\bar{u} = \sum_{s \in \mathcal{A}} p_s \quad (9)$$

220 The translation control term $\bar{w}(\dots)$ was set to 1 due to the absence of a translational
 221 regulatory network in the current system.

222 *Mechanistic resource competition model.* A key contribution of this study was the re-
 223 placement of the ad hoc resource depletion functions used previously [27] with a mech-
 224 anistic two-layer resource competition framework. The first layer described the allocation
 225 of transcriptional and translational machinery between competing genes. The free RNA
 226 polymerase and ribosome concentrations were computed algebraically from the total con-
 227 centrations $R_{X,T}$ and $R_{L,T}$ by accounting for the fraction of each machinery sequestered

228 by active genes:

$$R_{X,\text{free}} = \frac{R_{X,T}}{1 + \sum_j f_{X,j}} \quad (10)$$

$$R_{L,\text{free}} = \frac{R_{L,T}}{1 + \sum_j f_{L,j}} \quad (11)$$

229 where the sequestration fractions $f_{X,j}$ and $f_{L,j}$ were given by Michaelis-Menten-like ex-
230 pressions:

$$f_{X,j} = \frac{\mathcal{G}_j}{\tau_{X,j}K_X + (1 + \tau_{X,j})\mathcal{G}_j} \quad (12)$$

$$f_{L,j} = \frac{m_j}{\tau_{L,j}K_L + (1 + \tau_{L,j})m_j} \quad (13)$$

231 These expressions arose naturally from the same initiation-elongation kinetics used to
232 derive the transcription and translation rate laws and ensured that the allocation of ma-
233 chinery shifted dynamically as gene dosages and mRNA concentrations changed over
234 time.

235 The second layer described the irreversible depletion of consumable resources (nu-
236 cleotides, energy cofactors, amino acids, and related metabolites) that were not regener-
237 ated in the cell-free reaction. Two state variables, $\epsilon_X \in [0, 1]$ and $\epsilon_L \in [0, 1]$, represented
238 the fractional availability of transcriptional and translational consumable resources, re-
239 spectively. Their dynamics were governed by:

$$\frac{d\epsilon_X}{dt} = -\alpha_X \left(\sum_j l_{G,j} r_{X,j} \bar{u}_j \right) \epsilon_X \quad (14)$$

$$\frac{d\epsilon_L}{dt} = -\alpha_L \left(\sum_j l_{P,j} r_{L,j} \right) \epsilon_L \quad (15)$$

with initial conditions $\epsilon_X(0) = 1$ and $\epsilon_L(0) = 1$. Here, α_X and α_L are consumption rate constants (units: inverse flux), $l_{G,j}$ is the gene length in nucleotides, and $l_{P,j}$ is the protein length in amino acids. The depletion rates were proportional to the total transcription and translation flux weighted by the respective sequence lengths, reflecting the stoichiometric requirement that longer genes consume more nucleotides per transcript and longer proteins consume more amino acids per polypeptide. This formulation ensured that resource depletion was driven by the actual biosynthetic activity in the system rather than by a pre-determined time course, providing a mechanistic link between gene expression load and resource exhaustion.

Estimation of model parameters. Model parameter values were either taken from published studies, governed by experimental conditions, or estimated in this study. Unknown model parameters κ were estimated by minimizing the squared difference between model simulations and experimental measurements. The minimization problem was formulated as a multi-objective optimization problem with five objectives:

$$\min_{\kappa} \mathcal{O}_k = \sum_{i \in \mathcal{E}_k} \left(\hat{\mathcal{M}}_{ik} - \hat{x}_{ik}(\kappa) \right)^2 \quad k = 1, \dots, 5 \quad (16)$$

subject to:

$$\dot{x} = f(x, \kappa) \quad (17)$$

$$\mathcal{L} \leq \kappa \leq \mathcal{U} \quad (18)$$

$$x(t_0) = x_0 \quad (19)$$

The five objectives were: (1) Venus mRNA sum of squared errors (SSE) at 10 mM D-gluconate, (2) Venus protein SSE at 10 mM D-gluconate, (3) GntR mRNA SSE at 10 mM D-gluconate, (4) Venus protein SSE at 0 mM D-gluconate, and (5) a regularization term

penalizing unrealistic GntR protein accumulation. The inclusion of the 0 mM D-gluconate condition as a training objective constrained the repression strength of GntR. The GntR protein regularization objective was necessary because GntR protein was not experimentally measured; rather than using Venus protein data as a proxy for GntR protein (as was done previously), we allowed the resource competition model to naturally bound GntR protein levels through competition for shared translational resources.

The minimization problem was solved using `ParetoEnsembles.jl`, a multi-objective optimization package implemented in the Julia programming language. `ParetoEnsembles.jl` employed a cycling strategy that alternated between multi-objective exploration of the Pareto front and single-objective refinement. In the multi-objective phase, simulated annealing with Pareto ranking was used to broadly explore the parameter space. In the single-objective phase, the algorithm identified the worst-performing objective across the current ensemble and drilled down on that objective alone, thereby preventing any single data type from being poorly fit. After each cycle, the best parameter sets were reseeded into the next round of multi-objective exploration. This cycling strategy balanced diversity (exploring tradeoffs between objectives) with quality (ensuring adequate fit to each individual measurement type). A total of 25 parameters were estimated, reduced from 28 in the previous model: the four empirical resource depletion parameters (onset, slope, $T_{L,1/2}$, and translation capacity initial condition) were removed and replaced by two mechanistic parameters (α_X and α_L). The final ensemble consisted of $N = 7167$ parameter sets filtered for consistency across training objectives. The model equations were solved using the `DifferentialEquations.jl` package [39].

Morris sensitivity analysis. The importance of model parameters was quantified using Morris sensitivity analysis, a global sensitivity analysis method [40]. The Morris method quantifies the influence of parameters on a model performance function as the mean and variance of an elementary effect. The mean of the elementary effect (μ^*) represents the

284 direct influence of the parameter on system performance; a large absolute mean value
285 indicates a significant influence. Meanwhile, a large variance (σ) suggests that the influ-
286 ence of a parameter is non-linear or results from interactions with other parameters. The
287 Morris sensitivity coefficients were computed using the `GlobalSensitivity.jl` package
288 [41] over 200 trajectories, with parameter ranges derived from the ensemble distribution
289 (mean $\pm$ 3 standard deviations, clamped to the original bounds). Four model outputs
290 were evaluated: Venus mRNA at 6 hours, Venus protein at 12 hours, GntR mRNA at 6
291 hours, and GntR protein at 12 hours.

Results

Circuit architecture and components. The genetic circuit was composed of two genes, P70-GntR and mP70-Venus (Fig. 1). The promoters for each gene were derived from the P70 series developed by Noireaux and coworkers [34]; thus, expression from both promoters was responsive to the bacterial sigma factor 70 (σ_{70}). However, the mP70 promoter contained an additional GntR operator site between the -35 and -10 regions. The transcription factor GntR is an allosteric repressor responsive to D-gluconate. Numerous studies on bacterial regulators, including GntR, TetR, and ArgR, have shown that these protein families have a conserved N-terminal region that promotes binding to specific operator regions on the DNA via a helix-turn-helix (HTH) motif [28, 29, 42–45]. On the other hand, the C-terminal sequence varies for each member in the subfamily, giving rise to different ligand binding specificities. Studies on the gluconate operon of *B. subtilis*, *E. coli*, and *P. aeruginosa* have shown that D-gluconate acts as a de-repressor, relieving the repression of the GntR protein [28–32]. The fluorescent protein Venus was used as the reporter because it was previously shown to have good fluorescence properties in PURExpress [46]. All genes were encoded as PCR-amplified linear DNA, circumventing the plasmid preparation process.

GntR repression and de-repression by D-gluconate. The GntR protein repressed Venus expression in the absence of D-gluconate, and this repression was substantially relieved upon addition of the effector (Fig. 2). An mP70-Venus circuit was first expressed in the reconstituted cell-free system PURExpress to characterize the yield of the fluorescent protein and its mRNA transcript. *E. coli* RNAP holoenzyme saturated with σ_{70} , required for transcription, was added to the reaction mixture because PURExpress was not formulated for gene expression from an *E. coli* promoter. Transcriptional activity was maintained for 6 hours, after which the mRNA began to degrade, presumably due to the combined effect of the depletion of transcription resources and the degradation of mRNA

by ribonucleases (Fig. 2A, yellow circles). Venus protein concentration increased slowly, and while translation continued until the endpoint of measurement (12 h) to yield 2.1 μM , the translation rate decreased after approximately 5 hours (Fig. 2B, yellow circles). The saturating protein concentration suggested that the translational capacity of the cell-free system decreased over the course of the reaction, consistent with the progressive depletion of amino acids and energy cofactors [22, 47].

To characterize repression by GntR, two linear DNA fragments were expressed: *E. coli* GntR placed downstream of the P70 promoter (P70-GntR) and Venus placed downstream of the modified P70 promoter (mP70-Venus). Upon adding the P70-GntR gene, the mRNA transcript levels decreased, with the maximum Venus mRNA concentration at 6 hours decreasing from 44 nM to 11 nM (Fig. 2A, black circles). Venus protein expression was tightly repressed within 6.5 hours (Fig. 2B, black circles); the maximum protein yield at 12 hours decreased from 2.1 μM to 0.6 μM , a 3.5-fold change in the dynamic range. Transcript levels for GntR were more than an order of magnitude higher than those for Venus (Fig. 2C). This difference was attributed to the change in the promoter recognition sequence in the two genes; while the -35 region in both promoters was unchanged, the -10 region for mP70 was changed to *TATCAT* (from *GATAAT* in P70) to accommodate the GntR operator. Additionally, to confirm that the decrease in Venus protein yield was due to repression by GntR and not resource limitation owing to the addition of a second gene, the gene for a non-specific GntR variant from *P. aeruginosa* was added at the same dosage instead of the *E. coli* variant (Fig. S1). In this case, the Venus protein yield at 12 hours was 1.8 μM , indicating that while the yield decreased 14% (suggesting some level of TXTL resource competition), the repression in the previous case was indeed due to the *E. coli* GntR protein.

To study de-repression, two linear DNA fragments (P70-GntR and mP70-Venus) were expressed with the addition of 10 mM D-gluconate (approximately 10-fold the value of K_D

reported in the literature [32]) at the start of incubation (Fig. 2, grey circles). With the addition of the effector, the Venus mRNA transcript levels at 2 hours significantly increased to 7.7 nM (from 0.4 nM when D-gluconate was absent), confirming that D-gluconate prevented GntR from binding to its operator and relieved repression (Fig. 2A, grey circles). However, compared to the simple mP70-Venus circuit, Venus mRNA transcript levels were almost 2-fold lower (4-fold lower at 6 h), suggesting that repression was not entirely relieved. Despite the partial de-repression, Venus protein yield was recovered almost 80%, reaching 1.7 μ M at 12 hours (Fig. 2B, grey circles). The GntR mRNA transcript levels were, on average, almost 2-fold lower when D-gluconate was added, suggesting that D-gluconate either lowered the transcription rate or reduced mRNA stability (Fig. 2C). Taken together, GntR was functionally expressed in PURExpress, effectively interacted with its operator to repress transcription, and D-gluconate acted to relieve this repression.

Mechanistic model training. The mechanistic resource competition model captured the transcription and translation dynamics of the gluconate sensor circuit under the training conditions (Fig. 3). Model parameters were estimated from two experimental conditions: the de-repressed case (10 nM P70-GntR, 7 nM mP70-Venus, 10 mM D-gluconate) and the fully repressed case (10 nM P70-GntR, 7 nM mP70-Venus, 0 mM D-gluconate). The multi-objective/single-objective cycling strategy in ParetoEnsembles.jl produced an ensemble of $N = 7167$ parameter sets that captured the expression dynamics across 12 hours. The ensemble mean and 95% confidence bands for Venus mRNA closely tracked the experimental data, reproducing the rise to peak concentration near 6 hours followed by degradation (Fig. 3A). Venus protein simulations matched the experimental endpoint of approximately 1.7 μ M at 12 hours under 10 mM D-gluconate, with the ensemble mean reaching $1.82 \pm 0.19 \mu$ M (Fig. 3B). Despite the difference of more than an order of magnitude between Venus and GntR mRNA levels, the model effectively captured both measurements (Fig. 3C). Because GntR protein was not experimentally measured, its tra-

jectory was predicted rather than fit; the ensemble predicted a GntR protein concentration of $9.94 \pm 1.74 \mu\text{M}$ at 12 hours (coefficient of variation = 17.5%), bounded naturally by competition for shared translational resources rather than by an artificial constraint (Fig. 3D). The means and standard deviations of the 25 estimated model parameters are listed in Table 1.

Resource depletion dynamics. The mechanistic resource model revealed distinct depletion kinetics for transcriptional and translational consumable resources (Fig. 4). The transcriptional resource variable ϵ_X declined rapidly during the first 6 hours, reaching a value of 0.19 by 12 hours, indicating that approximately 81% of the consumable transcriptional resources (nucleotides and energy cofactors required for transcription) were consumed over the course of the reaction (Fig. 4). This rapid depletion was consistent with the observed peak and subsequent decline of mRNA concentrations near 6 hours. In contrast, the translational resource variable ϵ_L declined more gradually, reaching a value of 0.74 by 12 hours, corresponding to only 26% consumption of translational consumable resources (Fig. 4). The asymmetry between transcriptional and translational resource depletion was reflected in the estimated consumption rate constants: $\alpha_X = 2.3 \times 10^{-5}$ for transcriptional resources and $\alpha_L = 1.6 \times 10^{-4}$ for translational resources, indicating that translational resources were consumed approximately 7-fold faster per unit biosynthetic flux than transcriptional resources. However, because the total transcription flux (weighted by gene length) was much larger than the total translation flux (weighted by protein length), the net effect was a more rapid depletion of the transcriptional resource pool. The differential depletion rates suggested that transcriptional capacity was the primary bottleneck in the PURExpress system under the conditions tested, a finding that was consistent with the observation that mRNA levels peaked and declined while protein levels continued to rise, albeit at a decreasing rate.

Dose-response prediction. The model accurately predicted the dose-response relationship of Venus protein as a function of D-gluconate concentration without re-estimation of parameters (Fig. 5). The Venus concentration at 12 hours was measured for D-gluconate concentrations ranging from 0.0001 mM to 20 mM; above 20 mM, Venus protein expression rapidly declined (data not shown), suggesting inhibition of the PURExpress system at high effector concentrations. To simulate the dose-response relationship, the parameter ensemble trained on the 10 mM and 0 mM D-gluconate conditions was used, varying only the concentration of D-gluconate. The model captured the endpoint Venus levels within the limits of experimental error, yielding a characteristic sigmoidal curve (Fig. 5). The experimental data points fell within the 95% confidence bands of the ensemble prediction across four orders of magnitude of D-gluconate concentration. The estimated dissociation constant was $K_{\text{gluconate}} = 0.25 \pm 0.20$ mM, and the Hill coefficient was $n_{\text{gluconate}} = 1.92 \pm 0.86$. The Hill coefficient near 2 was consistent with cooperative binding of D-gluconate to the GntR dimer, which has been reported in the literature for GntR-family regulators. Compared to the previous model, which estimated $K_{\text{gluconate}} = 0.49$ mM and $n_{\text{gluconate}} = 1.63$ [27], the updated estimates reflected the inclusion of the 0 mM condition as a training constraint, which better anchored the lower asymptote of the dose-response curve. In terms of biosensor performance metrics, the circuit exhibited a dynamic range of approximately 3-fold (from $0.69 \mu\text{M}$ at baseline to $1.82 \mu\text{M}$ at saturation), with a limit of detection near 0.1 mM D-gluconate (the lowest concentration at which Venus protein was distinguishable from the repressed baseline). The response time was approximately 6–8 hours to reach 90% of the endpoint signal, limited by the combined kinetics of GntR expression, gluconate binding, and Venus accumulation in the cell-free system.

Discussion

This study demonstrated that a mechanistic resource competition model provided an improved and physically interpretable framework for describing the dynamics of a cell-free biosensor circuit. The two-layer formulation, comprising machinery allocation and consumable resource depletion, replaced the ad hoc logistic and exponential decay functions used in our earlier model [27] and simultaneously reduced the number of estimated parameters from 28 to 25. Importantly, the mechanistic formulation linked resource exhaustion to the actual biosynthetic load placed on the system, rather than to a predetermined time course, enabling the model to generalize across conditions with different gene dosages or effector concentrations.

The estimated resource depletion parameters provided quantitative insight into the capacity limitations of the PURExpress system. The finding that 81% of transcriptional resources were consumed by 12 hours, compared to only 26% of translational resources, identified transcription as the primary bottleneck under the conditions tested. This result has practical implications for circuit design: adding more genes or increasing promoter strength would further accelerate the depletion of an already limiting transcriptional resource pool, potentially reducing the operational lifetime of the sensor. The 7-fold difference in per-unit-flux consumption rates between translation (α_L) and transcription (α_X) suggested that the energetic or material cost per amino acid incorporated exceeded the cost per nucleotide incorporated, a finding that was qualitatively consistent with the higher ATP equivalents required for aminoacyl-tRNA charging and peptide bond formation compared to phosphodiester bond formation.

The machinery allocation layer of the resource model provided a natural mechanism for gene-gene coupling through competition for shared RNA polymerase and ribosome pools. Analysis of the ensemble revealed that approximately 40% of the total RNAP was sequestered by active genes at steady state, while ribosome sequestration was negli-

gible ($<0.5\%$) for this three-gene circuit. The RNAP allocation was constant over time (since gene concentrations were fixed), but the ribosome allocation shifted dynamically as mRNA concentrations changed. This asymmetry suggested that RNAP competition was the relevant machinery constraint for circuits of this scale, while ribosome competition would become significant only for circuits with more genes or higher mRNA concentrations. These findings were consistent with the resource competition framework of Gyorgy and Murray [20], who showed that shared machinery creates implicit coupling between genetic modules, and with the whole-cell resource allocation model of Weisse et al. [21]. Our framework extended these approaches to the cell-free context, where consumable resources impose an additional time-dependent constraint that is absent in living cells (which regenerate metabolites through metabolism). Concurrently, Han et al. developed a mathematical model of cell-free TX/TL with plasmid crosstalk that also captured resource competition effects [24]; our two-layer formulation differs from their approach in that we explicitly separated machinery allocation (algebraic, instantaneous) from consumable depletion (ODE, cumulative), enabling distinct physical interpretations for each resource constraint. Piorino et al. experimentally demonstrated that plasmid crosstalk in cell-free systems depends on DNA concentration and the type of transcriptional machinery [23], a finding that was consistent with our observation that RNAP allocation was the relevant machinery constraint for the circuit studied here. The distinction between renewable machinery resources and non-renewable consumable resources is unique to cell-free systems and represents a key contribution of this work.

This coupling was particularly important for predicting GntR protein levels: because GntR protein was not experimentally measured, the previous model had used Venus protein data as a proxy to constrain GntR expression. In the current model, GntR protein was naturally bounded by competition for shared resources, eliminating the need for this artificial constraint and yielding a predicted GntR protein concentration of $9.90 \pm 1.75 \mu\text{M}$

at 12 hours. This prediction was consistent with a back-of-envelope estimate: Venus accumulated to approximately 2.1 μM from 7 nM DNA template in the absence of GntR (Fig. 2A), and given GntR's higher DNA dosage (10 nM) and stronger promoter (P70 versus mP70), a naive scaling would predict GntR protein in the range of 15–30 μM . The model's lower estimate of approximately 10 μM reflected the dampening effect of shared resource competition, which redistributed translational capacity between the two genes.

Global sensitivity analysis using the Morris method confirmed that the resource consumption parameters α_X and α_L were the most influential parameters for all model outputs (Fig. 6). The transcriptional resource consumption rate α_X dominated the sensitivity of Venus mRNA, GntR mRNA, and both protein outputs, while α_L was the second most influential parameter for protein concentrations. Among the remaining parameters, the mRNA degradation modifiers and the translation time constants were the most sensitive, consistent with the observation that mRNA stability and translational kinetics are key determinants of protein yield in cell-free systems. The gluconate binding parameters ($n_{\text{gluconate}}$, $K_{\text{gluconate}}$) appeared in the top ten for Venus protein but not for the other outputs, confirming that these parameters affected Venus expression exclusively through the GntR-mediated repression pathway.

Analysis of parameter correlations across the ensemble revealed several structurally identifiable clusters that reflected mechanistic coupling within the model (Fig. 7). The translation time constants $\tau_{\text{protein,GntR}}$ and $\tau_{\text{protein,Venus}}$ were strongly positively correlated with each other ($\rho = 0.82$) and negatively correlated with the translation saturation constant K_L ($\rho = -0.85$ and -0.84 , respectively), reflecting the well-known trade-off between Michaelis constant and time constant in enzyme kinetic expressions. The pseudo-energy for Venus transcription ($\Delta G_{\text{Venus,RNAP}}$) was strongly anti-correlated with the Venus mRNA time constant ($\rho = -0.74$), indicating a compensatory relationship between promoter strength and initiation kinetics. The GntR mRNA degradation modifier was anti-correlated

with the transcriptional resource consumption rate α_X ($\rho = -0.72$), indicating that faster mRNA degradation could be compensated by slower resource depletion to maintain the same net mRNA output. These correlations provided useful information about the structural identifiability of the model: while individual parameter values within a correlated cluster could not be precisely determined, the constrained combinations of parameters that the data supported were biophysically meaningful.

The multi-objective/single-objective cycling strategy implemented in `ParetoEnsembles.jl` proved effective for navigating the tradeoffs inherent in fitting multiple data types simultaneously. The cycling approach balanced breadth (exploring the Pareto front across five objectives) with depth (drilling down on the worst-performing objective in each cycle), producing a large and well-diversified ensemble of 5464 parameter sets. The inclusion of the 0 mM D-gluconate condition as a training objective was critical for constraining the repression strength of GntR; without this constraint, the model could compensate for weak repression by adjusting other parameters, leading to unrealistic dose-response predictions. The resulting dissociation constant ($K_{\text{gluconate}} = 0.25 \pm 0.20$ mM) was lower than the value estimated previously ($K_{\text{gluconate}} = 0.49$ mM), reflecting the tighter constraint on the repressed state, and was in reasonable agreement with literature values for GntR-gluconate binding [32].

Several limitations of the current study should be noted. First, GntR protein was not directly measured, so the predicted GntR protein trajectory could not be experimentally validated. While the resource competition model provided a natural bound on GntR accumulation, direct measurement of GntR protein (e.g., via Western blot or mass spectrometry) would strengthen confidence in the model predictions. Second, the model did not capture the inhibitory effect of high D-gluconate concentrations (above 20 mM) on the PURExpress system, which was observed experimentally but not included in the model formulation. Incorporating an inhibition term for high effector concentrations would ex-

tend the predictive range of the model but was beyond the scope of this study. Third, the α_X and α_L parameters, while mechanistically motivated, were empirical consumption rate constants that aggregated many underlying biochemical processes (nucleotide depletion, energy cofactor exhaustion, byproduct accumulation) into single parameters. A more detailed model of the metabolic network underlying transcription and translation in PURExpress could replace these lumped parameters with explicit metabolite balances, as has been done for extract-based systems using metabolomic measurements [9, 48], but would require additional experimental measurements to constrain the expanded parameter space. Recent advances in LC/MS-based metabolite quantification in cell-free reactions [48] could provide the data needed to decompose the lumped α parameters into individual metabolite consumption terms in future work.

Conclusions

In this study, we developed and validated a mechanistic resource competition model for a D-gluconate responsive biosensor circuit operating in the reconstituted cell-free system PURExpress. The model incorporated a two-layer resource framework: machinery allocation described the distribution of RNA polymerase and ribosome between competing genes, while consumable resource depletion captured the irreversible exhaustion of nucleotides, energy cofactors, and amino acids through differential equations driven by cumulative biosynthetic flux. This mechanistic formulation replaced the ad hoc decay functions used previously, reduced the number of estimated parameters, and provided a physical interpretation of the resource limitations observed in cell-free reactions. The model accurately captured the transcription and translation kinetics under training conditions and, without re-estimation of parameters, predicted the dose-response relationship of Venus protein as a function of D-gluconate concentration over four orders of magnitude. The resource depletion analysis revealed that transcriptional resources were the primary

547 bottleneck in the PURExpress system, with 81% consumed by 12 hours compared to 26%
548 for translational resources. These findings provide quantitative design rules for allocating
549 transcriptional and translational capacity in synthetic gene circuits and demonstrate the
550 value of mechanistic resource models for improving the predictive capability of cell-free
551 biosensor frameworks.

Conflict of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions

J.V. directed the study. A.A., A.N., and H.L. conducted the cell-free experimental measurements. A.N. conducted qPCR experiments. J.V., A.M., and A.A. developed the mechanistic resource competition model and the parameter ensemble. A.M., A.A., and J.V. analyzed the model ensemble and generated figures for the manuscript. The manuscript was prepared and edited for publication by A.A., A.M., A.N., and J.V. All authors reviewed this manuscript.

Funding

The work described was supported by the Center on the Physics of Cancer Metabolism at Cornell University through Award Number 1U54CA210184-01 from the National Cancer Institute. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Cancer Institute or the National Institutes of Health. We also acknowledged the financial support to J.V. from the Robert Frederick Smith School of Chemical and Biomolecular Engineering, Cornell University.

Data Availability Statement

Model code is available under an MIT software license from the GitHub repository: <https://github.com/varnerlab/Gluconate-Sensor-Model-Paper>. The mRNA and protein measurements presented in this study are available in the `code/data` directory of the repository in comma separated value (CSV) format. The parameter estimation, simulation, and analysis scripts are provided in the `code/scripts` directory, along with instructions for

575 reproducing all results.

References

1. Swartz JR (2018) Expanding biological applications using cell-free metabolic engineering: an overview. *Metabolic engineering* 50: 156–172.
2. Silverman AD, Karim AS, Jewett MC (2020) Cell-free gene expression: an expanded repertoire of applications. *Nature reviews genetics* 21: 151–170.
3. Bundy BC, Hunt JP, Jewett MC, Swartz JR, Wood DW, et al. (2018) Cell-free biomanufacturing. *Current opinion in chemical engineering* 22: 177–183.
4. Vilkhovoy M, Adhikari A, Vadhin S, Varner JD (2020) The evolution of cell free biomanufacturing. *Processes* 8: 675.
5. Garenne D, Haines MC, Romantseva EF, Freemont P, Strychalski EA, et al. (2021) Cell-free gene expression. *Nature reviews methods primers* 1: 1–18.
6. Moore SJ, MacDonald JT, Freemont PS (2017) Cell-free synthetic biology for in vitro prototype engineering. *Biochemical Society Transactions* 45: 785–791.
7. Hunt AC, Rasor BJ, Seki K, Ekas HM, Warfel KF, et al. (2025) Cell-free gene expression: methods and applications. *Chemical Reviews* 125: 91–149.
8. Vilkhovoy M, Horvath N, Shih CH, Wayman JA, Calhoun K, et al. (2018) Sequence Specific Modeling of *E. coli* Cell-Free Protein Synthesis. *ACS Synth Biol* 7: 1844–1857.
9. Horvath N, Vilkhovoy M, Wayman J, Calhoun K, Swartz J, et al. (2020) Toward a genome scale sequence specific dynamic model of cell-free protein synthesis in *Escherichia coli*. *Metab Eng Comm* 10: e00113.
10. Karzbrun E, Shin J, Bar-Ziv RH, Noireaux V (2011) Coarse-grained dynamics of protein synthesis in a cell-free system. *Physical review letters* 106: 048104.
11. Soltani M, Davis BR, Ford H, Nelson JAD, Bundy BC (2018) Reengineering cell-free protein synthesis as a biosensor: Biosensing with transcription, translation, and protein-folding. *Biochemical Engineering Journal* 138: 165–171.

12. Gräwe A, Dreyer A, Vornholt T, Barteczko U, Buchholz L, et al. (2019) A paper-based, cell-free biosensor system for the detection of heavy metals and date rape drugs. *PloS one* 14: e0210940.
13. Voyvodic PL, Bonnet J (2020) Cell-free biosensors for biomedical applications. *Current opinion in biomedical engineering* 13: 9–15.
14. Jung JK, Alam KK, Verosloff MS, Capdevila DA, Desmau M, et al. (2020) Cell-free biosensors for rapid detection of water contaminants. *Nature biotechnology* 38: 1451–1459.
15. Pardee K, Green AA, Ferrante T, Cameron DE, DaleyKeyser A, et al. (2014) Paper-based synthetic gene networks. *Cell* 159: 940–954.
16. Wen KY, Cameron L, Chappell J, Jensen K, Bell DJ, et al. (2017) A cell-free biosensor for detecting quorum sensing molecules in *p. aeruginosa*-infected respiratory samples. *ACS synthetic biology* 6: 2293–2301.
17. Li Y, Lucci T, Dujovne MV, Jung JK, Capdevila DA, et al. (2025) A cell-free biosensor signal amplification circuit with polymerase strand recycling. *Nature Chemical Biology* 21: 949–958.
18. McSweeney MA, Patterson AT, Loeffler K, Cuellar Lelo de Larrea R, McNerney MP, et al. (2025) A modular cell-free protein biosensor platform using split T7 RNA polymerase. *Science Advances* 11: eado6280.
19. Voyvodic PL, Pandi A, Koch M, Conejero I, Valjent E, et al. (2019) Plug-and-play metabolic transducers expand the chemical detection space of cell-free biosensors. *Nature communications* 10: 1–8.
20. György A, Jimenez JI, Yazbek J, Huang HH, Chung H, et al. (2015) Isocost lines describe the cellular economy of genetic circuits. *Biophysical journal* 109: 639–646.
21. Weisse AY, Oyarzún DA, Danos V, Swain PS (2015) Mechanistic links between cellular trade-offs, gene expression, and growth. *Proceedings of the National Academy of*

Sciences 112: E1038–E1047.

22. Stögbauer T, Windhager L, Zimmer R, Rädler JO (2012) Experiment and mathematical modeling of gene expression dynamics in a cell-free system. *Integrative Biology* 4: 494–501.
23. Piorino F, Patterson AT, Han Y, Styczynski MP (2023) Plasmid crosstalk in cell-free expression systems. *ACS Synthetic Biology* 12: 2843–2856.
24. Han Y, Patterson AT, Piorino F, Styczynski MP (2025) A mathematical model of cell-free transcription-translation with plasmid crosstalk. *Synthetic Biology* 10: ysaf011.
25. Yurchenko A, Ozkul G, van Riel NAW, van Hest JCM, de Greef TFA (2024) Mechanism-based and data-driven modeling in cell-free synthetic biology. *Chemical Communications* 60: 6466–6475.
26. Ganesh RB, Maerkl SJ (2024) Towards self-regeneration: exploring the limits of protein synthesis in the PURE cell-free transcription-translation system. *ACS Synthetic Biology* 13: 2555–2566.
27. Adhikari A, Vilkhovoy M, Vadhin S, Lim HE, Varner JD (2020) Effective biophysical modeling of cell free transcription and translation processes. *Frontiers in bioengineering and biotechnology* 8: 539081.
28. Miwa Y, Fujita Y (1988) Purification and characterization of a repressor for the *Bacillus subtilis* gnt operon. *The Journal of biological chemistry* 263: 13252–7.
29. Fujita Y, Miwa Y (1989) Identification of an operator sequence for the *Bacillus subtilis* gnt operon. *The journal of biological chemistry* 264: 4201–6.
30. Peekhaus N, Conway T (1998) Positive and negative transcriptional regulation of the *Escherichia coli* gluconate regulon gene gntT by GntR and the cyclic AMP (cAMP)-cAMP receptor protein complex. *Journal of bacteriology* 180: 1777–1785.
31. Yoshida KI, Miwa Y, Ohmori H, Fujita Y (1995) Analysis of an insertional operator mutation (gntOi) that affects the expression level of the *Bacillus subtilis* gnt operon,

and characterization of *gntO* suppressor mutations. *Molecular and general genetics* MGG 248: 583–591.

32. Daddaoua A, Corral-Lugo A, Ramos JL, Krell T (2017) Identification of *gntR* as regulator of the glucose metabolism in *Pseudomonas aeruginosa*. *Environmental microbiology* 19: 3721–3733.

33. Ramachandran S, Fontanille P, Pandey A, Larroche C (2006) Gluconic acid: properties, applications and microbial production. *Food Technology and Biotechnology* 44: 185–195.

34. Shin J, Noireaux V (2010) Efficient cell-free expression with the endogenous *E. coli* RNA polymerase and sigma factor 70. *Journal of biological engineering* 4: 1–9.

35. Izu H, Adachi O, Yamada M (1997) Gene organization and transcriptional regulation of the *gntR* operon involved in gluconate uptake and catabolism of *Escherichia coli*. *Journal of molecular biology* 267: 778–793.

36. McClure WR (1980) Rate-limiting steps in RNA chain initiation. *Proceedings of the National Academy of Sciences* 77: 5634–5638.

37. Moon TS, Lou C, Tamsir A, Stanton BC, Voigt CA (2012) Genetic programs constructed from layered logic gates in single cells. *Nature* 491: 249–53.

38. Bintu L, Buchler NE, Garcia HG, Gerland U, Hwa T, et al. (2005) Transcriptional regulation by the numbers: models. *Current opinion in genetics & development* 15: 116–124.

39. Rackauckas C, Nie Q (2017) *DifferentialEquations.jl*—a performant and feature-rich ecosystem for solving differential equations in Julia. *Journal of Open Research Software* 5.

40. Morris MD (1991) Factorial sampling plans for preliminary computational experiments. *Technometrics* 33: 161–174.

41. Dixit VK, Rackauckas C (2022) *GlobalSensitivity.jl*: Performant and parallel global

sensitivity analysis with julia. *Journal of Open Source Software* 7: 4561.

42. Cuthbertson L, Nodwell JR (2013) The tetr family of regulators. *Microbiology and molecular biology reviews*: MMBR 77: 440–75.

43. Dimas RP, Jordan BR, Jiang XL, Martini C, Glavy JS, et al. (2019) Engineering dna recognition and allosteric response properties of tetr family proteins by using a module-swapping strategy. *Nucleic acids research* 47: 8913–8925.

44. Suvorova IA, Korostelev YD, Gelfand MS (2015) GntR family of bacterial transcription factors and their dna binding motifs: structure, positioning and co-evolution. *PloS one* 10: e0132618.

45. Cherney LT, Cherney MM, Garen CR, Lu GJ, James MNG (2008) Crystal structure of the arginine repressor protein in complex with the dna operator from mycobacterium tuberculosis. *Journal of molecular biology* 384: 1330–40.

46. Lentini R, Forlin M, Martini L, Del Bianco C, Spencer AC, et al. (2013) Fluorescent proteins and in vitro genetic organization for cell-free synthetic biology. *ACS synthetic biology* 2: 482–489.

47. Niederholtmeyer H, Xu L, Maerkl SJ (2013) Real-time mrna measurement during an in vitro transcription and translation reaction using binary probes. *ACS synthetic biology* 2: 411–417.

48. Vilkhovoy M, Dai D, Vadhin S, Adhikari A, Varner JD (2019) Absolute quantification of cell-free protein synthesis metabolism by reversed-phase liquid chromatography-mass spectrometry. *J Vis Exp* .

49. Garamella J, Marshall R, Rustad M, Noireaux V (2016) The all e. coli tx-tl toolbox 2.0: a platform for cell-free synthetic biology. *ACS synthetic biology* 5: 344–355.




Fig. 1: Genetic circuit architecture used in this study. The circles represent the promoters for each transcription unit. mP70 is the modified P70 promoter with the GntR operator included. The circuit components were added to the reaction as linear DNA. D-Gluconate was added as a de-repressor at the start of the reaction.




Fig. 2: Venus mRNA and protein levels for the three experimental conditions. Linear DNA used were: mP70-Venus (7 nM), P70-GntR (0 or 10 nM). **A:** Venus mRNA levels measured using qPCR. **B:** Venus protein levels measured on a plate reader (513 nm excitation/531 nm emission). **C:** GntR mRNA levels measured using qPCR. Yellow circles represent the unrepressed case, i.e., without the addition of GntR gene and D-gluconate. Grey circles represent the de-repressed case, i.e., with the addition of both GntR gene (10 nM) and D-gluconate (10 mM). Black circles represent the repressed case, i.e., with only the addition of GntR gene (10 nM). Error bars represent standard errors.




Fig. 3: Mechanistic resource competition model simulations versus experimental measurements for the de-repressed training condition. Linear DNA used were: mP70-Venus (7 nM), P70-GntR (10 nM). D-gluconate (10 mM) was added as the effector at the start of reaction. **A:** Venus mRNA. **B:** Venus protein. **C:** GntR mRNA. **D:** GntR protein prediction. Black circles represent experimental data with standard errors captured by error bars across at least three replicates. mRNA was measured using qPCR; Venus protein was measured on a plate reader (513 nm excitation/531 nm emission). The ParetoEnsembles.jl parameter ensemble ($N = 7167$) was used to calculate the mean (black line) and 95% confidence estimate of the model simulation (shaded region). GntR protein (panel D) was predicted by the model and not experimentally measured.




Fig. 4: Predicted trajectories of consumable resource availability over the 12-hour cell-free reaction. The transcriptional resource variable ϵ_X (representing the fractional availability of nucleotides and energy cofactors for transcription) and the translational resource variable ϵ_L (representing the fractional availability of amino acids and energy cofactors for translation) were computed from the mechanistic resource depletion model (Eqns. 14–15). By 12 hours, ϵ_X declined to 0.19 (81% depletion) while ϵ_L declined to 0.74 (26% depletion), indicating that transcriptional resources were the primary bottleneck. Lines represent the ensemble mean; shaded regions represent the 95% confidence interval across the parameter ensemble ($N = 7167$).




Fig. 5: Model prediction versus experimental measurements for the dose-response study. Linear DNA used were: mP70-Venus (7 nM), P70-GntR (10 nM). D-gluconate concentration was varied from 0.0001 to 20 mM. Black circles represent Venus protein measurements at the endpoint of the experiment (12 h). Error bars represent standard errors across at least three replicates. The black line represents the ensemble mean of the model prediction, and the shaded region represents the 95% confidence estimate across the parameter ensemble ($N = 7167$). All model parameters were fixed at the values estimated from the 10 mM and 0 mM D-gluconate training conditions; only D-gluconate concentration was varied.




Fig. 6: Global sensitivity analysis of the model parameters using the Morris method. The heatmap represents the normalized absolute mean elementary effect ($|\mu^*|$) of each parameter on four model outputs: Venus mRNA at 6 hours, Venus protein at 12 hours, GntR mRNA at 6 hours, and GntR protein at 12 hours. Values were normalized per output to the range $[0, 1]$ for visualization; darker colors indicate greater sensitivity. The resource consumption parameters α_X and α_L were the dominant parameters for all model outputs. Morris sensitivity coefficients were computed over 200 trajectories using parameter ranges derived from the ensemble distribution.




Fig. 7: Spearman rank correlation matrix of the estimated model parameters across the ensemble ($N = 7167$). Strong positive correlations (blue) and negative correlations (red) reveal parameter dependencies that reflect structural identifiability constraints. Notable correlated clusters include the translation time constants ($\tau_{\text{protein,GntR}}$, $\tau_{\text{protein,Venus}}$) and the translation saturation constant (K_L), which exhibited pairwise correlations $|\rho| > 0.82$, reflecting the well-known trade-off between Michaelis constant and time constant in enzyme kinetics. The mRNA degradation modifier for GntR was strongly anti-correlated with α_X ($\rho = -0.72$), indicating a compensatory relationship between mRNA stability and transcriptional resource depletion rate.

Table 1: Estimated model parameters for the D-gluconate sensor circuit. Values represent the ensemble mean $\pm$ standard deviation across $N = 7167$ parameter sets. The first group contains pseudo-energy parameters for promoter configurations, the second group contains binding and kinetic parameters, and the last two rows are the new resource consumption parameters introduced in this study.

Description	Parameter	Value	Units	Reference
RNA Polymerase concentration	R_X	0.13	μM	set by experiment
Ribosome concentration	R_L	2.0	μM	set by experiment
Transcription elongation rate	$\dot{v}_X$	60	nt/s	[49]
Translation elongation rate	$\dot{v}_L$	16.5	aa/s	[49]
Transcription saturation coeff.	K_X	0.054	μM	[36]
Transcription initiation time	$k_{\text{init},X}$	22	s	[36]
Translation initiation time	$k_{\text{init},L}$	1.5	s	[27]
Venus gene length	$l_{G,\text{Venus}}$	720	nt	from DNA sequence
GntR gene length	$l_{G,\text{GntR}}$	996	nt	from DNA sequence
cRNAP + P70 pseudo-energy	$\epsilon_{P70,cRNAP}$	0.027 ± 0.023	$k_B T$	—
hRNAP + P70 pseudo-energy	$\epsilon_{P70,hRNAP}$	-0.75 ± 0.85	$k_B T$	—
cRNAP + mP70 pseudo-energy	$\epsilon_{mP70,cRNAP}$	2.99 ± 1.34	$k_B T$	—
hRNAP + mP70 pseudo-energy	$\epsilon_{mP70,hRNAP}$	-0.28 ± 0.40	$k_B T$	—
GntR + mP70 operator	$\epsilon_{mP70,GntR}$	-1.26 ± 0.72	$k_B T$	—
Hill coeff. P70 + hRNAP	$n_{P70,hRNAP}$	1.35 ± 1.32	—	—
Dissoc. const. P70 + hRNAP	$K_{P70,hRNAP}$	0.52 ± 3.30	μM	—
Hill coeff. mP70 + hRNAP	$n_{mP70,hRNAP}$	1.93 ± 1.77	—	—
Dissoc. const. mP70 + hRNAP	$K_{mP70,hRNAP}$	4.38 ± 6.80	μM	—
Hill coeff. mP70 + GntR	$n_{mP70,GntR}$	1.40 ± 1.28	—	—
Dissoc. const. mP70 + GntR	$K_{mP70,GntR}$	0.83 ± 2.10	μM	—
TX time const. GntR	$\tau_{X,GntR}$	0.094 ± 0.108	—	—
TX time const. Venus	$\tau_{X,Venus}$	11.3 ± 13.4	—	—
TL time const. GntR	$\tau_{L,GntR}$	34.4 ± 26.8	—	—
TL time const. Venus	$\tau_{L,Venus}$	3.17 ± 3.04	—	—
Deg. modifier mRNA GntR	$\delta_{m,GntR}$	0.47 ± 0.16	—	—
Deg. modifier mRNA Venus	$\delta_{m,Venus}$	0.069 ± 0.092	—	—
Deg. modifier protein GntR	$\delta_{p,GntR}$	0.084 ± 0.14	—	—
Deg. modifier protein Venus	$\delta_{p,Venus}$	0.012 ± 0.049	—	—
Deg. modifier protein σ_{70}	$\delta_{p,\sigma_{70}}$	0.14 ± 0.18	—	—
Translation saturation const.	K_L	105 ± 143	μM	—
Hill coeff. gluconate-GntR	$n_{\text{gluconate}}$	1.92 ± 0.86	—	—
Dissoc. const. gluconate-GntR	$K_{\text{gluconate}}$	0.25 ± 0.20	mM	—
TX resource consumption rate	α_X	$(2.5 \pm 1.1) \times 10^{-5}$	$(\mu\text{M} \cdot \text{nt} \cdot \text{hr})^{-1}$	—
TL resource consumption rate	α_L	$(6.7 \pm 8.4) \times 10^{-5}$	$(\mu\text{M} \cdot \text{aa} \cdot \text{hr})^{-1}$	—

Supplemental Information




Fig. S1: Venus protein levels measured for two cases: addition of *P. aeruginosa* GntR gene and in the absence of GntR gene. Linear DNA used were: mP70-Venus (7 nM), P70-GntR (0 or 10 nM). The *P. aeruginosa* GntR variant does not bind to the *E. coli* GntR operator, serving as a control for resource competition effects.