
Variance-Corrected Multi-Asset Equity Simulation with Hybrid Hidden Markov Marginals

Abdulrahman Alswaidan

Robert Frederick Smith School of Chemical and Biomolecular Engineering
Cornell University, Ithaca, NY 14850
aa2725@cornell.edu

Jeffrey D. Varner*

Robert Frederick Smith School of Chemical and Biomolecular Engineering
Cornell University, Ithaca, NY 14850
jdv27@cornell.edu

Abstract

Synthetic multi-asset equity data must reproduce each asset’s return distribution and its relationship with the market. A common approach adds a synthetic full-return path to a market factor as though the synthetic path were a residual. Because that path already contains the asset’s total variance, the construction adds market variance a second time. We derived a correction that centers and rescales each synthetic path before adding the market factor. We tested the method on 423 non-market assets in a 424-asset United States equity and exchange-traded-fund universe. The correction preserved each generator’s target variance, recovered its calibrated market loading with low error, and retained heavy-tailed return distributions without fitting a second model to the regression residuals. On 416 complete asset histories held out from 2025, it increased the mean Kolmogorov–Smirnov pass rate over naive composition and brought the median ratio of synthetic to observed variance close to one. Its one-day left-tail 99% Value-at-Risk exceedance rate was similar to that of the best residual-fit method, although both exceeded the nominal rate. We generated the per-asset paths with a hidden Markov model whose jump-duration variant lengthened visits to extreme-return states. On a decade of daily data for a broad-market exchange-traded fund (ticker SPY), this variant improved the match to volatility clustering relative to the no-jump model while maintaining strong distributional fit. We selected it as a practical compromise, not because it outperformed every baseline. The multi-asset method still omits residual dependence between assets, which can overstate the benefit of daily rebalancing. Code and fitted per-asset models are released with the paper.

Keywords: hidden Markov model; jump-duration mechanism; single-index model; synthetic financial data; variance correction; Value-at-Risk coverage check; volatility clustering; heavy-tailed marginals.

*Corresponding author.

1 Introduction

Researchers and practitioners often use synthetic paths to stress-test risk models against unobserved market scenarios, validate portfolio optimization algorithms, and evaluate allocation strategies across many simulations [1, 2]. However, synthetic financial data often fail to capture important statistical features of real returns [3]. Daily excess growth rates for a broad-market exchange-traded fund (ticker SPY) from 2014–2024 illustrate three widely documented stylized facts [4, 5] (Fig. 1). Most observed returns lie close to zero, but extreme returns occur much more often than a Gaussian model predicts (Fig. 1a and Fig. 1b). Raw growth rates show little correlation across days (Fig. 1c). Absolute growth rates, in contrast, remain positively correlated over longer periods (Fig. 1d), so large and small changes cluster over time. Matching the marginal return distribution alone therefore does not reproduce how volatility evolves over time. Multi-asset simulations must also reproduce cross-asset dependence to evaluate portfolio risk and performance under different rebalancing strategies [6, 7]. Building a multi-asset simulation therefore starts with choosing a single-asset generator that reproduces each asset’s return distribution and volatility dynamics.

Single-asset generators differ in the return features they reproduce and in the effort required to fit them across a large asset universe. Generalized autoregressive conditional heteroskedasticity (GARCH) models capture volatility clustering, but may require heavy-tailed innovations to reproduce extreme returns [8, 9]. Stochastic-volatility models represent persistent changes in volatility, while jump-diffusion models represent abrupt extremes [10–12]. Jumps alone do not produce persistent volatility. Deep generative models can learn flexible return distributions and temporal patterns. Some generative adversarial networks (GANs) reproduce volatility clustering and other path-dependent features [13, 14]. Other evaluations of GANs and the time-series generative adversarial network (TimeGAN) miss the absolute-return autocorrelation associated with volatility clustering [15–17]. Hidden Markov models provide interpretable market regimes, but standard transition rules can leave high-volatility states too quickly to reproduce long clusters of large returns [18–20]. Choosing a single-asset model requires balancing distributional fit, volatility persistence, interpretability, and the effort required to fit the model across hundreds of assets. Portfolio simulation adds a separate requirement because independently fitted single-asset models do not specify how assets move together.

Multivariate hidden-state models allow assets to share market regimes, but become harder to fit as the number of assets grows [21]. Copula models combine separately fitted asset distributions with a model of cross-asset dependence and can capture nonlinear and tail relationships [6, 22, 23]. However, copula models require choosing a dependence structure, and standard static copulas do not describe how dependence changes across days. Factor models are easier to apply to large asset universes because each asset is linked to a small number of shared factors instead of modeling every relationship between pairs of assets [24–26]. The single-index model separates each asset’s return into a market component and an asset-specific residual. After fitting the model, the historical residuals are easy to calculate. The residuals can be resampled or used to fit a separate generator for each asset. Fitting residual generators is a direct solution, but adds a second fitting step for every asset. Generating residual paths independently also misses sector, style, and other relationships not explained by the market. Reusing an existing full-return generator avoids the second fit and preserves the fitted return distribution and temporal behavior. However, a full-return draw already contains the asset mean and market-related variance. Treating the full-return draw as a residual can shift the mean and count market variance twice. The problem is not how to calculate residuals. The problem is whether an existing full-return generator can be reused without shifting the mean or inflating the variance. Dependence among the residuals remains a separate problem.

In this study, we explored how to build a multi-asset generator from separately fitted single-asset components and a shared market path without shifting expected returns or overcounting market variance. We used a hidden Markov model (HMM) for each asset, with Laplace-quantile states, Student- t emissions, and counted transitions. Temporary episodes in the extreme states distinguished the hidden Markov model with jumps (HMM-WJ) from the hidden Markov model with no jumps (HMM-NJ). On a decade of SPY data, HMM-WJ reduced absolute-return autocorrelation error relative to HMM-NJ while retaining a high in-sample Kolmogorov–Smirnov pass rate. GARCH achieved a lower autocorrelation error but did not match the return distribution, so HMM-WJ served as a practical per-asset generator rather than a uniformly superior model. For multi-asset composition, we centered and rescaled each asset-level path before adding a shared SPY path. The correction removed the mean shift and prevented the market contribution to variance from being counted




Figure 1: **Empirical stylized facts of SPY daily excess growth rates (2014–2024).** Panel (a) shows the heavy-tailed return density; the Laplace maximum-likelihood fit tracks the empirical peak and tails while the Gaussian fit does not. Panel (b) shows departure from the diagonal in a normal quantile-quantile plot at both extremes. Panel (c) shows the autocorrelation function of raw excess growth rates. Individual correlations are small, although the Ljung–Box test detects joint serial dependence. Panel (d) shows the autocorrelation function of absolute excess growth rates, which decays slowly across the reported lags.

twice without fitting another model to each asset’s regression residuals. We tested the correction on 423 non-market assets in a 424-asset universe and set the jump probability to zero to isolate the composition rule. With all 2014–2024 parameters held fixed, the correction increased the mean Kolmogorov–Smirnov pass rate across 416 complete asset histories from 2025 and brought the median ratio of synthetic to observed variance from inflated under naive composition to near one. Independent residual generation did not reproduce residual dependence and could overstate returns from daily rebalancing.

2 Related Work

Single-asset return models treat changing volatility, market regimes, and extreme moves in different ways. Autoregressive conditional heteroskedasticity (ARCH) and generalized autoregressive conditional heteroskedasticity (GARCH) models allow conditional variance to change over time [8, 9]. Even with Gaussian innovations, the changing variance can produce an unconditional return distribution with excess kurtosis. Heavy-tailed innovations can improve the tail fit when Gaussian GARCH understates the frequency of extreme returns. Stochastic-volatility models make volatility a latent process [27, 10, 28]. Jump-diffusion models add discontinuous price moves [11, 12], but a jump component alone does not explain persistent volatility. Markov-switching ARCH and GARCH models combine discrete regimes with changing conditional variance [29, 30]. Hidden-state models represent returns with a small number of market regimes. In a hidden Markov model (HMM), each latent state has a separate return distribution. Gaussian-emission HMMs can reproduce heavy tails and weak raw-return autocorrelation, but may still miss volatility clustering [31, 19]. Hidden semi-Markov

models allow more flexible state durations and can improve volatility persistence [20]. An HMM does not create a jump process by itself. Rare extreme moves must come from the state-dependent return distributions or from a transition rule designed for such events. Deep generative models learn return distributions and temporal features without specifying a small set of market regimes or a parametric return process. Generative adversarial networks (GANs) are one class of deep generative model. Quant GANs have reproduced several financial time-series features, including volatility clustering and leverage effects [13]. Sig-Wasserstein GANs use path signatures to learn features that depend on the order of observations [14]. In contrast, evaluations of GAN paths, including paths from the time-series generative adversarial network (TimeGAN), have found too little autocorrelation in absolute returns [15–17]. The conflicting results show that flexible architectures do not guarantee a match to financial stylized facts. Model design, training, and evaluation all matter [1, 3].

Multi-asset generation also requires a model of cross-asset dependence. Multivariate GARCH models estimate time-varying covariance matrices. The dynamic conditional correlation model reduces the number of parameters by separating individual volatility models from correlation dynamics [32, 33]. Copulas separate each asset’s return distribution from the dependence model. Dynamic copulas allow cross-asset dependence to change over time, while vine copulas provide flexible high-dimensional constructions [34, 23]. Factor models reduce dimension by linking each asset to a small number of shared factors [24–26]. Shared-state HMMs and multivariate GANs can also generate joint market moves [21, 35]. A standard factor model can resample historical residuals or fit a separate residual generator for every asset. This paper addresses a narrower problem. We start with single-asset generators already fitted to full returns and ask whether they can be reused in a factor model. The regression residuals are easy to calculate. The difficulty is that a full-return draw already contains the asset mean and market-related variance. Using that draw as a residual can shift the mean and count part of the variance twice. The correction developed here removes the mean shift and variance double count without fitting a second generator for every asset. It does not model dependence among the remaining residuals.

3 Methods

We fitted a marginal generator for each asset and combined the generated paths with a variance-corrected single-index model. The dataset contained 424 United States-listed equities and exchange-traded funds observed from 2014 through 2024. The single-asset analysis focused on the broad-market exchange-traded fund with ticker SPY. It used 2,767 daily volume-weighted average prices for fitting, which yielded 2,766 growth rates, and the next 250 prices from 2025, which yielded 249 holdout growth rates. The multi-asset analysis used daily closing prices. Of the 423 non-market assets in the training universe, 416 had complete 2025 histories and were included in the multi-asset holdout evaluation. We defined the excess growth rate $G_{i,j}$ for ticker i between trading days $j - 1$ and j from prices $P_{i,j-1}$ and $P_{i,j}$.

$$G_{i,j} \equiv \left(\frac{1}{\Delta t} \right) \ln \left(\frac{P_{i,j}}{P_{i,j-1}} \right) - r_f. \quad (1)$$

Here, $\Delta t = 1/252$ is the fraction of a trading year spanned by one trading day, and r_f is a constant continuously compounded risk-free rate. We used $r_f = 0.043$ for the SPY training period and $r_f = 0.0421$ for its 2025 holdout, based on Treasury Separate Trading of Registered Interest and Principal of Securities (STRIPS) bond yields. The excess growth rate has units of year^{-1} and is time-additive under continuous compounding.

3.1 Per-asset hidden Markov marginal generator

For each asset, the hidden Markov model with jumps (HMM-WJ) converted the continuous return series into a sequence of N ordered return regimes. At each time step, the model first selected a regime and then drew a return from that regime’s distribution (Fig. 2). We represented the model as $\mathcal{M} = (\mathcal{S}, \mathcal{O}, \mathbf{T}, \mathbf{E}, \bar{\pi})$ [18]. The hidden state $S_t \in \mathcal{S} = \{1, \dots, N\}$ identifies the regime at time t , and the observed value is the excess growth rate G_t from Eq. (1). We omit the ticker index i when discussing one asset. The transition matrix $\mathbf{T}$ sets the probabilities of moving between regimes, $\mathbf{E}$ gives the return distribution within each regime, and $\bar{\pi}$ gives the initial regime probabilities. We ordered the regimes from the most negative to the most positive returns. Their boundaries were equally spaced quantiles of a Laplace distribution fitted to the asset’s in-sample returns. Specifically,

$Q_k = F_L^{-1}(k/N; \mu_L, b_L)$, where μ_L and b_L are the fitted location and scale. We set $Q_0 = -\infty$ and $Q_N = +\infty$ and assigned return G_t to state k when $Q_{k-1} < G_t \leq Q_k$. The Laplace distribution captured the sharp concentration of small price movements in Fig. 1a [36, 37]. We computed its closed-form quantiles for all 424 tickers [38].

The within-state emission followed a location-scale Student- t distribution.

$$G_t \mid S_t = k \sim \mu_k + \sigma_k \cdot t_\nu, \quad \nu = 5, \quad (2)$$

where t_ν denotes a standard Student- t random variable with ν degrees of freedom. For observations assigned to state k , μ_k is the sample mean and σ_k is the sample standard deviation. A standard t_ν has variance $\nu/(\nu - 2)$, so σ_k is a scale parameter rather than the emission standard deviation. At $\nu = 5$, the emission standard deviation is $\sigma_k \sqrt{5/3}$. We selected $\nu = 5$ from a sensitivity sweep. We used $\nu = \infty$ for the Normal baseline; Table 1 gives the explored range. Holding the rest of the model fixed, the Student- t emission raised simulated kurtosis toward the observed value and raised the Kolmogorov–Smirnov (KS) and Anderson–Darling (AD) pass rates relative to the Normal emission. We also compared HMM-WJ with a hidden semi-Markov model (HSMM) baseline adapted from Bulla and Bulla [20] (Table S3; Online Appendix S6). The emission and HSMM comparisons separated the effects of heavy-tailed emissions and the jump-duration mechanism. We estimated the transition matrix $\mathbf{T}$ by counting observed transitions, which avoided the initialization sensitivity of iterative expectation-maximization [38]. The counted matrix was concentrated near its diagonal, but individual states, including tail states, usually lasted only 1–2 days (Fig. S3 in Online Appendix S5). The counted transition probabilities caused the model to leave extreme-return states too quickly to reproduce the observed volatility clustering [20]: a chain fitted by counting reproduces the one-step state-pair frequencies exactly, and its absolute-return autocorrelation therefore decays geometrically at a rate set by the counted matrix. Online Appendix S4 derives the fitted generator’s stationary law and autocorrelation structure and quantifies both this rapid mixing and the correction added by the jump mechanism.

We added jump episodes to extend visits to extreme-return states. When no jump episode was active, a new episode began with probability ϵ . Its duration was $K \sim \text{Poisson}(\lambda)$, where λ is the mean episode length [39]. Because a Poisson draw can be zero, $K = 0$ caused no forced tail step. For $K > 0$, the model temporarily replaced ordinary transitions with draws from two tail sets. The set $\mathcal{S}_- = \{1, \dots, N_{\text{tail}}\}$ contained the most negative states, and $\mathcal{S}_+ = \{N - N_{\text{tail}} + 1, \dots, N\}$ contained the most positive states. At each forced step, the model selected the negative set with probability p_{neg} and the positive set otherwise. It made the tail-set choice independently at each step, so negative extreme returns occurred slightly more often than positive ones. Table 1 lists ϵ , λ , p_{neg} , and N_{tail} . Two closed-form quantities summarize the mechanism’s operating point: forced steps occupy a long-run fraction of approximately $\epsilon\lambda/(1 + \epsilon\lambda)$ of all steps, and a path of T steps contains at least one episode with probability approximately $1 - e^{-\epsilon T}$, so ϵ sets how many paths jump while λ sets how long an episode lasts (Online Appendix S4). We obtained the initial-state distribution $\bar{\pi}$ by solving $\bar{\pi}\mathbf{T} = \bar{\pi}$ with $\sum_k \bar{\pi}_k = 1$ [40, 41]. Because $\mathbf{T}$ was estimated by counting, $\bar{\pi}$ equals the vector of in-sample state frequencies up to end effects, so the model’s stationary growth-rate distribution is the frequency-weighted mixture of the state emission distributions. The solved $\bar{\pi}$ is exact for the ordinary transition matrix but only approximate for the jump-augmented process because it does not include the remaining jump duration. The full simulation procedure appears in Algorithm 1. Each step followed $\mathbf{T}$ or continued a jump episode and then drew a return from the selected state’s emission distribution.

We chose the jump probability ϵ and mean duration λ to match the SPY absolute-return autocorrelation and kurtosis [4, 5, 19]. Jump-diffusion models use Poisson events to add instantaneous price jumps [11, 12]. In HMM-WJ, a jump episode extended a visit to the tail states. We searched a grid of (ϵ, λ) values. At each grid point, we averaged squared errors in the autocorrelation function (ACF) of absolute returns through lag L and weighted squared errors in excess kurtosis over the jump-active paths. We minimized the objective $J(\epsilon, \lambda)$.

$$J(\epsilon, \lambda) = \frac{1}{|\mathcal{R}_J|} \sum_{r \in \mathcal{R}_J} \left[\sum_{\tau=1}^L (\text{ACF}_{\text{obs}}(\tau) - \text{ACF}_r(\tau))^2 + w_\kappa (\kappa_{\text{obs}} - \kappa_r)^2 \right], \quad (3)$$

where $\mathcal{R}_J$ is the set of jump-active simulated paths, $\text{ACF}_{\text{obs}}(\tau)$ and κ_{obs} are the empirical absolute-growth autocorrelation at lag τ and excess kurtosis, and $\text{ACF}_r(\tau)$ and κ_r are the corresponding




Figure 2: **Architecture of the hidden Markov model with jumps (HMM-WJ).** When no jump episode is active, the chain transitions according to the empirical transition matrix $\mathbf{T}$ with probability $1 - \epsilon$ (thin bidirectional arrows) or enters a Poisson jump episode with probability ϵ (dashed upward arrow to the Poisson clock). During an episode, each of the $K \sim \text{Poisson}(\lambda)$ forced steps independently draws the bottom tail set $\mathcal{S}_-$ (red, lowest-valued states) with probability p_{neg} or the top tail set $\mathcal{S}_+$ (blue, highest-valued states) otherwise. Ordinary transitions then resume. Small bell curves above each state show the state-conditional location-scale Student- t ($\nu = 5$) emission distribution. Tail sets are defined as the N_{tail} lowest- and highest-quantile states under the Laplace quantile partition.

Algorithm 1 Hidden Markov model with jumps (HMM-WJ) single-asset marginal generator.

Require: Fitted parameters $(\mathbf{T}, \mathbf{E}, \bar{\pi}, \epsilon, \lambda, p_{\text{neg}}, N_{\text{tail}})$; horizon T ; tail sets $\mathcal{S}_- = \{1, \dots, N_{\text{tail}}\}$ and $\mathcal{S}_+ = \{N - N_{\text{tail}} + 1, \dots, N\}$.

Ensure: State sequence $\{S_t\}_{t=1}^T$ and growth-rate path $\{G_t\}_{t=1}^T$.

- 1: Draw $S_1 \sim \bar{\pi}$; draw $G_1 \sim \mu_{S_1} + \sigma_{S_1} t_5$; $k \leftarrow 0$ {stationary start; k : remaining forced-step counter}
- 2: **for** $t = 2, \dots, T$ **do**
- 3: **if** $k > 0$ **then**
- 4: $\mathcal{J} \leftarrow \mathcal{S}_-$ w.p. p_{neg} , else $\mathcal{S}_+$; $S_t \sim \text{Uniform}(\mathcal{J})$; $k \leftarrow k - 1$ {forced tail step; sign drawn per step}
- 5: **else if** $\text{Uniform}(0, 1) > \epsilon$ **then**
- 6: $S_t \sim \mathbf{T}_{S_{t-1}, \cdot}$ {standard Markov transition}
- 7: **else**
- 8: $K \sim \text{Poisson}(\lambda)$ {trigger jump; K is the episode length}
- 9: **if** $K = 0$ **then**
- 10: $S_t \sim \mathbf{T}_{S_{t-1}, \cdot}$ { $K = 0$: no forced steps, standard transition}
- 11: **else**
- 12: $\mathcal{J} \leftarrow \mathcal{S}_-$ w.p. p_{neg} , else $\mathcal{S}_+$; $S_t \sim \text{Uniform}(\mathcal{J})$; $k \leftarrow \min(K - 1, T - t)$ {first forced step; K steps total, sign per step}
- 13: **end if**
- 14: **end if**
- 15: Draw $G_t \sim \mu_{S_t} + \sigma_{S_t} t_5$ {Eq. (2)}
- 16: **end for**

values for path r . We used κ for kurtosis and K for jump duration. We simulated 200 independent paths of 2,766 trading days at each grid point. A path with no jump reduces to the hidden Markov model with no jumps (HMM-NJ) and contains no information about (ϵ, λ) , so the objective included only paths with at least one jump [39]. Within those paths, λ controls both the height and the horizon of the slow autocorrelation component while the kurtosis penalty limits the total forced-step mass; ϵ acts mainly through the share of paths that contain an episode (Online Appendix S4). The value of w_κ set the relative weight of the kurtosis error. The search ranges, w_κ , L , and the remaining

HMM-WJ and SIM hyperparameters are reported in Table 1. We fixed the selected hyperparameters before running the generator comparisons and the holdout evaluation.

3.2 Variance-corrected multi-asset composition

The single-asset model generated each ticker separately and did not specify cross-asset dependence. For the multi-asset model, we added a simulated market path as a common factor. We used the single-index model (SIM) of Sharpe [24].

$$g_i(t) = \alpha_i + \beta_i g_m(t) + \varepsilon_i(t), \quad t = 1, \dots, T. \quad (4)$$

Here α_i is the intercept for asset i , g_m is the market growth-rate path, β_i is the asset's market loading, and ε_i is its asset-specific residual path. We use ε_i for the residual path and ϵ for the jump probability in the single-asset model. We estimated the intercept and market loading (α_i, β_i) from real data by ordinary least squares (OLS). The choice of ε_i determined the composition method. A standard SIM draws independent Gaussian residuals with the variance estimated by OLS. These Gaussian residuals preserve the fitted market relationship but do not retain the heavy tails and volatility regimes produced by the single-asset generator. We called the Gaussian-residual method *Gaussian SIM*.

Naive full-path composition. In the naive construction, a full synthetic return path $\tilde{g}_i$ from the single-asset generator served as the residual.

$$g_i = \alpha_i + \beta_i g_m + \tilde{g}_i. \quad (5)$$

The fitted intercept and market term already set the conditional mean of asset i , while the full-return draw generally has $\mathbb{E}[\tilde{g}_i] \neq 0$. Equation (5) therefore has conditional mean $\alpha_i + \mathbb{E}[\tilde{g}_i] + \beta_i g_m$, not $\alpha_i + \beta_i g_m$. We centered every full-return path before composition to remove the duplicate mean. Centering subtracted the realized path mean.

$$\bar{\tilde{g}}_i \equiv \frac{1}{T} \sum_{t=1}^T \tilde{g}_i(t), \quad \tilde{g}_i^c(t) \equiv \tilde{g}_i(t) - \bar{\tilde{g}}_i. \quad (6)$$

Therefore, $T^{-1} \sum_t \tilde{g}_i^c(t) = 0$ exactly. Subtracting a constant leaves both the sample variance and the sample covariance with g_m unchanged. We used the centered construction $g_i = \alpha_i + \beta_i g_m + \tilde{g}_i^c$ as the *naive* benchmark.

Centering corrected the mean but not the variance. The path variance $\sigma_{\text{gen},i}^2 = \text{Var}(\tilde{g}_i) = \text{Var}(\tilde{g}_i^c)$ is the *generator variance*. It already represents the asset's total variance. Adding the market term therefore adds variance a second time. We sampled the residual and market paths independently. Their covariance is zero in population and close to zero on a long realized path. We approximated the naive composed variance accordingly.

$$\text{Var}(g_i) \approx \beta_i^2 \sigma_m^2 + \sigma_{\text{gen},i}^2, \quad \text{Cov}(\tilde{g}_i^c, g_m) \approx 0, \quad (7)$$

where $\sigma_m^2 = \text{Var}(g_m)$ is the variance of the market path. The excess is $\beta_i^2 \sigma_m^2$, so the error is largest for assets with large market loadings.

Variance correction. We set $\sigma_{\text{gen},i}^2$ as the target total variance for asset i . The market term contributed $\beta_i^2 \sigma_m^2$, so the residual term could contribute only the remaining variance. We defined s_i as the nonnegative asset-specific residual scale, set $\varepsilon_i = s_i \tilde{g}_i^c$, and required $\text{Var}(\beta_i g_m + s_i \tilde{g}_i^c) = \sigma_{\text{gen},i}^2$. Under the zero-covariance assumption, solving for s_i gives the correction.

$$s_i^2 = 1 - \frac{\beta_i^2 \sigma_m^2}{\sigma_{\text{gen},i}^2} \quad (8)$$

The *market loading ratio* ρ_i is the market variance contribution divided by the target variance.

$$\rho_i \equiv \frac{\beta_i^2 \sigma_m^2}{\sigma_{\text{gen},i}^2}, \quad (9)$$

Under the zero-covariance assumption, ρ_i is the fraction of the target variance contributed by the market, and $s_i^2 = 1 - \rho_i$. At $\rho_i = 1$, the unconstrained rule leaves no asset-specific variance; for

$\rho_i > 1$, it has no real-valued solution. Values of ρ_i at or above one can arise when β_i is large, the generator variance is low, or the synthetic market is more volatile than the calibration market. We required the asset-specific draw to contribute at least a fraction f of the target variance. We set $\rho_i^{\max} = 1 - f$ and reduced the effective loading β_i^{eff} when $\rho_i > \rho_i^{\max}$. We applied the piecewise correction in Eq. (10).

$$\beta_i^{\text{eff}} = \begin{cases} \beta_i & \text{if } \rho_i \leq \rho_i^{\max} \\ \text{sign}(\beta_i) \sqrt{(1-f) \frac{\sigma_{\text{gen},i}^2}{\sigma_m^2}} & \text{if } \rho_i > \rho_i^{\max} \end{cases} \quad s_i^2 = \begin{cases} 1 - \rho_i & \text{if } \rho_i \leq \rho_i^{\max} \\ f & \text{if } \rho_i > \rho_i^{\max} \end{cases} \quad (10)$$

The correction retained the sign of the calibrated loading β_i . We used $f = 0.10$, so the asset-specific draw contributed at least 10% of the target variance for non-tracker assets (Table 1). Online Appendix S9.1 examines the substitution of a full-return draw for the residual: it derives what the construction preserves exactly and how the composed shape dilutes as ρ_i grows.

High- R^2 index trackers. The non-tracker branch targeted total generator variance. For an index-tracking fund, we instead targeted its fitted relationship with the market. The coefficient of determination R^2 measures the fraction of the asset's variance explained by the market. For example, SPY regressed on itself has $\beta = 1$, $R^2 = 1$, and no residual variance. QQQ and SPYG also had high R^2 against SPY. The non-tracker branch can change the fitted R^2 even when it preserves total variance. We therefore used a separate tracker branch when $R_{i,\text{real}}^2$ was at least R_{preserve}^2 . We set $R_{\text{preserve}}^2 = 0.80$, which separated the index trackers from individual stocks in our universe (Fig. S5). For the tracker branch, we computed the residual variance required to recover the calibrated R^2 under the zero-covariance assumption.

$$\sigma_{\varepsilon,\text{target}}^2 = \beta_i^2 \sigma_m^2 \cdot \frac{1 - R_{i,\text{real}}^2}{R_{i,\text{real}}^2} \quad (11)$$

We rescaled the generator draw to the target residual variance.

$$\varepsilon_i = \left(\sqrt{\frac{\sigma_{\varepsilon,\text{target}}^2}{\sigma_{\text{gen},i}^2}} \right) \tilde{g}_i^c. \quad (12)$$

When $R_{i,\text{real}}^2 = 1$, the target residual variance is zero. SPY regressed on itself therefore has no residual term. A scale factor greater than one increases the variance of the generator draw and can weaken the original marginal fit.

Complete composition rule. The tracker branch targeted the calibrated R^2 . The non-tracker branch targeted the generator variance and reduced β_i only when the market term would leave less than the required asset-specific variance. After selecting a branch, we added the scaled asset draw to the market term. The full procedure appears in Algorithm 2. If used, a residual-copula rank reorder was applied jointly after scaling and before composition. The reorder preserved the empirical variance of each residual path. Centering kept the path mean controlled by the intercept and market term. Under the zero-covariance assumption, OLS estimates from longer composed paths converge to the effective market loading and its intercept. The variance and loading properties are asymptotic. They do not specify exact finite-sample values or the complete return distribution. Online Appendix S9 gives the algebra. The construction adds only same-day market dependence; it does not reproduce lead-lag effects or market-driven volatility clustering.

Algorithm 2 Hybrid single-index model (SIM) composition with variance correction.

Require: Nonconstant $g_m, \tilde{g}_i \in \mathbb{R}^T, i = 1, \dots, N$; each $\tilde{g}_i$ is independent of g_m .

Require: Parameters $(\alpha_i, \beta_i, R_{i,\text{real}}^2)$; floor $f \in (0, 1)$; threshold $R_{\text{preserve}}^2 \in (0, 1]$.

Ensure: Composed paths $g_i \in \mathbb{R}^T$ and branch flags b_i .

```

1:  $\sigma_m^2 \leftarrow \text{Var}(g_m)$ .
2: for each asset  $i = 1, \dots, N$  do
3:    $\sigma_{\text{gen},i}^2 \leftarrow \text{Var}(\tilde{g}_i)$ ;  $\tilde{g}_i^c \leftarrow \tilde{g}_i - \bar{\tilde{g}}_i$ .
4:   if  $R_{i,\text{real}}^2 \geq R_{\text{preserve}}^2$  then
5:      $v_i \leftarrow \beta_i^2 \sigma_m^2 (1 - R_{i,\text{real}}^2) / R_{i,\text{real}}^2$ .
6:      $s_i \leftarrow \sqrt{v_i / \sigma_{\text{gen},i}^2}$ ;  $\beta_i^{\text{eff}} \leftarrow \beta_i$ ;  $b_i \leftarrow \text{R2-PRESERVE}$ .
7:   else
8:      $\rho_i \leftarrow \beta_i^2 \sigma_m^2 / \sigma_{\text{gen},i}^2$ 
9:     if  $\rho_i \leq 1 - f$  then
10:       $s_i \leftarrow \sqrt{1 - \rho_i}$ ;  $\beta_i^{\text{eff}} \leftarrow \beta_i$ ;  $b_i \leftarrow \text{VARIANCE}$ .
11:    else
12:       $s_i \leftarrow \sqrt{f}$ ;  $\beta_i^{\text{eff}} \leftarrow \text{sign}(\beta_i) \sqrt{(1 - f) \sigma_{\text{gen},i}^2 / \sigma_m^2}$ .
13:       $b_i \leftarrow \text{CLIPPED}$ .
14:    end if
15:  end if
16:   $\varepsilon_i \leftarrow s_i \tilde{g}_i^c$ .
17: end for
18: Optional: jointly rank-reorder  $\{\varepsilon_i\}_{i=1}^N$ .
19: for each asset  $i = 1, \dots, N$  do
20:    $g_i \leftarrow \alpha_i + \beta_i^{\text{eff}} g_m + \varepsilon_i$  {Eq. (4)}
21: end for

```

Table 1: Hidden Markov model with jumps (HMM-WJ) and single-index model (SIM) composition hyperparameters. Values used for the main SPY analysis, explored ranges, and locations of sensitivity checks. For the 423-ticker SIM composition, we set $\epsilon = 0$ instead of the SPY value. The cross-asset validation used per-ticker (ϵ^*, λ^*) values for NVDA, JNJ, and JPM (Table S9).

Symbol	Description	SPY value	Explored range
<i>State partition and emission</i>			
N	Laplace-quantile states	100	$\{30, \dots, 350\}$, Online Appendix S7
ν	Student- t emission degrees of freedom	5	$\{3, \dots, 30, \infty\}$
<i>Jump-duration override</i>			
ϵ	jump probability per step	10^{-4}	$[10^{-4}, 2.5 \times 10^{-2}]$, Eq. (3)
λ	mean jump-episode duration (steps)	90	$[10, 160]$, Eq. (3)
p_{neg}	probability a jump targets $\mathcal{S}_-$	0.52	fixed
N_{tail}	quantile states per tail set	5	fixed (at $N = 100$)
w_κ	kurtosis-penalty weight, Eq. (3)	0.20	fixed
L	autocorrelation lag window, Eq. (3)	25 days	fixed
<i>SIM composition</i>			
f	idiosyncratic variance floor	0.10	$\{0.05, \dots, 0.30\}$
R_{preserve}^2	branch-selection threshold	0.80	$\{0.70, \dots, 0.90\}$

Evaluation protocol. We evaluated the method on 424 United States equities and exchange-traded funds from Polygon.io. Each ticker had 2,767 daily closing prices from January 2014 through December 2024, which yielded 2,766 growth rates. This complete-case selection conditioned the analysis on assets that survived and retained data coverage throughout the training window. We used SPY as the market and composed the other 423 assets. For each asset, we fitted the state and emission model used in the SPY analysis (Table 1; $r_f = 0$). We set the jump probability to zero so that the experiment measured multi-asset composition without per-ticker jump calibration. We regressed each real asset on SPY to estimate $\alpha_i, \beta_i, R_{i,\text{real}}^2$, and residual variance. Across the sample, β_i ranged from 0.03 to 1.79, and $R_{i,\text{real}}^2$ ranged from 0.001 to 0.933 (Fig. S5).

We compared six ways to construct the asset-specific term. The naive method used the full JumpHMM draw without correction. The Gaussian SIM method used independent Gaussian residuals with variance estimated by OLS. The hybrid method used Algorithm 2. The JumpHMM-on-residuals method fitted a new JumpHMM to the OLS residual series after converting the residuals to a synthetic price path by cumulative compounding. The block-bootstrap method sampled the OLS residual series with the Politis–Romano stationary bootstrap [42] and a mean block length of $\sqrt{T} \approx 50$. The GARCH(1,1)- t method fitted a conditional-variance model to each residual series [9]. We excluded thirty non-stationary GARCH fits from the GARCH results. For each ticker, the naive and hybrid methods received the same JumpHMM draw, which isolated the effect of the variance correction. The other four methods generated their own residual draws. We did not use the optional cross-sectional rank reorder because the experiment tested each ticker’s marginal construction separately.

We evaluated each composed series on factor recovery, distributional fit, and variance preservation. For factor recovery, a new OLS regression produced the absolute errors $|\hat{\beta} - \beta|$ and $|\hat{R}^2 - R_{\text{real}}^2|$; lower values indicate closer recovery. For distributional fit, we used Kolmogorov–Smirnov (KS) and Anderson–Darling (AD) pass rates, Wasserstein-1 distance, excess kurtosis, and the Hill upper-tail index. Higher KS and AD pass rates and lower Wasserstein distance indicate a closer match; kurtosis and the Hill index describe tail behavior. For variance preservation, we used the composed-to-generator variance ratio. A ratio of one indicates equal variances. Online Appendix S3 gives the metric definitions. We ran 100 replications per ticker with seed 1234, producing 42,300 series per method. Pass rates used the significance level $\alpha = 0.05$.

For the holdout evaluation, we froze all marginal fits, residual models, and SIM parameters estimated from 2014–2024. We evaluated the methods against the 249 growth rates observed in 2025. Of the 423 non-market training assets, 416 had complete holdout histories; GARCH results covered 386 because its thirty non-stationary training fits remained excluded. For each replication, we generated a 249-day SPY path from the training-period SPY marginal and used the same simulated market path in all asset compositions. Observed 2025 SPY and asset returns were used only for scoring. We repeated the KS, AD, Wasserstein-1, variance, kurtosis, and one-day left-tail Value-at-Risk (VaR) evaluations with 100 paths per ticker. For each VaR threshold, we also used Kupiec’s unconditional-coverage test to compare the observed number of exceedances with the nominal rate [43]. Because goodness-of-fit tests have less power on 249 observations than on 2,766, we also compared each synthetic path with a randomly selected contiguous 249-day block from the training history. We first averaged replication-level results within ticker and then summarized across tickers so each asset received equal weight. No 2025 data were used for fitting. The 249-day training blocks provided a comparison at the same sample length as the holdout period.

4 Results

4.1 Single-asset generator validation

To determine whether the marginal generator reproduced the relevant SPY behavior, we compared it with the empirical distributional and temporal targets (Figs. 2 and 1). In both the in-sample (2014–2024, $T = 2,766$) and out-of-sample (2025, $T = 249$) windows, the real SPY series had heavy tails, and the Jarque–Bera test rejected normality. The series also showed persistent volatility clustering. Individual raw-return correlations were small, although the Ljung–Box test detected joint serial dependence (Table S1). Using these targets, we calibrated (ϵ, λ) for HMM-WJ against the in-sample SPY series using the grid search in Eq. (3) (Algorithm 1; search range in Table 1). The selected values were $(\epsilon, \lambda) = (10^{-4}, 90)$, although nearby grid points gave similar objective values. Under the counted transition matrix, ordinary state residence times were only 1–2 steps and the implied autocorrelation of absolute growth rates decayed to near zero within a week, whereas the selected jump duration had a mean of 90 steps (Fig. S3; Online Appendix S4). We then checked whether the results depended on the number of states. The primary analysis used $N = 100$ states. HMM-NJ pass rates moved by less than one percentage point across $N = 30$ to 200, whereas HMM-WJ traded its two metrics against each other as the partition changed, gaining autocorrelation accuracy and losing AD pass rate as the states grew narrower (Table S6). At $N = 350$, some states were never visited and their emission parameters could not be estimated (Online Appendix S7). We kept $N = 100$ as a midpoint of that tradeoff for the generator comparisons that followed.

Table 2: **In-sample and out-of-sample model comparison for SPY.** Results summarize 1,000 simulated paths. Values in parentheses are standard errors, and bold entries are best among the fitted generative models within each panel; the empirical bootstrap is excluded from this highlighting. Metric definitions and standard-error calculations are given in Online Appendix S3.

Model	KS (%)	AD (%)	κ	ACF-MAE	Coverage (%)	W_1	Hellinger
<i>In sample: 2014–2024 (2,766 days; observed $\kappa = 7.715$)</i>							
Bootstrap	100.0 (<0.1)	99.7 (0.2)	7.6 (0.08)	0.060 (<0.001)	100.0 (<0.1)	0.062 (0.001)	0.042 (<0.001)
Gaussian	0.0 (<0.1)	0.0 (<0.1)	−0.0 (<0.01)	0.060 (<0.001)	13.1 (0.1)	0.399 (0.001)	0.148 (<0.001)
Laplace	44.0 (1.6)	43.3 (1.6)	3.0 (0.02)	0.060 (<0.001)	37.4 (1.5)	0.138 (0.001)	0.072 (<0.001)
GARCH(1,1)	5.5 (0.7)	1.9 (0.4)	8.2 (0.37)	0.031 (0.001)	29.3 (1.4)	0.304 (0.006)	0.100 (0.001)
GRU	0.6 (0.2)	0.2 (0.1)	5.7 (0.16)	0.036 (<0.001)	17.2 (0.5)	0.421 (0.003)	0.134 (<0.001)
HSMM	82.0 (1.2)	42.5 (1.6)	4.8 (0.08)	0.059 (<0.001)	68.7 (1.1)	0.176 (0.001)	0.113 (<0.001)
HMM-NJ	99.3 (0.3)	98.2 (0.4)	8.1 (0.14)	0.059 (<0.001)	100.0 (<0.1)	0.081 (0.001)	0.072 (<0.001)
HMM-WJ	98.3 (0.4)	94.8 (0.7)	7.5 (0.09)	0.053 (<0.001)	100.0 (<0.1)	0.097 (0.001)	0.074 (<0.001)
<i>Out of sample: 2025 (249 days; observed $\kappa = 6.867$)</i>							
Bootstrap	97.4 (0.5)	98.5 (0.4)	6.0 (0.18)	0.043 (<0.001)	100.0 (<0.1)	0.232 (0.002)	0.205 (0.001)
Gaussian	62.2 (1.5)	46.3 (1.6)	−0.0 (<0.01)	0.043 (<0.001)	36.4 (1.9)	0.452 (0.002)	0.235 (0.001)
Laplace	88.0 (1.0)	92.9 (0.8)	2.7 (0.05)	0.043 (<0.001)	74.7 (1.0)	0.263 (0.002)	0.211 (0.001)
GARCH(1,1)	80.3 (1.3)	72.0 (1.4)	1.6 (0.07)	0.026 (<0.001)	96.0 (1.3)	0.507 (0.018)	0.232 (0.001)
GRU	71.8 (1.4)	44.8 (1.6)	2.9 (0.10)	0.033 (<0.001)	77.8 (2.9)	0.488 (0.005)	0.240 (0.001)
HSMM	96.2 (0.6)	96.7 (0.6)	4.1 (0.13)	0.042 (<0.001)	100.0 (0.2)	0.287 (0.002)	0.239 (0.001)
HMM-NJ	96.6 (0.6)	96.1 (0.6)	6.2 (0.15)	0.041 (<0.001)	100.0 (<0.1)	0.259 (0.003)	0.207 (0.001)
HMM-WJ	95.4 (0.7)	95.3 (0.7)	6.2 (0.15)	0.040 (<0.001)	100.0 (<0.1)	0.275 (0.005)	0.210 (0.001)

Model abbreviations are generalized autoregressive conditional heteroskedasticity (GARCH), gated recurrent unit (GRU), hidden semi-Markov model (HSMM), hidden Markov model with no jumps (HMM-NJ), and hidden Markov model with jumps (HMM-WJ). Kolmogorov–Smirnov (KS) and Anderson–Darling (AD) report pass rates at $\alpha = 0.05$; coverage is the fraction of empirical quantiles inside the synthetic 90% envelope. κ is simulated excess kurtosis. Mean absolute error in the autocorrelation function (ACF-MAE) is computed for $|G_t|$ through lag 252. Larger values are preferred for KS, AD, and coverage; smaller values are preferred for ACF-MAE, W_1 , and Hellinger distance; κ is compared with the observed value.

To determine whether HMM-WJ improved both distributional and temporal fit, we compared it with seven alternative generators on 1,000 paths of 2,766 trading days each (Table 2; Fig. 3). No baseline was best on both the return distribution and volatility clustering. Bootstrap resampling passed every distribution test but, like all independent sampling methods, did not reproduce clustering. Gaussian sampling passed none of the distribution tests. The GARCH(1,1) model had the lowest mean absolute error in the autocorrelation function (ACF-MAE) of 0.031 but rarely passed the distribution tests. The gated recurrent unit (GRU) [44] showed the same general pattern. The HSMM-style baseline adapted from Bulla and Bulla [20] reached an 82% KS pass rate but underestimated kurtosis by more than one third. Against these baselines, the two HMM variants gave the strongest overall distributional fit among the fitted parametric and latent-state generators. The HMM-NJ and HMM-WJ models had KS pass rates of 99.3% and 98.3%, respectively. The slightly lower HMM-WJ pass rates were consistent with jump episodes sometimes placing too much probability in the tails. Student- t emissions produced simulated kurtosis of 7.5, close to the observed 7.7, whereas Normal emissions underestimated it by about 29% (Table S3; Online Appendix S6). Both HMM variants also reproduced the observed skewness of -0.75 (Table S1; Online Appendix S4). That skewness arose from unequal time spent in negative and positive states rather than from an explicit skewness parameter.

The jump episodes were intended to improve temporal fit without sacrificing marginal fit. The HMM-WJ model reduced ACF-MAE from 0.059 for HMM-NJ to 0.053. About 24% of HMM-WJ paths contained at least one jump and showed more persistent autocorrelation in absolute returns; paths without a jump were equivalent to HMM-NJ. This share, the fraction of steps inside episodes (just under one percent), and the shape of the jump-path autocorrelation all matched the closed-form episode arithmetic derived in Online Appendix S4 (Table S2). We then tested whether the distributional fit carried beyond the training period using the held-out 249 days of 2025. The HMM-NJ and HMM-WJ models retained KS pass rates of 96.6% and 95.4%, with kurtosis close to the observed value (Table 2; Fig. S2). Tests on NVDA, JNJ, and JPM showed strong in-sample fit but more variable holdout performance (Table S9, Online Appendix S8). These comparisons support HMM-WJ as a compromise between distributional and temporal fit. We examined that compromise directly by varying the share of jump-active paths.




Figure 3: **Head-to-head in-sample model comparison for SPY** ($N = 100, 1,000$ simulated paths). *Panel (a)*: marginal density of excess growth rates with in-sample Kolmogorov–Smirnov (KS) pass rates annotated per generator. *Panel (b)*: autocorrelation function of $|G_t|$ at lags 1–252 (shaded bands: 10th–90th percentile across paths) per generator. *Panel (c)*: tail quantile-quantile plot at the 0.1st–99.9th percentile per generator.

To measure how jump-active paths changed the results, we resampled the fitted ensemble at different jump-path fractions f , the share of accepted paths that contain at least one jump, from $f = 0$, where all accepted paths reduce to HMM-NJ, to $f = 1$, where every accepted path contains a jump. This procedure held the fitted model fixed and changed only the mixture of accepted paths. As the jump share rose, the absolute-return autocorrelation error fell monotonically, simulated excess kurtosis declined through the observed value, and the KS and AD pass rates fell with it, while the Hill upper-tail index stayed flat because the Student- t ($\nu = 5$) emission sets the tail exponent independently of jump content. Increasing the jump share therefore traded fit to the distributional body for lower autocorrelation error without changing the tail index, the tradeoff derived in closed form in Online Appendix S4. We therefore used $f \approx 0.25$, the fitted ensemble’s own jump-path share, which kept simulated kurtosis close to the observed value while lowering the autocorrelation error and balanced distributional and temporal fit in the SPY analysis. The multi-asset experiment that followed set the jump probability to zero so that it could focus on how the paths were combined.

4.2 Variance-corrected multi-asset composition

To test whether the variance-corrected SIM preserved both generator variance and calibrated factor loading, we compared six composition methods across the 423-ticker universe on the aggregate scorecard (Table 3). Before comparing the methods, we measured the cross-term omitted from the variance correction. For the full-return generator, the universe-mean normalized cross-covariance with the market path was 1.53×10^{-4} , and the per-ticker values ranged from -0.0061 to 0.0061 (Fig. S1). These values supported the negligible-cross-term approximation used in Eq. (8).

Table 3: **Aggregate scorecard across 423 non-market tickers and 100 replications per ticker.** Errors, distances, and tail metrics are medians per method. Kolmogorov–Smirnov (KS) and Anderson–Darling (AD) pass fractions are pooled across replications. $|\hat{\alpha} - \alpha|$, $|\hat{\beta} - \beta|$, and $|\hat{R}^2 - R_{\text{real}}^2|$: absolute deviation between ordinary least-squares (OLS) recovery and calibration (intercept error has units of yr^{-1}). KS pass and AD pass: fraction of replications whose two-sample test against the real marginal exceeds $p = 0.05$. W_1 : Wasserstein-1 distance to the real marginal. κ : sample excess kurtosis. Hill: tail-index estimate on the upper 5% of $|g|$. SIM denotes the single-index model, and GARCH denotes generalized autoregressive conditional heteroskedasticity.

Method	$ \hat{\alpha} - \alpha $	$ \hat{\beta} - \beta $	$ \hat{R}^2 - R_{\text{real}}^2 $	KS pass (%)	AD pass (%)	W_1	κ	Hill
Naive	0.002	0.022	0.087	7.1	3.5	0.700	7.767	0.369
Gaussian SIM	0.048	0.017	0.008	0.6	0.5	0.682	1.427	0.217
Hybrid	0.002	0.018	0.021	73.6	64.6	0.249	7.165	0.365
JumpHMM-on-residuals	0.049	0.018	0.015	75.8	70.7	0.232	7.262	0.365
Block bootstrap	0.042	0.017	0.020	73.3	66.8	0.232	6.675	0.330
GARCH(1,1)- t	0.047	0.017	0.036	67.4	61.4	0.267	6.069	0.317

All six methods recovered the factor loading with similar accuracy: the median absolute β error ranged from 0.017 to 0.022 (Table 3). Path centering also kept the median absolute intercept error at 0.002 yr^{-1} for both methods that used full-return draws. The residual-fit methods retained random variation in the residual mean over each finite path. Their larger intercept errors therefore do not, by themselves, imply a worse fit to the return distribution.

The larger differences appeared in variance recovery and tail behavior (Table 3; Fig. S6). Gaussian SIM had the lowest median R^2 error because it targets R^2 directly. However, it almost never passed the distributional tests, and its median excess kurtosis was an order of magnitude below the observed value. Centered naive composition retained the shape of the full-return draw, but adding the market factor inflated its variance by $\beta_i^2 \sigma_m^2$. The hybrid method removed that extra variance. It passed 73.6% of KS comparisons and retained heavy tails, with median excess kurtosis of 7.2 and a Hill index of 0.36. Its median R^2 error fell between those of the naive and Gaussian methods because this branch preserves the generator variance rather than targeting the observed R^2 directly. The hybrid method reached that position without a separate residual generator, so we compared it with three residual-fit alternatives, which avoided double counting because the empirical OLS residuals already excluded the market component. Their KS pass rates ranged from 67.4% for GARCH(1,1)- t to 75.8% for JumpHMM-on-residuals. The hybrid rate was two percentage points below JumpHMM-on-residuals and slightly above block bootstrap. JumpHMM-on-residuals and block bootstrap had slightly lower Wasserstein distances. JumpHMM-on-residuals also matched the hybrid’s kurtosis and Hill index almost exactly, whereas the block-bootstrap and GARCH residual fits produced lighter tails. The correction substantially improved on naive composition and approached the best residual-fit method without requiring a second fit for each asset. We next tested whether this ordering persisted after all fitted parameters were frozen.

To test out-of-sample generalization without refitting or look-ahead bias, we froze the fitted models and SIM parameters before evaluating them against 416 complete asset histories from 2025 (Table 4). The hybrid method reached an 87.0% mean cross-ticker KS pass rate, against 83.5% for centered naive composition, and its median composed-to-observed variance ratio was 0.97, whereas naive composition remained inflated at 1.29. JumpHMM-on-residuals reached an 86.2% KS pass rate and a slightly lower median W_1 than the hybrid. Matched-length training blocks produced lower pass rates for every method. The higher 2025 pass rates did not arise solely from comparing samples of 249 rather than 2,766 observations. The average pass rates concealed substantial differences among tickers (Fig. 4). Most ticker-level rates were near the upper end of the range, but every method retained a lower tail of weak fits. For the hybrid method, many tickers improved relative to their matched-length training blocks, while a subset deteriorated out of sample. The held-out marginal improvement was broad but not uniform across assets, motivating a separate check of whether the resulting synthetic tails produced reliable risk thresholds.

Table 4: Frozen-parameter multi-asset evaluation on the 2025 holdout. Results cover 416 non-market tickers and 100 replications per ticker; generalized autoregressive conditional heteroskedasticity (GARCH) covers 386 tickers. Pass rates are first averaged within ticker and then across tickers. The *matched in-sample (IS)* columns compare each synthetic path with a randomly selected contiguous 249-day block from the 2014–2024 training history, matching the holdout sample length. W_1 and the variance ratio are medians across ticker-level medians; the variance ratio is synthetic variance divided by observed 2025 variance.

Method	KS pass (%)		AD pass (%)		W_1 (2025)	variance ratio
	matched IS	2025	matched IS	2025		
Naive	69.4	83.5	54.8	71.6	0.877	1.29
Gaussian SIM	57.4	73.9	44.9	64.6	0.866	0.97
Hybrid	78.8	87.0	66.2	79.7	0.716	0.97
JumpHMM-on-residuals	77.9	86.2	65.6	80.1	0.708	0.98
Block bootstrap	76.3	84.0	64.0	76.5	0.715	0.90
GARCH(1,1)- t	74.5	82.6	62.6	75.0	0.733	0.91




Figure 4: **Cross-ticker distribution of frozen-parameter out-of-sample marginal fit.** Panel (a) shows the distribution of per-ticker 2025 Kolmogorov–Smirnov (KS) pass rates across 100 replications; the dashed line marks 95%. Panel (b) compares each ticker’s hybrid KS pass rate against a matched-length training block with its pass rate against the 2025 holdout. The diagonal denotes equal performance.

4.3 VaR coverage

To test whether marginal fit translated into reliable tail thresholds, we used the frozen synthetic paths to estimate per-ticker one-day left-tail Value-at-Risk (VaR) thresholds and counted exceedances on the real 2025 histories (Table 5). At $\alpha = 0.95$, the hybrid and JumpHMM-on-residuals methods produced mean exceedance rates of 5.97% and 6.07%, respectively, against the 5% nominal rate. At $\alpha = 0.99$, both rates were 1.67%, against the 1% nominal rate. Naive composition was closest at the 99% level (1.21%), while Gaussian SIM had the largest exceedance rate (2.03%). The short holdout contains only about 2–3 expected 99% exceedances per ticker, so we interpret the corresponding Kupiec pass rates and cross-ticker dispersion as low-power checks rather than precise rankings. Taken together, the VaR results showed that stronger marginal fit did not automatically imply nominal tail coverage, motivating a closer look at where the variance correction was most stable.

Table 5: **Out-of-sample per-ticker one-day left-tail Value-at-Risk coverage across the six composition methods.** VaR thresholds at coverage level α were estimated from frozen-parameter synthetic paths and exceedances were counted against the real 2025 growth-rate history. *rate*: mean of the per-ticker mean exceedance rates (nominal: $1 - \alpha$). Cross-ticker standard deviation (SD) is reported in percentage points. *Kupiec pass*: mean within-ticker fraction of replications whose unconditional-coverage test p -value exceeds 0.05.

Method	$\alpha = 0.95$			$\alpha = 0.99$		
	rate (%)	SD (pp)	Kupiec pass (%)	rate (%)	SD (pp)	Kupiec pass (%)
Naive	4.37	2.15	65.8	1.21	0.76	78.9
Gaussian SIM	4.95	2.26	70.0	2.03	1.17	72.1
Hybrid	5.97	2.43	66.3	1.67	0.93	73.4
JumpHMM-on-residuals	6.07	2.59	67.0	1.67	0.95	74.5
Block bootstrap	6.13	2.56	63.7	1.84	1.01	73.4
GARCH(1,1)- t	5.79	2.47	62.4	1.80	1.02	72.0

4.4 Sensitivity analyses

To isolate how the variance correction changed marginal fit, we plotted each ticker’s KS pass rate and variance ratio $\text{Var}(g)/\sigma_{\text{gen}}^2$ against its calibrated β (Fig. 5). Under naive composition, the variance ratio followed $1 + \rho_i$ and exceeded generator variance by 50–75% at high β , and the naive KS pass rate fell toward zero above $\beta \approx 0.7$; the hybrid method held the ratio at one and retained substantially higher pass rates across the β range.




Figure 5: **Effect of the hybrid variance correction on each ticker’s Kolmogorov–Smirnov pass rate and variance ratio.** Each point is a per-ticker median over 100 replications; thick lines are binned medians. *Panel (a):* Kolmogorov–Smirnov pass rate against the real marginal at $\alpha = 0.05$ for naive composition (blue) and hybrid (orange). *Panel (b):* variance ratio $\text{Var}(g)/\sigma_{\text{gen}}^2$ for the same two methods. The dotted curve is the theoretical naive inflation $1 + \rho$ evaluated at the universe-median σ_{gen}^2 ; the dashed line marks a variance ratio of one.

The branch-selection rule assigned 421 tickers to the variance-preserving branch, two trackers to the R^2 -preserving branch, and none to the clipped branch (Figs. S5 and S4; Table S4). The two R^2 -preserving assignments were QQQ and SPYG, and both had a 100% KS pass rate; the highest- R^2 individual stock, BLK, fell about 0.16 below the $R^2_{\text{preserve}} = 0.80$ threshold. Clipping did not occur in the universe: $\rho_i = \beta_i^2 \sigma_m^2 / \sigma_{\text{gen},i}^2$ (Eq. (9)) stayed below $1 - f = 0.90$ for every ticker. Because the R^2 -preserving branch contained only two tickers, we tested it on a synthetic-tracker grid spanning $(\beta_{\text{true}}, R^2_{\text{true}}) \in \{0.8, 1.0, 1.2\} \times \{0.80, 0.85, 0.90, 0.95, 0.99\}$. Each cell had a 100% KS pass rate across replications and recovered its target loading and R^2 (Table S5). The correction removed the observed beta-related variance inflation. We then tested the clipping branch under conditions that would force it to activate.

To test whether the correction weakened as market exposure increased, we bucketed the universe by β quartile. The hybrid KS pass rate fell from 87.6% in the lowest quartile to 65.9% in the highest (Table S7). As β increased, the market term accounted for more of the composed variance, making the marginal more dependent on the market distribution. We tested the clipping rule directly by rescaling the market growth-rate path by $\gamma \in \{1, 2, 3\}$, holding the calibrated marginals fixed (Table S8). At $\gamma = 1$ no ticker crossed the threshold. At $\gamma = 2$ and 3, clipping applied to 76.4% and 95.2% of tickers. The clipping rule was unnecessary under observed market variance but activated as intended when market volatility was artificially increased. At the observed market variance, the floor f did not affect any branch assignment; varying R^2_{preserve} changed only the two tracker assignments identified above.

5 Discussion

Single-asset model choice involved a tradeoff between marginal and temporal fit. Relative to HMM-NJ, HMM-WJ reduced the autocorrelation error for absolute returns, although its in-sample KS and AD pass rates were slightly lower. The GARCH model achieved the lowest autocorrelation error of any generator but rarely passed the distributional tests, and the GRU showed a similar tradeoff. The adapted HSMM-style baseline passed the KS test more often but produced too little kurtosis. Bootstrap resampling passed the distributional tests by drawing directly from the observed returns, but it did not reproduce volatility clustering. We therefore interpret HMM-WJ as a compromise between distributional and temporal fit, not as a uniformly better single-asset model. Reading that compromise from the pass rates requires care, because the two-sample KS and AD tests assume independent observations, whereas the HMM, HSMM, GARCH, and block-bootstrap paths contain dependence. For that reason, we considered the tests alongside Wasserstein and Hellinger distances and the Hill tail index. Model resolution creates a second qualification. A sensitivity sweep over

$N \in \{30, 60, 90, 100, 150, 200\}$ showed little change across the middle of this range for the 2,766-day SPY sample, but shorter histories will support fewer well-populated states. We also tuned the jump parameters (ϵ, λ) on SPY and then applied them across the asset universe. Separate fits for NVDA, JNJ, and JPM achieved in-sample KS pass rates of at least 98.9% (Table S9), but three assets cannot establish that one jump calibration transfers to every return series. Finally, the Student- t emissions model heavy tails but not skewness directly. The fitted skewness instead came from unequal time spent in negative and positive states, so assets with different asymmetry may need a different state partition.

The central multi-asset question was whether a full-return generator could be reused without counting market variance twice. Naive composition added market variance to a draw that already represented the asset’s total variance. The correction removed that double count and passed 73.6% of the full-length in-sample KS comparisons (Table 3). After all fits were frozen, it raised the mean cross-ticker KS pass rate from 83.5% under centered naive composition to 87.0% on the 2025 holdout and kept the median variance ratio at 0.97 (Table 4). The improvement was broad but not uniform: every method retained a lower tail of tickers with weak fits, and some hybrid fits deteriorated relative to matched-length training blocks. A high average pass rate is therefore not a guarantee for every asset. Fitting the return generator directly to the OLS residuals is a competitive alternative on the same comparison, because residual-fit methods avoid double counting once the market component has been removed. The best such method, JumpHMM-on-residuals, had a slightly higher KS pass rate in sample, while the hybrid method had the higher rate in the holdout. The practical difference is reuse: the hybrid construction uses a full-return generator, whereas the residual method requires another fit for each asset. Neither approach achieved nominal tail coverage in the short holdout.

The short-holdout tail result narrowed the comparison further: the hybrid method’s one-day left-tail 99% VaR exceedance rate was 1.67%, the same rounded rate as JumpHMM-on-residuals and above the 1% nominal rate (Table 5). Several design choices also narrowed its interpretation. In sample, we used the observed SPY history as the market path. In the holdout experiment, we generated SPY from its frozen training-period model and reserved the observed 2025 path for scoring. The naive and hybrid methods shared the same JumpHMM draw for each ticker, while the residual methods used draws from their own fitted generators. We omitted the optional cross-sectional rank reorder and evaluated VaR for each ticker rather than for a portfolio. The variance correction also assumes $\text{Cov}(\tilde{g}_i^c, g_m) \approx 0$, which holds here because the asset and market paths were generated independently. A generator sampled from joint market history would require a covariance adjustment. In addition, the present construction adds only contemporaneous market dependence; it does not reproduce lead-lag effects or market-driven volatility clustering. The sensitivity analyses mark the same boundary from the other direction, by showing where the correction weakens inside the observed universe. Hybrid KS pass rates declined from 87.6% in the lowest β quartile to 65.9% in the highest because the market path determined more of the composed distribution as β increased (Table S7). The clipping rule did not activate in the observed universe, where every ticker had $\rho_i < 0.90$, but it activated for most tickers after market volatility was doubled or tripled (Table S8). Only two tracker assets used the R^2 -preserving branch (Table S4). The 249-day holdout also had much less power than the 2,766-day training comparison. Matched-length training blocks partly address that difference, but one year of test data cannot provide a precise ranking from either pass rates or left-tail 99% VaR counts. The evidence therefore supports per-asset variance correction and distributional evaluation, not portfolio-level dependence fidelity.

A rebalanced portfolio shows why that distinction matters. Independent residual draws remove the sector and style covariance present in real assets, and periodic rebalancing converts the excess cross-sectional dispersion into a diversification return that a real portfolio would not earn [45, 7, 46]. Online Appendix S2 derives that term for the composition rule used here and isolates the cross-asset covariance as the point where the construction departs from a real market. The problem affects every method that generates residuals independently, including the block-bootstrap and GARCH variants, so it is a property of the residual model rather than of any one generator. Removing it requires a joint residual model. A synchronized multivariate block bootstrap, residual factor model, or copula could restore cross-asset residual covariance. Multi-factor or lagged-factor versions could also capture dependence omitted by the current single-factor construction, with the variance correction retained as a compositional constraint. Per-asset jump calibration and stress tests involving highly volatile markets or leveraged assets would test the other transfer limits identified here. Separate univariate autoregressive or block-bootstrap models would not be enough: they can improve temporal behavior

within each asset without creating dependence between assets. Adjusting the level of the composed paths does not resolve this limitation, because rescaling or shifting a path leaves the cross-asset covariance untouched. The variance correction is therefore necessary for combining full-return generators with a market factor, but it is not a complete dependence model. Portfolio-level use requires joint residual dynamics.

6 Conclusion

In this study, we developed a variance-corrected single-index composition method that reuses full-return generators in a factor model without double counting their location or variance. Centering first prevented a full-return draw from shifting the calibrated intercept; the subsequent scale correction removed the variance inflation caused by adding the market factor. Across 423 non-market assets, it recovered the calibrated factor loading to a low median absolute error and produced heavy-tailed marginals. On 416 complete held-out asset histories from 2025, it improved the mean cross-ticker KS pass rate over centered naive composition and held the median variance ratio near one, while naive composition remained inflated. Its one-day left-tail 99% Value-at-Risk exceedance rate matched that of JumpHMM-on-residuals to the reported precision but stayed above the nominal rate. Those residual models, fitted directly to the OLS residual series, required a separate fit for each asset and produced a similar marginal pass rate. Solving the composition problem first required a per-asset generator that retained heavy tails and temporal structure, for which we used a hidden Markov marginal model with Laplace-quantile states, Student- t emissions, and direct transition counting. On SPY, its jump-duration variant reduced absolute-return autocorrelation error relative to the no-jump variant while keeping high KS and AD pass rates; GARCH reached a lower autocorrelation error but failed the distributional tests. The single-asset result therefore showed a measured tradeoff between distributional fit and volatility clustering, not uniform superiority. Taken together, these results separate two tasks that are easily conflated when synthesizing multi-asset market data: the marginal generator governs each asset’s distributional and temporal behavior, whereas the variance correction prevents the common market factor from double counting the generator’s total variance.

The current multi-asset implementation does not model cross-asset residual covariance. Independent residuals omit the sector and style dependence that links real assets, and a rebalanced portfolio built on them earns a diversification return that a real portfolio would not. A full joint residual model is needed before using these paths for portfolio-level return forecasts. The variance correction is a necessary compositional constraint rather than a complete dependence model. Portfolio-level synthetic market data would require combining it with joint residual dynamics.

Acknowledgments

The authors thank the eCornell artificial intelligence (AI) in Finance program participants for discussions that motivated the portfolio-level use of these paths, and Polygon.io for access to the 2014 to 2024 United States equity and exchange-traded-fund daily price histories used in the experiments. Computations were run on a single laptop; no shared-resource allocation was required.

Author contributions. A.A. designed the hybrid hidden Markov marginal generator, performed the single-asset experiments and validation analyses, and contributed to the manuscript. J.D.V. designed the variance-corrected single-index composition, performed the multi-asset experiments and downstream Value-at-Risk validation, released the code, and contributed to the manuscript.

Competing interests. The authors declare no competing interests.

Code and data availability. The experiment scripts, evaluation code (six-method comparison and Value-at-Risk coverage check), cached inputs and result summaries, and fitted per-asset marginals are released in the paper repository (<https://github.com/varnerlab/HMM-w-jumps-paper>); the underlying model-fitting library is released separately as JumpHMM.jl (<https://github.com/varnerlab/JumpHMM.jl>), pinned as a dependency of the experiment code.

Online Appendix

S1 Empirical cross-term diagnostic

The variance correction of Eq. (8) assumes $\text{Cov}(\tilde{g}_i^c, g_m) \approx 0$, allowing the variance derivation to omit the cross-term. This assumption does not follow automatically from the fitting procedure because each JumpHMM uses the asset’s full growth-rate history, including its market-correlated component. A remaining market loading in the generator draw would invalidate the closed-form scale. We tested the assumption by sampling $R = 100$ generator growth-rate paths $\tilde{g}_i^{(r)}$ per ticker from each generator and computing the normalized cross-covariance against the real return path for a broad-market exchange-traded fund (ticker SPY):

$$\text{cov}_i^{\text{norm}} = \frac{1}{R} \sum_{r=1}^R \frac{\text{Cov}((\tilde{g}_i^{(r)})^c, g_m)}{\sigma_{\text{gen},i} \sigma_m}. \quad (\text{S1})$$

Centering does not change covariance, so these values are numerically identical to those obtained from the raw draws. We plotted the per-ticker values as histograms (Fig. S1). Across the 423 tickers, the full-return JumpHMM generator (used by the hybrid and naive composition methods), the residual-fit JumpHMM generator (used by the JumpHMM-on-residuals composition method), and the Gaussian baseline had universe-mean normalized cross-covariances of 1.53×10^{-4} , 6.58×10^{-5} , and 2.26×10^{-4} , respectively. Their per-ticker values ranged from -0.0061 to 0.0061 . These values quantify the cross-term omitted in Eq. (8).




Figure S1: **Empirical normalized cross-covariance** $\text{Cov}(\tilde{g}_i^c, g_m)/(\sigma_{\text{gen},i} \sigma_m)$ **across the 423-ticker universe.** Histograms were stacked per generator, with each per-ticker value averaged over $R = 100$ generator growth-rate paths. Universe means were 1.53×10^{-4} , 6.58×10^{-5} , and 2.26×10^{-4} for the full-return JumpHMM, residual-fit JumpHMM, and Gaussian generators.

S2 Rebalancing under independent residuals

To determine what a portfolio-level use of these paths inherits from the independent-residual construction, this appendix derives the diversification return of a rebalanced portfolio under Eq. (4). The variance correction in Eq. (8) sets each synthetic asset’s second moment to its generator target. When that target matches the empirical second moment, the second-order formula for the diversification return is matched at the asset level as well. What the construction does not match is the cross-asset residual covariance, and the derivation below isolates where that omission enters.

The synthetic paths reproduce per-asset marginals and single-factor cross-sectional dependence. Each residual generator retains its own temporal structure, but the residual paths are generated

independently across assets at each date. The construction therefore omits cross-asset residual covariance. Consider a daily-rebalanced portfolio of N assets with constant target weights $w_i \geq 0$, $\sum_i w_i = 1$, reset to those weights at the start of each trading day. Throughout this derivation $g_i(t)$ is the annualized log-growth rate of Eq. (1), so the per-period log return on asset i is $g_i(t) \Delta t$ with $\Delta t = 1/252$ yr. The wealth process is:

$$W_t = W_{t-1} \sum_i w_i \exp(g_i(t) \Delta t), \quad (\text{S2})$$

so the per-step log return on the portfolio is $\log(W_t/W_{t-1}) = \log \sum_i w_i \exp(g_i(t) \Delta t)$. We suppress the time argument, writing g_i for $g_i(t)$. A second-order Taylor expansion of the exponential about zero is $\exp(g_i \Delta t) = 1 + g_i \Delta t + \frac{1}{2}(g_i \Delta t)^2 + O((g \Delta t)^3)$. Substituting it into each summand of $\sum_i w_i \exp(g_i \Delta t)$ gives:

$$\begin{aligned} \sum_i w_i \exp(g_i \Delta t) &= \sum_i w_i [1 + g_i \Delta t + \frac{1}{2}(g_i \Delta t)^2 + O((g \Delta t)^3)] \quad (\text{Taylor}) \\ &= \sum_i w_i + \Delta t \sum_i w_i g_i + \frac{1}{2}(\Delta t)^2 \sum_i w_i g_i^2 + O((g \Delta t)^3) \quad (\text{distribute}) \\ &= 1 + \Delta t \sum_i w_i g_i + \frac{1}{2}(\Delta t)^2 \sum_i w_i g_i^2 + O((g \Delta t)^3). \quad (\sum_i w_i = 1) \end{aligned} \quad (\text{S3})$$

A second-order Taylor expansion of $\log(1+x)$ about $x=0$ gives $\log(1+x) = x - \frac{1}{2}x^2 + O(x^3)$. Identifying $1+x$ with the right-hand side of Eq. (S3), the order- Δt correction is $x := \Delta t \sum_i w_i g_i + \frac{1}{2}(\Delta t)^2 \sum_i w_i g_i^2$, and squaring keeps only the leading $(\Delta t)^2$ term: $x^2 = (\Delta t)^2 (\sum_i w_i g_i)^2 + O((\Delta t)^3)$ because the cross term has an extra Δt factor. Substitution gives:

$$\begin{aligned} \log \frac{W_t}{W_{t-1}} &= \log(1 + x + O((g \Delta t)^3)) \\ &= x - \frac{1}{2}x^2 + O(x^3) \\ &= \Delta t \sum_i w_i g_i + \frac{1}{2}(\Delta t)^2 \sum_i w_i g_i^2 \\ &\quad - \frac{1}{2}(\Delta t)^2 \left(\sum_i w_i g_i \right)^2 + O((g \Delta t)^3) \\ &= \Delta t \sum_i w_i g_i + \frac{1}{2}(\Delta t)^2 \left[\sum_i w_i g_i^2 - \left(\sum_i w_i g_i \right)^2 \right] + O((g \Delta t)^3). \end{aligned} \quad (\text{S4})$$

Dividing through by Δt extracts the portfolio's annualized one-step log-growth rate $g_p(t) := \Delta t^{-1} \log(W_t/W_{t-1})$, the portfolio counterpart of the per-asset $g_i(t)$ (still suppressing the time argument):

$$g_p = \sum_i w_i g_i + \frac{1}{2} \Delta t \left(\sum_i w_i g_i^2 - \left(\sum_i w_i g_i \right)^2 \right) + O((\Delta t)^2). \quad (\text{S5})$$

To obtain expected annualized portfolio growth, we restored the time argument and took unconditional expectations of Eq. (S5) under stationarity. Define the per-asset annualized mean and variance:

$$\mu_i := \mathbb{E}(g_i(t)), \quad \sigma_i^2 := \text{Var}(g_i(t)), \quad (\text{S6})$$

the per-rate covariance $\Sigma_{ij} := \text{Cov}(g_i(t), g_j(t))$, and the per-rate portfolio variance $\sigma_p^2 := \text{Var}(\sum_i w_i g_i(t)) = w^\top \Sigma w$. The $O(\Delta t)$ correction in Eq. (S5) contains two squared quantities, $\sum_i w_i g_i(t)^2$ and $(\sum_i w_i g_i(t))^2$. Both expand by the second-moment identity, which holds for any random variable X with finite second moment:

$$\mathbb{E}(X^2) = \mathbb{E}(X)^2 + \text{Var}(X). \quad (\text{S7})$$

Applying this identity to $X = g_i(t)$ and summing against w_i gives:

$$\begin{aligned}\mathbb{E}\left(\sum_i w_i g_i(t)^2\right) &= \sum_i w_i \mathbb{E}(g_i(t)^2) && \text{(linearity)} \\ &= \sum_i w_i [\mu_i^2 + \sigma_i^2] && \text{(by Eq. (S7))} \\ &= \sum_i w_i \mu_i^2 + \sum_i w_i \sigma_i^2. && \text{(S8)}\end{aligned}$$

Applying the identity to $X = \sum_i w_i g_i(t)$, whose mean is $\sum_i w_i \mu_i$ and whose variance is σ_p^2 , gives:

$$\begin{aligned}\mathbb{E}\left(\left(\sum_i w_i g_i(t)\right)^2\right) &= \mathbb{E}\left(\sum_i w_i g_i(t)\right)^2 + \text{Var}\left(\sum_i w_i g_i(t)\right) && \text{(by Eq. (S7))} \\ &= \left(\sum_i w_i \mu_i\right)^2 + \sigma_p^2. && \text{(S9)}\end{aligned}$$

We substituted Eqs. (S8)–(S9) into the expectation of Eq. (S5) and grouped the variance and mean contributions as follows:

$$\begin{aligned}\mathbb{E}(g_p) &= \sum_i w_i \mu_i + \underbrace{\frac{1}{2} \Delta t \left(\sum_i w_i \sigma_i^2 - \sigma_p^2 \right)}_D \\ &\quad + \frac{1}{2} \Delta t \left[\sum_i w_i \mu_i^2 - \left(\sum_i w_i \mu_i \right)^2 \right] + O((\Delta t)^2),\end{aligned}\tag{S10}$$

the Booth-Fama identity for the daily-rebalanced portfolio in annualized growth-rate units [45, 46]. The middle term, D , is the diversification return: the $O(\Delta t)$ correction equal to half the difference between the weighted-average per-asset variance $\sum_i w_i \sigma_i^2$ and the portfolio variance σ_p^2 . The per-asset moments μ_i and σ_i^2 have units yr^{-1} and yr^{-2} respectively; the products $\Delta t \sigma_i^2$ and $\Delta t \sigma_p^2$ in Eq. (S10) are the annualized variances (yr^{-1}). The third term in Eq. (S10) is the cross-sectional dispersion of the per-asset annualized means, of order $\Delta t \mu^2 \sim 3 \times 10^{-5} \text{ yr}^{-1}$ at $\mu \sim 0.08 \text{ yr}^{-1}$ and negligible against the variance term $D \sim \Delta t \sigma^2 \sim 4 \times 10^{-2} \text{ yr}^{-1}$; we drop it. The realized annualized compound growth rate $\bar{g}_p = (T \Delta t)^{-1} \log(W_T/W_0) = T^{-1} \sum_t g_p(t)$ converges to $\mathbb{E}(g_p)$ by the ergodic theorem under any stationary residual process, so the long-run limit is fixed by the one-time-slice unconditional joint distribution of $(g_i(t))_i$ and does not, in itself, depend on the serial structure of the residual draw.

To rewrite D in a form that makes its non-negativity manifest, expand $\sigma_p^2 = \sum_i w_i^2 \sigma_i^2 + 2 \sum_{i < j} w_i w_j \Sigma_{ij}$ and use the identity $\sum_i w_i (1 - w_i) \sigma_i^2 = \sum_{i < j} w_i w_j (\sigma_i^2 + \sigma_j^2)$ that holds for any weights summing to one:

$$D = \frac{1}{2} \Delta t \left(\sum_i w_i \sigma_i^2 - \sigma_p^2 \right) = \frac{1}{2} \Delta t \sum_{i < j} w_i w_j \text{Var}(g_i - g_j) \geq 0, \tag{S11}$$

so D equals one-half the annualized weighted-average pairwise growth-rate-difference variance and is therefore non-negative, with equality only when every pair (i, j) comoves perfectly ($\text{Var}(g_i - g_j) = 0$) or all weight concentrates on a single asset. The rebalancing rule itself produces this effect: weights drift between rebalances, the rebalancer sells whatever has outperformed (its weight rose above target) and buys whatever has underperformed (its weight fell below target), and the trade systematically converts period-by-period cross-sectional dispersion into compounded growth. Markowitz [46] described this as the variance bonus of the long-run mean-variance allocation; Booth and Fama [45] called it the diversification return.

To separate market and residual contributions to the diversification return, we substituted the single-index decomposition $g_i(t) = \alpha_i + \beta_i g_m(t) + \varepsilon_i(t)$, with market variance $\sigma_m^2 := \text{Var}(g_m(t))$, per-asset residual variance $\sigma_{\varepsilon,i}^2 := \text{Var}(\varepsilon_i(t))$, and $\text{Cov}(g_m, \varepsilon_i) = 0$ (verified in Sec. S1), into Eq. (S11) gives:

$$\text{Var}(g_i - g_j) = (\beta_i - \beta_j)^2 \sigma_m^2 + \text{Var}(\varepsilon_i - \varepsilon_j), \tag{S12}$$

Because g_m and the residuals are uncorrelated, this relation splits the diversification return into market-beta-dispersion and idiosyncratic components:

$$D = \underbrace{\frac{1}{2} \Delta t \sigma_m^2 \sum_{i < j} w_i w_j (\beta_i - \beta_j)^2}_{D_{\text{mkt}}} + \underbrace{\frac{1}{2} \Delta t \sum_{i < j} w_i w_j \text{Var}(\varepsilon_i - \varepsilon_j)}_{D_\varepsilon}, \quad (\text{S13})$$

where the market-beta-dispersion component D_{mkt} equals $\frac{1}{2} \Delta t \sigma_m^2 [\sum_i w_i \beta_i^2 - (\sum_i w_i \beta_i)^2]$, the weighted cross-sectional β dispersion scaled by the annualized market variance $\Delta t \sigma_m^2$. Across the 423-ticker universe the empirical β distribution spans $[0.03, 1.79]$ with interquartile range $[0.81, 1.18]$ and the SPY annualized variance $\Delta t \sigma_m^2 \approx 0.04 \text{ yr}^{-1}$. The weighted variance of β is bounded by one quarter of its squared range, so D_{mkt} is at most 0.016 yr^{-1} , or 1.6 percentage points per year over this universe. For the idiosyncratic component D_ε , residuals were approximately orthogonal across assets at the same time slice. Therefore, $\text{Var}(\varepsilon_i - \varepsilon_j) \approx \sigma_{\varepsilon,i}^2 + \sigma_{\varepsilon,j}^2$, and D_ε scales with the weighted-average annualized idiosyncratic variance $\Delta t \sigma_{\varepsilon,i}^2$ per name.

The second-order value of D depends on same-time unconditional moments, not on serial dependence. Equation (8) sets $\sigma_{\text{gen},i}^2 = \beta_i^2 \sigma_m^2 + \sigma_{\varepsilon,i}^2$ to its empirical target. Any residual generator with the same unconditional second moments therefore gives the same D . This includes a stationary first-order autoregressive [AR(1)] residual generator $\varepsilon_i(t) = \phi \varepsilon_i(t-1) + \eta_i(t)$ with independent and identically distributed innovations η_i tuned so that $\text{Var}(\varepsilon_i) = \sigma_{\eta,i}^2 / (1 - \phi^2) = \sigma_{\varepsilon,i}^2$. The parameter ϕ leaves the marginal distribution at a single time slice unchanged. It therefore does not change $\mathbb{E}(g_p)$ in Eqs. (S10) and (S13), but it changes sampling variance. For a stationary AR(1) residual with variance $\sigma_{\varepsilon,i}^2$, the variance of its sample mean is approximately $T^{-1} \sigma_{\varepsilon,i}^2 (1 + \phi) / (1 - \phi)$. Under ergodicity, serial dependence changes per-path sampling noise, not the stationary mean. Serial dependence in the residual draw is therefore not a lever on the diversification return at the level of the second-order formula.

The direct structural difference is contemporaneous. The derivation assumes residuals are orthogonal across assets at each time slice, as they are in the independent per-asset generator. Real residuals share sector and style factors, which reduce $\text{Var}(\varepsilon_i - \varepsilon_j)$ below the independent-generator value $\sigma_{\varepsilon,i}^2 + \sigma_{\varepsilon,j}^2$. Matching each marginal variance does not restore this covariance. The synthetic D_ε will therefore exceed its real-market counterpart when the omitted covariance is positive. Quantifying that contribution requires a joint-residual experiment with the marginals and the allocation rule held fixed, which we did not run. Realized diversification returns on real equity portfolios are also attenuated relative to the second-order formula at all rebalancing horizons [7], so the synthetic excess is an upper bound on what an independent-residual generator would contribute in a live comparison.

S3 Validation metric definitions

To make the validation criteria reproducible, we defined the distributional and tail metrics reported in the main results. The two-sample Kolmogorov–Smirnov (KS) statistic between the empirical cumulative distribution functions (CDFs) of the observed and simulated growth-rate samples (G_{obs} of length n and G_{sim} of length m) is $D = \sup_x |F_n(x) - F_m(x)|$ [47, 48]; the test rejects when D exceeds the critical value at level $\alpha = 0.05$. The Anderson–Darling (AD) statistic places greater weight on the tails via the integral $A^2 = nm / (n + m) \int [F_n(x) - F_m(x)]^2 / [F(x)(1 - F(x))] dF(x)$, where F is the pooled empirical CDF [49]. The Wasserstein-1 distance between two empirical distributions of equal length is the average of sorted-quantile differences:

$$W_1 = \frac{1}{n} \sum_{i=1}^n |G_{\text{obs}}^{(i)} - G_{\text{sim}}^{(i)}|, \quad (\text{S14})$$

with $G^{(i)}$ the i th order statistic. The Hill upper-tail index is the classical order-statistic estimator:

$$\hat{\xi} = \frac{1}{k-1} \sum_{i=1}^{k-1} \log \left(\frac{X_{(i)}}{X_{(k)}} \right), \quad (\text{S15})$$

with $X_{(1)} \geq X_{(2)} \geq \dots$ the ordered upper-tail absolute returns; k is set to the upper 5% of the sample. For a Pareto-tailed distribution with shape α , $\hat{\xi} \approx 1/\alpha$. For the Hellinger calculation, we

binned both samples on a common grid and normalized the bin counts to probability masses p and q . Their distance is:

$$H(p, q) = \frac{1}{\sqrt{2}} \left\| \sqrt{p} - \sqrt{q} \right\|_2 \in [0, 1], \quad (\text{S16})$$

with $H = 0$ for identical probability masses and $H = 1$ for disjoint supports.

We abbreviate mean absolute error in the autocorrelation function as ACF-MAE.




Figure S2: **Statistical validation of the hidden Markov model with jumps (HMM-WJ)** ($N = 100$, $\alpha = 0.05$). Panels (a) and (d) condition on the $\sim 24\%$ of in-sample paths containing jump events. (a) In-sample KS p -values for 239 jump paths (pass rate 93.7%). (b) Out-of-sample KS p -values for all 1,000 paths (pass rate 95.4%). (c) Out-of-sample density fan chart; the observed density falls within the simulation envelope. (d) Observed and mean simulated in-sample autocorrelation function (ACF) of $|G_t|$ for jump paths.

Table S1: **Descriptive statistics and stylized-facts tests for SPY daily excess growth rates.** In-sample (IS): 2014–2024 ($T = 2,766$); out-of-sample (OoS): 2025 ($T = 249$). Gaussian and Laplace columns show maximum likelihood estimation (MLE) fits to the in-sample data. We used the Jarque–Bera (JB) normality test and the Ljung–Box (LB) autocorrelation test at lag 20. * denotes rejection at $\alpha = 0.05$.

Statistic	IS Observed	OoS Observed	Gaussian	Laplace
Mean (annualized, %)	6.31	11.60	6.31	14.19
Standard deviation of G_t (% yr ⁻¹)	214.50	220.62	214.46	205.93
Skewness	-0.753	-1.093	0	0
Excess Kurtosis	7.715	6.867	0	3
JB normality test	reject* ($p < 0.001$)	reject* ($p < 0.001$)	not used	not used
LB test on G_t (lag 20)	reject* ($p < 0.001$)	reject* ($p = 0.019$)	not used	not used
LB test on $ G_t $ (lag 20)	reject* ($p < 0.001$)	reject* ($p < 0.001$)	not used	not used

S4 Derivation of the single-asset generator

This appendix derives the probability law of the fitted single-asset generator and connects each fitted component to the empirical feature it controls. The hidden Markov model with jumps (HMM-WJ), its no-jump restriction (HMM-NJ), and their symbols follow the main-text definitions in Eq. (2) and Algorithm 1. Unless noted otherwise, numerical values evaluate the closed-form expressions at the fitted SPY model of Table 1: $N = 100$ states, Student- t degrees of freedom $\nu = 5$, jump probability $\epsilon = 10^{-4}$, mean episode duration $\lambda = 90$ steps, tail-selection probability $p_{\text{neg}} = 0.52$, tail-set size $N_{\text{tail}} = 5$, and the in-sample series of $T = 2,766$ daily excess growth rates.

S4.1 Generative law

The no-jump model generates a hidden state path $S_0, \dots, S_T$ on $\{1, \dots, N\}$ and observations $G_1, \dots, G_T$ with the joint density:

$$p(s_0, \dots, s_T, g_1, \dots, g_T) = \bar{\pi}_{s_0} \prod_{t=1}^T \mathbf{T}_{s_{t-1}, s_t} f_{s_t}(g_t), \quad (\text{S17})$$

where $\bar{\pi}_{s_0}$ is the initial probability of state s_0 , $\mathbf{T}_{s_{t-1}, s_t}$ is the probability of moving from state s_{t-1} to state s_t , and f_k is the emission density of state k . Under Eq. (2), f_k is the location-scale Student- t density:

$$f_k(g) = \frac{1}{\sigma_k} \psi_\nu\left(\frac{g - \mu_k}{\sigma_k}\right), \quad \psi_\nu(x) = \frac{\Gamma(\frac{\nu+1}{2})}{\sqrt{\nu\pi} \Gamma(\frac{\nu}{2})} \left(1 + \frac{x^2}{\nu}\right)^{-\frac{\nu+1}{2}}, \quad (\text{S18})$$

where ψ_ν is the density of a standard Student- t random variable with ν degrees of freedom, Γ is the gamma function, and (μ_k, σ_k) are the location and scale of state k . The factorization in Eq. (S17) states the structural property that drives everything that follows: given the state path, the observations are independent. Every form of serial dependence the generator can produce must therefore be carried by the state path [18].

The jump-augmented model enlarges the hidden state. Let $J_t \in \{0, 1, 2, \dots\}$ count the forced steps remaining after time t , and let u be the forced-step state distribution, which mixes uniform draws over the two tail sets:

$$u_k = \begin{cases} p_{\text{neg}}/N_{\text{tail}} & k \in \mathcal{S}_-, \\ (1 - p_{\text{neg}})/N_{\text{tail}} & k \in \mathcal{S}_+, \\ 0 & \text{otherwise.} \end{cases} \quad (\text{S19})$$

Writing $p_K(m) = e^{-\lambda} \lambda^m / m!$ for the Poisson probability of episode length m and $\mathbf{1}\{\cdot\}$ for the indicator function, the pair (S_t, J_t) evolves by the transition kernel:

$$\begin{aligned} & \Pr(S_t = k, J_t = j \mid S_{t-1} = s, J_{t-1} = i) \\ &= \begin{cases} u_k \mathbf{1}\{j = i - 1\} & i \geq 1, \\ (1 - \epsilon + \epsilon e^{-\lambda}) \mathbf{T}_{s,k} \mathbf{1}\{j = 0\} + \epsilon p_K(j + 1) u_k & i = 0, \end{cases} \end{aligned} \quad (\text{S20})$$

and the emission draw depends only on the first coordinate, $G_t \sim f_{S_t}$. The two cases read as follows. While an episode is active ($i \geq 1$), the next state is an independent draw from u and the counter decrements; the tail set and the state within it are re-drawn at every forced step. While no episode is active ($i = 0$), the chain either takes an ordinary transition (no trigger, or a trigger whose Poisson draw returns zero) or starts an episode of length $j + 1 \geq 1$ whose first forced step occurs immediately. The pair (S_t, J_t) is therefore itself a Markov chain, and the HMM-WJ generator is a hidden Markov model on the enlarged state space $\{1, \dots, N\} \times \{0, 1, 2, \dots\}$. Algorithm 1 implements Eq. (S20) with the episode counter truncated at the simulation horizon; it emits the stationary initial draw as the first observation and applies the kernel from the second step onward, which changes the law of a path only through the absent possibility of a trigger at the first step. When an episode ends, ordinary transitions resume from the last forced tail state, so the chain re-enters the body of the state space through the tail rows of $\mathbf{T}$.

S4.2 Estimators and the one-step empirical law

The fitting procedure consists of four closed-form estimators, and each one matches a specific empirical object exactly. The Laplace partition comes first. For observations $G_1, \dots, G_T$, the Laplace log-likelihood with location μ_L and scale b_L is:

$$\ell(\mu_L, b_L) = -T \ln(2b_L) - \frac{1}{b_L} \sum_{t=1}^T |G_t - \mu_L|. \quad (\text{S21})$$

For any fixed $b_L > 0$, maximizing ℓ over μ_L minimizes $\sum_t |G_t - \mu_L|$, so $\hat{\mu}_L$ is the sample median. Setting the b_L derivative to zero at $\hat{\mu}_L$ gives:

$$\begin{aligned} \frac{\partial \ell}{\partial b_L} &= -\frac{T}{b_L} + \frac{1}{b_L^2} \sum_{t=1}^T |G_t - \hat{\mu}_L| = 0 & (\text{first-order condition}) \\ \hat{b}_L &= \frac{1}{T} \sum_{t=1}^T |G_t - \hat{\mu}_L|, & (\text{solve for } b_L) \end{aligned} \quad (\text{S22})$$

the mean absolute deviation about the median. The state boundaries then invert the fitted Laplace cumulative distribution function (CDF) [36]. The Laplace CDF is piecewise exponential:

$$F_L(x) = \begin{cases} \frac{1}{2} \exp((x - \hat{\mu}_L)/\hat{b}_L) & x \leq \hat{\mu}_L, \\ 1 - \frac{1}{2} \exp(-(x - \hat{\mu}_L)/\hat{b}_L) & x > \hat{\mu}_L, \end{cases} \quad (\text{S23})$$

and setting $F_L(Q_k) = k/N$ and solving each branch for Q_k gives the closed-form boundaries:

$$\begin{aligned} \frac{k}{N} &= \frac{1}{2} e^{(Q_k - \hat{\mu}_L)/\hat{b}_L} \implies Q_k = \hat{\mu}_L + \hat{b}_L \ln \frac{2k}{N}, & (\text{lower branch}) \\ \frac{k}{N} &= 1 - \frac{1}{2} e^{-(Q_k - \hat{\mu}_L)/\hat{b}_L} \implies Q_k = \hat{\mu}_L - \hat{b}_L \ln \left(2 - \frac{2k}{N}\right). & (\text{upper branch}) \end{aligned} \quad (\text{S24})$$

The bins are equiprobable under the fitted Laplace, not under the empirical distribution, so the in-sample state counts are not uniform: across the 100 states, they range from 16 to 47 of the 2,766 observations. Every within-state mean μ_k and standard deviation σ_k was therefore computed from at least 16 observations, and no state required a fallback to the global moments.

The transition matrix is the maximum likelihood estimator (MLE) of a first-order Markov chain. Conditional on the assigned state sequence $s_1, \dots, s_T$, let n_{jk} count the transitions from state j to state k and let $n_{j\cdot} = \sum_k n_{jk}$ be the row total. The likelihood of the transition parameters factorizes over rows into multinomials, and maximizing the log-likelihood subject to each row summing to one, with multiplier c_j for row j , uses the Lagrangian:

$$\mathcal{L} = \sum_{j,k} n_{jk} \ln \mathbf{T}_{jk} + \sum_j c_j \left(1 - \sum_k \mathbf{T}_{jk}\right), \quad (\text{S25})$$

whose stationarity conditions solve in three steps:

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial \mathbf{T}_{jk}} &= \frac{n_{jk}}{\mathbf{T}_{jk}} - c_j = 0 & (\text{first-order condition}) \\ \mathbf{T}_{jk} &= \frac{n_{jk}}{c_j} & (\text{solve for } \mathbf{T}_{jk}) \\ c_j &= \sum_k n_{jk} = n_{j\cdot} & (\text{row constraint } \sum_k \mathbf{T}_{jk} = 1) \\ \hat{\mathbf{T}}_{jk} &= \frac{n_{jk}}{n_{j\cdot}}. & (\text{counted estimator}) \end{aligned} \quad (\text{S26})$$

The interior condition applies to cells with positive counts; for a cell with $n_{jk} = 0$ the log-likelihood is non-increasing in $\mathbf{T}_{jk}$, so its maximizer is the boundary value $\hat{\mathbf{T}}_{jk} = 0$, which the same formula returns. Multiplying $\hat{\mathbf{T}}_{jk}$ by the row frequency $n_{j\cdot}/(T-1)$ recovers $n_{jk}/(T-1)$, the empirical frequency of the state pair (j, k) . The counted estimator therefore reproduces the in-sample joint

law of (S_{t-1}, S_t) exactly; everything it asserts about longer horizons is the Chapman–Kolmogorov extension $\mathbf{T}^\tau$ of that one-step law [38, 41]. The emission estimator completes the pair: within each state, μ_k and σ_k are the sample mean and standard deviation of the assigned observations, and σ_k enters Eq. (S18) as the scale, so the emission standard deviation is $\sigma_k \sqrt{\nu/(\nu-2)}$. The aggregate consequence of this scale convention is quantified in Eq. (S30).

The initial distribution ties the pieces together. Let n_k count the occupancy of state k over the T observations, and let $\hat{\pi}_k = n_k/T$ be the occupancy frequency. The row total satisfies $n_{j\cdot} = n_j - \mathbf{1}\{s_T = j\}$, and the column total satisfies $n_{\cdot k} = n_k - \mathbf{1}\{s_1 = k\}$, because only the last observation opens no transition and only the first closes none. Substituting both identities gives:

$$\begin{aligned}
(\hat{\pi} \hat{\mathbf{T}})_k &= \sum_j \frac{n_j}{T} \cdot \frac{n_{jk}}{n_{j\cdot}} && \text{(definitions)} \\
&= \sum_j \frac{n_{j\cdot}}{T} \cdot \frac{n_{jk}}{n_{j\cdot}} + \sum_j \frac{\mathbf{1}\{s_T = j\}}{T} \cdot \frac{n_{jk}}{n_{j\cdot}} && (n_j = n_{j\cdot} + \mathbf{1}\{s_T = j\}) \\
&= \frac{n_{\cdot k}}{T} + \frac{\hat{\mathbf{T}}_{s_T, k}}{T} && \text{(sum transitions into } k\text{)} \\
&= \hat{\pi}_k + \frac{\hat{\mathbf{T}}_{s_T, k} - \mathbf{1}\{s_1 = k\}}{T}. && (n_{\cdot k} = n_k - \mathbf{1}\{s_1 = k\}) \quad (\text{S27})
\end{aligned}$$

The occupancy frequencies therefore satisfy the stationarity equation $\hat{\pi} \hat{\mathbf{T}} = \hat{\pi}$ up to a fixed-point residual bounded by $1/T$. For the fitted SPY model, the solved eigenvector $\hat{\pi}$ deviates from the occupancy frequencies by at most 3.7×10^{-4} , the same order as $1/T$, against occupancy values between 0.0058 and 0.017. The initial distribution is consequently not an abstract spectral object: $\hat{\pi}$ is, up to a correction of the order of $1/T$, the vector of in-sample state frequencies.

S4.3 Stationary distribution and moments

The stationary observation law of the no-jump generator follows directly from Eq. (S17). Marginalizing the state at a single time gives the mixture density and CDF:

$$p(g) = \sum_{k=1}^N \hat{\pi}_k f_k(g), \quad F_{\text{mix}}(x) = \sum_{k=1}^N \hat{\pi}_k \Psi_\nu \left(\frac{x - \mu_k}{\sigma_k} \right), \quad (\text{S28})$$

where Ψ_ν is the CDF of the standard Student- t with ν degrees of freedom. Every ingredient of Eq. (S28) is an empirical summary: the weights are the in-sample state frequencies, and the component locations and scales are the within-state sample moments. The generator's marginal is therefore a 100-component semiparametric estimate of the in-sample distribution, and reproducing the empirical marginal requires no further tuning. The mean is matched by construction. Weighting the within-state means by the occupancy frequencies telescopes into the grand mean:

$$\begin{aligned}
\sum_k \hat{\pi}_k \mu_k &= \sum_k \frac{n_k}{T} \cdot \frac{1}{n_k} \sum_{t: s_t=k} G_t && \text{(definitions of } \hat{\pi}_k, \mu_k\text{)} \\
&= \frac{1}{T} \sum_{t=1}^T G_t, && \text{(each } t \text{ counted once)} \quad (\text{S29})
\end{aligned}$$

and replacing $\hat{\pi}$ by $\bar{\pi}$ changes the value only through the $1/T$ correction of Eq. (S27): the model mean is 0.0636 yr^{-1} against the sample mean of 0.0631 yr^{-1} . The distributional match is equally direct. Evaluated at the fitted SPY model, the largest gap between the model CDF and the empirical CDF is $\sup_x |F_{\text{mix}}(x) - \hat{F}_T(x)| = 0.0045$, where $\hat{F}_T$ is the in-sample empirical CDF. The two-sample Kolmogorov–Smirnov (KS) critical value at significance level $\alpha = 0.05$ for two samples of 2,766 observations is $1.358 \sqrt{2/T} = 0.0365$, so the model-versus-data distance consumes about one eighth of the rejection threshold. Rejections of simulated paths are therefore driven by path-level sampling variation rather than by a misplaced marginal, which is consistent with the in-sample pass rates in Table S3.

The second moment makes the role of the emission scale convention explicit. Conditioning on the state and applying the law of total variance gives:

$$\begin{aligned}\text{Var}(G_t) &= \mathbb{E}[\text{Var}(G_t | S_t)] + \text{Var}(\mathbb{E}[G_t | S_t]) && \text{(law of total variance)} \\ &= \frac{\nu}{\nu-2} \sum_k \bar{\pi}_k \sigma_k^2 + \sum_k \bar{\pi}_k (\mu_k - \bar{\mu})^2, && \text{(Student-}t \text{ variance)}\end{aligned} \quad (\text{S30})$$

where $\bar{\mu} = \sum_k \bar{\pi}_k \mu_k$ is the stationary mean. Define the within-state variance $V_w = \sum_k \bar{\pi}_k \sigma_k^2$ and the between-state variance $V_b = \sum_k \bar{\pi}_k (\mu_k - \bar{\mu})^2$. The corresponding empirical decomposition of the sample variance is $V_b + V_w$, up to degrees-of-freedom corrections, so the model variance exceeds the sample variance by $2V_w/(\nu-2)$. At the SPY fit, $V_w = 0.17 \text{ yr}^{-2}$, only 3.6% of the total variance of 4.60 yr^{-2} : the quantile partition concentrates almost all dispersion between states rather than within them. The inflation is correspondingly small, a model variance of 4.72 against 4.60 yr^{-2} (1.3% in standard deviation, 2.17 against 2.14 yr^{-1}), and vanishes under Gaussian emissions (4.61 yr^{-2}).

The fourth moment explains the emission comparison in Table S3. Write $G_t = \mu_k + \sigma_k X$ within state k , where X is a standard Student- t draw, and let $\delta_k = \mu_k - \bar{\mu}$. Expanding the fourth central moment and dropping the odd-power terms, which vanish by the symmetry of ψ_ν , gives:

$$\begin{aligned}\mathbb{E}[(G_t - \bar{\mu})^4] &= \sum_k \bar{\pi}_k \mathbb{E}[(\delta_k + \sigma_k X)^4] && \text{(condition on the state)} \\ &= \sum_k \bar{\pi}_k [\delta_k^4 + 6 \delta_k^2 \sigma_k^2 \mathbb{E}(X^2) \\ &\quad + \sigma_k^4 \mathbb{E}(X^4)],\end{aligned} \quad (\text{S31})$$

with the standard Student- t moments, finite for $\nu > 4$:

$$\mathbb{E}(X^2) = \frac{\nu}{\nu-2}, \quad \mathbb{E}(X^4) = \frac{3\nu^2}{(\nu-2)(\nu-4)}. \quad (\text{S32})$$

At $\nu = 5$ these values are $5/3$ and 25 : the within-state kurtosis is 9, against 3 for Gaussian emissions. The excess kurtosis of the stationary mixture is $\mathbb{E}[(G_t - \bar{\mu})^4] / \text{Var}(G_t)^2 - 3$; evaluating it at the SPY fit gives 8.34 with Student- t emissions and 5.82 with Gaussian emissions, against an observed value of 7.72. The emission choice enters Eq. (S31) through both moments in Eq. (S32); the larger change by far is $\mathbb{E}(X^4)$ rising from 3 to 25, which acts on the σ_k^4 terms that the wide tail states dominate. The per-path simulated values reported in Table S3 (7.5 and 5.5) sit below these population values because the sample kurtosis of a heavy-tailed series of 2,766 observations is biased downward, but the population gap of 2.5 between the emission choices matches the simulated gap of 2.0 in both direction and size. The mixture also carries the asymmetry of the fitted state occupancies without an explicit skewness parameter: simulating 1,000 in-sample-length paths gives a mean sample skewness of -0.76 for HMM-NJ and -0.73 for HMM-WJ, against the observed -0.75 (Table S1). Because the states partition the Laplace quantiles of an asymmetric series, the occupancy-weighted mixture inherits that asymmetry through the μ_k and σ_k estimates.

S4.4 Dependence structure without jumps

The results stated so far concern single time slices; the dependence structure is where the no-jump generator falls short of the data, and the failure is structural rather than a matter of estimation. Within state k , the sojourn time is geometric: the chain remains for m steps with probability $\mathbf{T}_{kk}^{m-1}(1 - \mathbf{T}_{kk})$ and therefore stays for $1/(1 - \mathbf{T}_{kk})$ steps on average, between one and two days for every fitted state (Fig. S3c). The autocorrelation of absolute returns inherits this short memory. Let $a_k = \mathbb{E}[|G_t| | S_t = k]$ be the within-state mean absolute return and $\bar{a} = \sum_k \bar{\pi}_k a_k$ its stationary average. Because observations are conditionally independent given the states, the lag- τ product moment reduces to a state-pair sum:

$$\begin{aligned}\mathbb{E}[|G_t| |G_{t+\tau}|] &= \sum_{j,k} \Pr(S_t = j, S_{t+\tau} = k) a_j a_k && \text{(conditional independence)} \\ &= \sum_{j,k} \bar{\pi}_j (\mathbf{T}^\tau)_{jk} a_j a_k, && \text{(Chapman-Kolmogorov)}\end{aligned} \quad (\text{S33})$$

so the autocorrelation function (ACF) of $|G_t|$ is:

$$\text{ACF}_{|G|}(\tau) = \frac{\sum_{j,k} \bar{\pi}_j (\mathbf{T}^\tau)_{jk} a_j a_k - \bar{a}^2}{\text{Var}(|G_t|)}. \quad (\text{S34})$$

As τ grows, $\mathbf{T}^\tau$ converges to its rank-one limit, every row equal to $\bar{\pi}$, at the geometric rate $|\theta_2|^\tau$, where θ_2 is the second-largest eigenvalue of $\mathbf{T}$ in modulus; the ACF is therefore bounded by a constant multiple of $|\theta_2|^\tau$. For the fitted SPY matrix, $|\theta_2| = 0.253$, and $|\theta_2|^\tau$ falls below 0.01 from lag 4 onward. Evaluating Eq. (S33) gives a model ACF of 0.26 at lag 1, close to the observed 0.30 because the counted matrix pins the one-step state-pair law, but 0.021 at lag 3 and effectively zero from lag 5, while the observed ACF remains near 0.26 at lag 5 and 0.14 at lag 25. No first-order chain fitted by transition counting can do otherwise: matching the one-step law fixes $\mathbf{T}$, and the spectral gap of that matrix then forces geometric mixing within a week. This is the quantitative content of the main-text statement that the counted transition probabilities leave extreme-return states too quickly to reproduce volatility clustering, and it is the failure the jump mechanism corrects.

S4.5 Jump-episode arithmetic

The jump mechanism overlays the fast-mixing chain with rare, long episodes, and its operating characteristics follow from renewal arguments. On each step with no active episode, an episode with at least one forced step begins with probability $p_e = \epsilon(1 - e^{-\lambda})$, since a trigger whose Poisson draw returns zero is discarded; conditional on starting, the episode length is the zero-truncated Poisson draw with mean $\mathbb{E}[K \mid K \geq 1] = \lambda/(1 - e^{-\lambda})$. At $\lambda = 90$, the correction $e^{-\lambda}$ is below 10^{-39} , so $p_e = \epsilon$ and $\mathbb{E}[K \mid K \geq 1] = \lambda$ to machine precision. The long-run fraction of forced steps, denoted π_J , follows from the renewal-reward theorem. A renewal cycle consists of the free steps preceding an episode and the episode itself: the number of free trials up to and including the trigger is geometric with mean $1/p_e$, the triggering trial is itself the first forced step, and the episode contributes $\mathbb{E}[K \mid K \geq 1]$ forced steps in total:

$$\begin{aligned} \pi_J &= \frac{\mathbb{E}[K \mid K \geq 1]}{\frac{1 - p_e}{p_e} + \mathbb{E}[K \mid K \geq 1]} && \text{(forced steps over cycle length)} \\ &= \frac{p_e \mathbb{E}[K \mid K \geq 1]}{1 - p_e + p_e \mathbb{E}[K \mid K \geq 1]} && \text{(multiply through by } p_e) \\ &= \frac{\epsilon \lambda}{1 - \epsilon + \epsilon e^{-\lambda} + \epsilon \lambda} && (p_e \mathbb{E}[K \mid K \geq 1] = \epsilon \lambda) \\ &\approx \frac{\epsilon \lambda}{1 + \epsilon \lambda} = 0.0089. && \text{(drop } \epsilon e^{-\lambda}, \text{ then } \epsilon) \end{aligned} \quad (\text{S35})$$

Fewer than one step in a hundred is forced at the fitted values; the measured fraction across 400 simulated in-sample-length paths is 0.0088. The product $\epsilon \lambda$ is the quantity that sets this fraction, a point that matters for identification in Section S4.6.

At the path level, the same arithmetic determines how episodes are distributed across the ensemble. A simulated path contains no episode only if every one of its $T - 1$ transition steps fails to trigger one, so the jump-path probability at $T = 2,766$ is $1 - (1 - p_e)^{T-1} = 24.2\%$; the simulated ensemble gives 24.8% (99 of 400 paths), matching the $\sim 24\%$ jump-path fraction conditioned on in Fig. S2. Episodes begin at the long-run rate of one per expected renewal cycle, so a path carries about 0.274 episodes in expectation. The expected gap between episodes is $1/\epsilon = 10^4$ trading days, about 40 trading years, so the ensemble is a two-population mixture: roughly three quarters of the paths never trigger an episode and follow the HMM-NJ law exactly, while the remaining quarter carry a single episode (about one jump path in eight carries a second). Within a jump path, the forced fraction is $\lambda/T = 3.3\%$ for a single episode; the simulated mean is 3.6%, slightly higher through the multi-episode paths. Forced steps differ sharply from free ones. Under the forced-step distribution u of Eq. (S19), the conditional standard deviation is 5.38 yr^{-1} against the free-step 2.17 yr^{-1} , a variance ratio of 6.1, and the mean absolute return is $\bar{a}_J = \sum_k u_k a_k = 4.96 \text{ yr}^{-1}$ against $\bar{a} = 1.47 \text{ yr}^{-1}$. The stationary state marginal of the augmented chain is the mixture $(1 - \pi_J)$ times the free-phase distribution plus π_J times u ; the free phase relaxes from the post-episode tail state back to $\bar{\pi}$ at the rate $|\theta_2|$ within a few steps. Using $\bar{\pi}$ as the initial law therefore misallocates $O(\pi_J) \approx 1\%$ of probability mass, which quantifies the main-text remark that $\bar{\pi}$ is exact for the ordinary transition matrix but approximate for the jump-augmented process.

S4.6 Autocorrelation and kurtosis under jumps

The tuning objective of Eq. (3) scores the absolute-return ACF and the excess kurtosis, and both statistics admit closed-form approximations that show what (ϵ, λ) control. The ACF contribution rests on one more renewal quantity: the probability $h(\tau)$ that, given a step is forced, the step τ later belongs to the same episode. A uniformly chosen forced step lands in an episode with the length-biased law, an episode of length m covering m forced steps, and its position within that episode is uniform. Combining the two gives:

$$\begin{aligned} h(\tau) &= \sum_{m \geq 1} \frac{m \Pr(K = m \mid K \geq 1)}{\mathbb{E}[K \mid K \geq 1]} \cdot \frac{(m - \tau)_+}{m} \\ &= \frac{\mathbb{E}[(K - \tau)_+ \mid K \geq 1]}{\mathbb{E}[K \mid K \geq 1]} && \text{(sum over } m) \\ &\approx \left(1 - \frac{\tau}{\lambda}\right)_+, && (K \text{ concentrated near } \lambda) \end{aligned} \quad (\text{S36})$$

where $(x)_+ = \max(x, 0)$ and the approximation holds because the Poisson length concentrates within a few multiples of $\sqrt{\lambda} \approx 9.5$ around λ . The same-episode probability therefore decays almost linearly from one to zero over the horizon λ . The ACF follows by conditioning on the forced-step indicator $Z_t = \mathbf{1}\{\text{step } t \text{ is forced}\}$. Given the indicator path, every forced step's state is an independent draw from u , so two forced observations are conditionally independent; the dependence linking a forced observation to a later free one runs only through the episode's final state and, like the free-phase covariance itself, decays at the $|\theta_2|^\tau$ rate of Section S4.4. For lags beyond that scale, only the conditional means covary. Write $A_t = |G_t|$ for the absolute return. The conditional mean is $\mathbb{E}[A_t \mid Z_t] = \bar{a} + Z_t(\bar{a}_J - \bar{a})$, and for rare episodes the joint forced probability is $\Pr(Z_t = 1, Z_{t+\tau} = 1) \approx \pi_J h(\tau)$, so:

$$\begin{aligned} \text{Cov}(A_t, A_{t+\tau}) &\approx \text{Cov}(\mathbb{E}[A_t \mid Z_t], \mathbb{E}[A_{t+\tau} \mid Z_{t+\tau}]) && \text{(independence given } Z) \\ &= (\bar{a}_J - \bar{a})^2 \text{Cov}(Z_t, Z_{t+\tau}) && \text{(conditional mean in } Z) \\ &\approx (\bar{a}_J - \bar{a})^2 \pi_J h(\tau), && \text{(episodes rare)} \end{aligned} \quad (\text{S37})$$

so the jump mechanism adds an ACF component of height $\pi_J(\bar{a}_J - \bar{a})^2 / \text{Var}(A_t)$ that decays almost linearly to zero at lag λ , precisely the slow component the counted chain cannot produce. For a jump path carrying a single episode, the relevant forced fraction is λ/T rather than π_J , and evaluating the resulting expression at the fitted values, with $\text{Var}(A_t) = 3.01$ under the same conditioning, gives a predicted plateau of 0.13. Table S2 compares this prediction with the mean simulated jump-path ACF: the two differ by 0.013 at lag 5 and by less than 0.01 from lag 10 onward. At lags 1 to 3 the simulated values exceed the episode component (0.35 against 0.13 at lag 1) because the free-phase geometric component of Section S4.4 adds on top before it dies out.

The kurtosis contribution pulls in the opposite direction. Treating a jump path as a two-regime mixture with weights $(1 - \lambda/T, \lambda/T)$ and evaluating the fourth-moment expression of Eq. (S31) with the forced-step distribution u in place of $\bar{\pi}$ for the episode regime gives a population excess kurtosis of 7.20 for a single-episode jump path, against 8.34 with no jumps. Forced steps inflate the

Table S2: Episode-overlap prediction of the jump-path absolute-return autocorrelation function (ACF) against simulation. The prediction evaluates Eq. (S37) at the fitted SPY values with the single-episode forced fraction λ/T in place of π_J and the same-episode probability $h(\tau)$ of Eq. (S36). The simulated column is the mean ACF of $|G_t|$ over the 99 jump-active paths among 400 simulated in-sample-length paths. The free-phase component of Eq. (S33) is excluded from the prediction and has decayed below 0.01 at the lags shown.

Lag τ	$h(\tau)$	Predicted ACF	Simulated ACF
5	0.944	0.124	0.137
10	0.889	0.117	0.123
15	0.833	0.110	0.115
20	0.778	0.102	0.105
25	0.722	0.095	0.099

variance, by about 17% at the fitted values, faster than they inflate the fourth moment, because u draws uniformly within each tail set rather than extending the extreme tail. Adding jump content therefore lowers kurtosis toward and past the observed value while it raises the slow ACF component; the jump-share sweep reported in the main text traces the same tradeoff empirically, with the Hill tail index flat across the sweep because the tail exponent is set by ν , not by the jump content. The objective of Eq. (3) balances these two effects.

The same expressions show how the two jump parameters are identified, and where they are not. Within a single-episode jump path, the ACF component of Eq. (S37) has height proportional to λ/T and horizon λ , and the kurtosis shift depends on the same forced fraction: every statistic entering Eq. (3) for such a path is controlled by λ . The parameter ϵ moves the objective only through the incidence of jump paths, which the jump-active conditioning removes, and through the small fraction of multi-episode paths. This weak dependence is consistent with the tuned $\epsilon = 10^{-4}$ lying at the lower edge of the explored range for SPY and for all three cross-asset tickers, and with the Monte Carlo sensitivity of the tuned optima noted in Section S8. The objects entering every expression in this appendix, the fitted boundaries, transition matrix, and residence times, are plotted in Fig. S3.

S5 Transition matrix internals and partition diagnostics

To determine why the ordinary state process required a duration override, we inspected the fitted hidden Markov model with jumps (HMM-WJ) components for SPY at $N = 100$ (Fig. S3). The fitted Laplace CDF and the empirical CDF are shown with the 99 state boundaries Q_k . The transition matrix $\mathbf{T}$ has a near-diagonal band, with most off-diagonal mass within ± 5 states of the diagonal. Under the ordinary Markov transitions, every state has an expected residence time of only 1–2 steps. The fitted jump duration is $\lambda = 90$ steps, about two orders of magnitude longer. Every state had at least one in-sample observation. This residence-time gap, rather than missing states, motivated the duration override.




Figure S3: **Fitted hidden Markov model with jumps (HMM-WJ) internals for SPY with $N = 100$ states.** Panel (a) overlays the fitted Laplace cumulative distribution function (CDF) on the empirical CDF; the 99 vertical lines mark the equal-probability quantile boundaries Q_k that define the hidden states. Panel (b) displays the estimated transition matrix $\mathbf{T}$ in $\log_{10}$ color scale; the dominant near-diagonal band reflects short-range regime persistence. Panel (c) plots the expected natural residence time $1/(1 - T_{kk})$ for each state on a $\log_{10}$ scale, with the bottom tail set $\mathcal{S}_-$ (states 1–5, red) and the top tail set $\mathcal{S}_+$ (states 96–100, teal) highlighted. The dashed horizontal line marks the fitted mean jump duration $\lambda = 90$ steps, quantifying the two-order-of-magnitude gap between natural residence times and the jump-duration override.

S6 Emission distribution and semi-Markov comparison

To compare the emission and duration specifications, we evaluated three variants on SPY (Table S3): a hidden semi-Markov model (HSMM) baseline adapted from Bulla and Bulla [20] ($K = 8$ states, negative-binomial dwell times with a geometric fallback, and Student- t emissions), HMM-WJ with the original Gaussian emissions ($N = 100$), and HMM-WJ with Student- t emissions ($N = 100$). Both HMM-WJ variants had higher in-sample KS and AD pass rates than the HSMM-style baseline. The HMM-WJ variants use 100 states and the HSMM uses 8, so this comparison does not assign

the pass-rate difference solely to the duration model. Holding the rest of HMM-WJ fixed, changing the emission from Normal to Student- t ($\nu = 5$) increased simulated excess kurtosis from 5.5 to 7.5, against an observed value of 7.7. The Student- t emission also increased the KS and AD pass rates, while ACF-MAE and the distributional distances changed modestly. The emission choice therefore explained the kurtosis gain in this controlled HMM-WJ comparison, while it changed temporal fit little.

Table S3: **Three-way comparison of emission distributions and duration specifications for SPY.** Standard errors in parentheses; bold marks the best value per row. The HSMM is a hidden semi-Markov-style model adapted from Bulla and Bulla ($K = 8$ states, negative-binomial dwell times with a geometric fallback, and Student- t emissions) [20]; HMM-WJ: hidden Markov model with jumps ($N = 100$). All entries are computed from 1,000 simulated paths at significance level $\alpha = 0.05$. Kolmogorov–Smirnov (KS), Anderson–Darling (AD), and mean absolute error in the autocorrelation function (ACF-MAE) are defined in Section S3.

Metric	HSMM (Student- t)	HMM-WJ (Gaussian)	HMM-WJ (Student- t)
States	$K = 8$	$N = 100$	$N = 100$
<i>In-sample (2,766 trading days, 2014–2024)</i>			
KS pass rate (%)	82.0 (1.2)	97.0 (0.5)	98.3 (0.4)
AD pass rate (%)	42.5 (1.6)	90.5 (0.9)	94.8 (0.7)
Excess kurtosis	4.8 (0.08)	5.5 (0.03)	7.5 (0.09)
ACF-MAE	0.059 (<0.001)	0.052 (<0.001)	0.053 (<0.001)
Wasserstein-1	0.176 (0.001)	0.101 (0.002)	0.097 (0.001)
Hellinger	0.113 (<0.001)	0.076 (<0.001)	0.074 (<0.001)
<i>Out-of-sample (249 trading days, 2025)</i>			
KS pass rate (%)	96.2 (0.6)	95.0 (0.7)	95.4 (0.7)
AD pass rate (%)	96.7 (0.6)	95.9 (0.6)	95.3 (0.7)
Excess kurtosis	4.1 (0.13)	4.9 (0.08)	6.2 (0.15)
ACF-MAE	0.042 (<0.001)	0.040 (<0.001)	0.040 (<0.001)
Wasserstein-1	0.287 (0.002)	0.275 (0.005)	0.275 (0.005)
Hellinger	0.239 (0.001)	0.212 (0.001)	0.210 (0.001)

Table S4: **Hybrid composition method broken out by branch.** N_{tickers} : number of tickers assigned to each branch by the $R^2_{\text{preserve}} = 0.80$ threshold. The clipped variance-preserving branch does not trigger anywhere in the universe ($\rho_i < 0.90$ for all tickers) and is omitted. The two trackers in the R^2 -preserving branch (QQQ and SPYG) had median absolute errors of 0.005 for β and 0.001 for R^2 , with a KS pass rate of 100%.

Branch	N_{tickers}	$ \hat{\beta} - \beta $	$ \hat{R}^2 - R^2_{\text{real}} $	KS pass (%)	W_1	κ
hybrid	421	0.018	0.021	73.5	0.250	7.142
r2-preserve	2	0.005	0.001	100.0	0.099	11.525

Table S5: **Synthetic-tracker grid for the R^2 -preserving branch.** The observed universe places only two tickers on this branch, so we generated synthetic trackers at each $(\beta_{\text{true}}, R^2_{\text{true}})$ cell, composed them with Algorithm 2, and recovered $(\hat{\beta}, \hat{R}^2)$ by ordinary least squares. Entries are medians over 100 replications per cell with cross-replication standard deviations in parentheses; the Kolmogorov–Smirnov (KS) pass rate is the fraction of replications whose two-sample test against the target marginal exceeds $p = 0.05$. Every cell was assigned to the R^2 -preserving branch.

β_{true}	R^2_{true}	$\hat{\beta}$	$\hat{R}^2$	KS pass (%)
0.8	0.80	0.8006 (0.0072)	0.7998 (0.0044)	100.0
0.8	0.85	0.8006 (0.0059)	0.8497 (0.0041)	100.0
0.8	0.90	0.8004 (0.0048)	0.8999 (0.0028)	100.0
0.8	0.95	0.7995 (0.0033)	0.9500 (0.0013)	100.0
0.8	0.99	0.7999 (0.0016)	0.9901 (0.0003)	100.0
1.0	0.80	1.0003 (0.0089)	0.8017 (0.0052)	100.0
1.0	0.85	0.9994 (0.0084)	0.8497 (0.0039)	100.0
1.0	0.90	0.9999 (0.0065)	0.8999 (0.0026)	100.0
1.0	0.95	1.0005 (0.0036)	0.9502 (0.0013)	100.0
1.0	0.99	1.0000 (0.0018)	0.9900 (0.0003)	100.0
1.2	0.80	1.2007 (0.0126)	0.8001 (0.0048)	100.0
1.2	0.85	1.2011 (0.0105)	0.8503 (0.0043)	100.0
1.2	0.90	1.1996 (0.0081)	0.8994 (0.0027)	100.0
1.2	0.95	1.1995 (0.0052)	0.9498 (0.0014)	100.0
1.2	0.99	1.1998 (0.0025)	0.9900 (0.0003)	100.0

S7 State-resolution sensitivity

To test whether the results depended on state resolution, we swept the partition over $N \in \{30, 60, 90, 100, 150, 200, 350\}$ and re-evaluated the in-sample Kolmogorov–Smirnov (KS) and Anderson–Darling (AD) pass rates and the mean absolute error in the absolute-return autocorrelation function (ACF-MAE) for the hidden Markov model with no jumps (HMM-NJ) and the hidden Markov model with jumps (HMM-WJ), holding the tail-set size N_{tail} fixed (Table S6). HMM-NJ was insensitive to the partition: its pass rates moved by less than one percentage point and its ACF-MAE was unchanged at the reported precision, as expected for a model with no duration override whose behavior could depend on the width of a tail state. HMM-WJ traded the two metrics against each other. Because N_{tail} is fixed, a finer partition makes the tail set a narrower quantile band, so each forced step lands further out, and ACF-MAE improved from 0.057 at $N = 30$ to 0.045 at $N = 200$. The same narrowing left fewer observations behind each tail state’s emission parameters, and the AD pass rate, which weights tail discrepancies heavily, moved the other way, from 97.1% at $N = 30$ to 82.8% at $N = 200$. The intermediate values were not monotone, which is consistent with re-running the (ϵ, λ) grid search on a finite path ensemble at each N . At $N = 350$ several quantile bins were never visited in the historical record, leaving the corresponding emission parameters undefined. We adopted $N = 100$ for all subsequent experiments as a midpoint that held the KS pass rate above 98% and the AD pass rate within a few percentage points of its best value while capturing part of the ACF-MAE improvement.

Table S6: **Sensitivity of in-sample fit to the number of states N .** All entries are computed from 1,000 simulated in-sample-length paths for SPY at significance level $\alpha = 0.05$, with the tail-set size N_{tail} and the remaining hyperparameters of Table 1 held fixed and the (ϵ, λ) grid search re-run at each N for the hidden Markov model with jumps (HMM-WJ). Standard errors in parentheses: binomial for pass rates, bootstrap ($B = 500$) for the mean absolute error in the absolute-return autocorrelation function (ACF-MAE). HMM-NJ denotes the hidden Markov model with no jumps. Values at $N = 100$ reproduce the corresponding entries of Table 2.

Model	N	KS pass (%)	AD pass (%)	ACF-MAE
HMM-NJ	30	99.3 (0.3)	97.8 (0.5)	0.059 (<0.001)
HMM-NJ	60	99.5 (0.2)	98.5 (0.4)	0.059 (<0.001)
HMM-NJ	90	99.3 (0.3)	98.6 (0.4)	0.059 (<0.001)
HMM-NJ	100	99.3 (0.3)	98.2 (0.4)	0.059 (<0.001)
HMM-NJ	150	99.1 (0.3)	98.7 (0.4)	0.059 (<0.001)
HMM-NJ	200	99.2 (0.3)	98.3 (0.4)	0.059 (<0.001)
HMM-WJ	30	98.4 (0.4)	97.1 (0.5)	0.057 (<0.001)
HMM-WJ	60	96.7 (0.6)	91.4 (0.9)	0.053 (<0.001)
HMM-WJ	90	98.3 (0.4)	95.2 (0.7)	0.055 (<0.001)
HMM-WJ	100	98.3 (0.4)	94.8 (0.7)	0.053 (<0.001)
HMM-WJ	150	97.7 (0.5)	91.7 (0.9)	0.052 (<0.001)
HMM-WJ	200	96.0 (0.6)	82.8 (1.2)	0.045 (0.001)




Figure S4: **Branch assignment in calibrated $(\beta, R^2_{\text{real}})$ space.** The $R^2_{\text{preserve}} = 0.80$ threshold (dashed line) separates tracker assets from single stocks. In this universe, only QQQ and SPYG exceed the threshold, and both are assigned to the R^2 -preserving branch. The remaining 421 tickers populate the variance-preserving branch; the clipped variance-preserving branch is unused because the market loading ratio $\rho_i = \beta_i^2 \sigma_m^2 / \sigma_{\text{gen},i}^2$ stays below 0.90 for every ticker.

Table S7: **Aggregate metrics by β quartile, across composition methods.** Quartile cutoffs are computed from the calibrated β distribution across the 423-ticker universe. The hybrid KS pass rate declines from 87.6% in Q1 to 65.9% in Q4 as the market loading ratio ρ_i grows; the naive KS pass rate falls from 24.6% to 1.3% over the same range.

β bucket	Method	$ \hat{\beta} - \beta $	KS pass (%)	W_1	κ	$\text{Var}(g)$
Q1 (low β)	Naive	0.018	24.6	0.403	8.971	18.1
	Gaussian SIM	0.015	0.0	0.620	0.622	14.0
	Hybrid	0.016	87.6	0.177	8.369	14.6
	JumpHMM-on-residuals	0.015	73.2	0.204	7.059	14.3
	Block bootstrap	0.015	71.3	0.208	6.286	13.9
	GARCH(1,1)- t	0.015	57.8	0.253	5.255	13.4
Q2	Naive	0.021	1.9	0.621	7.510	25.1
	Gaussian SIM	0.016	0.2	0.653	1.509	18.0
	Hybrid	0.017	74.9	0.231	6.889	19.0
	JumpHMM-on-residuals	0.017	82.7	0.200	7.799	18.7
	Block bootstrap	0.016	79.8	0.197	7.213	18.0
	GARCH(1,1)- t	0.015	76.4	0.227	6.470	17.8
Q3	Naive	0.023	0.7	0.792	7.390	33.3
	Gaussian SIM	0.017	1.4	0.688	1.947	23.4
	Hybrid	0.018	66.1	0.279	6.904	24.4
	JumpHMM-on-residuals	0.018	78.3	0.242	7.454	24.0
	Block bootstrap	0.017	75.0	0.236	6.888	22.8
	GARCH(1,1)- t	0.016	71.0	0.263	6.265	21.9
Q4 (high β)	Naive	0.029	1.3	0.965	7.233	51.2
	Gaussian SIM	0.022	0.6	0.825	2.054	36.3
	Hybrid	0.022	65.9	0.339	6.739	37.1
	JumpHMM-on-residuals	0.022	69.1	0.313	6.842	37.0
	Block bootstrap	0.021	67.2	0.311	6.334	35.3
	GARCH(1,1)- t	0.021	64.5	0.361	6.064	34.7

Table S8: **Clipping-branch stress test under market-volatility scaling.** The market growth-rate path was rescaled by $\gamma \in \{1, 2, 3\}$ with the calibrated marginals held fixed, and the full hybrid composition method re-evaluated on the 423-ticker universe. As γ rises, the share of clipped tickers grows.

γ	Clipped (%)
1	0.0
2	76.4
3	95.2

S8 Cross-asset generalization

To test whether the single-asset procedure generalized beyond SPY, we repeated the fitting and evaluation on three stocks: NVDA (semiconductors, high volatility), JNJ (health care, low volatility), and JPM (financials, moderate volatility). Per-ticker tune outputs at ($n_{\text{paths}} = 200$, seed = 1234) were reproducible with these settings (Table S9). The objective combines ACF and kurtosis errors with fixed weights. Because kurtosis varies across finite samples of heavy-tailed paths, the selected grid point can change with the Monte Carlo sample. The reported (ϵ^* , λ^*) values are reproducible with the stated path count and seed, but they are not unique per-ticker optima. At the selected parameter values, in-sample HMM-WJ KS pass rates were 98.9% for NVDA, 99.4% for JNJ, and 99.1% for JPM. Out-of-sample KS pass rates on the 249-day 2025 hold-out were 97.7%, 74.0%, and 79.2%, respectively. Thus the strong in-sample fit transferred unevenly across assets, especially on the short holdout.

Table S9: **Cross-asset generalization of the hidden Markov model with jumps (HMM-WJ) across three single-stock tickers spanning sector and volatility regimes.** Per-ticker tune outputs (ϵ^*, λ^*) at $n_{\text{paths}} = 200$ and seed = 1234, and the corresponding HMM-WJ Kolmogorov–Smirnov (KS) and Anderson–Darling (AD) pass rates on 1,000 simulated paths at $\alpha = 0.05$ against each ticker’s in-sample (IS; 2014–2024) and out-of-sample (OoS; 2025) growth-rate history. Values in parentheses are binomial standard errors (SEs) for pass rates. The (ϵ^*, λ^*) values are reported as the parameter values used for the IS / OoS metrics rather than as canonical per-ticker optima.

Ticker	ϵ^*	λ^*	IS KS (%)	IS AD (%)	OoS KS (%)	OoS AD (%)
NVDA (semiconductors)	10^{-4}	70	98.9 (0.3)	96.7 (0.6)	97.7 (0.5)	96.1 (0.6)
JNJ (health care)	10^{-4}	70	99.4 (0.2)	96.3 (0.6)	74.0 (1.4)	61.3 (1.5)
JPM (financials)	10^{-4}	80	99.1 (0.3)	95.7 (0.6)	79.2 (1.3)	77.7 (1.3)




Figure S5: **Calibrated R^2 distribution across the 423 non-SPY tickers.** Histogram of ordinary least-squares (OLS) R^2 against SPY over the 2014–2024 window. The two tickers crossing the $R^2_{\text{preserve}} = 0.80$ threshold (dashed line) are index exchange-traded funds (QQQ, SPYG); the highest- R^2 individual stock (BLK, 0.645) falls ~ 0.16 below the threshold. The R^2 -preserving branch therefore contains only QQQ and SPYG in this universe.




Figure S6: **Heavy-tail preservation: excess kurtosis and Hill tail index vs calibrated β .** Each point is a per-ticker median over 100 replications. *Left:* excess kurtosis, y -axis clipped to $[-2, 20]$ for readability. The dashed line marks the median real-data excess kurtosis across the universe ($\kappa_{\text{real}} \approx 13$); the hybrid and naive composition methods (orange and blue) track the generator’s kurtosis, while the Gaussian single-index model (SIM) collapses to $\kappa \approx 1$. *Right:* Hill tail-index estimator on the upper 5% of $|g|$. The hybrid and naive medians remain near 0.37 throughout the β range; Gaussian SIM drops to ~ 0.22 . The hybrid’s median being below κ_{real} reflects the JumpHMM marginal’s own kurtosis rather than a composition-rule distortion: hybrid tracks naive (both retain the generator’s tails) while Gaussian SIM does not.

S9 SIM composition derivations and properties verification

To verify the two closed-form targets used by the variance-corrected single-index composition, we derived the variance scale of Eq. (8), the residual-variance target of Eq. (11), and the resulting large- T OLS coefficients and composed variance. To derive Eq. (8), first center the full-return draw as in Eq. (6), then set $\varepsilon_i = s_i \tilde{g}_i^c$ for $s_i \in [0, 1]$ and require that the composed series have the same variance as the generator path:

$$\text{Var}(g_i) = \text{Var}(\alpha_i + \beta_i g_m + \varepsilon_i) = \text{Var}(\beta_i g_m + \varepsilon_i) = \sigma_{\text{gen},i}^2, \quad (\text{S38})$$

where α_i drops out because it is constant. Expanding the sum gives:

$$\text{Var}(\beta_i g_m + \varepsilon_i) = \beta_i^2 \text{Var}(g_m) + 2\beta_i \text{Cov}(g_m, \varepsilon_i) + \text{Var}(\varepsilon_i). \quad (\text{S39})$$

Substituting $\varepsilon_i = s_i \tilde{g}_i^c$ gives $\text{Var}(\varepsilon_i) = s_i^2 \sigma_{\text{gen},i}^2$ and $\text{Cov}(g_m, \varepsilon_i) = s_i \text{Cov}(g_m, \tilde{g}_i^c)$. Under the assumption $\text{Cov}(\tilde{g}_i^c, g_m) \approx 0$, the cross term vanishes and the variance condition becomes $\beta_i^2 \sigma_m^2 + s_i^2 \sigma_{\text{gen},i}^2 = \sigma_{\text{gen},i}^2$. Solving for s_i^2 gives Eq. (8).

To derive Eq. (11), use the same zero-cross-covariance assumption and write the population R^2 identity for Eq. (4):

$$R^2 = \frac{\beta_i^2 \sigma_m^2}{\beta_i^2 \sigma_m^2 + \sigma_{\varepsilon,i}^2}. \quad (\text{S40})$$

Solving this identity for the residual variance gives Eq. (11). Because $\bar{\varepsilon}_i = 0$ after centering and scaling, the sample mean of a composed path obeys $\bar{g}_i = \alpha_i + \beta_i^{\text{eff}} \bar{g}_m$. Hence, the finite-sample OLS intercept was:

$$\hat{\alpha}_i = \bar{g}_i - \hat{\beta}_i \bar{g}_m = \alpha_i - (\hat{\beta}_i - \beta_i^{\text{eff}}) \bar{g}_m. \quad (\text{S41})$$

Under the zero-cross-covariance assumption, in the large- T limit OLS therefore gives $\hat{\alpha}_i \rightarrow \alpha_i$ and $\hat{\beta}_i \rightarrow \beta_i^{\text{eff}}$. Here $\beta_i^{\text{eff}} = \beta_i$ unless Eq. (10) clips the market loading. On the variance-preserving branch, the composed variance is $\sigma_{\text{gen},i}^2$. Without clipping, $\text{Var}(g_i) = \beta_i^2 \sigma_m^2 + s_i^2 \sigma_{\text{gen},i}^2 = \sigma_{\text{gen},i}^2$. Under clipping, Eq. (10) sets β_i^{eff} to give the same variance. On the R^2 -preserving branch, the composed variance is $\beta_i^2 \sigma_m^2 / R_{i,\text{real}}^2$, the value implied by the calibrated $(\beta_i, R_{i,\text{real}}^2)$ rather than by the generator marginal. On the ordinary-stock branch, the floor guarantees only that $\text{Var}(\varepsilon_i) \geq f \sigma_{\text{gen},i}^2$; it does not guarantee that adding the market term preserves the generator’s tail distribution. The KS pass rate fell from 87.6% in the lowest-beta quartile to 65.9% in the highest-beta quartile (Table S7).

S9.1 Role of the centered full-return draw as the residual

The hybrid composition places $\varepsilon_i = s_i \tilde{g}_i^c$, a centered and rescaled draw of asset i 's full return, in the residual slot of Eq. (4). On its face this is a category mismatch: the residual of a single-index decomposition is the idiosyncratic component of the return, orthogonal to the market, while $\tilde{g}_i^c$ is distributed like the centered total return, market-driven component included. This appendix states what the substitution preserves exactly, what it preserves approximately, and how the approximation degrades as the market loading grows.

The residual slot must supply three things: a zero mean, so that the intercept and market term alone set the conditional mean; the variance $\sigma_{\text{gen},i}^2 - \beta_i^2 \sigma_m^2$ not already contributed by the market term; and the asset-specific distributional shape, the heavy tails and volatility clustering that motivated the single-asset generator. Centering meets the first requirement exactly (Eq. (6)), the scale of Eq. (8) meets the second exactly, and the large- T limits $\hat{\alpha}_i \rightarrow \alpha_i$ and $\hat{\beta}_i \rightarrow \beta_i^{\text{eff}}$ hold because the draw is generated independently of the simulated market path (Sec. S1), so the recovered factor structure is exact as well. Any zero-mean, correctly scaled, market-independent draw satisfies the first two requirements; the full-return draw is chosen for the third, because it carries the per-asset marginal that the single-asset validation certified. Only the shape requirement is met approximately, and two exact statements bracket the approximation.

At $\rho_i = 0$ the substitution is not an approximation: Eq. (8) gives $s_i = 1$, the market term contributes no variance, and the composed series is the generator draw itself, so the composed distribution equals the validated single-asset marginal. For $\rho_i > 0$ the composed return is the sum of two independent centered draws, and the survival of the generator's shape follows in closed form. In population, conditional on the fitted parameters, write the deviation of the composed return from its mean as $X_m + X_i$, where $X_m = \beta_i g_m^c$ is the market contribution, g_m^c is the market path centered at its mean, and $X_i = s_i \tilde{g}_i^c$ is the scaled asset draw; their variances are $v_m = \rho_i \sigma_{\text{gen},i}^2$ and $v_i = (1 - \rho_i) \sigma_{\text{gen},i}^2$. Independence factors every cross term in the fourth moment:

$$\begin{aligned} \mathbb{E}[(X_m + X_i)^4] &= \mathbb{E}(X_m^4) + 4 \mathbb{E}(X_m^3) \mathbb{E}(X_i) \\ &\quad + 6 \mathbb{E}(X_m^2) \mathbb{E}(X_i^2) + 4 \mathbb{E}(X_m) \mathbb{E}(X_i^3) \\ &\quad + \mathbb{E}(X_i^4) \quad (\text{independence}) \\ &= \mathbb{E}(X_m^4) + 6 v_m v_i + \mathbb{E}(X_i^4). \quad (\mathbb{E}X_m = \mathbb{E}X_i = 0) \quad (\text{S42}) \end{aligned}$$

Writing κ_m and κ_{gen} for the excess kurtosis of the market path and of the generator draw, the component fourth moments are $\mathbb{E}(X_m^4) = (\kappa_m + 3) v_m^2$ and $\mathbb{E}(X_i^4) = (\kappa_{\text{gen}} + 3) v_i^2$, where rescaling by s_i leaves excess kurtosis unchanged. The excess kurtosis of the composed return follows by dividing Eq. (S42) by the squared total variance:

$$\begin{aligned} \kappa(X_m + X_i) &= \frac{(\kappa_m + 3) v_m^2 + 6 v_m v_i + (\kappa_{\text{gen}} + 3) v_i^2}{(v_m + v_i)^2} - 3 \\ &= \frac{\kappa_m v_m^2 + \kappa_{\text{gen}} v_i^2 + 3(v_m + v_i)^2}{(v_m + v_i)^2} - 3 \quad (\text{collect terms}) \\ &= \kappa_m \rho_i^2 + \kappa_{\text{gen}} (1 - \rho_i)^2. \quad (v_m + v_i = \sigma_{\text{gen},i}^2) \quad (\text{S43}) \end{aligned}$$

The generator's excess kurtosis survives composition with weight $(1 - \rho_i)^2$ and the market path's enters with weight ρ_i^2 ; a simulation check with independent generator draws matched Eq. (S43) to within 0.03 in mean excess kurtosis at $\rho_i \in \{0.1, 0.3, 0.5\}$. For a low-loading asset the composed tails are essentially the generator's: at $\rho_i = 0.10$ the generator's kurtosis retains weight 0.81. As ρ_i grows the composed shape moves away from the asset's own marginal, which is the pattern in the documented diagnostics: the hybrid tracks the generator's kurtosis across the β range while the Gaussian SIM does not (Fig. S6), and the hybrid KS pass rate declines from the lowest to the highest β quartile (Table S7).

The residual-fit methods provide the exact-orthogonal alternative: they fit a generator to each asset's OLS residual series, contain no market component to double count, and are compared with the hybrid in Table 3; fitting JumpHMM to residuals requires converting each residual series to a synthetic price path by cumulative compounding and a second fit per asset. In summary, $\tilde{g}_i^c$ is not a residual in the orthogonal-decomposition sense; it is a carrier of the validated marginal shape. The substitution fixes the mean, the variance, and the recovered factor loadings exactly, reproduces the asset's marginal

exactly in the $\rho_i \rightarrow 0$ limit, and dilutes the generator’s shape with weight $(1 - \rho_i)^2$ as the market share grows; the clipped and R^2 -preserving branches of Algorithm 2 handle the region where the dilution is largest.

References

- [1] Samuel A. Assefa, Danial Dervovic, Mahmoud Mahfouz, Robert E. Tillman, Prashant Reddy, and Manuela Veloso. Generating synthetic data in finance: Opportunities, challenges and pitfalls. In *Proceedings of the First ACM International Conference on AI in Finance*, pages 1–8, 2020. doi: 10.1145/3383455.3422554.
- [2] James Jordon, Lukasz Szpruch, Florimond Houssiau, Mirko Bottarelli, Giovanni Cherubin, Carsten Maple, Samuel N. Cohen, and Adrian Weller. Synthetic data: What, why and how? *arXiv preprint arXiv:2205.03257*, 2022.
- [3] Michael Stenger, André Bauer, Thomas Prantl, Robert Leppich, Nathaniel Hudson, Kyle Chard, Ian Foster, and Samuel Kounev. Thinking in categories: A survey on assessing the quality for time series synthesis. *Journal of Data and Information Quality*, 16(2):1–32, 2024. doi: 10.1145/3666006.
- [4] Benoit Mandelbrot. The variation of certain speculative prices. *The Journal of Business*, 36(4): 394–419, 1963.
- [5] R. Cont. Empirical properties of asset returns: Stylized facts and statistical issues. *Quantitative Finance*, 1(2):223–236, 2001. doi: 10.1080/713665670.
- [6] Paul Embrechts, Alexander J. McNeil, and Daniel Straumann. Correlation and dependence in risk management: Properties and pitfalls. In *Risk Management: Value at Risk and Beyond*, pages 176–223. Cambridge University Press, 2002. doi: 10.1017/CBO9780511615337.008.
- [7] Paul Bouchev, Vassilii Nemtchinov, and Ting-Kam Leonard Wong. Volatility harvesting in theory and practice. *Journal of Wealth Management*, 18(3):89–100, 2015. doi: 10.3905/jwm.2015.18.3.089.
- [8] Robert F. Engle. Autoregressive conditional heteroscedasticity with estimates of the variance of United Kingdom inflation. *Econometrica*, 50(4):987–1007, 1982.
- [9] Tim Bollerslev. Generalized autoregressive conditional heteroskedasticity. *Journal of Econometrics*, 31(3):307–327, 1986. doi: 10.1016/0304-4076(86)90063-1.
- [10] Steven L. Heston. A closed-form solution for options with stochastic volatility with applications to bond and currency options. *The Review of Financial Studies*, 6(2):327–343, 1993. doi: 10.1093/rfs/6.2.327.
- [11] Robert C. Merton. Option pricing when underlying stock returns are discontinuous. *Journal of Financial Economics*, 3(1-2):125–144, 1976. doi: 10.1016/0304-405X(76)90022-2.
- [12] S. G. Kou. A jump-diffusion model for option pricing. *Management Science*, 48(8):1086–1101, 2002. doi: 10.1287/mnsc.48.8.1086.166.
- [13] Magnus Wiese, Robert Knobloch, Ralf Korn, and Peter Kretschmer. Quant GANs: Deep generation of financial time series. *Quantitative Finance*, 20(9):1419–1440, 2020. doi: 10.1080/14697688.2020.1730426.
- [14] Hao Ni, Lukasz Szpruch, Marc Sabate-Vidales, Baoren Xiao, Magnus Wiese, and Shujian Liao. Sig-wasserstein GANs for time series generation. In *Proceedings of the Second ACM International Conference on AI in Finance*, pages 1–8. ACM, 2021. doi: 10.1145/3490354.3494393.
- [15] Shuntaro Takahashi, Yu Chen, and Kumiko Tanaka-Ishii. Modeling financial time-series with generative adversarial networks. *Physica A: Statistical Mechanics and Its Applications*, 527: 121261, 2019. doi: 10.1016/j.physa.2019.121261.

- [16] Jinsung Yoon, Daniel Jarrett, and Mihaela van der Schaar. Time-series generative adversarial networks. In *Advances in Neural Information Processing Systems*, volume 32, pages 5509–5519, 2019.
- [17] Sohyeon Kwon and Yongjae Lee. Can GANs learn the stylized facts of financial time series? In *Proceedings of the 5th ACM International Conference on AI in Finance*, pages 126–133, 2024. doi: 10.1145/3677052.3698661.
- [18] L. Rabiner and B. Juang. An introduction to hidden Markov models. *IEEE ASSP Magazine*, 3(1):4–16, 1986. doi: 10.1109/MASSP.1986.1165342.
- [19] Tobias Rydén, Timo Teräsvirta, and Stefan Åsbrink. Stylized facts of daily return series and the hidden Markov model. *Journal of Applied Econometrics*, 13(3):217–244, 1998. doi: 10.1002/(SICI)1099-1255(199805/06)13:3<217::AID-JAE476>3.0.CO;2-V.
- [20] Jan Bulla and Ingo Bulla. Stylized facts of financial time series and hidden semi-Markov models. *Computational Statistics & Data Analysis*, 51(4):2192–2209, 2006. doi: 10.1016/j.csda.2006.07.021.
- [21] José G. Dias, Jeroen K. Vermunt, and Sofia Ramos. Mixture hidden Markov models in finance research. In *Advances in Data Analysis, Data Handling and Business Intelligence*, pages 451–459. Springer, 2010. doi: 10.1007/978-3-642-01044-6_41.
- [22] Umberto Cherubini, Elisa Luciano, and Walter Vecchiato. *Copula Methods in Finance*. John Wiley & Sons, 2004. ISBN 9780470863459.
- [23] Kjersti Aas, Claudia Czado, Arnoldo Frigessi, and Henrik Bakken. Pair-copula constructions of multiple dependence. *Insurance: Mathematics and Economics*, 44(2):182–198, 2009.
- [24] William F. Sharpe. A simplified model for portfolio analysis. *Management Science*, 9(2):277–293, 1963. doi: 10.1287/mnsc.9.2.277.
- [25] Stephen A. Ross. The arbitrage theory of capital asset pricing. *Journal of Economic Theory*, 13(3):341–360, 1976. doi: 10.1016/0022-0531(76)90046-6.
- [26] Eugene F. Fama and Kenneth R. French. Common risk factors in the returns on stocks and bonds. *Journal of Financial Economics*, 33(1):3–56, 1993. doi: 10.1016/0304-405X(93)90023-5.
- [27] Elias M. Stein and Jeremy C. Stein. Stock price distributions with stochastic volatility: An analytic approach. *The Review of Financial Studies*, 4(4):727–752, 1991.
- [28] Sangjoon Kim, Neil Shephard, and Siddhartha Chib. Stochastic volatility: Likelihood inference and comparison with ARCH models. *The Review of Economic Studies*, 65(3):361–393, 1998.
- [29] James D. Hamilton and Raul Susmel. Autoregressive conditional heteroskedasticity and changes in regime. *Journal of Econometrics*, 64(1–2):307–333, 1994. doi: 10.1016/0304-4076(94)90067-1.
- [30] Franc Klaassen. Improving GARCH volatility forecasts with regime-switching GARCH. *Empirical Economics*, 27(2):363–394, 2002. doi: 10.1007/s001810100100.
- [31] James D. Hamilton. A new approach to the economic analysis of nonstationary time series and the business cycle. *Econometrica*, 57(2):357–384, 1989. doi: 10.2307/1912559.
- [32] Robert Engle. Dynamic conditional correlation: A simple class of multivariate generalized autoregressive conditional heteroskedasticity models. *Journal of Business & Economic Statistics*, 20(3):339–350, 2002. doi: 10.1198/073500102288618487.
- [33] Luc Bauwens, Sébastien Laurent, and Jeroen V. K. Rombouts. Multivariate GARCH models: A survey. *Journal of Applied Econometrics*, 21(1):79–109, 2006. doi: 10.1002/jae.842.
- [34] Andrew J. Patton. Modelling asymmetric exchange rate dependence. *International Economic Review*, 47(2):527–556, 2006. doi: 10.1111/j.1468-2354.2006.00387.x.

- [35] Matteo Rizzato, Julien Wallart, Christophe Geissler, Nicolas Morizet, and Noureddine Boumlaik. Generative adversarial networks applied to synthetic financial scenarios generation. *Physica A: Statistical Mechanics and its Applications*, 623:128899, 2023. doi: 10.1016/j.physa.2023.128899.
- [36] Samuel Kotz, Tomasz J. Kozubowski, and Krzysztof Podgórski. *The Laplace Distribution and Generalizations: A Revisit with Applications to Communications, Economics, Engineering, and Finance*. Birkhauser, 2001. ISBN 9780817641665.
- [37] David Toth and Bruce Jones. Against the norm: Modeling daily stock returns with the Laplace distribution. *arXiv preprint arXiv:1906.10325*, 2019.
- [38] Jeff A. Bilmes. A gentle tutorial of the EM algorithm and its application to parameter estimation for Gaussian mixture and hidden Markov models. Technical Report TR-97-021, International Computer Science Institute, 1998.
- [39] Paul Glasserman. *Monte Carlo Methods in Financial Engineering*. Springer, 2003. doi: 10.1007/978-0-387-21617-1.
- [40] James D. Hamilton. Regime switching models. *The New Palgrave Dictionary of Economics*, pages 11421–11426, 2018. doi: 10.1057/978-1-349-95189-5_2459.
- [41] Charles M. Grinstead and J. Laurie Snell. *Introduction to Probability*. American Mathematical Society, 1997.
- [42] Dimitris N. Politis and Joseph P. Romano. The stationary bootstrap. *Journal of the American Statistical Association*, 89(428):1303–1313, 1994. doi: 10.1080/01621459.1994.10476870.
- [43] Paul H. Kupiec. Techniques for verifying the accuracy of risk measurement models. *The Journal of Derivatives*, 3(2):73–84, 1995. doi: 10.3905/jod.1995.407942.
- [44] Kyunghyun Cho, Bart van Merriënboer, Caglar Gulcehre, Dzmitry Bahdanau, Fethi Bougares, Holger Schwenk, and Yoshua Bengio. Learning phrase representations using RNN encoder-decoder for statistical machine translation. In *Proceedings of the 2014 Conference on Empirical Methods in Natural Language Processing (EMNLP)*, pages 1724–1734. Association for Computational Linguistics, 2014. doi: 10.3115/v1/D14-1179.
- [45] David G. Booth and Eugene F. Fama. Diversification returns and asset contributions. *Financial Analysts Journal*, 48(3):26–32, 1992. doi: 10.2469/faj.v48.n3.26.
- [46] Harry M. Markowitz. Investment for the long run: New evidence for an old rule. *Journal of Finance*, 31(5):1273–1286, 1976. doi: 10.1111/j.1540-6261.1976.tb03213.x.
- [47] A. N. Kolmogorov. Sulla determinazione empirica di una legge di distribuzione. *Giornale dell’Istituto Italiano degli Attuari*, 4:83–91, 1933.
- [48] N. Smirnov. Table for estimating the goodness of fit of empirical distributions. *The Annals of Mathematical Statistics*, 19(2):279–281, 1948.
- [49] Theodore W. Anderson and Donald A. Darling. Asymptotic theory of certain “goodness of fit” criteria based on stochastic processes. *The Annals of Mathematical Statistics*, 23(2):193–212, 1952.