
Variance-Corrected Multi-Asset Equity Simulation with Hybrid Hidden Markov Marginals

Abdulrahman Alswaidan

Robert Frederick Smith School of Chemical and Biomolecular Engineering
Cornell University, Ithaca, NY 14850
aa2725@cornell.edu

Jeffrey D. Varner*

Robert Frederick Smith School of Chemical and Biomolecular Engineering
Cornell University, Ithaca, NY 14850
jdv27@cornell.edu

Abstract

Synthetic multi-asset equity data must reproduce each asset’s return distribution and its relationship with the market. A common approach adds a synthetic full-return path to a market factor as though the synthetic path were a residual. Because that path already contains the asset’s total variance, the construction adds market variance a second time. We derived a correction that centers and rescales each synthetic path before adding the market factor. We tested the method on 423 non-market assets in a 424-asset US equity and exchange-traded-fund universe. The correction preserved each generator’s target variance, recovered its calibrated market loading with low error, and retained heavy-tailed return distributions without fitting a second model to the regression residuals. On 416 complete asset histories held out from 2025, it increased the mean Kolmogorov–Smirnov pass rate from 83.5% to 87.0% and reduced the median ratio of synthetic to observed variance from 1.29 to 0.97. Its one-day left-tail 99% Value-at-Risk exceedance rate was similar to that of the best residual-fit method, although both exceeded the nominal rate. We generated the per-asset paths with a hidden Markov model whose jump-duration variant lengthened visits to extreme-return states. On a decade of daily SPY data, this variant improved the match to volatility clustering relative to the no-jump model while maintaining strong distributional fit. We selected it as a practical compromise, not because it outperformed every baseline. The multi-asset method still omits residual dependence between assets, which can overstate the benefit of daily rebalancing. Code and fitted per-asset models are released with the paper.

Keywords: hidden Markov model; jump-duration mechanism; single-index model; synthetic financial data; variance correction; Value-at-Risk coverage check; volatility clustering; heavy-tailed marginals.

*Corresponding author.

1 Introduction

Synthetic financial data often fail to reproduce the statistical properties of real returns. These paths are used to stress-test risk models against unobserved market scenarios, validate portfolio optimization algorithms, and evaluate allocation strategies across many simulated scenarios [1, 2]. Empirical equity excess growth rates exhibit three statistical regularities a generative model must reproduce: a heavy-tailed marginal distribution, weak autocorrelation of raw growth rates, and persistent volatility clustering, in which large changes are followed by large changes and small changes by small changes [3–5]. All three patterns are visible in SPY daily excess growth rates from 2014–2024 (Fig. 1). The empirical density is sharply concentrated near zero but places substantially more probability in the tails than a Gaussian model, as shown by the density and quantile-quantile comparisons (Fig. 1a and Fig. 1b). Raw growth rates have little serial correlation (Fig. 1c), whereas the positive, slowly decaying autocorrelation of their absolute values (Fig. 1d) shows that their magnitudes remain temporally dependent. These regularities are often called *stylized facts*. They show why matching the distribution of individual returns is not enough: raw returns should remain nearly uncorrelated, while the sizes of those returns should remain correlated over time.

Existing generators typically reproduce some of these regularities but not others. The generalized autoregressive conditional heteroskedasticity (GARCH) family [6, 7] captures volatility clustering but requires fat-tailed innovations to reproduce heavy tails. Stochastic-volatility and jump-diffusion models [8–10] produce heavy tails, but the jump component itself does not create persistent volatility. Deep generative models [11–13] can learn complex marginals but often miss the absolute-return autocorrelation associated with volatility clustering. Standard hidden Markov models (HMMs) [14] provide a regime-switching structure but return to central regimes too quickly to dwell in tail states long enough to reproduce the empirical clustering profile [15, 16]. We therefore sought a model that retained distributional fit, reduced absolute-return autocorrelation error, and could be fitted separately across a large asset universe.

We addressed the problem in two stages. First, we built a per-asset generator based on a hidden Markov model. It divided returns into Laplace-quantile states, used Student- t draws within each state, and estimated transitions by direct counting. Its jump-duration variant (HMM-WJ) added temporary episodes in the extreme-return states. On a decade of SPY data, HMM-WJ reduced the error in absolute-return autocorrelation from 0.059 to 0.053 relative to the no-jump model while retaining a 98.3% in-sample Kolmogorov–Smirnov pass rate. The GARCH model achieved a lower autocorrelation error of 0.031 but did not match the return distribution. We therefore used HMM-WJ as a practical per-asset generator, not as a uniformly superior model.

Our main contribution concerns the second stage: combining those per-asset paths with a shared market path. Treating a full synthetic return path as a single-index residual creates an unintended mean shift and counts market variance twice. We removed the path mean and derived a closed-form scale correction that preserved target variance and recovered the calibrated market loading across a 424-asset universe. With all 2014–2024 parameters held fixed, the correction increased the mean Kolmogorov–Smirnov pass rate from 83.5% to 87.0% on 416 complete asset histories from 2025 and reduced the median variance ratio from 1.29 to 0.97. The method does not, however, reproduce residual dependence between assets. Independent residuals created too much cross-sectional dispersion and could overstate the return from daily rebalancing. A location adjustment corrected the average return level but could not restore the missing dependence. This distinction between per-asset fit, variance preservation, and cross-asset dependence defines both the contribution and the limit of the method.




Figure 1: **Empirical stylized facts of SPY daily excess growth rates (2014–2024).** Panel (a) shows the heavy-tailed return density; the Laplace maximum-likelihood fit tracks the empirical peak and tails while the Gaussian fit does not. Panel (b) shows departure from the diagonal in a normal quantile-quantile plot at both extremes. Panel (c) shows the autocorrelation function of raw excess growth rates. Individual correlations are small, although the Ljung–Box test detects joint serial dependence. Panel (d) shows the autocorrelation function of absolute excess growth rates, which decays slowly across the reported lags.

2 Related Work

Models of equity-market stylized facts follow either a bottom-up or top-down approach. Bottom-up agent-based models [17–22] derive return dynamics from simulated interactions among heterogeneous agents. They are demanding to calibrate and expensive to simulate at scale. Top-down models instead fit statistical structures directly to observed growth-rate series. Top-down generators trace to Bachelier’s diffusion model, later formalized as geometric Brownian motion and used by Black-Scholes-Merton for option pricing [23–26]. These diffusion baselines are tractable but fail on the empirical regularities that drive risk and pricing. Mandelbrot [3] showed that returns have heavier tails than a Gaussian model allows. Fama [27] reviewed evidence that raw returns have near-zero autocorrelation. Schwert [28] documented persistence in monthly absolute residuals, a pattern later described by Cont [4] as volatility clustering. The autoregressive conditional heteroskedasticity (ARCH) and generalized ARCH (GARCH) models of Engle and Bollerslev [6, 7] reproduced clustering but required a heavy-tailed noise term to produce heavy-tailed marginals. Neither model includes discrete regimes or sudden jumps. Stochastic-volatility models [29, 8] make volatility a latent process to capture clustering, but estimating that process requires filtering or Markov chain Monte Carlo (MCMC) methods [30]; jump-diffusion models [9, 10] inject Poisson events into the price path, producing heavy tails by construction, but the jump component alone does not produce persistent volatility.

Hidden Markov models embed regime and jump behavior directly in a latent discrete chain. This approach follows Hamilton’s [31] two-state regime-switching model, later applied to volatility forecasting [32] and equity trading [33]. Ryden et al. [15] found that Gaussian-emission HMMs

matched heavy tails and the absence of return autocorrelation but did not reproduce volatility clustering. Bulla and Bulla [16] modeled clustering with hidden semi-Markov models (HSMMs) that replaced the geometric dwell-time distribution of every state with a parametric (negative-binomial) alternative. Likelihood-fitted HMM and HSMM formulations commonly use the Baum–Welch expectation-maximization (EM) algorithm [34], which is iterative and can be sensitive to initialization. Deep generative alternatives learn the marginal directly but can miss the volatility-clustering signature: Takahashi et al. [11] found that generative adversarial network (GAN) architectures failed to reproduce the absolute-return autocorrelation function, and Kwon and Lee [13] reported the same failure for TimeGAN [12]. Recent surveys [1, 2, 5] agree that stylized-facts reproduction remains the primary quality criterion. Our marginal component also used a latent-state chain, but it differed from the HSMM. It kept the empirically counted transition matrix for the central states and added an event-driven override that forces the chain into the tail states for a Poisson-distributed duration on a subset of time steps.

Building a multi-asset path requires a dependence model in addition to the separate single-asset generators. Mixture hidden Markov models [35] share latent state across assets but are costly to estimate in high dimensions. Copula approaches [36–38] separate marginals from cross-asset dependence but require choosing a copula family and, in their standard static form, treat successive time slices as independent. Factor models instead reduce a firm’s growth rate to one or more shared factors [39, 40]. The single-index model (SIM) of Sharpe [41] uses one shared factor and decomposes each asset’s growth rate into a market term and an idiosyncratic residual. This structure can use a single-factor market generator directly. We derived a closed-form correction that preserves the generator variance while recovering the calibrated market loading.

A statistical match does not guarantee that synthetic paths will be useful in a downstream financial task. We therefore combined goodness-of-fit tests for the return distribution [42–44] with autocorrelation diagnostics for temporal structure [4]. We also evaluated Value-at-Risk coverage and a daily-rebalanced allocation example. Booth and Fama [45] described the problem that arises in the latter setting: paths whose residuals are drawn independently across assets can create a *diversification return* that is muted in real markets, where common sector and style factors correlate residuals across assets. We measured Value-at-Risk performance with Kupiec’s unconditional-coverage test and used a location adjustment to measure, but not solve, the return-level effect of the missing residual dependence.

3 Methods

We fitted a marginal generator for each asset and then combined the generated paths with a variance-corrected single-index model. The dataset contained 424 United States-listed equities and exchange-traded funds observed from 2014 through 2024. The single-asset analysis focused on the SPDR S&P 500 ETF (SPY). It used 2,767 daily volume-weighted average prices for fitting, which yielded 2,766 growth rates, and the next 250 prices from 2025, which yielded 249 holdout growth rates. The multi-asset analysis used daily closing prices. Of the 423 non-market assets in the training universe, 416 had complete 2025 histories and entered the multi-asset holdout evaluation. We computed the excess growth rate $G_{i,j}$ for ticker i between trading days $j - 1$ and j from price $P_{i,j}$ as follows:

$$G_{i,j} \equiv \left(\frac{1}{\Delta t} \right) \ln \left(\frac{P_{i,j}}{P_{i,j-1}} \right) - r_f. \quad (1)$$

where $\Delta t = 1/252$ is the fraction of a trading year spanned by one trading day, and r_f is a constant continuously compounded risk-free rate. We used $r_f = 0.043$ for the SPY training period and $r_f = 0.0421$ for its 2025 holdout, based on Treasury STRIPS (Separate Trading of Registered Interest and Principal of Securities) bond yields. The excess growth rate has units of year^{-1} and is time-additive under continuous compounding.

3.1 Per-asset hidden Markov marginal generator

For each asset, HMM-WJ converted the continuous return series into a sequence of N ordered return regimes. At each time step, the model first selected a regime and then drew a return from that regime’s distribution (Fig. 2). We defined the model as follows: $\mathcal{M} = (\mathcal{S}, \mathcal{O}, \mathbf{T}, \mathbf{E}, \bar{\pi})$ [14]. The hidden state $S_t \in \mathcal{S} = \{1, \dots, N\}$ identifies the regime at time t , and the observed value is the excess growth rate

G_t from Eq. (1). We suppress the ticker index i when discussing one asset. The transition matrix $\mathbf{T}$ sets the probabilities of moving between regimes, $\mathbf{E}$ gives the return distribution within each regime, and $\bar{\pi}$ gives the initial regime probabilities.

We ordered the regimes from the most negative to the most positive returns. Their boundaries were equally spaced quantiles of a Laplace distribution fitted to the asset’s in-sample returns. Specifically, $Q_k = F_L^{-1}(k/N; \mu_L, b_L)$, where μ_L and b_L are the fitted location and scale. We set $Q_0 = -\infty$ and $Q_N = +\infty$ and assigned return G_t to state k when $Q_{k-1} < G_t \leq Q_k$. The Laplace distribution captured the sharp concentration of small price movements in Fig. 1a [46, 47]. Its closed-form quantiles also made the same partition inexpensive to compute across all 424 tickers [48].

The within-state emission followed a location-scale Student- t distribution:

$$G_t \mid S_t = k \sim \mu_k + \sigma_k \cdot t_\nu, \quad \nu = 5, \quad (2)$$

where t_ν denotes a standard Student- t random variable with ν degrees of freedom. For observations assigned to state k , μ_k is the sample mean and σ_k is the sample standard deviation. A standard t_ν has variance $\nu/(\nu - 2)$, so σ_k is a scale parameter rather than the emission standard deviation. At $\nu = 5$, that standard deviation is $\sigma_k \sqrt{5/3}$. We selected $\nu = 5$ from a sensitivity sweep ($\nu = \infty$ recovers the Normal baseline; explored range in Table 1). At $\nu = 5$, simulated kurtosis was 7.5 against the observed 7.7, compared with 5.5 under Normal emissions. The corresponding Student- t KS and AD pass rates were 98.3% and 94.8%. The full comparison against an HSMM-style baseline adapted from Bulla and Bulla [16] is given in Table S3 of Sec. S5.

With the regimes and within-regime distributions fixed, we estimated the transition matrix $\mathbf{T}$ by counting observed transitions. This avoided the initialization sensitivity of iterative expectation-maximization [48]. The counted matrix was strongly concentrated near its diagonal, so a state usually persisted for only 1–2 days, including in the tails (Fig. S5 in Online Appendix S4). As a result, the ordinary model left extreme-return states too quickly to reproduce the observed volatility clustering [16].

We addressed that problem with occasional jump episodes. Whenever no jump episode was active, a new episode began with probability ϵ . Its duration was $K \sim \text{Poisson}(\lambda)$, where λ is the mean episode length [49]. Because a Poisson draw can be zero, $K = 0$ caused no forced tail step. For $K > 0$, the model temporarily replaced ordinary transitions with draws from one of two tail sets: $\mathcal{S}_- = \{1, \dots, N_{\text{tail}}\}$ for the most negative states and $\mathcal{S}_+ = \{N - N_{\text{tail}} + 1, \dots, N\}$ for the most positive states. It selected the negative set with probability p_{neg} and the positive set otherwise. The model made this choice independently at each forced step, allowing negative extreme returns to occur slightly more often than positive ones. Table 1 gives all four jump settings.

We obtained the initial-state distribution $\bar{\pi}$ by solving $\bar{\pi}\mathbf{T} = \bar{\pi}$ with $\sum_k \bar{\pi}_k = 1$ [50, 51]. This distribution is exact for the ordinary transition matrix but only approximate for the jump-augmented process because it does not include the remaining jump duration. Algorithm 1 gives the full simulation procedure. Each step either followed $\mathbf{T}$ or continued a jump episode and then drew a return from the selected state’s emission distribution.

We chose the jump probability ϵ and mean duration λ so that simulated paths matched two features of the SPY history: volatility clustering and kurtosis [3, 4, 15]. The jump mechanism follows the Poisson event idea used in jump-diffusion models [9, 10], but here an event extends a visit to the tail states rather than adding an instantaneous price jump. We searched a grid of (ϵ, λ) values. For each jump-active path, the first part of the objective measured the squared difference between the observed and simulated autocorrelation functions of absolute returns through lag L . The second part measured the squared difference in excess kurtosis. We then averaged both errors over the jump-active paths. The calibration objective was:

$$J(\epsilon, \lambda) = \frac{1}{|\mathcal{R}_J|} \sum_{r \in \mathcal{R}_J} \left[\sum_{\tau=1}^L (\text{ACF}_{\text{obs}}(\tau) - \text{ACF}_r(\tau))^2 + w_\kappa (\kappa_{\text{obs}} - \kappa_r)^2 \right], \quad (3)$$

where $\mathcal{R}_J$ is the set of jump-active simulated paths, $\text{ACF}_{\text{obs}}(\tau)$ and κ_{obs} are the empirical absolute-growth autocorrelation at lag τ and excess kurtosis, and $\text{ACF}_r(\tau)$ and κ_r are the corresponding values for path r . We use κ for kurtosis to avoid collision with the jump-duration variable K . We simulated 200 independent paths of 2,766 trading days at each grid point. A path with no jump reduces to HMM-NJ and contains no information about (ϵ, λ) , so the objective included only paths




Figure 2: **Architecture of the hybrid hidden Markov model with a jump-duration mechanism (HMM-WJ).** When no jump episode is active, the chain transitions according to the empirical transition matrix $\mathbf{T}$ with probability $1 - \epsilon$ (thin bidirectional arrows) or enters a Poisson jump episode with probability ϵ (dashed upward arrow to the Poisson clock). During an episode, each of the $K \sim \text{Pois}(\lambda)$ forced steps independently draws the bottom tail set $\mathcal{S}_-$ (red, lowest-valued states) with probability p_{neg} or the top tail set $\mathcal{S}_+$ (blue, highest-valued states) otherwise, before normal Markovian evolution resumes. Small bell curves above each state depict the state-conditional location-scale Student- t ($\nu = 5$) emission distribution. Tail sets are defined as the N_{tail} lowest- and highest-quantile states under the Laplace quantile partition.

Algorithm 1 HMM-WJ single asset marginal generator.

Require: Fitted parameters $(\mathbf{T}, \mathbf{E}, \bar{\pi}, \epsilon, \lambda, p_{\text{neg}}, N_{\text{tail}})$; horizon T ; tail sets $\mathcal{S}_- = \{1, \dots, N_{\text{tail}}\}$ and $\mathcal{S}_+ = \{N - N_{\text{tail}} + 1, \dots, N\}$.
Ensure: State sequence $\{S_t\}_{t=0}^T$ and growth-rate path $\{G_t\}_{t=1}^T$.
1: Draw $S_0 \sim \bar{\pi}$; $k \leftarrow 0$ { k : remaining forced-step counter}
2: **for** $t = 1, \dots, T$ **do**
3: **if** $k > 0$ **then**
4: $\mathcal{J} \leftarrow \mathcal{S}_-$ w.p. p_{neg} , else $\mathcal{S}_+$; $S_t \sim \text{Uniform}(\mathcal{J})$; $k \leftarrow k - 1$ {forced tail step; sign drawn per step}
5: **else if** $\text{Uniform}(0, 1) > \epsilon$ **then** {standard Markov transition}
6: $S_t \sim \mathbf{T}_{S_{t-1}, \cdot}$
7: **else** {trigger jump; K is the episode length}
8: $K \sim \text{Poisson}(\lambda)$
9: **if** $K = 0$ **then** { $K = 0$: no forced steps, standard transition}
10: $S_t \sim \mathbf{T}_{S_{t-1}, \cdot}$
11: **else**
12: $\mathcal{J} \leftarrow \mathcal{S}_-$ w.p. p_{neg} , else $\mathcal{S}_+$; $S_t \sim \text{Uniform}(\mathcal{J})$; $k \leftarrow \min(K - 1, T - t)$ {first forced step; K steps total, sign per step}
13: **end if**
14: **end if**
15: Draw $G_t \sim \mu_{S_t} + \sigma_{S_t} t_5$ {Eq. (2)}
16: **end for**

with at least one jump [49]. The kurtosis weight w_κ balanced the kurtosis and ACF terms. The search ranges, w_κ , L , and the remaining HMM-WJ and SIM hyperparameters are reported in Table 1. The fitted model was evaluated against the marginal and temporal statistics reported in Section 4.

3.2 Variance-corrected multi-asset composition

The single-asset model describes each ticker separately. It does not say how several tickers should move together. For the multi-asset extension, we added one common source of movement: a market

path. We used the single-index model (SIM) of Sharpe [41] as follows:

$$g_i(t) = \alpha_i + \beta_i g_m(t) + \varepsilon_i(t), \quad t = 1, \dots, T. \quad (4)$$

Here α_i is the intercept for asset i , g_m is the market growth-rate path, β_i is the asset's market loading, and ε_i is its asset-specific residual path. The symbol ε_i denotes a residual path; it is unrelated to the jump probability ϵ used by the single-asset model. We estimated the intercept and market loading (α_i, β_i) from real data by ordinary least squares (OLS). The remaining modeling choice is ε_i . A standard SIM draws independent Gaussian residuals with the variance estimated by the real-data regression. This preserves the fitted market relationship but discards the heavy tails and volatility regimes produced by the single-asset generator. We called this baseline *Gaussian SIM*.

Why direct construction fails. A tempting alternative is to use a full synthetic return path $\tilde{g}_i$ from the single-asset generator as the residual:

$$g_i = \alpha_i + \beta_i g_m + \tilde{g}_i. \quad (5)$$

This raw substitution double counts location as well as variance. The calibrated intercept already fixes the conditional location of asset i , whereas the full-return draw generally has $\mathbb{E}[\tilde{g}_i] \neq 0$. Equation (5) therefore has conditional mean $\alpha_i + \mathbb{E}[\tilde{g}_i] + \beta_i g_m$, not $\alpha_i + \beta_i g_m$.

All full-return constructions in our comparison first removed the realized path mean. We defined the centered generator path as follows:

$$\bar{\tilde{g}}_i \equiv \frac{1}{T} \sum_{t=1}^T \tilde{g}_i(t), \quad \tilde{g}_i^c(t) \equiv \tilde{g}_i(t) - \bar{\tilde{g}}_i. \quad (6)$$

Therefore, $T^{-1} \sum_t \tilde{g}_i^c(t) = 0$ exactly. Subtracting a constant leaves both the sample variance and the sample covariance with g_m unchanged. We therefore used the centered direct construction $g_i = \alpha_i + \beta_i g_m + \tilde{g}_i^c$ as the *naive* variance comparator. Centering isolates the variance-composition question: the naive method has the correct residual location but no variance correction.

The path variance $\sigma_{\text{gen},i}^2 = \text{Var}(\tilde{g}_i) = \text{Var}(\tilde{g}_i^c)$ is the *generator variance*. It already represents the asset's total variance. Adding the market term therefore contributes variance a second time. Because $\tilde{g}_i$ was sampled independently of g_m , the naive composed variance was as follows:

$$\text{Var}(g_i) = \beta_i^2 \sigma_m^2 + \sigma_{\text{gen},i}^2, \quad \text{Cov}(\tilde{g}_i^c, g_m) \approx 0, \quad (7)$$

where $\sigma_m^2 = \text{Var}(g_m)$ is the variance of the market path. The excess is $\beta_i^2 \sigma_m^2$, so the error is largest for assets with high market exposure.

Variance-preserving correction. We treated the generator variance for asset i , $\sigma_{\text{gen},i}^2$, as the total variance budget for the composed asset. The market term uses part of that budget. We scaled the single-asset draw so that it uses only the part that remains. We defined s_i as the asset-specific residual scale factor, set $\varepsilon_i = s_i \tilde{g}_i^c$, and required $\text{Var}(\beta_i g_m + s_i \tilde{g}_i^c) = \sigma_{\text{gen},i}^2$. Solving for s_i gives the correction:

$$s_i^2 = 1 - \frac{\beta_i^2 \sigma_m^2}{\sigma_{\text{gen},i}^2} \quad (8)$$

We defined the *market loading ratio* as follows:

$$\rho_i \equiv \frac{\beta_i^2 \sigma_m^2}{\sigma_{\text{gen},i}^2}, \quad (9)$$

This ratio is the fraction of the variance budget used by the market. The residual receives the remaining fraction, so $s_i^2 = 1 - \rho_i$. This correction is impossible when $\rho_i \geq 1$ because the market term alone uses the entire variance budget. This case can occur when the market loading β_i is large, the per-asset generator has low variance, or the synthetic market is more volatile than the calibration market. In that case we reserved a minimum fraction f , the idiosyncratic variance floor, for the asset-specific draw and reduced the effective market loading β_i^{eff} . Equivalently, the maximum allowable market loading ratio is $\rho_i^{\text{max}} = 1 - f$. The resulting rule was:

$$\beta_i^{\text{eff}} = \begin{cases} \beta_i & \text{if } \rho_i \leq \rho_i^{\text{max}} \\ \text{sign}(\beta_i) \sqrt{(1-f) \frac{\sigma_{\text{gen},i}^2}{\sigma_m^2}} & \text{if } \rho_i > \rho_i^{\text{max}} \end{cases} \quad s_i^2 = \begin{cases} 1 - \rho_i & \text{if } \rho_i \leq \rho_i^{\text{max}} \\ f & \text{if } \rho_i > \rho_i^{\text{max}} \end{cases} \quad (10)$$

The sign of the calibrated market loading β_i is retained. We used $f = 0.10$, which guarantees that at least 10% of the variance budget remains asset-specific (Table 1).

High- R^2 index trackers. For an ordinary stock, our main target was total variance. For an index-tracking fund, preserving its relationship with the market was more important. The coefficient of determination R^2 measures the fraction of the asset’s variance explained by the market. For example, SPY regressed on itself has $\beta = 1$, $R^2 = 1$, and no residual variance. The QQQ and SPYG funds also have high R^2 against SPY. Applying the ordinary-stock rule to these trackers can preserve total variance while changing their market relationship. We therefore used a separate branch when the calibrated coefficient of determination, $R_{i,\text{real}}^2$, is at least the preservation threshold R_{preserve}^2 . In our universe, 0.80 separated the index trackers from individual stocks (Fig. S3). On this branch, we computed the target residual variance needed to recover the calibrated R^2 as follows:

$$\sigma_{\varepsilon,\text{target}}^2 = \beta_i^2 \sigma_m^2 \cdot \frac{1 - R_{i,\text{real}}^2}{R_{i,\text{real}}^2} \quad (11)$$

We rescaled the generator draw to this target as follows:

$$\varepsilon_i = \left(\sqrt{\frac{\sigma_{\varepsilon,\text{target}}^2}{\sigma_{\text{gen},i}^2}} \right) \tilde{g}_i^c. \quad (12)$$

When $R_{i,\text{real}}^2 = 1$, the target residual variance is zero, as it should be for SPY against itself. If the scale factor in Eq. (12) exceeds one, the procedure stretches the generator draw. We flagged this case because it can weaken the original marginal fit.

Complete composition rule. Algorithm 2 puts these steps together. For a high- R^2 tracker, it preserves the calibrated R^2 . For other assets, it preserves the generator variance and clips beta only when the market term would leave too little asset-specific variance. It then adds the scaled asset draw to the market term. An optional rank reorder can add dependence among the residual draws without changing their individual variances. Centering kept the path mean controlled by the intercept and market term. Under the zero-covariance assumption, an OLS regression on a long composed path therefore recovered the effective market loading and its corresponding intercept. The selected branch also fixed the target variance. These are guarantees about variance and factor loading, not about the complete return distribution. Online Appendix S8 gives the algebra. The construction adds only same-day market dependence; it does not reproduce lead-lag effects or market-driven volatility clustering.

Table 1: **HMM-WJ and SIM composition hyperparameters.** Values used for the main SPY analysis, with the sensitivity-sweep or grid-search span and the location of the corresponding check where one exists. The 423-ticker SIM composition universe holds $\epsilon = 0$ instead of the SPY value; per-ticker (ϵ^* , λ^*) for the NVDA, JNJ, and JPM cross-asset validation are given in Table S7.

Symbol	Description	SPY value	Explored range
<i>State partition and emission</i>			
N	Laplace-quantile states	100	$\{30, \dots, 350\}$, Online Appendix S6
ν	Student- t emission degrees of freedom	5	$\{3, \dots, 30, \infty\}$
<i>Jump-duration override</i>			
ϵ	jump probability per step	10^{-4}	$[10^{-4}, 2.5 \times 10^{-2}]$, Eq. (3)
λ	mean jump-episode duration (steps)	90	$[10, 160]$, Eq. (3)
p_{neg}	probability a jump targets $\mathcal{S}_-$	0.52	fixed
N_{tail}	quantile states per tail set	5	fixed (at $N = 100$)
w_κ	kurtosis-penalty weight, Eq. (3)	0.20	fixed
L	ACF lag window, Eq. (3)	25 days	fixed
<i>SIM composition</i>			
f	idiosyncratic variance floor	0.10	$\{0.05, \dots, 0.30\}$
R_{preserve}^2	branch-selection threshold	0.80	$\{0.70, \dots, 0.90\}$

Algorithm 2 Hybrid SIM composition with variance correction.

Require: Market growth-rate path $g_m \in \mathbb{R}^T$ with sample variance σ_m^2 ; per-asset generator growth-rate paths $\tilde{g}_i \in \mathbb{R}^T$ for $i = 1, \dots, N$; calibrated SIM parameters $(\alpha_i, \beta_i, R_{i,\text{real}}^2)$; idiosyncratic variance floor $f \in (0, 1)$; R^2 -preservation threshold $R_{\text{preserve}}^2 \in (0, 1)$.

Ensure: Composed paths $g_i \in \mathbb{R}^T$ and per-asset construction flags.

```
1: for each asset  $i = 1, \dots, N$  do
2:   Compute generator variance  $\sigma_{\text{gen},i}^2 \leftarrow \text{Var}(\tilde{g}_i)$ .
3:   Center the full-return draw:  $\tilde{g}_i^c \leftarrow \tilde{g}_i - T^{-1} \sum_{t=1}^T \tilde{g}_i(t)$ .
4:   if  $R_{i,\text{real}}^2 \geq R_{\text{preserve}}^2$  then
5:      $\sigma_{\varepsilon,\text{target}}^2 \leftarrow \beta_i^2 \sigma_m^2 (1 - R_{i,\text{real}}^2) / R_{i,\text{real}}^2$  {Eq. (11)}
6:      $\varepsilon_i \leftarrow \left( \sqrt{\sigma_{\varepsilon,\text{target}}^2 / \sigma_{\text{gen},i}^2} \right) \tilde{g}_i^c$ 
7:      $\beta_i^{\text{eff}} \leftarrow \beta_i$ ; branch  $\leftarrow R^2$ -preserving
8:   else
9:      $\rho_i \leftarrow \beta_i^2 \sigma_m^2 / \sigma_{\text{gen},i}^2$  {Eq. (9)}
10:    if  $\rho_i \leq 1 - f$  then
11:       $s_i^2 \leftarrow 1 - \rho_i$ ;  $\beta_i^{\text{eff}} \leftarrow \beta_i$ ; branch  $\leftarrow$  variance-preserving
12:    else
13:       $s_i^2 \leftarrow f$ ;  $\beta_i^{\text{eff}} \leftarrow \text{sign}(\beta_i) \sqrt{(1 - f) \sigma_{\text{gen},i}^2 / \sigma_m^2}$ ; branch  $\leftarrow$  clipped variance-preserving
14:    end if
15:     $\varepsilon_i \leftarrow s_i \tilde{g}_i^c$ 
16:  end if
17:  (Optional) apply cross-sectional rank reorder to  $\varepsilon_i$ .
18:   $g_i \leftarrow \alpha_i + \beta_i^{\text{eff}} g_m + \varepsilon_i$  {Eq. (4)}
19: end for
```

Evaluation protocol. We evaluated the method on 424 US equities and exchange-traded funds from Polygon.io. We selected tickers with a complete sequence of 2,767 daily closing prices from January 2014 through December 2024, which yielded 2,766 growth rates. This complete-case selection conditions the analysis on assets that survived and retained data coverage throughout the training window. We used SPY as the market and treated the other 423 tickers as assets to compose. For each asset, we fitted the same state and emission model used in the SPY analysis (Table 1; $r_f = 0$). To focus the experiment on multi-asset composition rather than per-ticker jump calibration, we set the jump probability to zero. We also regressed each real asset series on SPY to estimate the intercept α_i , market loading β_i , coefficient of determination $R_{i,\text{real}}^2$, and residual variance. Across the universe, β ranged from 0.03 to 1.79 and $R_{i,\text{real}}^2$ ranged from 0.001 to 0.933 (Fig. S3).

We compared six ways to construct the asset-specific term. The naive method used the full JumpHMM draw without correction. The Gaussian SIM method used independent Gaussian residuals with variance estimated by OLS. The hybrid method used Algorithm 2. The JumpHMM-on-residuals method fitted a new JumpHMM to the OLS residual series after converting the residuals to a synthetic price path by cumulative compounding. The block-bootstrap method sampled the OLS residual series with the Politis–Romano stationary bootstrap [52] and a mean block length of $\sqrt{T} \approx 50$. The GARCH(1,1)- t method fitted a conditional-variance model to each residual series [7]. We excluded thirty non-stationary fits from the GARCH results only. For each ticker, the naive and hybrid methods received the same JumpHMM draw. This pairing isolated the effect of the variance correction. The other four methods generated their own residual draws. We did not use the optional cross-sectional rank reorder because the experiment tested each ticker’s marginal construction separately.

We asked three questions about each composed series. First, did a new OLS regression recover the calibrated market loading and R^2 ? We measured this with the absolute errors $|\hat{\beta} - \beta|$ and $|\hat{R}^2 - R_{\text{real}}^2|$, where lower values are better. Second, did the synthetic and real returns have similar distributions? We used Kolmogorov–Smirnov (KS) and Anderson–Darling (AD) pass rates, Wasserstein-1 distance, excess kurtosis, and the Hill upper-tail index. Higher KS and AD pass rates and lower Wasserstein distance indicate a closer match; kurtosis and the Hill index describe tail behavior. Third, did the

composed path preserve generator variance? For this question, a composed-to-generator variance ratio of one is ideal. Online Appendix S3 gives the metric definitions. We ran 100 replications per ticker with seed 1234, producing 42,300 series per method. Pass rates used the significance level $\alpha = 0.05$.

We then froze all marginal fits, residual models, and SIM parameters estimated from 2014–2024 and evaluated the methods against the 249 growth rates observed in 2025. Of the 423 non-market training assets, 416 had complete holdout histories; GARCH results covered 386 because its thirty non-stationary training fits remained excluded. For each replication, we generated a 249-day SPY path from the training-period SPY marginal and used that common simulated market path in all asset compositions. Observed 2025 SPY and asset returns were used only for scoring. We repeated the KS, AD, Wasserstein-1, variance, kurtosis, and one-day left-tail VaR evaluations with 100 paths per ticker. Because goodness-of-fit tests have less power on 249 observations than on 2,766, we also compared each synthetic path with a randomly selected contiguous 249-day block from the training history. We first averaged replication-level results within ticker and then summarized across tickers so each asset received equal weight.

4 Results

4.1 Single-asset generator validation

We first tested whether the marginal generator in Fig. 2 reproduced the empirical SPY targets shown in Fig. 1. In both the in-sample (2014–2024, $T = 2,766$) and out-of-sample (2025, $T = 249$) windows, the real SPY series had heavy tails, and the Jarque-Bera test rejected normality. The series also showed persistent volatility clustering. Individual raw-return correlations were small, although the Ljung–Box test detected joint serial dependence (Table S1). We calibrated (ϵ, λ) for HMM-WJ against the in-sample SPY series using the grid search in Eq. (3) (Algorithm 1; search range in Table 1). The selected values were $(\epsilon, \lambda) = (10^{-4}, 90)$, although nearby grid points gave similar objective values. Under the counted transition matrix, ordinary state residence times were only 1–2 steps, whereas the selected jump duration had a mean of 90 steps (Fig. S5). The primary analysis used $N = 100$ states. Across $N = 60$ to 200, the KS and AD pass rates varied by less than one percentage point for both HMM-NJ and HMM-WJ. At $N = 350$, some states were never visited and their emission parameters could not be estimated (Online Appendix S6). The results were stable across a broad range of state resolutions, so we kept $N = 100$ for the generator comparisons that followed.

With the model calibrated, we compared HMM-WJ with seven alternative generators on 1,000 paths of 2,766 trading days each (Table 2; Fig. 3). No baseline was best on both the return distribution and volatility clustering. Bootstrap resampling passed every distribution test but, like all independent sampling methods, did not reproduce clustering. Gaussian sampling passed none of the distribution tests. The GARCH(1,1) model had the lowest mean absolute autocorrelation error (ACF-MAE) of 0.031 but rarely passed the distribution tests. The gated recurrent unit (GRU) [53] showed the same general pattern. The HSMM-style baseline adapted from Bulla and Bulla [16] reached an 82% KS pass rate but underestimated kurtosis by more than one third.

Among the fitted parametric and latent-state generators, the two HMM variants gave the strongest overall distributional fit. The HMM-NJ and HMM-WJ models had KS pass rates of 99.3% and 98.3%, respectively. The slightly lower HMM-WJ pass rates were consistent with jump episodes sometimes placing too much probability in the tails. Student- t emissions produced simulated kurtosis of 7.5, close to the observed 7.7, whereas Normal emissions underestimated it by about 29% (Table S3; Online Appendix S5). Both HMM variants also reproduced the observed skewness of -0.75 . That skewness arose from unequal time spent in negative and positive states rather than from an explicit skewness parameter.

The jump episodes improved temporal fit. The HMM-WJ model reduced ACF-MAE from 0.059 for HMM-NJ to 0.053. About 24% of HMM-WJ paths contained at least one jump and showed more persistent autocorrelation in absolute returns; paths without a jump were equivalent to HMM-NJ. On the held-out 249 days of 2025, HMM-NJ and HMM-WJ retained KS pass rates of 96.6% and 95.4%, with kurtosis close to the observed value (Table 2; Fig. S4). Tests on NVDA, JNJ, and JPM showed strong in-sample fit but more variable holdout performance (Table S7, Online Appendix S7).

Table 2: **In-sample and out-of-sample model comparison for SPY**. Results summarize 1,000 simulated paths. Values in parentheses are standard errors, and bold entries are best among the fitted generative models within each panel; the empirical bootstrap is excluded from this highlighting. Metric definitions and standard-error calculations are given in Online Appendix S3.

Model	KS (%)	AD (%)	κ	ACF-MAE	Cov. (%)	W_1	Hellinger
<i>In sample: 2014–2024 (2,766 days; observed $\kappa = 7.715$)</i>							
Bootstrap	100.0 (<0.1)	99.7 (0.2)	7.6 (0.08)	0.060 (<0.001)	100.0 (<0.1)	0.062 (0.001)	0.042 (<0.001)
Gaussian	0.0 (<0.1)	0.0 (<0.1)	−0.0 (<0.01)	0.060 (<0.001)	13.1 (0.1)	0.399 (0.001)	0.148 (<0.001)
Laplace	44.0 (1.6)	43.3 (1.6)	3.0 (0.02)	0.060 (<0.001)	37.4 (1.5)	0.138 (0.001)	0.072 (<0.001)
GARCH(1,1)	5.5 (0.7)	1.9 (0.4)	8.2 (0.37)	0.031 (0.001)	29.3 (1.4)	0.304 (0.006)	0.100 (0.001)
GRU	0.6 (0.2)	0.2 (0.1)	5.7 (0.16)	0.036 (<0.001)	17.2 (0.5)	0.421 (0.003)	0.134 (<0.001)
HSMM	82.0 (1.2)	42.5 (1.6)	4.8 (0.08)	0.059 (<0.001)	68.7 (1.1)	0.176 (0.001)	0.113 (<0.001)
HMM-NJ	99.3 (0.3)	98.2 (0.4)	8.1 (0.14)	0.059 (<0.001)	100.0 (<0.1)	0.081 (0.001)	0.072 (<0.001)
HMM-WJ	98.3 (0.4)	94.8 (0.7)	7.5 (0.09)	0.053 (<0.001)	100.0 (<0.1)	0.097 (0.001)	0.074 (<0.001)
<i>Out of sample: 2025 (249 days; observed $\kappa = 6.867$)</i>							
Bootstrap	97.4 (0.5)	98.5 (0.4)	6.0 (0.18)	0.043 (<0.001)	100.0 (<0.1)	0.232 (0.002)	0.205 (0.001)
Gaussian	62.2 (1.5)	46.3 (1.6)	−0.0 (<0.01)	0.043 (<0.001)	36.4 (1.9)	0.452 (0.002)	0.235 (0.001)
Laplace	88.0 (1.0)	92.9 (0.8)	2.7 (0.05)	0.043 (<0.001)	74.7 (1.0)	0.263 (0.002)	0.211 (0.001)
GARCH(1,1)	80.3 (1.3)	72.0 (1.4)	1.6 (0.07)	0.026 (<0.001)	96.0 (1.3)	0.507 (0.018)	0.232 (0.001)
GRU	71.8 (1.4)	44.8 (1.6)	2.9 (0.10)	0.033 (<0.001)	77.8 (2.9)	0.488 (0.005)	0.240 (0.001)
HSMM	96.2 (0.6)	96.7 (0.6)	4.1 (0.13)	0.042 (<0.001)	100.0 (0.2)	0.287 (0.002)	0.239 (0.001)
HMM-NJ	96.6 (0.6)	96.1 (0.6)	6.2 (0.15)	0.041 (<0.001)	100.0 (<0.1)	0.259 (0.003)	0.207 (0.001)
HMM-WJ	95.4 (0.7)	95.3 (0.7)	6.2 (0.15)	0.040 (<0.001)	100.0 (<0.1)	0.275 (0.005)	0.210 (0.001)

KS and AD report pass rates at $\alpha = 0.05$; coverage is the fraction of empirical quantiles inside the synthetic 90% envelope. κ is simulated excess kurtosis. ACF-MAE is computed for $|G_t|$ through lag 252. Larger values are preferred for KS, AD, and coverage; smaller values are preferred for ACF-MAE, W_1 , and Hellinger distance; κ is compared with the observed value.

These comparisons support HMM-WJ as a compromise between distributional and temporal fit. We examined that compromise directly by varying the share of jump-active paths.

To measure how jump-active paths changed the results, we resampled the fitted ensemble at different jump-path fractions f (Fig. 4). Here f is the share of accepted paths that contain at least one jump. We varied f from 0, where all accepted paths reduce to HMM-NJ, to 1, where every accepted path contains a jump. This procedure held the fitted model fixed and changed only the mixture of accepted paths. The absolute-return autocorrelation error fell monotonically from 0.059 at $f = 0$ to 0.037 at $f = 1$, moving from the no-jump value toward the GARCH value as the jump share rose. Over the same sweep, simulated excess kurtosis declined from 7.9 to 6.7, matching the observed value near $f \approx 0.14$ and falling below it thereafter. The two-sample KS and AD pass rates also declined. The Hill upper-tail index stayed near 0.35, against an observed 0.32, across the whole sweep because the




Figure 3: **Head-to-head in-sample model comparison for SPY** ($N = 100, 1,000$ simulated paths). *Panel (a)*: marginal density of excess growth rates with in-sample KS pass rates annotated per generator. *Panel (b)*: autocorrelation function of $|G_t|$ at lags 1–252 (shaded bands: 10th–90th percentile across paths) per generator. *Panel (c)*: tail Q-Q plot at the 0.1st–99.9th percentile region per generator.

Student- t ($\nu = 5$) emission sets the tail exponent independently of jump content. Increasing the jump share therefore traded fit to the distributional body for lower autocorrelation error without changing the tail index.

The value $f \approx 0.25$ used throughout kept simulated kurtosis within 2% of the observed value and lowered the autocorrelation error about a quarter of the way from the no-jump value toward the all-jump endpoint. The selected mixture therefore provided a compromise between marginal and temporal fit before the fitted marginals were carried into the multi-asset composition experiment.

4.2 Variance-corrected multi-asset composition

The single-asset results defined the tradeoff in the marginal generator. We then tested whether the variance-corrected SIM preserved each asset’s variance and factor loading across the 423-ticker universe, comparing the six composition methods on the aggregate scorecard (Table 3). Before comparing the methods, we measured the cross-term omitted from the variance correction. For the full-return generator, the universe-mean normalized cross-covariance with the market path was 1.53×10^{-4} , and the per-ticker values ranged from -0.0061 to 0.0061 (Fig. S1). These values supported the negligible-cross-term approximation used in Eq. (8).




Figure 4: **Jump-mix enrichment frontier for SPY** (in-sample, $N = 100$, pool of 4,000 simulated paths). Resampling the fitted HMM-WJ ensemble to a target jump-path fraction f (the share of accepted paths containing at least one jump) traces the temporal-marginal tradeoff. Both panels share the horizontal axis: absolute-return autocorrelation error (ACF-MAE of $|G_t|$ over lags 1–252); lower values indicate stronger volatility clustering. Panel (a): simulated excess kurtosis. Panel (b): Hill upper-tail index ξ , the raw estimator equal to the reciprocal of the tail exponent. Red dashed lines mark the observed SPY values; the star marks the $f \approx 0.25$ value used throughout. Endpoints $f = 0$ and $f = 1$ correspond to the no-jump generator and an all-jump ensemble.

Table 3: **Aggregate scorecard across 423 non-market tickers and 100 replications per ticker.** Errors, distances, and tail metrics are medians per method. Pass fractions for KS and AD are pooled across replications. $|\hat{\alpha} - \alpha|$, $|\hat{\beta} - \beta|$, and $|\hat{R}^2 - R_{\text{real}}^2|$: absolute deviation between OLS recovery and calibration (intercept error has units of yr^{-1}). KS pass and AD pass: fraction of replications whose two-sample test against the real marginal exceeds $p = 0.05$. W_1 : Wasserstein-1 distance to the real marginal. κ : sample excess kurtosis. Hill: tail-index estimate on the upper 5% of $|g|$.

Method	$ \hat{\alpha} - \alpha $	$ \hat{\beta} - \beta $	$ \hat{R}^2 - R_{\text{real}}^2 $	KS pass (%)	AD pass (%)	W_1	κ	Hill
Naive	0.002	0.022	0.087	7.1	3.5	0.700	7.767	0.369
Gaussian SIM	0.048	0.017	0.008	0.6	0.5	0.682	1.427	0.217
Hybrid	0.002	0.018	0.021	73.6	64.6	0.249	7.165	0.365
JumpHMM-on-residuals	0.049	0.018	0.015	75.8	70.7	0.232	7.262	0.365
Block bootstrap	0.042	0.017	0.020	73.3	66.8	0.232	6.675	0.330
GARCH(1,1)- t	0.047	0.017	0.036	67.4	61.4	0.267	6.069	0.317

All six methods recovered the factor loading with similar accuracy: the median absolute β error ranged from 0.017 to 0.022 (Table 3). Path centering also kept the median absolute intercept error at 0.002 yr^{-1} for both methods that used full-return draws. The residual-fit methods retained random variation in the residual mean over each finite path. Their larger intercept errors therefore do not, by themselves, imply a worse fit to the return distribution.

The larger differences appeared in variance recovery and tail behavior. Gaussian SIM had the lowest median R^2 error because it targets R^2 directly. However, it almost never passed the distributional tests, and its median excess kurtosis was an order of magnitude below the observed value (Fig. S2). Centered naive composition retained the shape of the full-return draw, but adding the market factor inflated its variance by $\beta_i^2 \sigma_m^2$. The hybrid method removed that extra variance. It passed 73.6% of KS comparisons and retained heavy tails, with median excess kurtosis of 7.2 and a Hill index of 0.36. Its median R^2 error fell between those of the naive and Gaussian methods because this branch preserves the generator variance rather than targeting the observed R^2 directly.

The three residual-fit methods avoided double counting by fitting their generators to the empirical OLS residuals, whose variance already excludes the market component. Their KS pass rates ranged from 67.4% for GARCH(1,1)- t to 75.8% for JumpHMM-on-residuals. The hybrid rate was two percentage points below JumpHMM-on-residuals and slightly above block bootstrap. JumpHMM-on-residuals and block bootstrap had slightly lower Wasserstein distances. JumpHMM-on-residuals also matched the hybrid’s kurtosis and Hill index almost exactly, whereas the block-bootstrap and GARCH residual fits produced lighter tails. The correction substantially improved on naive composition and approached the best residual-fit method without requiring a second fit for each asset. We next tested whether this ordering persisted after all fitted parameters were frozen.

We froze the fitted models and SIM parameters before evaluating them against 416 complete asset histories from 2025 (Table 4). The hybrid method reached an 87.0% mean cross-ticker KS pass rate, against 83.5% for centered naive composition, and its median composed-to-observed variance ratio was 0.97, whereas naive composition remained inflated at 1.29. JumpHMM-on-residuals reached an 86.2% KS pass rate and a slightly lower median W_1 than the hybrid. Matched-length training blocks produced lower pass rates for every method. The higher 2025 pass rates did not arise solely from comparing samples of 249 rather than 2,766 observations.

The average pass rates concealed substantial differences among tickers (Fig. 5). Most ticker-level rates were near the upper end of the range, but every method retained a lower tail of weak fits. For the hybrid method, many tickers improved relative to their matched-length training blocks, while a subset deteriorated out of sample. The held-out marginal improvement was broad but not uniform across assets, motivating a separate check of whether the resulting synthetic tails produced reliable risk thresholds.

Table 4: **Frozen-parameter multi-asset evaluation on the 2025 holdout.** Results cover 416 non-market tickers and 100 replications per ticker; GARCH covers 386 tickers. Pass rates are first averaged within ticker and then across tickers. *matched IS* compares each synthetic path with a randomly selected contiguous 249-day block from the 2014–2024 training history, matching the holdout sample length. W_1 and the variance ratio are medians across ticker-level medians; the variance ratio is synthetic variance divided by observed 2025 variance.

Method	KS pass (%)		AD pass (%)		W_1 (2025)	variance ratio
	matched IS	2025	matched IS	2025		
Naive	69.4	83.5	54.8	71.6	0.877	1.29
Gaussian SIM	57.4	73.9	44.9	64.6	0.866	0.97
Hybrid	78.8	87.0	66.2	79.7	0.716	0.97
JumpHMM-on-residuals	77.9	86.2	65.6	80.1	0.708	0.98
Block bootstrap	76.3	84.0	64.0	76.5	0.715	0.90
GARCH(1,1)- t	74.5	82.6	62.6	75.0	0.733	0.91




Figure 5: **Cross-ticker distribution of frozen-parameter out-of-sample marginal fit.** Panel (a) shows the distribution of per-ticker 2025 KS pass rates across 100 replications; the dashed line marks 95%. Panel (b) compares each ticker’s hybrid KS pass rate against a matched-length training block with its pass rate against the 2025 holdout. The diagonal denotes equal performance.

4.3 VaR coverage

We used the frozen synthetic paths to estimate per-ticker one-day left-tail Value-at-Risk (VaR) thresholds and counted exceedances on the real 2025 histories (Table 5). At $\alpha = 0.95$, the hybrid and JumpHMM-on-residuals methods produced mean exceedance rates of 5.97% and 6.07%, respectively, against the 5% nominal rate. At $\alpha = 0.99$, both rates were 1.67%, against the 1% nominal rate. Naive composition was closest at the 99% level (1.21%), while Gaussian SIM had the largest exceedance rate (2.03%). The short holdout contains only about 2–3 expected 99% exceedances per ticker, so we interpret the corresponding Kupiec pass rates and cross-ticker dispersion as low-power checks rather than precise rankings. Taken together, the VaR results showed that stronger marginal fit did not automatically imply nominal tail coverage, motivating a closer look at where the variance correction was most stable.

Table 5: **Out-of-sample per-ticker one-day left-tail Value-at-Risk coverage across the six composition methods.** VaR thresholds at coverage level α were estimated from frozen-parameter synthetic paths and exceedances were counted against the real 2025 growth-rate history. *rate*: mean of the per-ticker mean exceedance rates (nominal: $1 - \alpha$). *SD*: cross-ticker standard deviation, in percentage points. *Kupiec pass*: mean within-ticker fraction of replications whose unconditional-coverage test *p*-value exceeds 0.05.

Method	$\alpha = 0.95$			$\alpha = 0.99$		
	rate (%)	SD (pp)	Kupiec pass (%)	rate (%)	SD (pp)	Kupiec pass (%)
Naive	4.37	2.15	65.8	1.21	0.76	78.9
Gaussian SIM	4.95	2.26	70.0	2.03	1.17	72.1
Hybrid	5.97	2.43	66.3	1.67	0.93	73.4
JumpHMM-on-residuals	6.07	2.59	67.0	1.67	0.95	74.5
Block bootstrap	6.13	2.56	63.7	1.84	1.01	73.4
GARCH(1,1)- <i>t</i>	5.79	2.47	62.4	1.80	1.02	72.0

4.4 Sensitivity analyses

To isolate how the variance correction changed marginal fit, we plotted each ticker’s KS pass rate and variance ratio $\text{Var}(g)/\sigma_{\text{gen}}^2$ against its calibrated β (Fig. 6). Under naive composition, the variance ratio followed $1 + \rho_i$ and exceeded generator variance by 50–75% at high β , and the naive KS pass rate fell toward zero above $\beta \approx 0.7$; the hybrid method held the ratio at one and retained substantially higher pass rates across the β range.

The branch-selection rule assigned 421 tickers to the variance-preserving branch, two trackers to the R^2 -preserving branch, and none to the clipped branch (Figs. S3 and S6; Table S4). The two R^2 -preserving assignments were QQQ and SPYG, and both had a 100% KS pass rate; the highest- R^2 individual stock, BLK, fell about 0.16 below the $R^2_{\text{preserve}} = 0.80$ threshold. Clipping did not occur in the universe: $\rho_i = \beta_i^2 \sigma_m^2 / \sigma_{\text{gen},i}^2$ (Eq. (9)) stayed below $1 - f = 0.90$ for every ticker. Because the R^2 -preserving branch contained only two tickers, we tested it on a synthetic-tracker grid spanning $(\beta_{\text{true}}, R^2_{\text{true}}) \in \{0.8, 1.0, 1.2\} \times \{0.80, 0.85, 0.90, 0.95, 0.99\}$. Each cell had a 100% KS pass rate across replications. The correction removed the observed beta-related variance inflation. We then tested the clipping branch under conditions that would force it to activate.

Bucketing the universe by β quartile, the hybrid KS pass rate fell from 87.6% in the lowest quartile to 65.9% in the highest (Table S5). As β increased, the market term accounted for more of the composed




Figure 6: **Hybrid’s variance correction and its KS-pass-rate consequence, per ticker vs calibrated β .** Each point is a per-ticker median over 100 replications; thick lines are binned medians. *Left (consequence)*: Kolmogorov–Smirnov pass rate against the real marginal at $\alpha = 0.05$ for naive composition (blue) and hybrid (orange). *Right (cause)*: variance ratio $\text{Var}(g)/\sigma_{\text{gen}}^2$ for the same two methods. The dotted curve is the theoretical naive inflation $1 + \rho$ evaluated at the universe-median σ_{gen}^2 ; the dashed line marks a variance ratio of one.

variance, making the marginal more dependent on the market distribution. We tested the clipping rule directly by rescaling the market growth-rate path by $\gamma \in \{1, 2, 3\}$, holding the calibrated marginals fixed (Table S6). At $\gamma = 1$ no ticker crossed the threshold. At $\gamma = 2$ and 3, clipping applied to 76.4% and 95.2% of tickers. The clipping rule was unnecessary under observed market variance but activated as intended when market volatility was artificially increased. At the observed market variance, the floor f did not affect any branch assignment; varying R_{preserve}^2 changed only the two tracker assignments identified above.

4.5 Daily-rebalanced portfolio check

To test a use that depends on joint cross-asset behavior, we applied a 20-asset target-weight allocator to the hybrid-composed 2025 paths. Switching the same target weights from static allocation to daily rebalancing increased the median annualized growth rate from 1.97% to 29.69%. Adding the allocation signal increased it by only another 0.13 percentage points, to 29.83%. The same allocator on the real 2025 history returned 11.50% (Table S2). This comparison was consistent with omitted residual covariance creating a diversification-return bias, but it did not isolate how much of the gap that omission caused. A constant log-wealth shift set the synthetic ensemble median to the prespecified 8% annual return prior, but it did not restore the missing covariance.

5 Discussion

The single-asset analysis supplied the return generator used in the multi-asset construction. Relative to HMM-NJ, HMM-WJ reduced the autocorrelation error for absolute returns, although its in-sample KS and AD pass rates were slightly lower. The GARCH model achieved the lowest autocorrelation error of any generator but rarely passed the distributional tests, and the GRU showed a similar tradeoff. The adapted HSMM-style baseline passed the KS test more often but produced too little kurtosis. Bootstrap resampling passed the distributional tests by drawing directly from the observed returns, but it did not reproduce volatility clustering. We therefore interpret HMM-WJ as a compromise between distributional and temporal fit, not as a uniformly better single-asset model.

The distributional pass rates also need cautious interpretation. The two-sample KS and AD tests assume independent observations, whereas the HMM, HSMM, GARCH, and block-bootstrap paths contain dependence. For that reason, we considered the tests alongside Wasserstein and Hellinger distances and the Hill tail index. Model resolution creates a second qualification. A sensitivity sweep over $N \in \{30, 60, 90, 100, 150, 200\}$ showed little change across the middle of this range for the 2,766-day SPY sample, but shorter histories will support fewer well-populated states. We also tuned the jump parameters (ϵ, λ) on SPY and then applied them across the asset universe. Separate fits for NVDA, JNJ, and JPM achieved in-sample KS pass rates of at least 98.9%, but three assets cannot establish that one jump calibration transfers to every return series. Finally, the Student- t emissions model heavy tails but not skewness directly. The fitted skewness instead came from unequal time spent in negative and positive states, so assets with different asymmetry may need a different state partition.

The main multi-asset result was the variance correction. Naive composition added market variance to a draw that already represented the asset’s total variance. The correction removed that double count and passed 73.6% of the full-length in-sample KS comparisons. After all fits were frozen, it raised the mean cross-ticker KS pass rate from 83.5% under centered naive composition to 87.0% on the 2025 holdout and kept the median variance ratio at 0.97. The improvement was broad but not uniform: every method retained a lower tail of tickers with weak fits, and some hybrid fits deteriorated relative to matched-length training blocks. A high average pass rate is therefore not a guarantee for every asset.

Fitting the return generator directly to OLS residuals is a competitive alternative. Residual-fit methods avoid double counting because the market component has already been removed. The best such method, JumpHMM-on-residuals, had a slightly higher KS pass rate in sample, while the hybrid method had the higher rate in the holdout. The practical difference is reuse: the hybrid construction can use a generator already fitted to full returns, whereas the residual method requires another fit for each asset. Neither approach achieved nominal tail coverage in the short holdout. The hybrid method’s one-day left-tail 99% VaR exceedance rate was 1.67%, the same rounded rate as JumpHMM-on-residuals and above the 1% nominal rate.

Several design choices narrow this comparison. In sample, we used the observed SPY history as the market path. In the holdout experiment, we generated SPY from its frozen training-period model and reserved the observed 2025 path for scoring. The naive and hybrid methods shared the same JumpHMM draw for each ticker, while the residual methods used draws from their own fitted generators. We omitted the optional cross-sectional rank reorder and evaluated VaR for each ticker rather than for a portfolio. The variance correction also assumes $\text{Cov}(\tilde{g}_i^c, g_m) \approx 0$, which holds here because the asset and market paths were generated independently. A generator sampled from joint market history would require a covariance adjustment. In addition, the present construction adds only contemporaneous market dependence; it does not reproduce lead-lag effects or market-driven volatility clustering.

The sensitivity results mark the boundary of the observed evidence. Hybrid KS pass rates declined from 87.6% in the lowest β quartile to 65.9% in the highest because the market path determined more of the composed distribution as β increased. The clipping rule did not activate in the observed universe, where every ticker had $\rho_i < 0.90$, but it activated for most tickers after market volatility was doubled or tripled. Only two tracker assets used the R^2 -preserving branch. The 249-day holdout also had much less power than the 2,766-day training comparison. Matched-length training blocks partly address that difference, but one year of test data cannot provide a precise ranking from either pass rates or left-tail 99% VaR counts. The evidence therefore supports per-asset variance correction and distributional evaluation, not portfolio-level dependence fidelity.

The portfolio experiment shows why that distinction matters. Independent residual draws removed the sector and style covariance present in real assets [45]. Daily rebalancing then converted the excess cross-sectional dispersion into a spurious return premium. This problem affected every method that generated residuals independently, including the block-bootstrap and GARCH variants. In a 20-asset target-weight example, the synthetic median annualized growth rate exceeded the realized 2025 return by about 18 percentage points. Moving from static weights to daily rebalancing changed the median by 27.73 percentage points, whereas adding the allocation signal changed it by only 0.13 percentage points [45, 54, 55]. The apparent allocation gain was therefore almost entirely a diversification-return bias.

A joint residual model could address this limitation. A synchronized multivariate block bootstrap, residual factor model, or copula could restore cross-asset residual covariance. Multi-factor or lagged-factor versions could also capture dependence omitted by the current single-factor construction, with the variance correction retained as a compositional constraint. Per-asset jump calibration and stress tests involving highly volatile markets or leveraged assets would test the other transfer limits identified here. Separate univariate autoregressive or block-bootstrap models would not be enough: they can improve temporal behavior within each asset without creating dependence between assets.

The current location adjustment does not resolve this central limitation. We calibrated a constant shift in log-wealth to an external 8% annual long-run return prior [56, 57], not to the realized test-year result. The shift preserved path ranks and adjusted the return level, although it changed drawdown timing and depth as wealth accumulated. It did not restore the missing residual covariance. The variance correction is therefore necessary for combining full-return generators with a market factor, but it is not a complete dependence model. Portfolio-level use requires joint residual dynamics.

6 Conclusion

In this study, we developed a variance-corrected method for composing per-asset generators through a single-index model. Centering first prevented a full-return draw from shifting the calibrated intercept; the subsequent scale correction removed the variance inflation caused by adding the market factor. Across 423 non-market assets, it recovered the calibrated factor loading to a low median absolute error and produced heavy-tailed marginals. On 416 complete held-out asset histories from 2025, it improved the mean cross-ticker KS pass rate from 83.5% under centered naive composition to 87.0% and held the median variance ratio at 0.97. Its one-day left-tail 99% Value-at-Risk exceedance rate was 1.67%, the same rounded rate as JumpHMM-on-residuals but above the 1% nominal rate. Those residual models, fitted directly to the OLS residual series, required a separate fit for each asset and produced a similar marginal pass rate.

The per-asset component underpinning this construction was a hidden Markov marginal model with Laplace-quantile states, Student- t emissions, and direct transition counting. On SPY, its jump-duration

variant reduced absolute-return autocorrelation error relative to the no-jump variant while keeping high KS and AD pass rates; GARCH reached a lower autocorrelation error but failed the distributional tests. The single-asset result therefore showed a measured tradeoff between distributional fit and volatility clustering, not uniform superiority. Taken together, these results separate two tasks that are easily conflated when synthesizing multi-asset market data: the marginal generator governs each asset’s distributional and temporal behavior, whereas the variance correction prevents the common market factor from double counting the generator’s total variance.

The current multi-asset implementation does not model cross-asset residual covariance. In a 20-asset allocator, independent residuals inflated the synthetic median daily-rebalanced return well above the realized 2025 outcome, almost entirely through the static-to-daily-rebalance diversification return rather than the allocation signal. A location-shift correction set to a documented 8% annual long-run prior adjusted the return level without restoring the missing covariance. A full joint residual model is needed before using these paths for portfolio-level return forecasts. The variance correction is a necessary compositional constraint rather than a complete dependence model. Portfolio-level synthetic market data would require combining it with joint residual dynamics.

Acknowledgments

The authors thank the eCornell AI in Finance program participants for discussions that motivated the downstream-use bias-correction analysis, and Polygon.io for access to the 2014 to 2024 US equity and ETF daily price histories used in the experiments. Computations were run on a single laptop; no shared-resource allocation was required.

Author contributions. A.A. designed the hybrid hidden Markov marginal generator, performed the single-asset experiments and validation analyses, and contributed to the manuscript. J.D.V. designed the variance-corrected single-index composition, performed the multi-asset experiments and downstream Value-at-Risk validation, developed the bias-correction methodology, released the code, and contributed to the manuscript.

Competing interests. The authors declare no competing interests.

Code and data availability. The experiment scripts, evaluation code (six-method comparison, Value-at-Risk coverage check, and bias correction), cached inputs and result summaries, and fitted per-asset marginals are released in the paper repository (<https://github.com/varnerlab/HMM-w-jumps-paper>); the underlying model-fitting library is released separately as JumpHMM.jl (<https://github.com/varnerlab/JumpHMM.jl>), pinned as a dependency of the experiment code.

Online Appendix

S1 Empirical cross-term diagnostic

The variance correction of Eq. (8) assumes $\text{Cov}(\tilde{g}_i^c, g_m) \approx 0$, allowing the variance derivation to omit the cross-term. This assumption does not follow automatically from the fitting procedure because each JumpHMM uses the asset’s full growth-rate history, including its market-correlated component. A remaining market loading in the generator draw would invalidate the closed-form scale. We tested the assumption by sampling $R = 100$ generator growth-rate paths $\tilde{g}_i^{(r)}$ per ticker from each generator and computing the normalized cross-covariance against the real SPY path:

$$\text{cov}_i^{\text{norm}} = \frac{1}{R} \sum_{r=1}^R \frac{\text{Cov}((\tilde{g}_i^{(r)})^c, g_m)}{\sigma_{\text{gen},i} \sigma_m}. \quad (\text{S1})$$

Centering does not change covariance, so these values are numerically identical to those obtained from the raw draws. Per-ticker values were plotted as histograms in Fig. S1. Across the 423 tickers, the full-return JumpHMM generator (used by the hybrid and naive composition methods), the residual-fit JumpHMM generator (used by the JumpHMM-on-residuals composition method), and the Gaussian baseline had universe-mean normalized cross-covariances of 1.53×10^{-4} , 6.58×10^{-5} , and 2.26×10^{-4} , respectively. Their per-ticker values ranged from -0.0061 to 0.0061 . These values quantify the cross-term omitted in Eq. (8).




Figure S1: **Empirical normalized cross-covariance** $\text{Cov}(\tilde{g}_i^c, g_m)/(\sigma_{\text{gen},i} \sigma_m)$ **across the 423-ticker universe.** Histograms were stacked per generator, with each per-ticker value averaged over $R = 100$ generator growth-rate paths. Universe means were 1.53×10^{-4} , 6.58×10^{-5} , and 2.26×10^{-4} for the full-return JumpHMM, residual-fit JumpHMM, and Gaussian generators.

S2 Bias correction for daily-rebalanced backtests

This appendix derives the bias correction in Eq. (S14). Under daily rebalancing, the raw synthetic ensemble had a median annualized growth rate about 22 percentage points above the long-run prior. The variance correction in Eq. (8) first sets each synthetic asset’s second moment to its generator target. When that target matches the empirical second moment, the second-order formula for the diversification return is also matched at the asset level. The remaining difference can reflect omitted cross-asset residual covariance, higher-order Taylor terms, calibration drift, and the documented attenuation of realized diversification returns [54]. The location-shift correction absorbs these effects as a single empirical drag on log-wealth.

The synthetic paths reproduce per-asset marginals and single-factor cross-sectional dependence. Each residual generator retains its own temporal structure, but the residual paths are generated

independently across assets at each date. The construction therefore omits cross-asset residual covariance. Consider a daily-rebalanced portfolio of N assets with constant target weights $w_i \geq 0$, $\sum_i w_i = 1$, reset to those weights at the start of each trading day. Throughout this derivation $g_i(t)$ is the annualized log-growth rate of Eq. (1), so the per-period log return on asset i is $g_i(t) \Delta t$ with $\Delta t = 1/252$ yr. The wealth process is:

$$W_t = W_{t-1} \sum_i w_i \exp(g_i(t) \Delta t), \quad (\text{S2})$$

so the per-step log return on the portfolio is $\log(W_t/W_{t-1}) = \log \sum_i w_i \exp(g_i(t) \Delta t)$. We suppress the time argument, writing g_i for $g_i(t)$. A second-order Taylor expansion of the exponential about zero is $\exp(g_i \Delta t) = 1 + g_i \Delta t + \frac{1}{2}(g_i \Delta t)^2 + O((g \Delta t)^3)$. Substituting it into each summand of $\sum_i w_i \exp(g_i \Delta t)$ gives:

$$\begin{aligned} \sum_i w_i \exp(g_i \Delta t) &= \sum_i w_i \left[1 + g_i \Delta t + \frac{1}{2}(g_i \Delta t)^2 + O((g \Delta t)^3) \right] \quad (\text{Taylor}) \\ &= \sum_i w_i + \Delta t \sum_i w_i g_i + \frac{1}{2}(\Delta t)^2 \sum_i w_i g_i^2 + O((g \Delta t)^3) \quad (\text{distribute}) \\ &= 1 + \Delta t \sum_i w_i g_i + \frac{1}{2}(\Delta t)^2 \sum_i w_i g_i^2 + O((g \Delta t)^3). \quad (\sum_i w_i = 1) \end{aligned} \quad (\text{S3})$$

A second-order Taylor expansion of $\log(1+x)$ about $x=0$ gives $\log(1+x) = x - \frac{1}{2}x^2 + O(x^3)$. Identifying $1+x$ with the right-hand side of Eq. (S3), the order- Δt correction is $x := \Delta t \sum_i w_i g_i + \frac{1}{2}(\Delta t)^2 \sum_i w_i g_i^2$, and squaring keeps only the leading $(\Delta t)^2$ term: $x^2 = (\Delta t)^2 (\sum_i w_i g_i)^2 + O((\Delta t)^3)$ because the cross term has an extra Δt factor. Substitution gives:

$$\begin{aligned} \log \frac{W_t}{W_{t-1}} &= \log \left(1 + x + O((g \Delta t)^3) \right) \\ &= x - \frac{1}{2}x^2 + O(x^3) \\ &= \Delta t \sum_i w_i g_i + \frac{1}{2}(\Delta t)^2 \sum_i w_i g_i^2 \\ &\quad - \frac{1}{2}(\Delta t)^2 \left(\sum_i w_i g_i \right)^2 + O((g \Delta t)^3) \\ &= \Delta t \sum_i w_i g_i + \frac{1}{2}(\Delta t)^2 \left[\sum_i w_i g_i^2 - \left(\sum_i w_i g_i \right)^2 \right] + O((g \Delta t)^3). \end{aligned} \quad (\text{S4})$$

Dividing through by Δt extracts the portfolio's annualized one-step log-growth rate $g_p(t) := \Delta t^{-1} \log(W_t/W_{t-1})$, the portfolio counterpart of the per-asset $g_i(t)$ (still suppressing the time argument):

$$g_p = \sum_i w_i g_i + \frac{1}{2} \Delta t \left(\sum_i w_i g_i^2 - \left(\sum_i w_i g_i \right)^2 \right) + O((\Delta t)^2). \quad (\text{S5})$$

Restoring the time argument, we now take unconditional expectations of Eq. (S5) under stationarity. Define the per-asset annualized mean and variance:

$$\mu_i := \mathbb{E}(g_i(t)), \quad \sigma_i^2 := \text{Var}(g_i(t)), \quad (\text{S6})$$

the per-rate covariance $\Sigma_{ij} := \text{Cov}(g_i(t), g_j(t))$, and the per-rate portfolio variance $\sigma_p^2 := \text{Var}(\sum_i w_i g_i(t)) = w^\top \Sigma w$. The $O(\Delta t)$ correction in Eq. (S5) contains two squared quantities, $\sum_i w_i g_i(t)^2$ and $(\sum_i w_i g_i(t))^2$. Both expand by the second-moment identity, which holds for any random variable X with finite second moment:

$$\mathbb{E}(X^2) = \mathbb{E}(X)^2 + \text{Var}(X). \quad (\text{S7})$$

Applying this identity to $X = g_i(t)$ and summing against w_i gives:

$$\begin{aligned}\mathbb{E}\left(\sum_i w_i g_i(t)^2\right) &= \sum_i w_i \mathbb{E}(g_i(t)^2) && \text{(linearity)} \\ &= \sum_i w_i [\mu_i^2 + \sigma_i^2] && \text{(by Eq. (S7))} \\ &= \sum_i w_i \mu_i^2 + \sum_i w_i \sigma_i^2. && \text{(S8)}\end{aligned}$$

Applying the identity to $X = \sum_i w_i g_i(t)$, whose mean is $\sum_i w_i \mu_i$ and whose variance is σ_p^2 , gives:

$$\begin{aligned}\mathbb{E}\left(\left(\sum_i w_i g_i(t)\right)^2\right) &= \mathbb{E}\left(\sum_i w_i g_i(t)\right)^2 + \text{Var}\left(\sum_i w_i g_i(t)\right) && \text{(by Eq. (S7))} \\ &= \left(\sum_i w_i \mu_i\right)^2 + \sigma_p^2. && \text{(S9)}\end{aligned}$$

We substituted Eqs. (S8)–(S9) into the expectation of Eq. (S5) and grouped the variance and mean contributions as follows:

$$\begin{aligned}\mathbb{E}(g_p) &= \sum_i w_i \mu_i + \underbrace{\frac{1}{2} \Delta t \left(\sum_i w_i \sigma_i^2 - \sigma_p^2 \right)}_D \\ &\quad + \frac{1}{2} \Delta t \left[\sum_i w_i \mu_i^2 - \left(\sum_i w_i \mu_i \right)^2 \right] + O((\Delta t)^2),\end{aligned}\tag{S10}$$

the Booth-Fama identity for the daily-rebalanced portfolio in annualized growth-rate units [45, 55]. The middle term, D , is the diversification return: the $O(\Delta t)$ correction equal to half the difference between the weighted-average per-asset variance $\sum_i w_i \sigma_i^2$ and the portfolio variance σ_p^2 . The per-asset moments μ_i and σ_i^2 have units yr^{-1} and yr^{-2} respectively; the products $\Delta t \sigma_i^2$ and $\Delta t \sigma_p^2$ in Eq. (S10) are the annualized variances (yr^{-1}). The third term in Eq. (S10) is the cross-sectional dispersion of the per-asset annualized means, of order $\Delta t \mu^2 \sim 3 \times 10^{-5} \text{ yr}^{-1}$ at $\mu \sim 0.08 \text{ yr}^{-1}$ and negligible against the variance term $D \sim \Delta t \sigma^2 \sim 4 \times 10^{-2} \text{ yr}^{-1}$; we drop it. The realized annualized compound growth rate $\bar{g}_p = (T \Delta t)^{-1} \log(W_T/W_0) = T^{-1} \sum_t g_p(t)$ converges to $\mathbb{E}(g_p)$ by the ergodic theorem under any stationary residual process, so the long-run limit is fixed by the one-time-slice unconditional joint distribution of $(g_i(t))_i$ and does not, in itself, depend on the serial structure of the residual draw.

To rewrite D in a form that makes its non-negativity manifest, expand $\sigma_p^2 = \sum_i w_i^2 \sigma_i^2 + 2 \sum_{i < j} w_i w_j \Sigma_{ij}$ and use the identity $\sum_i w_i (1 - w_i) \sigma_i^2 = \sum_{i < j} w_i w_j (\sigma_i^2 + \sigma_j^2)$ that holds for any weights summing to one:

$$D = \frac{1}{2} \Delta t \left(\sum_i w_i \sigma_i^2 - \sigma_p^2 \right) = \frac{1}{2} \Delta t \sum_{i < j} w_i w_j \text{Var}(g_i - g_j) \geq 0, \tag{S11}$$

so D equals one-half the annualized weighted-average pairwise growth-rate-difference variance and is therefore non-negative, with equality only when every pair (i, j) comoves perfectly ($\text{Var}(g_i - g_j) = 0$) or all weight concentrates on a single asset. The rebalancing rule itself produces this effect: weights drift between rebalances, the rebalancer sells whatever has outperformed (its weight rose above target) and buys whatever has underperformed (its weight fell below target), and the trade systematically converts period-by-period cross-sectional dispersion into compounded growth. Markowitz [55] described this as the variance bonus of the long-run mean-variance allocation; Booth and Fama [45] called it the diversification return.

Substituting the single-index decomposition $g_i(t) = \alpha_i + \beta_i g_m(t) + \varepsilon_i(t)$, with market variance $\sigma_m^2 := \text{Var}(g_m(t))$, per-asset residual variance $\sigma_{\varepsilon,i}^2 := \text{Var}(\varepsilon_i(t))$, and $\text{Cov}(g_m, \varepsilon_i) = 0$ (verified in Sec. S1), into Eq. (S11) gives:

$$\text{Var}(g_i - g_j) = (\beta_i - \beta_j)^2 \sigma_m^2 + \text{Var}(\varepsilon_i - \varepsilon_j), \tag{S12}$$

Because g_m and the residuals are uncorrelated, this relation splits the diversification return into market-beta-dispersion and idiosyncratic components:

$$D = \underbrace{\frac{1}{2} \Delta t \sigma_m^2 \sum_{i < j} w_i w_j (\beta_i - \beta_j)^2}_{D_{\text{mkt}}} + \underbrace{\frac{1}{2} \Delta t \sum_{i < j} w_i w_j \text{Var}(\varepsilon_i - \varepsilon_j)}_{D_\varepsilon}, \quad (\text{S13})$$

where the market-beta-dispersion component D_{mkt} equals $\frac{1}{2} \Delta t \sigma_m^2 [\sum_i w_i \beta_i^2 - (\sum_i w_i \beta_i)^2]$, the weighted cross-sectional β dispersion scaled by the annualized market variance $\Delta t \sigma_m^2$. Across the 423-ticker universe the empirical β distribution spans $[0.03, 1.79]$ with interquartile range $[0.81, 1.18]$ and the SPY annualized variance $\Delta t \sigma_m^2 \approx 0.04 \text{ yr}^{-1}$. The weighted variance of β is bounded by one quarter of its squared range, so D_{mkt} is at most 0.016 yr^{-1} , or 1.6 percentage points per year over this universe. For the idiosyncratic component D_ε , residuals were approximately orthogonal across assets at the same time slice. Therefore, $\text{Var}(\varepsilon_i - \varepsilon_j) \approx \sigma_{\varepsilon,i}^2 + \sigma_{\varepsilon,j}^2$, and D_ε scales with the weighted-average annualized idiosyncratic variance $\Delta t \sigma_{\varepsilon,i}^2$ per name.

The second-order value of D depends on same-time unconditional moments, not on serial dependence. Equation (8) sets $\sigma_{\text{gen},i}^2 = \beta_i^2 \sigma_m^2 + \sigma_{\varepsilon,i}^2$ to its empirical target. Any residual generator with the same unconditional second moments therefore gives the same D . This includes a stationary AR(1) residual generator $\varepsilon_i(t) = \phi \varepsilon_i(t-1) + \eta_i(t)$ with i.i.d. innovations η_i tuned so that $\text{Var}(\varepsilon_i) = \sigma_{\eta,i}^2 / (1 - \phi^2) = \sigma_{\varepsilon,i}^2$. The parameter ϕ leaves the marginal distribution at a single time slice unchanged. It therefore does not change $\mathbb{E}(g_p)$ in Eqs. (S10) and (S13), but it changes sampling variance. For a stationary AR(1) residual with variance σ_ε^2 , the variance of its sample mean is approximately $T^{-1} \sigma_\varepsilon^2 (1 + \phi) / (1 - \phi)$. Under ergodicity, serial dependence changes per-path sampling noise, not the stationary mean. The raw median's ~ 22 -percentage-point excess over an external long-run growth prior is therefore not caused by i.i.d. sampling *across time* at the level of the second-order formula.

The direct structural difference is contemporaneous. The derivation assumes residuals are orthogonal across assets at each time slice, as they are in the independent per-asset generator. Real residuals share sector and style factors, which reduce $\text{Var}(\varepsilon_i - \varepsilon_j)$ below the independent-generator value $\sigma_{\varepsilon,i}^2 + \sigma_{\varepsilon,j}^2$. Matching each marginal variance does not restore this covariance. The synthetic D_ε will therefore exceed its real-market counterpart when the omitted covariance is positive. Measuring that contribution requires a joint-residual experiment, which we did not run. The gap may also include higher-moment corrections to the Taylor expansion (heavy-tailed JumpHMM marginals have non-trivial third and fourth cumulants), the time-varying volatility and correlation across the 2014–2024 calibration window relative to a long-run prior, and the documented attenuation of realized diversification returns on real equity portfolios at all rebalancing horizons [54]. The location-shift correction of Eq. (S14) treats this composite shortfall as a single empirical drag b to be absorbed and is agnostic to its decomposition.

Table S1: Descriptive statistics and stylized-facts tests for SPY daily excess growth rates. In-sample (IS): 2014–2024 ($T = 2,766$); out-of-sample (OoS): 2025 ($T = 249$). Gaussian and Laplace columns show maximum likelihood estimation (MLE) fits to the in-sample data. We used the Jarque–Bera (JB) normality test and the Ljung–Box (LB) autocorrelation test at lag 20. * denotes rejection at $\alpha = 0.05$.

Statistic	IS Observed	OoS Observed	Gaussian	Laplace
Mean (annualized, %)	6.31	11.60	6.31	14.19
Std. dev. of G_t (% yr^{-1})	214.50	220.62	214.46	205.93
Skewness	−0.753	−1.093	0	0
Excess Kurtosis	7.715	6.867	0	3
JB normality test	reject* ($p < 0.001$)	reject* ($p < 0.001$)	n/a	n/a
LB test on G_t (lag 20)	reject* ($p < 0.001$)	reject* ($p = 0.019$)	n/a	n/a
LB test on $ G_t $ (lag 20)	reject* ($p < 0.001$)	reject* ($p < 0.001$)	n/a	n/a

To measure the bias, we applied a 20-asset, long-only target-weight allocator to the hybrid-composed universe. We calibrated the assets over 2014–2024, deployed the allocator in 2025, limited the universe to names with calibrated (α, β) and $\max \beta = 1.21$, and assumed a 5-basis-point round-trip

cost. We then decomposed the path-mean annualized growth rate $\bar{g}_p \equiv (T \Delta t)^{-1} \log(W_{p,T}/W_{p,0})$ into static and rebalancing-driven components (Table S2). The static-to-daily-rebalance switch increased the median by 27.73 percentage points. Adding a signal after daily rebalancing changed it by another 0.13 percentage points. The same allocator on the real 2025 growth-rate history produced 11.5%/yr, about 18 percentage points below the synthetic median. The synthetic residuals were independent across tickers and therefore omitted the contemporaneous covariance present in real markets. Determining how much of the gap this omission caused would require a joint-residual experiment with the marginals and allocator held fixed.

Table S2: Diversification-return decomposition on a 20-asset target-weight allocator deployed against the hybrid-composed 2025 universe. Median path-mean annualized growth rate $\bar{g}_p$ across the synthetic ensemble. Equal-weight buy-and-hold serves as a passive baseline; the target-weight allocation is reported under static (no intra-year rebalance) and daily-rebalanced regimes, with and without a signal on top of the target weights. The static-to-daily increase of +27.73 pp on the same target weights isolates the diversification return of Eq. (S13). The same allocator deployed against the real 2025 growth-rate history is shown for comparison.

Allocation rule (synthetic 2025 universe)	$\bar{g}_p$ (%/yr)
Equal-weight buy-and-hold	7.68
Target-weight, static (no rebalance)	1.97
Target-weight, daily rebalanced	29.69
Target-weight, daily rebalanced + signal	29.83
<i>Real 2025 history, same allocator</i>	11.50




Figure S2: Heavy-tail preservation: excess kurtosis and Hill tail index vs calibrated β . Each point is a per-ticker median over 100 replications. *Left:* excess kurtosis, y -axis clipped to $[-2, 20]$ for readability. The dashed line marks the median real-data excess kurtosis across the universe ($\kappa_{\text{real}} \approx 13$); the hybrid and naive composition methods (orange and blue) track the generator's kurtosis, while Gaussian SIM collapses to $\kappa \approx 1$. *Right:* Hill tail-index estimator on the upper 5% of $|g|$. The hybrid and naive medians remain near 0.37 throughout the β range; Gaussian SIM drops to ~ 0.22 . The hybrid's median being below κ_{real} reflects the JumpHMM marginal's own kurtosis rather than a composition-rule distortion: hybrid tracks naive (both retain the generator's tails) while Gaussian SIM does not.

The correction subtracts a constant drift in log-wealth so that the ensemble median annualized growth rate matches an external prior $\bar{g}^{\text{prior}}$. At each time t , it multiplies every path by the same factor and therefore preserves cross-path wealth ranks. It also shifts every log-return increment by the same constant, which leaves the variance of those increments unchanged. For path p , we applied the following correction:

$$W_{p,t}^{\text{corr}} = W_{p,t} \cdot \exp(-b \cdot t \Delta t), \quad b = \text{median}_p \bar{g}_p - \bar{g}^{\text{prior}}, \quad (\text{S14})$$

with Δt the integration step in years and $b, \bar{g}_p, \bar{g}^{\text{prior}}$ all in the same per-year growth-rate units. The correction is a linear detrend in log-wealth and remains multiplicative in wealth. It changes drawdown dates and depths, so it adjusts the return level rather than preserving drawdowns. It does preserve terminal-wealth ranks and the variance of log-return increments. We set $\bar{g}^{\text{prior}} = 8\%/yr$ before examining the holdout result. This value was based on external long-run equity-return estimates and an allowance for fees, turnover, and market impact [56, 57]. We used this documented prior rather than the realized test year to avoid circular calibration. Matching the realized year would force the synthetic median to equal the test sample. In this deployment, the real-2025 outcome of 11.5%/yr fell just above the median of the corrected ensemble.




Figure S3: **Calibrated R^2 distribution across the 423 non-SPY tickers.** Histogram of OLS R^2 against SPY over the 2014–2024 window. The two tickers crossing the $R^2_{\text{preserve}} = 0.80$ threshold (dashed line) are index ETFs (QQQ, SPYG); the highest- R^2 individual stock (BLK, 0.645) falls ~ 0.16 below the threshold. The R^2 -preserving branch therefore contains only QQQ and SPYG in this universe.

Several alternative fixes were considered and rejected. A path-filtering rule that drops synthetic paths above a $\bar{g}_p$ threshold turns the threshold itself into a tuning target and removes only a subset of the bias’s consequences while leaving the underlying bias intact. Inflating per-trade transaction costs to absorb the diversification return would require round-trip costs of ~ 70 basis points versus the 1–10 basis-point range assumed for liquid US equity, shifting the misrepresentation from the synthetic generator to its cost model. Drifting the calibrated parameters between calibration and deployment to widen the predictive envelope, with drift scale varied across $\{0, 1, 2, 3, 5\}$ multiples of the OLS standard error, left the median Sharpe ratio essentially unchanged across the sweep, so standard-error-scale parameter drift did not absorb the bias. Imposing an AR(1) on the residual draw would conflict with the copula reorder step because reordering destroys temporal structure. Preserving both structures would require a block-ordered copula sampler and therefore a new residual model. The per-ticker block bootstrap [52] preserves each ticker’s temporal residual structure but not covariance across tickers. A joint bootstrap or residual copula would be required for that purpose. The block bootstrap also replaces the JumpHMM marginal rather than correcting it. A multiplicative haircut on the raw-versus-static deviation, $W^{\text{corr}} = W_{\text{static}} \cdot (W_{\text{raw}}/W_{\text{static}})^k$, hits the same target at k chosen to match the prior, but it changes the rank ordering of paths and inherits the static allocation’s drawdown pattern.

The location-shift correction preserves percentile ranks and the relative ranking of strategies evaluated on the same paths. It also preserves centered moments of terminal log growth, including variance, skewness, and kurtosis. Because it changes the drawdown distribution, we evaluated drawdowns on the uncorrected ensemble. The correction does not repair residual autocorrelation, absolute Sharpe under daily rebalancing, or sensitivity to rebalancing frequency. Those targets require a residual model that preserves both temporal and cross-sectional dependence.

S3 Validation metric definitions

The distributional and tail metrics reported in the main results were defined as follows. The two-sample Kolmogorov-Smirnov statistic between the empirical cumulative distribution functions (CDFs) of the observed and simulated growth-rate samples (G_{obs} of length n and G_{sim} of length m) is $D = \sup_x |F_n(x) - F_m(x)|$ [42, 43]; the test rejects when D exceeds the critical value at level $\alpha = 0.05$. The Anderson-Darling statistic places greater weight on the tails via the integral $A^2 = nm/(n+m) \int [F_n(x) - F_m(x)]^2 / [F(x)(1-F(x))] dF(x)$, where F is the pooled empirical CDF [44]. The Wasserstein-1 distance between two empirical distributions of equal length is the average of sorted-quantile differences:

$$W_1 = \frac{1}{n} \sum_{i=1}^n |G_{\text{obs}}^{(i)} - G_{\text{sim}}^{(i)}|, \quad (\text{S15})$$

with $G^{(i)}$ the i th order statistic. The Hill upper-tail index is the classical order-statistic estimator:

$$\hat{\xi} = \frac{1}{k-1} \sum_{i=1}^{k-1} \log \left(\frac{X_{(i)}}{X_{(k)}} \right), \quad (\text{S16})$$

with $X_{(1)} \geq X_{(2)} \geq \dots$ the ordered upper-tail absolute returns; k is set to the upper 5% of the sample. For a Pareto-tailed distribution with shape α , $\hat{\xi} \approx 1/\alpha$. For the Hellinger calculation, we binned both samples on a common grid and normalized the bin counts to probability masses p and q . Their distance is:

$$H(p, q) = \frac{1}{\sqrt{2}} \left\| \sqrt{p} - \sqrt{q} \right\|_2 \in [0, 1], \quad (\text{S17})$$

with $H = 0$ for identical probability masses and $H = 1$ for disjoint supports.




Figure S4: **Statistical validation of HMM-WJ** ($N = 100$, $\alpha = 0.05$). Panels (a) and (d) condition on the $\sim 24\%$ of in-sample paths containing jump events. (a) In-sample KS p -values for 239 jump paths (pass rate 93.7%). (b) Out-of-sample KS p -values for all 1,000 paths (pass rate 95.4%). (c) Out-of-sample density fan chart; the observed density falls within the simulation envelope. (d) Observed and mean simulated in-sample ACF of $|G_t|$ for jump paths.

S4 Transition matrix internals and partition diagnostics

Fig. S5 shows the fitted HMM-WJ components for SPY with $N = 100$ states. The fitted Laplace CDF and the empirical CDF are shown with the 99 state boundaries Q_k . The transition matrix $\mathbf{T}$ has a near-diagonal band, with most off-diagonal mass within ± 5 states of the diagonal. Under the ordinary Markov transitions, every state has an expected residence time of only 1–2 steps. The fitted jump duration is $\lambda = 90$ steps, about two orders of magnitude longer. Every state had at least one in-sample observation.




Figure S5: **Fitted HMM-WJ internals for SPY with $N = 100$ states.** Panel (a) overlays the fitted Laplace CDF on the empirical CDF; the 99 vertical lines mark the equal-probability quantile boundaries Q_k that define the hidden states. Panel (b) displays the estimated transition matrix $\mathbf{T}$ in $\log_{10}$ color scale; the dominant near-diagonal band reflects short-range regime persistence. Panel (c) plots the expected natural residence time $1/(1 - T_{kk})$ for each state on a $\log_{10}$ scale, with the bottom tail set $\mathcal{S}_-$ (states 1–5, red) and the top tail set $\mathcal{S}_+$ (states 96–100, teal) highlighted. The dashed horizontal line marks the fitted mean jump duration $\lambda = 90$ steps, quantifying the two-order-of-magnitude gap between natural residence times and the jump-duration override.

S5 Emission distribution and semi-Markov comparison

To compare the emission and duration specifications, we evaluated three variants on SPY (Table S3): an HSMM-style baseline adapted from Bulla and Bulla [16] ($K = 8$ states, negative-binomial dwell times with a geometric fallback, and Student- t emissions), HMM-WJ with the original Gaussian emissions ($N = 100$), and HMM-WJ with Student- t emissions ($N = 100$). Both HMM-WJ variants had higher in-sample KS and AD pass rates than the HSMM-style baseline. The HMM-WJ variants use 100 states and the HSMM uses 8, so this comparison does not assign the pass-rate difference solely to the duration model. Holding the rest of HMM-WJ fixed, changing the emission from Normal to Student- t ($\nu = 5$) increased simulated excess kurtosis from 5.5 to 7.5, against an observed value of 7.7. The Student- t emission also increased the KS and AD pass rates, while ACF-MAE and the distributional distances changed modestly.

Table S3: **Three-way comparison of emission distributions and duration specifications for SPY.** Standard errors in parentheses; bold marks the best value per row. The HSMM is a hidden semi-Markov-style model adapted from Bulla and Bulla ($K = 8$ states, negative-binomial dwell times with a geometric fallback, and Student- t emissions) [16]; HMM-WJ: hybrid hidden Markov model with jump-duration override ($N = 100$). All entries are computed from 1,000 simulated paths at significance level $\alpha = 0.05$.

Metric	HSMM (Student- t)	HMM-WJ (Gaussian)	HMM-WJ (Student- t)
States	$K = 8$	$N = 100$	$N = 100$
<i>In-sample (2,766 trading days, 2014–2024)</i>			
KS pass rate (%)	82.0 (1.2)	97.0 (0.5)	98.3 (0.4)
AD pass rate (%)	42.5 (1.6)	90.5 (0.9)	94.8 (0.7)
Excess kurtosis	4.8 (0.08)	5.5 (0.03)	7.5 (0.09)
ACF-MAE	0.059 (< 0.001)	0.052 (< 0.001)	0.053 (< 0.001)
Wasserstein-1	0.176 (0.001)	0.101 (0.002)	0.097 (0.001)
Hellinger	0.113 (< 0.001)	0.076 (< 0.001)	0.074 (< 0.001)
<i>Out-of-sample (249 trading days, 2025)</i>			
KS pass rate (%)	96.2 (0.6)	95.0 (0.7)	95.4 (0.7)
AD pass rate (%)	96.7 (0.6)	95.9 (0.6)	95.3 (0.7)
Excess kurtosis	4.1 (0.13)	4.9 (0.08)	6.2 (0.15)
ACF-MAE	0.042 (< 0.001)	0.040 (< 0.001)	0.040 (< 0.001)
Wasserstein-1	0.287 (0.002)	0.275 (0.005)	0.275 (0.005)
Hellinger	0.239 (0.001)	0.212 (0.001)	0.210 (0.001)

Table S4: **Hybrid composition method broken out by branch.** N_{tickers} : number of tickers assigned to each branch by the $R^2_{\text{preserve}} = 0.80$ threshold. The clipped variance-preserving branch does not trigger anywhere in the universe ($\rho_i < 0.90$ for all tickers) and is omitted. The two trackers in the R^2 -preserving branch (QQQ and SPYG) had median absolute errors of 0.005 for β and 0.001 for R^2 , with a KS pass rate of 100%.

Branch	N_{tickers}	$ \hat{\beta} - \beta $	$ \hat{R}^2 - R^2_{\text{real}} $	KS pass (%)	W_1	κ
hybrid	421	0.018	0.021	73.5	0.250	7.142
r2-preserve	2	0.005	0.001	100.0	0.099	11.525

S6 State-resolution sensitivity

We swept the partition resolution over $N \in \{30, 60, 90, 100, 150, 200, 350\}$ and re-evaluated the in-sample KS and AD pass rates for HMM-NJ and HMM-WJ. Pass rates varied little across $N \in \{60, \dots, 200\}$. At $N = 30$ both models lost several percentage points on the AD test. At $N = 350$ several quantile bins were never visited in the historical record, leaving the corresponding emission parameters undefined. We adopted $N = 100$ for all subsequent experiments.




Figure S6: **Branch assignment in calibrated $(\beta, R^2_{\text{real}})$ space.** The $R^2_{\text{preserve}} = 0.80$ threshold (dashed line) separates tracker assets from single stocks. In this universe, only QQQ and SPYG exceed the threshold, and both are assigned to the R^2 -preserving branch. The remaining 421 tickers populate the variance-preserving branch; the clipped variance-preserving branch is unused because the market loading ratio $\rho_i = \beta_i^2 \sigma_m^2 / \sigma_{\text{gen},i}^2$ stays below 0.90 for every ticker.

Table S5: **Aggregate metrics by β quartile, across composition methods.** Quartile cutoffs are computed from the calibrated β distribution across the 423-ticker universe. The hybrid KS pass rate declines from 87.6% in Q1 to 65.9% in Q4 as the market loading ratio ρ_i grows; the naive KS pass rate falls from 24.6% to 1.3% over the same range.

β bucket	Method	$ \hat{\beta} - \beta $	KS pass (%)	W_1	κ	Var(g)
Q1 (low β)	Naive	0.018	24.6	0.403	8.971	18.1
	Gaussian SIM	0.015	0.0	0.620	0.622	14.0
	Hybrid	0.016	87.6	0.177	8.369	14.6
	JumpHMM-on-residuals	0.015	73.2	0.204	7.059	14.3
	Block bootstrap	0.015	71.3	0.208	6.286	13.9
	GARCH(1,1)- t	0.015	57.8	0.253	5.255	13.4
Q2	Naive	0.021	1.9	0.621	7.510	25.1
	Gaussian SIM	0.016	0.2	0.653	1.509	18.0
	Hybrid	0.017	74.9	0.231	6.889	19.0
	JumpHMM-on-residuals	0.017	82.7	0.200	7.799	18.7
	Block bootstrap	0.016	79.8	0.197	7.213	18.0
	GARCH(1,1)- t	0.015	76.4	0.227	6.470	17.8
Q3	Naive	0.023	0.7	0.792	7.390	33.3
	Gaussian SIM	0.017	1.4	0.688	1.947	23.4
	Hybrid	0.018	66.1	0.279	6.904	24.4
	JumpHMM-on-residuals	0.018	78.3	0.242	7.454	24.0
	Block bootstrap	0.017	75.0	0.236	6.888	22.8
	GARCH(1,1)- t	0.016	71.0	0.263	6.265	21.9
Q4 (high β)	Naive	0.029	1.3	0.965	7.233	51.2
	Gaussian SIM	0.022	0.6	0.825	2.054	36.3
	Hybrid	0.022	65.9	0.339	6.739	37.1
	JumpHMM-on-residuals	0.022	69.1	0.313	6.842	37.0
	Block bootstrap	0.021	67.2	0.311	6.334	35.3
	GARCH(1,1)- t	0.021	64.5	0.361	6.064	34.7

Table S6: **Clipping-branch stress test under market-volatility scaling.** The market growth-rate path was rescaled by $\gamma \in \{1, 2, 3\}$ with the calibrated marginals held fixed, and the full hybrid composition method re-evaluated on the 423-ticker universe. As γ rises, the share of clipped tickers grows.

γ	Clipped (%)
1	0.0
2	76.4
3	95.2

S7 Cross-asset generalization

We repeated the fitting and evaluation procedure on three stocks: NVDA (semiconductors, high volatility), JNJ (health care, low volatility), and JPM (financials, moderate volatility). Per-ticker tune outputs at ($n_{\text{paths}} = 200$, seed = 1234) are reported in Table S7. The objective combines ACF and kurtosis errors with fixed weights. Because kurtosis varies across finite samples of heavy-tailed paths, the selected grid point can change with the Monte Carlo sample. The reported (ϵ^*, λ^*) values are reproducible with the stated path count and seed, but they are not unique per-ticker optima. At the selected parameter values, in-sample HMM-WJ KS pass rates were 98.9% for NVDA, 99.4% for JNJ, and 99.1% for JPM. Out-of-sample KS pass rates on the 249-day 2025 hold-out were 97.7%, 74.0%, and 79.2%, respectively.

Table S7: **Cross-asset generalization of HMM-WJ across three single-stock tickers spanning sector and volatility regimes.** Per-ticker tune outputs (ϵ^*, λ^*) at $n_{\text{paths}} = 200$ and seed = 1234, and the corresponding HMM-WJ pass rates on 1,000 simulated paths at $\alpha = 0.05$ against each ticker’s in-sample (2014–2024) and out-of-sample (2025) growth-rate history. Standard errors in parentheses (binomial SE for pass rates). The (ϵ^*, λ^*) values are reported as the parameter values used for the IS / OoS metrics rather than as canonical per-ticker optima.

Ticker	ϵ^*	λ^*	IS KS (%)	IS AD (%)	OoS KS (%)	OoS AD (%)
NVDA (semiconductors)	10^{-4}	70	98.9 (0.3)	96.7 (0.6)	97.7 (0.5)	96.1 (0.6)
JNJ (health care)	10^{-4}	70	99.4 (0.2)	96.3 (0.6)	74.0 (1.4)	61.3 (1.5)
JPM (financials)	10^{-4}	80	99.1 (0.3)	95.7 (0.6)	79.2 (1.3)	77.7 (1.3)

S8 SIM composition derivations and properties verification

The variance-corrected single-index composition states two closed-form results, the variance scale of Eq. (8) and the residual-variance target of Eq. (11). The derivation below gives the large- T OLS coefficients and composed variance. To derive Eq. (8), first center the full-return draw as in Eq. (6), then set $\varepsilon_i = s_i \tilde{g}_i^c$ for $s_i \in (0, 1]$ and require that the composed series have the same variance as the generator path:

$$\text{Var}(g_i) = \text{Var}(\alpha_i + \beta_i g_m + \varepsilon_i) = \text{Var}(\beta_i g_m + \varepsilon_i) = \sigma_{\text{gen},i}^2, \quad (\text{S18})$$

where α_i drops out because it is constant. Expanding the sum gives:

$$\text{Var}(\beta_i g_m + \varepsilon_i) = \beta_i^2 \text{Var}(g_m) + 2\beta_i \text{Cov}(g_m, \varepsilon_i) + \text{Var}(\varepsilon_i). \quad (\text{S19})$$

Substituting $\varepsilon_i = s_i \tilde{g}_i^c$ gives $\text{Var}(\varepsilon_i) = s_i^2 \sigma_{\text{gen},i}^2$ and $\text{Cov}(g_m, \varepsilon_i) = s_i \text{Cov}(g_m, \tilde{g}_i^c)$. Under the assumption $\text{Cov}(\tilde{g}_i^c, g_m) \approx 0$, the cross term vanishes and the variance condition becomes $\beta_i^2 \sigma_m^2 + s_i^2 \sigma_{\text{gen},i}^2 = \sigma_{\text{gen},i}^2$. Solving for s_i^2 gives Eq. (8).

To derive Eq. (11), write the population R^2 identity for Eq. (4):

$$R^2 = \frac{\beta_i^2 \sigma_m^2}{\beta_i^2 \sigma_m^2 + \sigma_{\varepsilon,i}^2}. \quad (\text{S20})$$

Solving this identity for the residual variance gives Eq. (11). Because $\bar{\varepsilon}_i = 0$ after centering and scaling, the sample mean of a composed path obeys $\bar{g}_i = \alpha_i + \beta_i^{\text{eff}} \bar{g}_m$. Hence, the finite-sample OLS

intercept was:

$$\hat{\alpha}_i = \bar{g}_i - \hat{\beta}_i \bar{g}_m = \alpha_i - (\hat{\beta}_i - \beta_i^{\text{eff}}) \bar{g}_m. \quad (\text{S21})$$

Under the zero-cross-covariance assumption, in the large- T limit OLS therefore gives $\hat{\alpha}_i \rightarrow \alpha_i$ and $\hat{\beta}_i \rightarrow \beta_i^{\text{eff}}$. Here $\beta_i^{\text{eff}} = \beta_i$ unless Eq. (10) clips the market loading. On the variance-preserving branch, the composed variance is $\sigma_{\text{gen},i}^2$. Without clipping, $\text{Var}(g_i) = \beta_i^2 \sigma_m^2 + s_i^2 \sigma_{\text{gen},i}^2 = \sigma_{\text{gen},i}^2$. Under clipping, Eq. (10) sets β_i^{eff} to give the same variance. On the R^2 -preserving branch, the composed variance is $\beta_i^2 \sigma_m^2 / R_{i,\text{real}}^2$, the value implied by the calibrated $(\beta_i, R_{i,\text{real}}^2)$ rather than by the generator marginal. The floor guarantees only that $\text{Var}(\varepsilon_i) \geq f \sigma_{\text{gen},i}^2$; it does not guarantee that adding the market term preserves the generator's tail distribution. In Table S5, the KS pass rate fell from 87.6% in the lowest-beta quartile to 65.9% in the highest-beta quartile.

References

- [1] Samuel A. Assefa, Danial Dervovic, Mahmoud Mahfouz, Robert E. Tillman, Prashant Reddy, and Manuela Veloso. Generating synthetic data in finance: Opportunities, challenges and pitfalls. In *Proceedings of the First ACM International Conference on AI in Finance*, pages 1–8, 2020. doi: 10.1145/3383455.3422554.
- [2] James Jordon, Lukasz Szpruch, Florimond Houssiau, Mirko Bottarelli, Giovanni Cherubin, Carsten Maple, Samuel N. Cohen, and Adrian Weller. Synthetic data: What, why and how? *arXiv preprint arXiv:2205.03257*, 2022.
- [3] Benoit Mandelbrot. The variation of certain speculative prices. *The Journal of Business*, 36(4): 394–419, 1963.
- [4] R. Cont. Empirical properties of asset returns: Stylized facts and statistical issues. *Quantitative Finance*, 1(2):223–236, 2001. doi: 10.1080/713665670.
- [5] Michael Stenger, André Bauer, Thomas Prantl, Robert Leppich, Nathaniel Hudson, Kyle Chard, Ian Foster, and Samuel Kounev. Thinking in categories: A survey on assessing the quality for time series synthesis. *Journal of Data and Information Quality*, 16(2):1–32, 2024. doi: 10.1145/3666006.
- [6] Robert F. Engle. Autoregressive conditional heteroscedasticity with estimates of the variance of United Kingdom inflation. *Econometrica*, 50(4):987–1007, 1982.
- [7] Tim Bollerslev. Generalized autoregressive conditional heteroskedasticity. *Journal of Econometrics*, 31(3):307–327, 1986. doi: 10.1016/0304-4076(86)90063-1.
- [8] Steven L. Heston. A closed-form solution for options with stochastic volatility with applications to bond and currency options. *The Review of Financial Studies*, 6(2):327–343, 1993. doi: 10.1093/rfs/6.2.327.
- [9] Robert C. Merton. Option pricing when underlying stock returns are discontinuous. *Journal of Financial Economics*, 3(1-2):125–144, 1976. doi: 10.1016/0304-405X(76)90022-2.
- [10] S. G. Kou. A jump-diffusion model for option pricing. *Management Science*, 48(8):1086–1101, 2002. doi: 10.1287/mnsc.48.8.1086.166.
- [11] Shuntaro Takahashi, Yu Chen, and Kumiko Tanaka-Ishii. Modeling financial time-series with generative adversarial networks. *Physica A: Statistical Mechanics and Its Applications*, 527: 121261, 2019. doi: 10.1016/j.physa.2019.121261.
- [12] Jinsung Yoon, Daniel Jarrett, and Mihaela van der Schaar. Time-series generative adversarial networks. In *Advances in Neural Information Processing Systems*, volume 32, pages 5509–5519, 2019.
- [13] Sohyeon Kwon and Yongjae Lee. Can GANs learn the stylized facts of financial time series? In *Proceedings of the 5th ACM International Conference on AI in Finance*, pages 126–133, 2024. doi: 10.1145/3677052.3698661.

- [14] L. Rabiner and B. Juang. An introduction to hidden Markov models. *IEEE ASSP Magazine*, 3(1):4–16, 1986. doi: 10.1109/MASSP.1986.1165342.
- [15] Tobias Rydén, Timo Teräsvirta, and Stefan Åsbrink. Stylized facts of daily return series and the hidden Markov model. *Journal of Applied Econometrics*, 13(3):217–244, 1998. doi: 10.1002/(SICI)1099-1255(199805/06)13:3<217::AID-JAE476>3.0.CO;2-V.
- [16] Jan Bulla and Ingo Bulla. Stylized facts of financial time series and hidden semi-Markov models. *Computational Statistics & Data Analysis*, 51(4):2192–2209, 2006. doi: 10.1016/j.csda.2006.07.021.
- [17] Thomas Lux and Michele Marchesi. Scaling and criticality in a stochastic multi-agent model of a financial market. *Nature*, 397:498–500, 1999. doi: 10.1038/17290.
- [18] Rama Cont and Jean-Philippe Bouchaud. Herd behavior and aggregate fluctuations in financial markets. *Macroeconomic Dynamics*, 4(2):170–196, 2000. doi: 10.1017/S1365100500015029.
- [19] J. Dooyne Farmer and Shareen Joshi. The price dynamics of common trading strategies. *Journal of Economic Behavior & Organization*, 49(2):149–171, 2002. doi: 10.1016/S0167-2681(02)00065-3.
- [20] Blake LeBaron. Agent-based computational finance. In Leigh Tesfatsion and Kenneth L. Judd, editors, *Handbook of Computational Economics*, volume 2, pages 1187–1233. Elsevier, 2006. doi: 10.1016/S1574-0021(05)02024-1.
- [21] W. Brian Arthur. Foundations of complexity economics. *Nature Reviews Physics*, 3:136–145, 2021. doi: 10.1038/s42254-020-00273-3.
- [22] J. Dooyne Farmer. Quantitative agent-based models: A promising alternative for macroeconomics. *Oxford Review of Economic Policy*, 41(2):571–590, 2025. doi: 10.1093/oxrep/graf027.
- [23] Louis Bachelier. Théorie de la spéculation. *Annales Scientifiques de l’École Normale Supérieure*, 17:21–86, 1900.
- [24] M. F. M. Osborne. Brownian motion in the stock market. *Operations Research*, 7(2):145–173, 1959.
- [25] Paul A. Samuelson. Rational theory of warrant pricing. *Industrial Management Review*, 6(2):13–32, 1965.
- [26] Fischer Black and Myron Scholes. The pricing of options and corporate liabilities. *Journal of Political Economy*, 81(3):637–654, 1973. doi: 10.1086/260062.
- [27] Eugene F. Fama. Efficient capital markets: A review of theory and empirical work. *The Journal of Finance*, 25(2):383–417, 1970. doi: 10.2307/2325486.
- [28] G. William Schwert. Why does stock market volatility change over time? *The Journal of Finance*, 44(5):1115–1153, 1989. doi: 10.1111/j.1540-6261.1989.tb02647.x.
- [29] Elias M. Stein and Jeremy C. Stein. Stock price distributions with stochastic volatility: An analytic approach. *The Review of Financial Studies*, 4(4):727–752, 1991.
- [30] Sangjoon Kim, Neil Shephard, and Siddhartha Chib. Stochastic volatility: Likelihood inference and comparison with ARCH models. *The Review of Economic Studies*, 65(3):361–393, 1998.
- [31] James D. Hamilton. A new approach to the economic analysis of nonstationary time series and the business cycle. *Econometrica*, 57(2):357–384, 1989. doi: 10.2307/1912559.
- [32] Alessandro Rossi and Giampiero M. Gallo. Volatility estimation via hidden Markov models. *Journal of Empirical Finance*, 13(2):203–230, 2006. doi: 10.1016/j.jempfin.2005.09.003.
- [33] Nguyet Nguyen. Hidden Markov model for stock trading. *International Journal of Financial Studies*, 6(2):36, 2018. doi: 10.3390/ijfs6020036.

- [34] Leonard E. Baum, Ted Petrie, George Soules, and Norman Weiss. A maximization technique occurring in the statistical analysis of probabilistic functions of Markov chains. *The Annals of Mathematical Statistics*, 41(1):164–171, 1970. doi: 10.1214/aoms/1177697196.
- [35] José G. Dias, Jeroen K. Vermunt, and Sofia Ramos. Mixture hidden Markov models in finance research. In *Advances in Data Analysis, Data Handling and Business Intelligence*, pages 451–459. Springer, 2010. doi: 10.1007/978-3-642-01044-6_41.
- [36] Paul Embrechts, Alexander J. McNeil, and Daniel Straumann. Correlation and dependence in risk management: Properties and pitfalls. In *Risk Management: Value at Risk and Beyond*, pages 176–223. Cambridge University Press, 2002. doi: 10.1017/CBO9780511615337.008.
- [37] Umberto Cherubini, Elisa Luciano, and Walter Vecchiato. *Copula Methods in Finance*. John Wiley & Sons, 2004. ISBN 9780470863459.
- [38] Kjersti Aas, Claudia Czado, Arnoldo Frigessi, and Henrik Bakken. Pair-copula constructions of multiple dependence. *Insurance: Mathematics and Economics*, 44(2):182–198, 2009.
- [39] Stephen A. Ross. The arbitrage theory of capital asset pricing. *Journal of Economic Theory*, 13(3):341–360, 1976. doi: 10.1016/0022-0531(76)90046-6.
- [40] Eugene F. Fama and Kenneth R. French. Common risk factors in the returns on stocks and bonds. *Journal of Financial Economics*, 33(1):3–56, 1993. doi: 10.1016/0304-405X(93)90023-5.
- [41] William F. Sharpe. A simplified model for portfolio analysis. *Management Science*, 9(2):277–293, 1963. doi: 10.1287/mnsc.9.2.277.
- [42] A. N. Kolmogorov. Sulla determinazione empirica di una legge di distribuzione. *Giornale dell’Istituto Italiano degli Attuari*, 4:83–91, 1933.
- [43] N. Smirnov. Table for estimating the goodness of fit of empirical distributions. *The Annals of Mathematical Statistics*, 19(2):279–281, 1948.
- [44] Theodore W. Anderson and Donald A. Darling. Asymptotic theory of certain “goodness of fit” criteria based on stochastic processes. *The Annals of Mathematical Statistics*, 23(2):193–212, 1952.
- [45] David G. Booth and Eugene F. Fama. Diversification returns and asset contributions. *Financial Analysts Journal*, 48(3):26–32, 1992. doi: 10.2469/faj.v48.n3.26.
- [46] Samuel Kotz, Tomasz J. Kozubowski, and Krzysztof Podgórski. *The Laplace Distribution and Generalizations: A Revisit with Applications to Communications, Economics, Engineering, and Finance*. Birkhauser, 2001. ISBN 9780817641665.
- [47] David Toth and Bruce Jones. Against the norm: Modeling daily stock returns with the Laplace distribution. *arXiv preprint arXiv:1906.10325*, 2019.
- [48] Jeff A. Bilmes. A gentle tutorial of the EM algorithm and its application to parameter estimation for Gaussian mixture and hidden Markov models. Technical Report TR-97-021, International Computer Science Institute, 1998.
- [49] Paul Glasserman. *Monte Carlo Methods in Financial Engineering*. Springer, 2003. doi: 10.1007/978-0-387-21617-1.
- [50] James D. Hamilton. Regime switching models. *The New Palgrave Dictionary of Economics*, pages 11421–11426, 2018. doi: 10.1057/978-1-349-95189-5_2459.
- [51] Charles M. Grinstead and J. Laurie Snell. *Introduction to Probability*. American Mathematical Society, 1997.
- [52] Dimitris N. Politis and Joseph P. Romano. The stationary bootstrap. *Journal of the American Statistical Association*, 89(428):1303–1313, 1994. doi: 10.1080/01621459.1994.10476870.

- [53] Kyunghyun Cho, Bart van Merriënboer, Caglar Gulcehre, Dzmitry Bahdanau, Fethi Bougares, Holger Schwenk, and Yoshua Bengio. Learning phrase representations using RNN encoder-decoder for statistical machine translation. In *Proceedings of the 2014 Conference on Empirical Methods in Natural Language Processing (EMNLP)*, pages 1724–1734. Association for Computational Linguistics, 2014. doi: 10.3115/v1/D14-1179.
- [54] Paul Bouchey, Vassilii Nemtchinov, and Ting-Kam Leonard Wong. Volatility harvesting in theory and practice. *Journal of Wealth Management*, 18(3):89–100, 2015. doi: 10.3905/jwm.2015.18.3.089.
- [55] Harry M. Markowitz. Investment for the long run: New evidence for an old rule. *Journal of Finance*, 31(5):1273–1286, 1976. doi: 10.1111/j.1540-6261.1976.tb03213.x.
- [56] Aswath Damodaran. Equity risk premiums: Determinants, estimation and implications. NYU Stern School of Business working paper, updated annually, 2025. <https://pages.stern.nyu.edu/~adamodar/>.
- [57] Roger G. Ibbotson and Duff and Phelps. *SBBI Yearbook: Stocks, Bonds, Bills, and Inflation*. Duff and Phelps / Kroll, annual edition, 2024.